

A FATAL GAS EMBOLISM FOLLOWING THE USE OF INTRAOPERATIVE HYDROGEN PEROXIDE AS AN IRRIGATION FLUID

Z. Hussain-Khan¹, A. A. Soleimani^{*1} and M. Farzan²

1) Department of Anesthesiology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

2) Department of Orthopedic Surgery, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

Abstract- Hydrogen peroxide, an oxidizing agent used in surgery for its antiseptic properties, has been associated with life-threatening complications. We report a case of suspected gas embolism following irrigation of a tibial bone cyst with hydrogen peroxide in a five year-old boy under general anesthesia, who died despite resuscitative measures and a long stay in the intensive care unit.

Acta Medica Iranica, 42(2): 151-153; 2004

Key words: Hydrogen peroxide, oxygen embolism

INTRODUCTION

Hydrogen peroxide (H₂O₂) is frequently used for irrigation, cleansing and dressing of surgical wounds. H₂O₂ decomposes rapidly as a result of the action of catalases in blood and tissues, releasing molecular oxygen and water. There are some reports of cases of gas embolism due to H₂O₂ administration (1-3). Here, we report a case of gas embolism in a patient that led to death after a prolonged stay in the intensive care unit. We wish to warn against the potential dangers of using H₂O₂ during surgery.

Case reports

A five year-old boy presented with a tibial bone cyst. Preoperative vital signs and laboratory data were all within the normal range. After employing routine standard monitoring, the patient underwent an uneventful induction of anesthesia with fentanyl 1

µg/kg, thiopental sodium 5 mg/kg and atracurium in a dose of 0.5 mg/kg. Anesthesia was maintained with a combination of halothane 0.5%, N₂O and oxygen each in a concentration of 50% and incremental intravenous fentanyl. The patient was placed in supine

position and the table maintained in a neutral position. After preparing and draping the surgical field, a Jamshidi needle was inserted into the cyst cavity by the surgeon with the aid of fluoroscopy. After drainage of some fairly clear fluid from the cyst cavity, 10 ml of 30% H₂O₂ solution was forcibly flushed into the cavity via the same needle in order to irrigate the cystic cavity. A few seconds after the injection of H₂O₂, severe bradycardia and hypotension, accompanied by difficulties in ventilation ensued. Ventilatory pressure raised suddenly and hemoglobin saturation of oxygen (SpO₂) decreased abruptly to about 40%. There was also a reduction in end-tidal CO₂. N₂O was turned off immediately and the lungs were ventilated manually, but with difficulty, utilizing one hundred percent oxygen. The patient was placed in the Trendelenburg and left lateral position and repeated doses of atropine 0.25 mg followed by epinephrine 0.1 mg/kg (1: 1000) were injected together with rinsing of the

Received: 3 June. 2003, Revised: ---, Accepted: 24 Dec. 2003

* Corresponding Author:


A. A. Soleimani, Department of Anesthesiology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran
Tel: +98 21 6935061, Fax: +98 21 6438634
E-mail: aasolei@yahoo.com

Fatal gas embolism

surgical field with saline solution. Meanwhile, a needle was inserted into the heart of the patient with air gushing out (the central venous pressure had not been established prior to surgery because of a short surgical procedure). Subsequently, the initial stony bag gave way to a comparatively soft bag with resultant improvement in oxygenation culminating in an SpO₂ level of 90%. At this moment, the heart rate was about 130 beats per minute and the blood pressure was 90/60 mmHg. Dexamethasone 8 mg was given intravenously to counteract ensuing cerebral edema secondary to the hypoxemic episode. The patient was sent to the intensive care unit and mechanical ventilation (CMV mode with FiO₂=100%) was initiated. Nearly an hour later, the patient developed a generalized seizure which was controlled with diazepam 5 mg intravenously. A neurologist was consulted and intravenous phenobarbital 30 mg three times a day was started to control further seizures. No more seizures occurred. Nutritional support was started on the third day. Unfortunately, the patient never regained his consciousness and died after a protracted stay of 18 days in the ICU.

DISCUSSION

H₂O₂ is a clear and colorless solution with a mildly piquant or acidic odor. It is unstable and decomposes violently when in direct contact with rough surfaces or trace metals (such as Cu and Fe), dirt, heat, light and biological materials such as blood (4). H₂O₂ was primarily used for bleaching straw hats. At present, it has wide-spread usages, ranging from food processing to energy production in space crafts. In medicine it is frequently used for irrigation and dressing of surgical wounds (the broad spectrum effect is responsible for its wide-spread common usage as an effective antiseptic and disinfectant agent). The blood and liver of mammals contain an enzyme, catalase, that catalyzes the decomposition of H₂O₂ into water and oxygen (5,6). It is oxygen that kills bacteria (4).



In 1967, intravenous H₂O₂ was used in pigs. It was demonstrated that a solution of H₂O₂ 3% diluted 50/ 50 with NaCl 0.9% perfused in the right ventricle,

led to the rapid formation of bubbles with subsequent pulmonary and systemic embolization, severe methemoglobinemia and death (7). In a review of literature, we can find reports of sudden cardiovascular collapse and pulmonary embolism due to H₂O₂ application. Loeb reported a case of severe oxygen embolism after wound irrigation with H₂O₂ in a patient undergoing surgical dressing of a large thigh trauma under general anesthesia (1). In another report by Despound et al, an oxygen venous embolism occurred during lumbar discectomy after the use of H₂O₂ (2). There is also a report of a similar event occurring in a patient undergoing bilateral pulmonary lobectomy in whom the thoracic cavity had been irrigated with H₂O₂ at the end of the procedure (3). In the present case, although a transoesophageal echocardiography or precordial Doppler was not used to exactly diagnose and pinpoint gas embolism, the air gushing out after inserting the needle in the heart, and also the timing of the symptoms and clinical presentation, such as hypotension, bradycardia and a decrease in end-tidal CO₂ and oxygen saturation occurring abruptly after H₂O₂ irrigation, strongly and unequivocally pointed to the diagnosis of a gas embolism. Noteworthy is the fact that clinical and pathophysiological features of air and oxygen emboli are almost identical, including the management protocol. Unfortunately, medical management is not always successful and despite all efforts, some patients may die. Indeed, the present case, clearly demonstrated once again that H₂O₂ application is not totally free of danger and fatal repercussions. One milliliter of a 30% H₂O₂ solution will yield 100 ml of oxygen gas in tissue when it decomposes (8,9). This gas can enter the circulation and cause severe gas embolism. So we need to be more cautious while using H₂O₂. In our opinion, it is better to avoid the routine-use of H₂O₂, and restrict its use in highly indicated cases in whom its merits outweigh its potential hazards.

REFERENCES

1. Loeb T, Loubert G, Templier F, Pasteyer J. Iatrogenic gas embolism following surgical lavage of a wound with hydrogen peroxide. *Ann Fr Anesth Reanim* 2000; 19(2): 108-110.

2. Despond O, Fiset P. Oxygen venous embolism after the use of hydrogen peroxide during lumbar discectomy. *Can J Anaesth* 1997; 44(4): 410-413.
3. Konrad C, Schuupfer G, Wietlisbach M, Gerber H. Pulmonary embolism and hydrogen peroxide. *Can J Anaesth* 1997; 44 (3): 338-339.
4. Stoker S. H. Chemical calculations. The mole concept and chemical formulas. In: Stoker S. H. Introduction to chemical principles. 6th ed. New Jersey Prentice Hall: 1999. p. 272-325.
5. Brown TL, LeMay EH, Bursten BE. Chemical kinetics. In: Brown TL, LeMay EH, Bursten BE. Chemistry: the central science. 8th ed. New Jersey Prentice Hall: 2000 .p. 509-558.
6. Jones L, Atkins P. Chemistry molecules, matter and change. 4th ed. New York W H Freeman and Company: 1999. p. 885-917.
7. Fuson RL, Kylstra JA, Hochstein P, Salzman HA. Intravenous hydrogen peroxide infusion as a mean of extrapulmonary oxygenation. *Clin Res* 1967; 15: 74.
8. Green S. Oxygenation therapy unproven treatments for cancer and AIDS. *Sci Rev Alt Med* 1998; 2: 6-12.
9. Mendham J, Denney RC, Barnes JD, Thomas M J K. Titrimetric analysis. In: VOGEL'S textbook of Quantitative chemical analysis. 6 th ed. England Prentice Hall 2000. p. 312-460.