

1939

A classification of the proteolytic micrococci isolated from dairy products

Horace Hammerton Harned
Iowa State College

Follow this and additional works at: <https://lib.dr.iastate.edu/rtd>

 Part of the [Microbiology Commons](#)

Recommended Citation

Harned, Horace Hammerton, "A classification of the proteolytic micrococci isolated from dairy products " (1939). *Retrospective Theses and Dissertations*. 13703.
<https://lib.dr.iastate.edu/rtd/13703>

This Dissertation is brought to you for free and open access by the Iowa State University Capstones, Theses and Dissertations at Iowa State University Digital Repository. It has been accepted for inclusion in Retrospective Theses and Dissertations by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

INFORMATION TO USERS

This manuscript has been reproduced from the microfilm master. UMI films the text directly from the original or copy submitted. Thus, some thesis and dissertation copies are in typewriter face, while others may be from any type of computer printer.

The quality of this reproduction is dependent upon the quality of the copy submitted. Broken or indistinct print, colored or poor quality illustrations and photographs, print bleedthrough, substandard margins, and improper alignment can adversely affect reproduction.

In the unlikely event that the author did not send UMI a complete manuscript and there are missing pages, these will be noted. Also, if unauthorized copyright material had to be removed, a note will indicate the deletion.

Oversize materials (e.g., maps, drawings, charts) are reproduced by sectioning the original, beginning at the upper left-hand corner and continuing from left to right in equal sections with small overlaps.

ProQuest Information and Learning
300 North Zeeb Road, Ann Arbor, MI 48106-1346 USA
800-521-0600

UMI[®]

NOTE TO USERS

This reproduction is the best copy available.

UMI[®]

A CLASSIFICATION OF THE PROTEOLYTIC
MICROCOCCI ISOLATED FROM
DAIRY PRODUCTS

by

Horace H. Harned

A Thesis Submitted to the Graduate Faculty
for the Degree of

DOCTOR OF PHILOSOPHY

Major Subject Dairy Bacteriology

Approved:

Signature was redacted for privacy.


In charge of Major work

Signature was redacted for privacy.

Head of Major Department

Signature was redacted for privacy.

Dean of Graduate College



Iowa State College
1939

UMI Number: DP12737

UMI[®]

UMI Microform DP12737

Copyright 2005 by ProQuest Information and Learning Company.

All rights reserved. This microform edition is protected against
unauthorized copying under Title 17, United States Code.

ProQuest Information and Learning Company
300 North Zeeb Road
P.O. Box 1346
Ann Arbor, MI 48106-1346

QTR 121
H229c

TABLE OF CONTENTS

INTRODUCTION	6
STATEMENT OF PROBLEM	7
HISTORICAL	8
METHODS	14
Detection of Proteolytic Micrococci	14
Milk agar plates	14
Gelatine liquefaction	14
General Characteristics	14
Action on milk	14
pH and titratable acidity	14
Nitrate reduction	15
Nitrogen assimilation and ammonia production	15
Carbon dioxide	15
Hemolysis	15
Lipolysis	16
Diastatic action	16
Carbohydrate fermentation	16
Thermal death point	16
Acetylmethylcarbinol plus diacetyl	16
Volatile acidity	16
Hydrogen sulfide	17
Indol	17
Pathogenicity	17

T 6 1 5 4

Protein Breakdown in Skimmilk	17
General Action of Proteolytic Micrococci on Butter	18
EXPERIMENTAL	19
Isolation of Proteolytic Micrococci from Dairy Products	19
Chromogenesis	22
Influence of Various Factors on Chromogenesis ..	22
Carbon dioxide	22
Growth on potato	23
pH of the medium	23
Incubation in water	24
Heating at 55°C.	24
Incubation temperature	24
Darkness	25
Chromogenesis as a Means of Separating Genera ..	25
General Growth and Viability of Proteolytic Micrococci	26
General Characteristics of Proteolytic Micrococci	26
Action on litmus milk	29
Titratable acidity produced in skimmilk ...	29
Growth on agar slants	30
Growth in beef broth	30
Nitrate reduction	30

Nitrogen assimilation and ammonia production	30
Gelatine liquefaction	31
Carbon dioxide production	31
Hemolysis	31
Lipolysis	31
Diastatic action	31
Carbohydrate fermentation	32
Thermal death point	32
Acetylmethylcarbinol plus diacetyl, volatile acidity, hydrogen sulfide and indol production	32
Pathogenicity, spore formation and motility	33
Protein Breakdown in Skimmilk by Proteolytic Micrococci	33
General Action of Proteolytic Micrococci on Unsalted Butter	38
Action of orange micrococci	39
Action of yellow micrococci	44
Action of white micrococci	48
Species of Proteolytic Micrococci	52
Orange Group	52
Yellow Group	53
White Group	54

Key to the Identification of Proteolytic Micrococci	55
DISCUSSION OF RESULTS	57
SUMMARY AND CONCLUSIONS	60
ACKNOWLEDGMENTS	65
LITERATURE CITED	66

INTRODUCTION

In studying changes in dairy products, bacteriologists have given the micrococci relatively little attention since it is generally thought that these organisms produce changes slowly and therefore are of little importance. The usual classifications of the micrococci are based on such characteristics as gelatine liquefaction, nitrate reduction and the fermentation of various carbohydrates. These characteristics are of little practical value to the dairy bacteriologist. The action of the organisms on the constituents of milk, however, may be of considerable importance.

Various micrococci found in milk and milk derivatives are capable of attacking the proteins of milk. The extent to which the milk is proteolyzed varies greatly with the particular organism. In some instances proteolysis can be detected by the appearance of the milk while in other instances chemical analyses are necessary to detect changes.

The ability of certain micrococci to break down proteins is important since many dairy products contain enough proteins to make such changes undesirable from the standpoint of quality; however, such changes may be of importance in cheese ripening. The micrococci, therefore, should be seriously considered by the dairy bacteriologist.

STATEMENT OF PROBLEM

The objects of the work herein reported are:

1. To isolate proteolytic micrococci from various dairy products and to study their general characteristics.
2. To determine the effect of various types of proteolytic micrococci on milk and butter.
3. To classify the proteolytic micrococci in dairy products and to identify them.

HISTORICAL

The name Micrococcus was first applied by Hallier (18) to indicate the growth form of a mold and, later, by Cohn (8) to all spherical bacteria. In 1872, Cohn (8) published his conception of the systematic relationships of bacteria and for the first time the coccus types were placed in a separate tribe on an equal basis with other tribes of bacteria. The genus name Micrococcus was used for this group which included all spherical bacteria; a definite description of the genus was given. In 1875, Cohn (10) proposed a more complete classification of bacteria than anyone else had attempted up to this time. His classification was based on the assumption that the various types were degenerates of the blue-green algae. Having in mind the close relationship between the algae and the bacteria, he arranged those species of bacteria which he supposed were closely related to species of blue-green algae in the same genus with the algae so that many genera contained both bacteria and algae. The streptococci and sarcina were distinguished by their respective chain and packet formations.

The genus Micrococcus of Migula (31) and of Chester (7) was a kind of residual genus including all cocci which do not occur in well marked chains or packets. One of the outstanding characteristics of the group is the ability to produce

pigments, and many investigators have used this character as a starting point for classification.

In 1908, Dudgeon (12) studied the relationships of a group of staphylococci with particular emphasis on the use of chromogenesis for separating the individual species. Working with 46 strains of the orange type and 71 strains of the white type, he observed their liquefying power, their reducing power and their acid producing power in 11 carbohydrate media. His conclusion was that all the staphylococci studied were members of one species, although only a few of the strains were identical in all characters studied.

The Winslows (35), in 1908, made a comprehensive study of the coccus group. A large number of cultures was studied and divisions were made on the basis of habitat, chromogenesis and cell division. They divided the family Cocceaceae into two sub-families, the Paracocceaceae which included the parasitic types and the Metacocceaceae which included the non-parasitic or free living forms. The Paracocceaceae were further divided into five genera on the basis of cell division and pigment production. The old genus Staphylococcus was divided into two new genera, namely, Aurococcus which included the orange pigment producers, and Albococcus which included the porcelain white or non-pigment producers. The Metacocceaceae were divided into three genera, principally on the basis of cell formation and pigment production.

Orla-Jenson (32), in 1909, proposed a classification of

bacteria based primarily on physiological characters, the most important of which was acid production in various carbohydrate media. Although the conventional morphological grouping of the cocci was recognized, only the physiological characteristics were used in his classification. His classification of the coccus group is as follows:

Family Acidobacteriaceae
Genus Streptococcus
Genus Micrococcus
Genus Sarcina

In 1917, Buchanan (5) classified the micrococci on the basis of their physiology and habitat. The genera were separated on the basis of cell division, habitat, and chromogenesis. The tribe Micrococceae was divided into three genera:

Genus Micrococcus
Genus Rhodococcus
Genus Sarcina

Using the cultures of the American Museum of Natural History Collection, Kligler (26), in 1915, confirmed the work of the Winslows (35) and concluded that the white and orange micrococci were usually parasitic strains while the yellow and red micrococci were more often saprophytic.

In 1920, Winslow and his associates (36), after further study of the micrococci, were doubtful of the validity of the genus Aurococcus and genus Albococcus, described in 1908. Since the organisms in the two genera were similar, these investigators combined the groups into the single genus Staphylococcus.

Hucker (24) reported an exhaustive study of the micrococci and was convinced that chromogenesis of the members of this group is a very important character for use in classification but that groups should not be divided into genera on the basis of chromogenesis as the chief diagnostic feature. A study of the constancy of various characters of this group showed that chromogenesis, nitrate reduction, ability to utilize ammonium salts as the only source of nitrogen, gelatine liquefaction, and action on milk are tests whereby the natural species of the group can be separated.

In 1920, the committee on Classification of Bacterial Types, appointed by the Society of American Bacteriologists, (37), arranged the Micrococceae into three genera:

Genus Micrococcus
Genus Sarcina
Genus Rhodococcus

This is similar to Buchanan's classification (5).

Bergey (5), in his Manual of Determinative Bacteriology, divided the Micrococceae into five genera:

Genus Staphylococcus
Genus Gaffkya
Genus Micrococcus
Genus Sarcina
Genus Rhodococcus

For over 50 years research workers in the field of dairy industry have known that micrococci were plentiful in dairy products. Gorini (17), in 1901, reported micrococci as the most common type of organism in fresh milk. He thought the proteolytic micrococci in the mammalian glands were advan-

tageous in the digesting of milk and in the ripening of cheeses. In hard cheeses the organisms reduce casein to more digestible forms during ripening and at the same time promote the development of lactic acid organisms which contribute to flavor development. Liquefying, chromogenic micrococci were found by von Froudenreich (16) to predominate in milk. Orla-Jensen (32) reported many micrococci and serratias in milk and termed them tetracocci. The sources of the tetracocci were cow manure, soil, dust, the coat of the animal and the skin of the milker. According to Orla-Jensen, the organisms play an important part in the ripening of cheeses.

In an extensive study on the microflora of milk, Harding and Wilson (21) found that 75 per cent of the organisms present in freshly drawn milk were micrococci. In a similar study, Evans (14) noted that the predominant flora in fresh cows' milk consisted of Gram positive micrococci. Copeland and Olson (11) found mostly Gram positive micrococci in aseptically drawn milk, some of which liquefied gelatine. Breed (4) reported that the predominant flora in milk consisted of Gram positive micrococci, while Knowles (26) found that 48 of 68 cultures isolated from freshly drawn foremilk were micrococci.

Hucker (23), in examining microscopic sections of old cheddar cheese, found that micrococci predominated. The

characteristic clumping of this group of organisms showed that in their growth and reproduction they probably broke down the surrounding medium for a source of energy. Frazier and Rupp (15) reported the micrococci in milk as mostly acidoproteolytic.

METHODS

Detection of Proteolytic Micrococci

Milk agar plates

Proteolytic micrococci were detected on agar plates by the addition of 5 ml. of sterile skim milk to 100 ml. of beef infusion agar at the time of plating.

Gelatine liquefaction

Tubes of standard nutrient gelatine were inoculated and incubated at 37°C. for 5 days. The cultures were then placed in a refrigerator at 10°C. for 4 hours. After the holding period, the tubes were examined for liquefaction. Tests were also made by incubating the tubes at 15° to 18°C. for 15 days.

General Characteristics

Action on milk

The action of proteolytic micrococci on milk was determined by inoculating tubes of litmus milk with the organisms. The tubes were examined after 48 hours at 37°C.

pH and titratable acidity

The pH values were obtained electrometrically with the quinhydrone electrode. The titratable acidity of milk cul-

tures was obtained by titrating 20 ml. portions with N/10 sodium hydroxide, using phenolphthalein. The results were calculated as lactic acid.

Nitrate reduction

Nitrate reduction was determined by a method adopted by the committee on technique of the Society of American Bacteriologists (13).

Nitrogen assimilation and ammonia production

Tests for the assimilation of amino and ammonia nitrogen and the production of ammonia from peptone and nitrates were those described by Tucker (14).

Carbon dioxide

The production of carbon dioxide was measured by means of Aldridge tubes (15).

Hemolysis

Hemolysis was studied by streaking cultures on blood agar plates. The plates were prepared by adding 4 ml. of the following mixture to 100 ml. of beef infusion agar: 2 ml. aseptically drawn cow's blood and 2 ml. 5 per cent aqueous sodium citrate solution

Lipolysis

Lipolysis was investigated by the modified Nile blue sulfate and natural fat techniques as suggested by Long (29).

Diastatic action

Diastatic action was determined by the method suggested by Allen (1).

Carbohydrate fermentation

The fermentation of carbohydrates was studied by growing organisms in broths containing various materials and also bromocresol purple as an indicator.

Thermal death point

The thermal death point of organisms was determined by heating cotton stoppered tubes containing broth cultures for 15 minutes at various temperatures in a water bath.

Acetylmethylcarbinol plus diacetyl

Acetylmethylcarbinol plus diacetyl was determined qualitatively by the Voges-Proskauer reaction (55) and quantitatively by the procedure employed by Michaelian and Hamner (30).

Volatile acidity

Volatile acidity was determined by the method used by

Michaelian and Hammer (30).

Hydrogen sulfide

Hydrogen sulfide was determined with the method used by Bailey and Lacy (2).

Indol

The production of indol was determined by the Ehrlich-Bohme method adopted by the committee on bacteriological technique of the Society of American Bacteriologists (35).

Pathogenicity

Pathogenicity was tested by inoculating guinea pigs.

Protein Breakdown in Skimmilk

The total soluble nitrogen produced by cultures of proteolytic micrococci was obtained by precipitation of milk cultures of the organisms with acetic acid and then filtering through paper. The nitrogen in both the filtrates and the precipitates were determined by the Kjeldahl method. Amino nitrogen was obtained with the Van Slyke gasometric method (34). Nitrogen soluble and insoluble in trichloroacetic acid, phosphotungstic acid and ethyl alcohol was fractionated according to the procedure used by Lane and Hammer (28).

General Action of Proteolytic Micrococci on Butter

Churnings were made from sterile cream inoculated with milk cultures of the proteolytic micrococci. The butter was examined organoleptically at various periods during the holding.

EXPERIMENTAL

Isolation of Proteolytic Micrococci
from Dairy Products

Proteolytic micrococci were isolated from milk, butter, cheese (cheddar) and ice cream. The sources of the samples of these materials and the numbers from each are as follows:

<u>SOURCES</u>	<u>Number of samples examined</u>			
	<u>Milk</u>	<u>Butter</u>	<u>Cheese</u>	<u>Ice Cream</u>
Market Milk Department	136			
Iowa State College				
Ames, Iowa				
Dairy Department				
Mississippi State College				
State College, Mississippi:				
Barn No. 1	148			
Barn No. 2	164			
Dairy of H. H. Harned	82			
State College, Mississippi:				
A & M Cooperative Creamery:		42		46
State College, Mississippi:				
Grocery Stores			36	
Starkville, Mississippi				

All samples of milk were taken from fresh morning milk; each sample (about 10 ml.) was collected in a sterile test tube. The samples of butter were cut from one-pound prints with sterile spatulas and held in petri dishes. The samples of cheese were obtained with sterile triers and held in petri dishes. The samples of ice cream were obtained with sterile

spotulas, and allowed to melt in petri dishes at room temperature.

The samples of the four dairy products were plated with beef infusion milk agar in the following manner: The milk and ice cream were diluted with water as usual. The butter was melted at 45°C. and dilutions were made with warm water. The cheese were ground in a mortar with 10 ml. of 2 per cent aqueous sodium citrate until a homogeneous liquid was obtained; the necessary dilutions were made with water. All plates were incubated at 57°C. for 48 to 72 hours. Those colonies which were smooth and round, and which were encircled with a clear area characteristic of proteolytic organisms, were streaked on beef infusion milk agar plates for purification and for identification as micrococci. Special care was exercised to avoid duplicating organisms from a sample. In order to determine the morphological characters, smears were made and stained. After an organism had been identified as a Micrococcus by microscopic examination, it was transferred to an agar slant as a stock culture for detailed study. From 704 samples of dairy products, 426 proteolytic micrococci were isolated. The general sources of the organisms are shown in the following summary:

	<u>Milk</u>	<u>Butter</u>	<u>Cheese</u>	<u>Ice Cream</u>
Number of samples examined:	580	42	36	46
:	:	:	:	:
Number of samples yielding:	:	:	:	:
proteolytic micrococci	340	36	20	30

The sources of the individual cultures that were studied in detail are given in Table 1.

Table 1.
Sources of the proteolytic micrococci

Source	Total cultures	Culture numbers
Milk	340	All numbers not listed under butter, cheese and ice cream.
Butter	36	21, 36, 44, 49, 53, 132, 135, 134, 135, 178, 179, 180, 181, 210, 211, 212, 213, 214, 215, 250, 251, 252, 253, 254, 255, 256, 257, 329, 330, 331, 332, 354, 355, 356, 357.
Cheese	20	6, 96, 97, 98, 99, 100, 101, 126, 127, 174, 175, 176, 177, 243, 244, 245, 246, 247, 248, 249.
Ice Cream	30	27, 54, 55, 56, 57, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 169, 170, 171, 172, 173, 322, 323, 324, 325, 326, 327, 328, 341.

Chronogenesis

The pigments produced by the proteolytic micrococci encountered were red, orange, yellow or white (non-pigmented). In this investigation the red micrococci were omitted since only a few were isolated and these grew very slowly at both room temperature and 37°C. The slow development of the red organisms may account for the fact that so few were isolated as most of the colonies were picked from plates that were incubated not more than 72 hours. The cultures studied included 53 orange micrococci, 115 yellow and 258 white. On beef infusion agar, the cultures were very consistent in their pigment production.

Influence of Various Factors on Chronogenesis

Carbon dioxide

Orange cultures 4, 7, 12, 13, 15, 19, 20, 22, 30, 31, 36, 37, 39, 49, 55, and yellow cultures 59, 61, 64, 69, 72, 74, 81, 88, 91, 92, 94, 95, and 97 were streaked on beef infusion agar plates. The plates were placed in a Novy jar and carbon dioxide was allowed to flow into the jar until the air was largely displaced. After 5 and 10 days incubation periods at room temperature the cultures were compared with duplicate plates streaked at the same time but incubated under aerobic conditions. With cultures 7, 13, 15, 31, 36, and 49 held under carbon dioxide, the orange

pigment slightly lighter in color than in the controls, but most of the cultures appeared to produce as much pigment under carbon dioxide as in air. No change in the pigment production was noted with the yellow cultures.

Growth on potato

Orange cultures 4, 7, 12, 13, 15, 19, 20, 22, 30, 31, 36, 37, 39, 49, and 53 and yellow cultures 59, 61, 64, 69, 72, 74, 81, 83, 91, 92, 94, 95, and 97 were streaked on potato to compare the growth with that on beef infusion agar slants. The cultures were held at room temperature. Cultures 7, 13, 36, and 49 showed more pigmentation on potato than on beef infusion agar, while cultures 31, 39, and 47 appeared to have a deeper color on beef infusion agar than on potato. The yellow cultures showed about the same pigmentation on the two media.

pH of the medium

The pH of the medium appeared to affect the amount of pigment produced. The test organisms employed were those used in studying the growth on potato. When agars having values of 6.6, 7.0 and 7.6, respectively, were employed and the cultures held at room temperature, pigmentation was most pronounced on the agar having a pH of 7.0 with all cultures studied but was definite in all cultures.

Incubation in water

Orange cultures 12, 13, 15, 19, 20, 22, 30, 31, 36 and 49, and yellow cultures 59, 61, 64, 69, 72, 74, 81, 88, 97 and 103 were each placed in a tube of sterile water for 6 days at room temperature and then streaked on agar slants. The cultures were incubated at both 20° and 37°C. for 48 hours and then compared with cultures not placed in water. No difference in pigmentation was noted between the two types of cultures.

Heating at 55°C.

Orange cultures 12, 13, 15, 19, 20, 22, 30, 31, 39 and 49, and yellow cultures 59, 61, 64, 69, 72, 74, 81, 88, 97 and 103 were diluted in tubes of water and placed in a water bath at 55°C. for 10 minutes. The cultures were streaked on beef infusion agar and held at 37°C. for 48 hours. With orange cultures 39 and 49 there appeared to be a deeper color produced by the heated cultures than by unheated control cultures, but in the majority of comparisons there was no noticeable change due to heating. No change in pigmentation was noted with the yellow cultures that were heated.

Incubation temperature

Agar plates streaked with orange cultures 4, 7, 12, 13, 15, 19, 20, 22, 30, 31, 36, 37, 39, 49 and 53 and with

yellow cultures 59, 61, 64, 69, 72, 74, 81, 83, 91, 93, 94, 95, and 97 were incubated at 21°C. and also at 37°C. The orange cultures showed a deeper pigment production at 21°C. than at 37°C., whereas the yellow cultures showed no variation with the incubation temperatures.

Darkness

The orange and yellow cultures used in studying the effect of incubation temperature were streaked on duplicate tubes of agar. One set of tubes was placed in closed mailing tubes, while the other set was held in a room having sunlight most of the day. After 10 days the orange cultures incubated in darkness were much lighter in color than those incubated in the light room. With the yellow cultures little difference in pigmentation was noted, regardless of the amount of light present.

Chronogenesis as a Means of Separating Genera

Many workers have made use of chronogenesis as a diagnostic feature in separating genera. Usually it is an easy character to determine and the colors are relatively constant. However, orange culture 53 changed from orange to white on agar but later the orange color reappeared; this color change occurred twice over a 2-year period. In litmus milk cultures an orange ring regularly formed at the surface while agar

streak cultures were white with an orange periphery.

Since the chromogenic properties of the proteolytic micrococci were relatively constant, chromogenesis is apparently a satisfactory means of dividing relatively large numbers of organisms into groups. In this study the chromogenic properties were employed to divide the organisms into the orange, yellow and white groups.

General Growth and Viability of Proteolytic Micrococci

All of the 426 cultures of proteolytic micrococci studied grew well on beef infusion milk agar, although the amount of growth varied considerably among individual cultures.

In general, the cultures were viable for relatively long periods. Litmus milk cultures held 3 to 4 months and agar slant cultures held 10 to 12 months at room temperature remained viable. Agar slant cultures held for long periods were easily revived by adding a small amount of beef broth to the cultures, holding over night and then streaking on agar.

General Characteristics of Proteolytic Micrococci

The general characteristics, grouping and identification of the organisms studied are given in Table 2. The

General characteristics grouping and

Cult- ure no.	Form	Size microns	Gram stain	Action on milk 37°C. 48 hrs.	Titrat- able acidity in milk 37°C. 48 hrs.	Growth on agar slants	Growth in beef broth	Nitrate reduction	Amino N. assimilation	Ammonia N. assimilation
---------------------	------	-----------------	---------------	--	---	--------------------------------	-------------------------------	----------------------	--------------------------	----------------------------

1	singly	0.6 to		acid	0.30 to					
48	to pairs	1.0	+	coag. ²	0.36	abun-	turbid	+	-	-
49	clusters	av. 0.8		dig. ³	av. 0.33	dant				
50	and clusters	0.8 to 1.4	+	inert	0.28	abun-	turbid	-	-	+
51		av. 1.1			av. 0.26	dant				
52	singly	0.9	var. ¹	acid	0.39	abun-	turbid	-	-	-
53	clusters			coag.		dant				
54	and singly	0.6 to 1.0	+	acid	0.22 to 0.26	abun-	turbid	+	-	+
55	fours	av. 0.8			av. 0.24	dant				

54	singly	1.0 to		acid	0.1 to					
95	to pairs	1.6	var.	coag.	0.20	abun-	turbid	-	+	+
96	clusters	av. 1.3		dig.	av. 0.18	dant				
113	singly	1.0 to			0.14 to					
114	to pairs	1.4	var.	inert	0.16	abun-	turbid	-	-	+
137	clusters	av. 1.2			av. 0.15	dant				
138	singly	0.8 to		acid	0.14 to					
168	to pairs	1.2	var.	coag.	0.18	abun-	turbid	+	-	-
169	clusters	av. 1.0		dig.	av. 0.16	dant				
170	singly	0.9 to			0.12 to					
171	to pairs	1.3	var.	inert	0.14	abun-	turbid	+	-	-
172	clusters	av. 1.2			av. 0.13	dant				

1 = variable 2 = coagulation 3 = digestion

1 from dairy products

Glycerol	Normal	Classification
Inulin	Normal	
Lactose	Normal	
Levulose	Normal	
Maltose	Normal	
Mannitol	Normal	
Raffinose	Normal	
Salicin	Normal	
Sorbitol	Normal	
Sucrose	Normal	
Xylose	Normal	

Normal
 10 min
 20 min
 30 min
 45 min
 1 hr
 2 hr
 3 hr
 4 hr
 5 hr
 6 hr
 7 hr
 8 hr
 9 hr
 10 hr
 11 hr
 12 hr
 13 hr
 14 hr
 15 hr
 16 hr
 17 hr
 18 hr
 19 hr
 20 hr
 21 hr
 22 hr
 23 hr
 24 hr
 25 hr
 26 hr
 27 hr
 28 hr
 29 hr
 30 hr
 31 hr
 32 hr
 33 hr
 34 hr
 35 hr
 36 hr
 37 hr
 38 hr
 39 hr
 40 hr
 41 hr
 42 hr
 43 hr
 44 hr
 45 hr
 46 hr
 47 hr
 48 hr
 49 hr
 50 hr
 51 hr
 52 hr
 53 hr
 54 hr
 55 hr
 56 hr
 57 hr
 58 hr
 59 hr
 60 hr
 61 hr
 62 hr
 63 hr
 64 hr
 65 hr
 66 hr
 67 hr
 68 hr
 69 hr
 70 hr
 71 hr
 72 hr
 73 hr
 74 hr
 75 hr
 76 hr
 77 hr
 78 hr
 79 hr
 80 hr
 81 hr
 82 hr
 83 hr
 84 hr
 85 hr
 86 hr
 87 hr
 88 hr
 89 hr
 90 hr
 91 hr
 92 hr
 93 hr
 94 hr
 95 hr
 96 hr
 97 hr
 98 hr
 99 hr
 100 hr
 101 hr
 102 hr
 103 hr
 104 hr
 105 hr
 106 hr
 107 hr
 108 hr
 109 hr
 110 hr
 111 hr
 112 hr
 113 hr
 114 hr
 115 hr
 116 hr
 117 hr
 118 hr
 119 hr
 120 hr
 121 hr
 122 hr
 123 hr
 124 hr
 125 hr
 126 hr
 127 hr
 128 hr
 129 hr
 130 hr
 131 hr
 132 hr
 133 hr
 134 hr
 135 hr
 136 hr
 137 hr
 138 hr
 139 hr
 140 hr
 141 hr
 142 hr
 143 hr
 144 hr
 145 hr
 146 hr
 147 hr
 148 hr
 149 hr
 150 hr
 151 hr
 152 hr
 153 hr
 154 hr
 155 hr
 156 hr
 157 hr
 158 hr
 159 hr
 160 hr
 161 hr
 162 hr
 163 hr
 164 hr
 165 hr
 166 hr
 167 hr
 168 hr
 169 hr
 170 hr
 171 hr
 172 hr
 173 hr
 174 hr
 175 hr
 176 hr
 177 hr
 178 hr
 179 hr
 180 hr
 181 hr
 182 hr
 183 hr
 184 hr
 185 hr
 186 hr
 187 hr
 188 hr
 189 hr
 190 hr
 191 hr
 192 hr
 193 hr
 194 hr
 195 hr
 196 hr
 197 hr
 198 hr
 199 hr
 200 hr
 201 hr
 202 hr
 203 hr
 204 hr
 205 hr
 206 hr
 207 hr
 208 hr
 209 hr
 210 hr
 211 hr
 212 hr
 213 hr
 214 hr
 215 hr
 216 hr
 217 hr
 218 hr
 219 hr
 220 hr
 221 hr
 222 hr
 223 hr
 224 hr
 225 hr
 226 hr
 227 hr
 228 hr
 229 hr
 230 hr
 231 hr
 232 hr
 233 hr
 234 hr
 235 hr
 236 hr
 237 hr
 238 hr
 239 hr
 240 hr
 241 hr
 242 hr
 243 hr
 244 hr
 245 hr
 246 hr
 247 hr
 248 hr
 249 hr
 250 hr
 251 hr
 252 hr
 253 hr
 254 hr
 255 hr
 256 hr
 257 hr
 258 hr
 259 hr
 260 hr
 261 hr
 262 hr
 263 hr
 264 hr
 265 hr
 266 hr
 267 hr
 268 hr
 269 hr
 270 hr
 271 hr
 272 hr
 273 hr
 274 hr
 275 hr
 276 hr
 277 hr
 278 hr
 279 hr
 280 hr
 281 hr
 282 hr
 283 hr
 284 hr
 285 hr
 286 hr
 287 hr
 288 hr
 289 hr
 290 hr
 291 hr
 292 hr
 293 hr
 294 hr
 295 hr
 296 hr
 297 hr
 298 hr
 299 hr
 300 hr
 301 hr
 302 hr
 303 hr
 304 hr
 305 hr
 306 hr
 307 hr
 308 hr
 309 hr
 310 hr
 311 hr
 312 hr
 313 hr
 314 hr
 315 hr
 316 hr
 317 hr
 318 hr
 319 hr
 320 hr
 321 hr
 322 hr
 323 hr
 324 hr
 325 hr
 326 hr
 327 hr
 328 hr
 329 hr
 330 hr
 331 hr
 332 hr
 333 hr
 334 hr
 335 hr
 336 hr
 337 hr
 338 hr
 339 hr
 340 hr
 341 hr
 342 hr
 343 hr
 344 hr
 345 hr
 346 hr
 347 hr
 348 hr
 349 hr
 350 hr
 351 hr
 352 hr
 353 hr
 354 hr
 355 hr
 356 hr
 357 hr
 358 hr
 359 hr
 360 hr
 361 hr
 362 hr
 363 hr
 364 hr
 365 hr
 366 hr
 367 hr
 368 hr
 369 hr
 370 hr
 371 hr
 372 hr
 373 hr
 374 hr
 375 hr
 376 hr
 377 hr
 378 hr
 379 hr
 380 hr
 381 hr
 382 hr
 383 hr
 384 hr
 385 hr
 386 hr
 387 hr
 388 hr
 389 hr
 390 hr
 391 hr
 392 hr
 393 hr
 394 hr
 395 hr
 396 hr
 397 hr
 398 hr
 399 hr
 400 hr
 401 hr
 402 hr
 403 hr
 404 hr
 405 hr
 406 hr
 407 hr
 408 hr
 409 hr
 410 hr
 411 hr
 412 hr
 413 hr
 414 hr
 415 hr
 416 hr
 417 hr
 418 hr
 419 hr
 420 hr
 421 hr
 422 hr
 423 hr
 424 hr
 425 hr
 426 hr
 427 hr
 428 hr
 429 hr
 430 hr
 431 hr
 432 hr
 433 hr
 434 hr
 435 hr
 436 hr
 437 hr
 438 hr
 439 hr
 440 hr
 441 hr
 442 hr
 443 hr
 444 hr
 445 hr
 446 hr
 447 hr
 448 hr
 449 hr
 450 hr
 451 hr
 452 hr
 453 hr
 454 hr
 455 hr
 456 hr
 457 hr
 458 hr
 459 hr
 460 hr
 461 hr
 462 hr
 463 hr
 464 hr
 465 hr
 466 hr
 467 hr
 468 hr
 469 hr
 470 hr
 471 hr
 472 hr
 473 hr
 474 hr
 475 hr
 476 hr
 477 hr
 478 hr
 479 hr
 480 hr
 481 hr
 482 hr
 483 hr
 484 hr
 485 hr
 486 hr
 487 hr
 488 hr
 489 hr
 490 hr
 491 hr
 492 hr
 493 hr
 494 hr
 495 hr
 496 hr
 497 hr
 498 hr
 499 hr
 500 hr
 501 hr
 502 hr
 503 hr
 504 hr
 505 hr
 506 hr
 507 hr
 508 hr
 509 hr
 510 hr
 511 hr
 512 hr
 513 hr
 514 hr
 515 hr
 516 hr
 517 hr
 518 hr
 519 hr
 520 hr
 521 hr
 522 hr
 523 hr
 524 hr
 525 hr
 526 hr
 527 hr
 528 hr
 529 hr
 530 hr
 531 hr
 532 hr
 533 hr
 534 hr
 535 hr
 536 hr
 537 hr
 538 hr
 539 hr
 540 hr
 541 hr
 542 hr
 543 hr
 544 hr
 545 hr
 546 hr
 547 hr
 548 hr
 549 hr
 550 hr
 551 hr
 552 hr
 553 hr
 554 hr
 555 hr
 556 hr
 557 hr
 558 hr
 559 hr
 560 hr
 561 hr
 562 hr
 563 hr
 564 hr
 565 hr
 566 hr
 567 hr
 568 hr
 569 hr
 570 hr
 571 hr
 572 hr
 573 hr
 574 hr
 575 hr
 576 hr
 577 hr
 578 hr
 579 hr
 580 hr
 581 hr
 582 hr
 583 hr
 584 hr
 585 hr
 586 hr
 587 hr
 588 hr
 589 hr
 590 hr
 591 hr
 592 hr
 593 hr
 594 hr
 595 hr
 596 hr
 597 hr
 598 hr
 599 hr
 600 hr
 601 hr
 602 hr
 603 hr
 604 hr
 605 hr
 606 hr
 607 hr
 608 hr
 609 hr
 610 hr
 611 hr
 612 hr
 613 hr
 614 hr
 615 hr
 616 hr
 617 hr
 618 hr
 619 hr
 620 hr
 621 hr
 622 hr
 623 hr
 624 hr
 625 hr
 626 hr
 627 hr
 628 hr
 629 hr
 630 hr
 631 hr
 632 hr
 633 hr
 634 hr
 635 hr
 636 hr
 637 hr
 638 hr
 639 hr
 640 hr
 641 hr
 642 hr
 643 hr
 644 hr
 645 hr
 646 hr
 647 hr
 648 hr
 649 hr
 650 hr
 651 hr
 652 hr
 653 hr
 654 hr
 655 hr
 656 hr
 657 hr
 658 hr
 659 hr
 660 hr
 661 hr
 662 hr
 663 hr
 664 hr
 665 hr
 666 hr
 667 hr
 668 hr
 669 hr
 670 hr
 671 hr
 672 hr
 673 hr
 674 hr
 675 hr
 676 hr
 677 hr
 678 hr
 679 hr
 680 hr
 681 hr
 682 hr
 683 hr
 684 hr
 685 hr
 686 hr
 687 hr
 688 hr
 689 hr
 690 hr
 691 hr
 692 hr
 693 hr
 694 hr
 695 hr
 696 hr
 697 hr
 698 hr
 699 hr
 700 hr
 701 hr
 702 hr
 703 hr
 704 hr
 705 hr
 706 hr
 707 hr
 708 hr
 709 hr
 710 hr
 711 hr
 712 hr
 713 hr
 714 hr
 715 hr
 716 hr
 717 hr
 718 hr
 719 hr
 720 hr
 721 hr
 722 hr
 723 hr
 724 hr
 725 hr
 726 hr
 727 hr
 728 hr
 729 hr
 730 hr
 731 hr
 732 hr
 733 hr
 734 hr
 735 hr
 736 hr
 737 hr
 738 hr
 739 hr
 740 hr
 741 hr
 742 hr
 743 hr
 744 hr
 745 hr
 746 hr
 747 hr
 748 hr
 749 hr
 750 hr
 751 hr
 752 hr
 753 hr
 754 hr
 755 hr
 756 hr
 757 hr
 758 hr
 759 hr
 760 hr
 761 hr
 762 hr
 763 hr
 764 hr
 765 hr
 766 hr
 767 hr
 768 hr
 769 hr
 770 hr
 771 hr
 772 hr
 773 hr
 774 hr
 775 hr
 776 hr
 777 hr
 778 hr
 779 hr
 780 hr
 781 hr
 782 hr
 783 hr
 784 hr
 785 hr
 786 hr
 787 hr
 788 hr
 789 hr
 790 hr
 791 hr
 792 hr
 793 hr
 794 hr
 795 hr
 796 hr
 797 hr
 798 hr
 799 hr
 800 hr
 801 hr
 802 hr
 803 hr
 804 hr
 805 hr
 806 hr
 807 hr
 808 hr
 809 hr
 810 hr
 811 hr
 812 hr
 813 hr
 814 hr
 815 hr
 816 hr
 817 hr
 818 hr
 819 hr
 820 hr
 821 hr
 822 hr
 823 hr
 824 hr
 825 hr
 826 hr
 827 hr
 828 hr
 829 hr
 830 hr
 831 hr
 832 hr
 833 hr
 834 hr
 835 hr
 836 hr
 837 hr
 838 hr
 839 hr
 840 hr
 841 hr
 842 hr
 843 hr
 844 hr
 845 hr
 846 hr
 847 hr
 848 hr
 849 hr
 850 hr
 851 hr
 852 hr
 853 hr
 854 hr
 855 hr
 856 hr
 857 hr
 858 hr
 859 hr
 860 hr
 861 hr
 862 hr
 863 hr
 864 hr
 865 hr
 866 hr
 867 hr
 868 hr
 869 hr
 870 hr
 871 hr
 872 hr
 873 hr
 874 hr
 875 hr
 876 hr
 877 hr
 878 hr
 879 hr
 880 hr
 881 hr
 882 hr
 883 hr
 884 hr
 885 hr
 886 hr
 887 hr
 888 hr
 889 hr
 890 hr
 891 hr
 892 hr
 893 hr
 894 hr
 895 hr
 896 hr
 897 hr
 898 hr
 899 hr
 900 hr
 901 hr
 902 hr
 903 hr
 904 hr
 905 hr
 906 hr
 907 hr
 908 hr
 909 hr
 910 hr
 911 hr
 912 hr
 913 hr
 914 hr
 915 hr
 916 hr
 917 hr
 918 hr
 919 hr
 920 hr
 921 hr
 922 hr
 923 hr
 924 hr
 925 hr
 926 hr
 927 hr
 928 hr
 929 hr
 930 hr
 931 hr
 932 hr
 933 hr
 934 hr
 935 hr
 936 hr
 937 hr
 938 hr
 939 hr
 940 hr
 941 hr
 942 hr
 943 hr
 944 hr
 945 hr
 946 hr
 947 hr
 948 hr
 949 hr
 950 hr
 951 hr
 952 hr
 953 hr
 954 hr
 955 hr
 956 hr
 957 hr
 958 hr
 959 hr
 960 hr
 961 hr
 962 hr
 963 hr
 964 hr
 965 hr
 966 hr
 967 hr
 968 hr
 969 hr
 970 hr
 971 hr
 972 hr
 973 hr
 974 hr
 975 hr
 976 hr
 977 hr
 978 hr
 979 hr
 980 hr
 981 hr
 982 hr
 983 hr
 984 hr
 985 hr
 986 hr
 987 hr
 988 hr
 989 hr
 990 hr
 991 hr
 992 hr
 993 hr
 994 hr
 995 hr
 996 hr
 997 hr
 998 hr
 999 hr
 1000 hr

+	65	<u>MICROCOCCUS AUREUS</u>
-	60	<u>MICROCOCCUS V</u>
+	60	<u>MICROCOCCUS W</u>
+	60	<u>MICROCOCCUS X</u>
-	65	<u>MICROCOCCUS FLAVUS</u>
-	65	<u>MICROCOCCUS FLAVESCENS</u>
-	65	<u>MICROCOCCUS CITREUS</u>
-	65	<u>MICROCOCCUS CONJUGATUS</u>

Cult- ure no.	Form	Size microns	Gram Stain	Action on milk 37°C. 48 hrs.	Titrat- able acidity in milk 37°C. 48 hrs.	Growth on agar slants	Growth in beef broth	Nitrate reduction	Amino N. assimilation	Ammonic N. assimilation
---------------------	------	-----------------	---------------	--	---	--------------------------------	-------------------------------	----------------------	--------------------------	----------------------------

169	:singly	:1.0 to	:	:acid	:2.20 to	:	:	:	:	:
	to	:pairs	: 1.4	: +	:coag.	:2.8	:abun-	:turbid:	+	-
224	:clusters	:av. 1.2	:	:dig.	:av.2.50	:dant	:	:	:	:
225	:	:0.7 to	:	:acid	:0.16 to	:	:	:	:	:
	to	:singly	: 1.0	: +	:coag.	:0.20	:moder-	:turbid:	+	-
241	:clusters	:av. 0.9	:	:dig.	:av.0.18	:ate	:	:	:	:
242	:singly	:0.6 to	:	:acid	:0.60 to	:	:	:	:	:
	to	:pairs	: 1.0	: +	:coag.	:0.64	:slight	:turbid:	-	-
320	:clusters	:av. 0.8	:	:dig.	:av.0.52	:	:	:	:	:
321	:singly	:0.6 to	:	:acid	:0.60 to	:	:	:	:	:
	to	:pairs	: 1.2	: +	:coag.	:0.62	:abun-	:turbid:	+	-
405	:clusters	:av. 0.9	:	:dig.	:av.0.61	:dant	:	:	:	:
406	:singly	:1.0 to	:	:	:0.12 to	:	:	:	:	:
	to	:pairs	: 1.6	: +	:inert	:0.18	:slight	:turbid:	+	-
417	:clusters	:av. 1.3	:	:	:av.0.15	:	:	:	:	:
418	:	:1.5 to	:	:	:0.12 to	:	:	:	:	:
	to	:singly	: 2.0	: +	:alka-	:0.16	:moder-	:turbid:	-	-
424	:fours	:av. 1.8	:	:line	:av.0.14	:ate	:	:	:	:
425	:	:1.0 to	:	:	:0.12 to	:	:	:	:	:
	and	:singly	: 1.4	: +	:alka-	:0.14	:slight	:turbid:	-	-
426	:clusters	:av. 1.2	:	:line	:av.0.13	:	:	:	:	:

Table 2. (continued)

Nitrate reduction	Amino N. assimilation	Ammonic N. assimilation	Ammonia from peptone	Ammonia from nitrates	Gelatine liquefaction	Carbon dioxide equiv. 0.1 N. acid per 10 ml. milk	Hemolysis	Lipolysis	Diastatic action	Arabinose	Dextrine	Dextrose	Galactose	Glycerol	Inulin	Lactose	Levulose	Maltose	Mannitol	Raffinose
White group																				
+	-	-	+	-	crateriform	2.8 to 3.2 av. 3.0	-	-	-	-	+	+	+	-	-	+	+	+	-	-
+	-	-	+	-	crateriform	6.2 to 6.8 av. 6.5	-	-	-	-	+	+	+	-	-	+	+	+	-	-
-	-	-	+	-	stratiform	4.0 to 4.8 av. 4.4	-	-	-	-	+	+	+	+	-	+	+	+	+	-
+	-	-	+	-	stratiform	3.0 to 3.4 av. 3.2	-	-	-	-	+	+	+	+	-	+	+	+	-	-
+	-	-	+	-	stratiform	0.1 to 0.3 av. 0.2	-	-	-	-	+	+	+	+	+	+	+	+	-	-
-	-	-	+	-	stratiform	6.4 to 6.8 av. 6.6	-	-	-	-	+	+	+	+	-	+	+	+	-	-
-	-	-	+	-			-	-	-	-	-	-	-	-	-	+	-	-	-	-

Galactose	Glycerol	Inulin	Lactose	Levulose	Maltose	Mannitol	Raffinose	Salicin	Sorbitol	Sucrose	Xylose	Thermal death point (°C. for 15 minutes)	Classification
+	-	-	+	+	+	-	-	-	-	+	-	65	<u>Micrococcus albus</u>
-	-	-	+	+	-	-	-	-	-	-	-	55	<u>Micrococcus cremoris-viscosi</u>
+	+	-	+	+	+	+	-	+	+	+	-	55	<u>Micrococcus freudenreichii</u>
+	+	-	+	+	+	-	-	-	-	+	-	60	<u>Micrococcus caseolyticus</u>
+	+	+	+	+	+	-	-	-	-	+	+	65	<u>Micrococcus epidermidis</u>
+	+	-	-	+	+	-	-	-	-	-	+	55	<u>Micrococcus y</u>
-	-	-	+	-	-	-	-	-	-	-	+	55	<u>Micrococcus z</u>

data show the following relationships:

Action on litmus milk

Changes in litmus milk inoculated with proteolytic micrococci were of five general types:

1. No noticeable change other than an accumulation of cells at the bottom of tubes. This condition was most common with organisms of the yellow group.
2. Producing acid reaction without coagulation. This change was produced by organisms in all groups.
3. Producing alkaline reaction with digestion of protein. The change was produced only by organisms of the yellow and white groups.
4. Producing acid with coagulation and slight protein digestion. The change was noted with all groups of organisms.
5. Producing acid with coagulation and considerable protein digestion. This condition was most commonly produced by members of the orange group.

Although all the cultures were definitely proteolytic on milk agar, the change produced in the appearance of litmus milk by certain of them did not suggest proteolysis.

Titratable acidity produced in skimmilk

With 20 representative cultures titratable acidities varied from 0.12 to 0.66 per cent, calculated as lactic acid. In general, relatively high acidities were produced by members of the orange and white groups while comparatively low acidities were obtained with members of the yellow group. With a few white cultures, however, the

acidities were very low.

Growth on agar slants

All of the cultures studied grew well on beef infusion agar slants. In general, the yellow organisms grew more luxuriantly than those of the orange or white groups.

Growth in beef broth

The organisms regularly produced turbidity in beef broth with the formation of sediment.

Nitrate reduction

A majority of cultures studied reduced nitrates. With the orange group, 50 of 53 cultures were positive; with the yellow group 55 of 115 were positive; and with the white group, 168 of 258 were positive.

Nitrogen assimilation and ammonia production

The assimilation of amino nitrogen was negative with all but a few organisms of the yellow group. Ammonia nitrogen was assimilated by many organisms of the orange and yellow groups; with the white group, the results were negative in all trials.

All of the organisms produced ammonia from peptone, whereas none were capable of producing ammonia from nitrates.

Gelatin liquefaction

All of the cultures liquefied gelatine with the exception of a few in the white group. In general, liquefaction was more rapid and extensive with members of the orange groups than with members of the yellow or white groups.

Carbon dioxide production

Members of the yellow group generally produced larger amounts of carbon dioxide than members of either the orange or white groups.

Hemolysis

Hemolysis was shown by 49 cultures of the orange group and 17 cultures of the yellow group; in the white group, none were hemolytic.

Lipolysis

Lipolytic action by the organisms was limited to the orange group in which all of the cultures gave positive results. Apparently none of the yellow or white organisms studied were lipolytic according to the methods employed.

Diastatic action

The production of diastase was exhibited by 1 culture in the orange group and 96 cultures in the yellow group.

None of the organisms in the white group gave positive results.

Carbohydrate fermentation

In general, most of the cultures studied fermented dextrose, galactose, lactose, levulose and maltose. The fermentation of dextrine and xylose was confined principally to members of the white group, while glycerol, mannitol, salicin, sorbitol and sucrose were fermented by some cultures of the orange and white groups. Arabinose and raffinose were not fermented.

Thermal death point

In determining the thermal death points of the organisms, cotton stoppered tubes containing broth cultures were suspended in a water bath at 55°, 60°, 65°, and 70°C. for 15 minutes. All of the organisms were killed at temperatures ranging from 55° to 65°C.

Acetylmethylcarbinol plus diacetyl, volatile acidity, hydrogen sulfide and indol production

Acetylmethylcarbinol plus diacetyl was not produced from milk, with or without added citric acid, by any of the cultures.

Appreciable amounts of volatile acid were not produced by the organisms. Tests for hydrogen sulfide and

indol were negative in all instances.

Pathogenicity, spore formation and motility

To determine the pathogenicity of 50 representative cultures of proteolytic micrococci, the organisms were grown in lactose broth and 2 ml. of a culture was inoculated intraperitoneally into a guinea pig. At the same time guinea pigs were inoculated with cultures classified as Staphylococcus by the Bacteriology Department of the Ohio State University. The animals inoculated with proteolytic micrococci were not affected, whereas several of the cultures from Ohio produced death in 12 to 24 hours.

None of the proteolytic micrococci produced spores and all of the organisms were apparently non-motile.

Protein Breakdown in Skimmilk by Proteolytic
Micrococci

Four representative cultures of proteolytic micrococci were selected for the study of protein breakdown in skimmilk. These were cultures 59 (orange), 98 and 110 (yellow) and 170 (white).

Skimmilk, in approximately 200 ml. portions, was sterilized in pint bottles after the weight of each bottle and its contents had been recorded. Several bottles were then inoculated with each culture to be studied and incubated at 37°C. or 21°C. After 2, 4 and 7 day incubation

periods, the serum from each culture was analysed. The serum from a culture was obtained by adding distilled water to make up to the original weight, adding 3 ml. of glacial acetic acid and heating in a bath of boiling water with frequent agitation. After cooling, the culture was filtered through paper. The following determinations were made on 5 ml. portions of the serum: Total nitrogen, amino nitrogen, and the nitrogen soluble and insoluble in trichloroacetic acid, ethyl alcohol and phosphotungstic acid. The values were expressed as milliliters of N/10 acid equivalent to the nitrogen in 5 ml. of serum.

Three trials were carried out with the four organisms. The data for one of the trials are presented in Table 3; the data for the other trials are not given since they are practically identical with those given in Table 5.

All cultures of the test organisms produced protein breakdown in skim milk although the extent of proteolysis varied considerably with the individual cultures. At 37°C., the total nitrogen in the serum of skim milk inoculated with the organisms increased with continued holding. After 7 days, the values with some cultures were relatively high, especially with culture 170. At 21°C., the values for total nitrogen also increased with continued holding and in some instances the increases were greater than at 37°C. Although there were irregularities with the different incubation periods, in general, cultures 170 and 98 produced

Table 3.

Protein breakdown in skimmilk by certain proteolytic micrococci

Culture no.	Incubation time (days)	ml. of 0.1 normal acid equiv. to nitrogen of 5 ml. of skimmilk serum	Total N.	Nitrogen fractionated into sol. and insol. portions with				* Amino N.		
				Trichloroacetic acid	Ethyl alcohol	Phosphotungstic acid				
				Sol.	Insol.	Sol.	Insol.	Sol.	Insol.	
Held at 37°C.										
39		2.9	2.9	0.2	1.3	1.9	1.3	2.0	1.75	
170		8.0	7.5	1.2	4.5	3.7	1.4	6.5	1.92	
98	2	2.6	2.4	0.5	1.4	1.4	1.0	1.7	1.14	
110		2.7	1.8	0.6	1.6	1.3	1.3	1.5	1.35	
Control		2.6	2.6	0.5	1.3	0.6	1.2	1.5	0.92	
39		3.0	2.8	0.4	1.3	1.9	1.4	2.0	1.83	
170		11.2	9.9	1.0	6.8	4.8	2.4	8.8	3.35	
98	4	3.5	2.8	0.3	1.2	2.5	1.4	1.9	1.42	
110		3.0	2.5	0.5	1.2	2.1	1.7	1.8	1.89	
Control		2.7	2.5	0.4	1.3	1.3	1.0	1.9	0.64	
39		3.8	3.4	0.6	1.6	1.9	1.3	2.5	1.93	
170		11.5	11.0	0.5	6.2	4.8	2.2	8.7	3.59	
98	7	7.4	6.6	0.5	4.3	2.8	2.6	4.7	3.14	
110		4.1	3.3	0.6	2.5	1.3	1.9	2.1	1.92	
Control		2.5	2.4	0.5	1.3	1.4	1.1	1.5	0.71	

Table 3. (continued)

Culture no.	Incubation time (days)	:ml. of 0.1 normal acid equiv. to nitrogen of 5 ml. of skimmilk serum									
		Total N.	Nitrogen fractionated into sol. and insol. portions with				* Amino N.				
			Trichloroacetic acid	Ethyl alcohol	Phosphotungstic acid	Sol.	Insol.	Sol.	Insol.	Sol.	Insol.
Held at 21°C.											
39		3.3	2.5	0.6	1.7	2.0	1.5	1.9	1.73		
170		5.7	5.7	0.4	2.5	2.9	1.0	4.2	2.00		
98	2	2.4	2.1	0.4	1.2	1.4	1.0	1.7	1.50		
110		2.5	2.4	0.2	1.4	1.4	1.4	1.5	1.50		
Control		2.7	2.6	0.5	1.5	1.2	1.2	1.5	0.85		
39		4.8	4.5	0.4	2.0	3.0	1.5	3.5	2.14		
170		8.4	3.1	0.5	4.4	4.1	1.1	7.5	2.28		
98	4	2.7	2.4	0.3	1.6	1.3	1.5	1.5	1.57		
110		3.0	2.7	0.5	1.7	1.3	1.4	2.0	1.64		
Control		2.8	2.2	0.4	1.5	1.2	1.4	1.5	0.73		
39		5.7	5.5	0.4	2.9	3.0	1.2	4.7	2.32		
170		11.5	9.8	1.4	6.1	5.5	2.1	9.8	3.57		
98	7	2.8	2.5	0.5	1.5	1.5	0.3	1.6	1.59		
110		9.4	9.5	0.1	6.2	3.1	3.0	6.1	2.93		
Control		2.5	2.5	0.1	1.2	1.5	0.9	1.5	0.71		

* Calculated from milligrams of amino nitrogen as obtained by the Van Slyke gasometric method.

more breakdown at the higher temperature, while cultures 39 and 110 produced more breakdown at the lower one.

The distribution of nitrogen in the various fractions was much the same with the different organisms and with the two incubation temperatures. The fractions soluble in trichloroacetic acid were relatively large and increased with continued incubation, while the values for the fraction insoluble in trichloroacetic acid, as would be expected, were regularly low even after 7 days incubation. The amounts of nitrogen soluble and insoluble in ethyl alcohol were more or less the same, and each of the amounts generally increased with continued incubation. When phosphotungstic acid was used as the precipitant, most of the nitrogen was contained in the insoluble fractions; the soluble fractions remained about the same during incubation, whereas the insoluble fractions generally increased.

The amounts of amino nitrogen increased with continued incubation at both temperatures. There was a very general relationship between the values obtained for total nitrogen and those obtained for amino nitrogen; when the values for total nitrogen were high, those for amino nitrogen were also high.

General Action of Proteolytic Micrococci
on Unsalted Butter

The general action of 58 cultures of proteolytic micrococci on butter was studied by churning small portions of sterile cream inoculated with test organisms; this was done by agitating in a motor driven shaking machine. The cream was prepared by placing 500 ml. portions in 1-quart glass top fruit jars and heating in an autoclave at 15 pounds pressure for 30 minutes on 2 consecutive days. A culture of each test organism was prepared by inoculating 100 ml. of sterile skim milk and incubating at 37°C. for 48 hours. At the time of churning all of a milk culture was added to a jar of cream. The relatively large inoculation was used to insure many test organisms in the butter since washing of the butter would remove many of them. From four to six jars of cream constituted a series, one of the jars being left uninoculated as a control.

The butter was washed three times with sterile water and worked after each washing; the working after the last washing was intended to give a good distribution of the moisture droplets. All the equipment used in handling the butter was sterile. The butter was left unsalted so that the conditions for growth would be more satisfactory than if salt were added. Each churning was then divided and held in sterile petri dishes at about 4°C. and 20°C. At several intervals during the holding periods of from 1 to

8 weeks the butter was scored and criticized.*

The control butter in each series was treated in the same manner as the butter containing the test organisms. In general, the control butter kept reasonably well, although occasionally it developed mold growth or an off flavor, presumably as a result of a contamination. When this occurred the entire series of churnings was discarded. The uniformity of scores on the control butter and the fresh butter made from cream inoculated with the test organisms can be explained by the fact that the cream used in all the churnings came from the same source and that the cream was sterilized rather than pasteurized. The heat treatment of the cream caused an oily flavor in the butter, but this was considered preferable to the use of pasteurized cream which might contain considerable numbers of organisms in addition to the test organism.

Action of orange micrococci

Fourteen cultures of the orange micrococci were studied from the standpoint of their action on butter. The data on one of these cultures, which was considered representative, are given in Table 4. Culture No. 49 brought about relatively rapid changes in the flavor of butter. After 4 to 7 weeks at 4°C. or after 1 to 2 weeks at 20°C. definite flavor defects

* Scores and criticisms were given by Prof. F. H. Herzer of the Dairy Department, Mississippi State College.

Table 4.

General action of orange culture No. 49 on butter

Churning no.	Temp. held (°C.)	Period for develop- ment of definite defect (weeks)	Score and general criticism on butter		Decrease in score
			At churning	After storage	
1	4	6	90 oily, flat	88 tallowy sl* rancid	2
	20	2	90 oily, flat	85 rancid	5
2	4	7	90 oily, flat	86.5 rancid	3.5
	20	2	90 oily, flat	87.5 sl. rancid	2.5
3	4	6	90 oily, flat	87 sl. rancid	3
	20	1	90 oily, flat	88 cheesy	2
4	4	4	90 oily, flat	87.5 sl. rancid	2.5
	20	1	90 oily, flat	87.5 sl. rancid	2.5
5	4	6	90 oily, flat	88 sl. rancid	2
	20	1	90 oily, flat	88 sl. rancid	2

* Sl. = slightly

were apparent. A rancid flavor generally developed in the samples although in churning No. 5 a distinct cheesy flavor was evident at 20°C. The scores on the butter decreased from 2 to 5 points between the time of churning and the end of the holding periods.

Table 5 summarizes the data on the cultures of orange micrococci. The results show that relatively rapid changes were brought about in the flavor of butter by the strains of orange micrococci employed. When the samples were held at 4°C., rancid, cheesy or pungent flavors commonly developed after 4 to 7 weeks, while with samples held at 20°C. the same general defects were conspicuous after 1 to 5 weeks. A few of the cultures failed to produce a rancid or putrid defect at 4°C., which presumably was due to a slow growth at relatively low temperature. Decreases in the scores of samples during the holding varied from 1.5 to 5 points. The decreases in score were frequently greater with the relatively short holding at 20°C. than with the longer holding at 4°C.

As a group the orange proteolytic micrococci are lipolytic so that action on both the protein and fat of butter would be expected if the holding temperature and other conditions permitted growth. Various organisms which attack protein and fat give both cheesy and rancid flavors in butter, although commonly a rancid flavor tends to predominate.

Table 5.

General action of orange proteolytic micrococci on butter

Culture no.	No. of chummings	Temp. held (°C.)	Period for development of definite defect (weeks)	Average score and general criticism on butter		Decrease in score
				At churning	After storage	
7	3	4	5	90 oily, flat	86 rancid cheesy	4
		20	5	90 oily, flat	86 rancid	4
8	1	4	5	90 oily, flat	87 rancid	3
		20	1	90 oily, flat	85.5 bitter	4.5
10	3	4	6	90 oily, flat	88 oily cheesy	2
		20	3	90 oily, flat	85 rancid	5
12	1	4	4	90 oily, flat	86 oily bitter	4
		20	1	90 oily, flat	85 rancid pungent	5
13	4	4	6	90 oily, flat	86 rancid	4
		20	1	90 oily, flat	85 pungent	5
18	2	4	7	90 oily, flat	88 bitter tallowy	2
		20	2	90 oily, flat	88 sl.* rancid	2
30	5	4	6	90 oily, flat	88 sl. rancid	2
		20	3	90 oily, flat	85 rancid pungent	5

Table 5. (continued)

Culture no.	No. of churnings	Temp. held (°C.)	Period for development of definite defect (weeks)	Average score and general criticism on butter		Decrease in score
				At churning	After storage	
31	2	4	4	90 oily, flat	86 sl. rancid	2
		20	2	90 oily, flat	85 putrid	5
33	5	4	6	90 oily, flat	87 sl. rancid	3
		20	1	90 oily, flat	88 sl. rancid	2
34	2	4	8	90 oily, flat	83.5 oily	1.5
		20	1	90 oily, flat	86 rancid	4
36	6	4	6	90 oily, flat	87 sl. rancid	3
		20	2	90 oily, flat	86 rancid	4
39	7	4	6	90 oily, flat	85.5 oily storage	1.5
		20	2	90 oily, flat	85 pungent	5
43	2	4	8	90 oily, flat	83.5 oily storage	1.5
		20	2	90 oily, flat	86 putrid	4
49	5	4	6	90 oily, flat	88 tallowy sl. rancid	3
		20	2	90 oily, flat	86 rancid	4

* Sl. = Slightly

151

Action of yellow micrococci

The action of the yellow micrococci on butter was studied with 11 cultures. The data on one representative culture are given in Table 6. Culture No. 63 produced a variety of changes in the flavor of butter. After 4 to 8 weeks at 4°C. or 2 to 5 weeks at 20°C., flavor defects were conspicuous and included musty, bitter, cheesy, tallowy and rancid flavors. The score on the butter decreased from 0.5 to 3.0 points between the time of churning and the end of the holding.

Table 7 summarizes the results obtained with the 11 yellow cultures. The data show that changes were produced in the flavor of butter by most of the yellow strains studied although the changes occurred relatively slowly. When samples were held at 4°C., the most common defects were cheesy, musty or bitter flavors which developed after 4 to 8 weeks. With the samples held at 20°C., similar defects were detected in from 2 to 5 weeks. Decreases in the scores of the samples held at both temperatures varied from 0.5 to 3.5 points; in some cases the decreases in score were considerably greater at 20°C. than at 4°C. In general, members of the yellow group were relatively slow in producing definite flavor defects in butter compared to members of the orange group (Table 5). Although in three churnings a slight rancid flavor was detected in the butter, the group as a whole is nonlipolytic. The chief defect in butter caused by members of this group was a cheesy flavor which was probably due to the protein breakdown.

Table 6.

General action of yellow culture No. 63 on butter

Churning no.	Temp. held (°C.)	Period for develop- ment of definite defect (weeks)	Score and general criticism on butter		Decrease in score
			at churning	after storage	
1	4	5	90 oily, flat	87.5 bitter rusty	2.5
	20	2	90 oily, flat	87.5 oily rusty	2.5
2	4	8	90 oily, flat	89 sl.* tallowy	1
	20	5	90 oily, flat	87 sl. rancid	3
3	4	5	90 oily, flat	88 bitter	2
	20	4	90 oily, flat	87.5 cheesy	2.5
4	4	4	90 oily, flat	88.5 storage	0.5
	20	3	90 oily, flat	87 musty	3
5	4	7	90 oily, flat	88.5 oily	0.5
	20	4	90 oily, flat	88 oily	2

sl. = slightly

Table 7.

General action of yellow proteolytic microcci on butter

Cul- ture no.	No. of churn- ings	Temp. held (°C.)	Period for development of definite defect (weeks)	Average score and general criticism on butter		Decrease in score
				At churning	After storage	
65	5	4	8	90 oily, flat	88.5 oily	1.5
		20	4	90 oily, flat	86.5 cheesy musty	3.5
67	4	4	6	90 oily	88 oily cheesy	2
		20	4	90 oily	88 oily musty	2
74	2	4	4	90 oily, flat	83.5 sl.* rancid	1.5
		20	2	90 oily, flat	86.5 rancid	3.5
85	6	4	7	90 oily, flat	88 sl. rancid	2
		20	5	90 oily, flat	87 cheesy musty	3
97	4	4	7	90 oily, flat	87.5 cheesy sl.	2.5
		20	4	90 oily, flat	88 cheesy	2
98	5	4	6	90 oily, flat	88 musty sl.	2
		20	3	90 oily, flat	88 cheesy	2
99	2	4	7	90 oily	88.5 sl. musty	1.5
		20	3	90 oily	88 cheesy	2

* Sl. = Slightly

Table 7 (continued)

Cul- ture no.	No. of churn- ings	Temp. held (°C.)	Period for development of definite defect (weeks)	Average score and general criticism on butter		Decrease in score
				At churning	After storage	
110	1	4	6	90 oily, flat	88.5 storage	1.5
		20	3	90 oily, flat	87.5 cheesy musty	2.5
116	1	4	4	90 oily, flat	89 oily	1
		20	4	90 oily, flat	87 cheesy	3
154	1	4	5	90 grass flavor	89.5 oily	0.5
		20	5	90 grass flavor	87 bitter	3
157	1	4	7	90 oily	89 oily	1
		20	2	90 oily	88.5 oily sl.* rancid	1.5

*Sl. = slightly

Action of white micrococci

Fourteen cultures of the white micrococci were studied from the standpoint of their action on butter. The data on one of these cultures, which was considered representative, are given in Table 8. Butter containing culture No. 175 commonly developed a cheesy flavor after 4 to 7 weeks at 4°C. or after 1 to 3 weeks at 20°C. The scorings were rather uniform and the decreases in score during holding ranged from 1 to 3 points.

Table 9 summarizes the data obtained with the 14 white cultures. The results indicate that the organisms brought about comparatively slow changes in the flavor of butter at 4°C., while relatively rapid changes were noted at 20°C. At 4°C., oily or bitter flavors developed after 4 to 6 weeks, whereas at 20°C. most of the samples developed a cheesy flavor in 1 to 4 weeks. Decreases in the scores of samples during holding varied from 0.5 to 4 points. In general, the defects caused by this group were not as pronounced as those brought about by the orange cultures (Table 5.), and were more comparable with those caused by the yellow cultures (Table 7.).

The principal defect in butter caused by the white proteolytic micrococci was a cheesy or bitter flavor which presumably was due to protein decomposition in the butter. No lipolytic cultures were studied in this group and in only a few cases was the butter found to have a slightly rancid flavor.

Table 8.

General action of white culture No. 175 on butter

Churning no.	Temp. held (°C.)	Period for develop- ment of definite defect (weeks)	Score and general criticism on butter		Decrease in score
			At churning	After storage	
1	4	4	90 oily, flat	88.5 musty	1.5
	20	1	90 oily, flat	88 bitter	2
2	4	4	90 oily, flat	88 oily bitter	2
	20	3	90 oily, flat	88 cheesy	2
3	4	7	90 oily, flat	88.5 sl.* tallowy	1.5
	20	2	90 oily, flat	88.5 cheesy	1.5
4	4	5	90 oily, flat	88 storage	2
	20	3	90 oily, flat	88 cheesy	2
5	4	6	90 oily, flat	88.5 cheesy	1.5
	20	3	90 oily, flat	87.5 oily sl. cheesy	2.5
6	4	6	90 oily, flat	89 oily	1
	20	2	90 oily, flat	87 oily cheesy	3

* Sl. = Slightly

Table 9.

General action of white proteolytic micrococci on butter

Cul- ture no.	No. of churn- ings	Temp. held (°C.)	Period for the development of definite defect (weeks)	Average score and general criticism on butter		Decrease in score		
				At churning	After storage			
175	6	4	6	90	oily, flat	83	oily	2
		20	2	90	oily, flat	87.5	cheesy	2.5
200	2	4	6	90	oily, flat	89	oily storage	1.5
		20	2	90	oily, flat	88.5	oily	1
250	5	4	6	90	oily, flat	89	oily	1
		20	2	90	oily, flat	88.5	oily bitter	1.5
260	3	4	5	90	oily, flat	89	sl. bitter	1
		20	4	90	oily, flat	88	cheesy	2
274	3	4	4	90	oily, flat	89	oily	1
		20	2	90	oily, flat	87	sl. rancid	3
286	2	4	4	90	oily, flat	89	oily	1
		20	2	90	oily, flat	87.5	cheesy	2.5
289	1	4	4	90	oily, flat	88.5	oily bitter	1.5
		20	1	90	oily, flat	88	oily bitter	2

* SL. = Slightly

Table 9. (continued)

Cul- ture no.	No. of churn- ings	Temp. held (°C.)	Period for the development of definite defect (weeks)	Average score and general criticism on butter		Decrease in score		
				At churning	After storage			
310	1	4	4	90	oily, flat	89.5	oily	0.5
		20	5	90	oily, flat	87	flat sl* cheesy	3
315	4	4	6	90	oily, flat	87.5	cheesy	2.5
		20	2	90	oily, flat	87	oily bitter	3
317	2	4	4	90	oily, flat	85	oily storage	2
		20	1	90	oily, flat	87.5	cheesy	2.5
325	4	4	5	90	oily, flat	88	oily cheesy	2
		20	3	90	oily, flat	87.5	sl. rancid	2.5
328	2	4	4	90	oily, flat	87	musty bitter	3
		20	2	90	oily, flat	86	musty cheesy	4
355	1	4	6	90	oily, flat	86.5	oily	0.5
		20	2	90	oily, flat	87.5	cheesy	2.5
412	2	4	5	90	oily, flat	85	oily sl. rancid	2
		20	5	90	oily, flat	87.5	tainted	2.5

* Sl. = slightly

Species of Proteolytic Micrococci

An attempt was made to identify, on a species basis, the 426 cultures of proteolytic micrococci isolated from various dairy products. All the organisms were put in the genus Micrococcus, although some of them are included in the genus Staphylococcus by Bergey (3). This arrangement is in agreement with Hucker (25) who has considered the advisability of placing the organisms that have been variously described as micrococci, staphylococci and rhodococci in the single genus Micrococcus. It meets the problem of classifying the organisms of this general group that are so widely distributed in dairy products better than a division into two or more genera. The characters given for the various species by Bergey (3) were used in the identification.

Orange Group

Forty-eight of the 53 cultures of orange micrococci were identified as Micrococcus aureus; of these cultures, 39 were isolated from milk, 6 from butter, 1 from cheese and 2 from ice cream.

Five of the orange cultures could not be identified on the basis of the species described by Bergey (3). Some investigators would probably consider them M. aureus, but they differ from M. aureus in so many respects that a distinction seems advisable. The 5 cultures appear to repre-

sent three species which are designated Micrococcus v, Micrococcus w and Micrococcus x in Table 2.

Micrococcus v

Two strains of this organism were isolated from milk. They differ from M. aureus in their action on litmus milk, nitrate reduction and carbohydrate fermentation and from Micrococcus w and Micrococcus x in action on litmus milk, ammonia nitrogen assimilation and carbohydrate fermentation.

Micrococcus w

One strain of this culture was isolated from milk. It differs from M. aureus in nitrate reduction, diastatic action and its ability to ferment dextrine, maltose, salicin and sorbitol, and from Micrococcus x in action on litmus milk, nitrate reduction, ammonia nitrogen assimilation and carbohydrate fermentation.

Micrococcus x

Two strains of this culture were isolated from milk. They differ from M. aureus in action on litmus milk, ammonia nitrogen assimilation, hemolysis and ability to ferment maltose, salicin and sorbitol.

Yellow Group

Forty-one cultures of the 115 yellow micrococci were

identified as Micrococcus flavus; of these, 35 were isolated from milk, 2 from butter and 4 from ice cream. Eighteen cultures were identified as Micrococcus flavescens; 12 of these were isolated from milk and 6 from cheese. Twenty-four cultures were identified as Micrococcus citreus; of these, 6 were obtained from milk, 4 from butter, 2 from cheese and 12 from ice cream. Thirty-two cultures were identified as Micrococcus conglomeratus; all of these were isolated from milk.

White Group

Fifty-six of the 258 white micrococci were identified as Micrococcus albus; 37 of these were isolated from milk, 10 from butter, 4 from cheese and 5 from ice cream. Seventeen cultures were identified as Micrococcus cremoris-viscosi; all of these were isolated from milk. Eighty-eight cultures were identified as Micrococcus freudenreichii; of these 73 were isolated from milk, 8 from butter and 7 from cheese. Eighty-four cultures were identified as Micrococcus caseolyticus; 68 of these were obtained from milk, 8 from butter and 8 from ice cream. Eleven cultures were identified as Micrococcus epidermidis; all of these were isolated from milk.

Eight of the white cultures could not be identified on the basis of the species described by Bergey (3). These cultures appear to represent two species which are designated

Micrococcus y and Micrococcus z in Table 2.

Micrococcus y

Six cultures were obtained from milk. They differ from the identified species of micrococci in their action on litmus milk and the fermentation of certain carbohydrates and from Micrococcus z in their ability to ferment certain carbohydrates.

Micrococcus z

Two cultures were isolated from milk. They differ from the identified species of micrococci in their action on milk and the fermentation of certain carbohydrates.

Key to the Identification of Proteolytic
Micrococci

- A. Orange pigment produced
 - 1. Nitrates reduced
 - a. Acid in litmus milk without coagulation
Micrococcus x
 - aa. Acid and coagulation in litmus milk
Micrococcus aureus
 - 2. Nitrates not reduced
 - a. Acid and coagulation in litmus milk
Micrococcus w
 - aa. No change in litmus milk
Micrococcus v
- B. Yellow pigment produced
 - 1. Nitrates reduced

- a. Acid reaction in litmus milk
Micrococcus citreus
- aa. No change in litmus milk
Micrococcus conglomeratus
- 2. Nitrates not reduced
 - a. Acid reaction in litmus milk
Micrococcus flavus
 - aa. No change in litmus milk
Micrococcus flavescens
- C. White pigment produced
 - 1. Nitrates reduced
 - a. Acid reaction in litmus milk
 - b. Ferment sucrose
 - c. Ferment glycerol
Micrococcus caseolyticus
 - cc. Do not ferment glycerol
Micrococcus albus
 - bb. Do not ferment sucrose
Micrococcus cremoris-viscosi
 - aa. No change in litmus milk
Micrococcus epidermidis
 - 2. Nitrates not reduced
 - a. Alkaline reaction in litmus milk
 - b. Ferment dextrose
Micrococcus y
 - bb. Do not ferment dextrose
Micrococcus z
 - aa. Acid reaction in litmus milk
Micrococcus freudenreichii

DISCUSSION OF RESULTS

The relatively large numbers of proteolytic micrococci encountered in detailed studies on the distribution of these organisms in various dairy products is in agreement with the observations of investigators who had repeatedly obtained proteolytic micrococci in attempts to isolate other types of organisms that attack protein of milk. They also suggest that the organisms may be of importance from the standpoint of various changes in dairy products. Although many investigators have considered the micrococci a more or less inert group, producing changes rather slowly, various cultures were obtained which brought about comparatively rapid changes in milk and unsalted butter. Extensive proteolysis was produced in milk by certain of the organisms in relatively short periods and butter made from cream inoculated with the organisms showed definite flavor defects rather early.

The defects produced in butter indicated both proteolysis and lipolysis by the organisms. The fact that some of the organisms studied were isolated from cheese suggests they may be important in the cheese ripening process, since proteolysis and lipolysis appear to be of significance in the flavor development of various cheeses.

In detecting proteolysis by micrococci, it appears that culturing on milk agar plates is of greater value than cultur-

ing in litmus milk, since certain cultures which gave very evident proteolysis on milk agar did not produce changes in the appearance of litmus milk justifying the conclusion that the organisms were proteolytic. With some cultures other types of changes in milk, such as acid development, tended to obscure proteolysis and possibly to retard it. Some of the proteolytic micrococci did not liquefy gelatine. This deviation from a relationship that might be expected is in agreement with observations reported for organisms belonging to various genera.

The relatively minor effect of various factors on the color production of the cultures on agar emphasizes the value of chromogenesis in the classification of the organisms.

The viability of the cultures on agar and in milk parallels the general resistance of the micrococci to various unfavorable conditions that has been noted in investigations on these organisms. It indicates that contamination of dairy products with material containing these organisms may be greatly delayed without the organisms being destroyed. The general viability greatly simplifies the carrying of cultures.

The cultures studied were all placed in the genus Micrococcus since, in general, the differences among them did not appear to justify division into two or more genera. An important point in this connection was the failure of representative cultures to kill guinea pigs with which they were inoculated, since the genus Staphylococcus is characterized

by Bergey (3) as "usually parasitic". There is precedent for including all the cultures in the genus Micrococcus in Hucker's treatment (25) of this general group.

While most of the characters that have been used by various investigators in classifying the organisms of the genus Micrococcus are of importance, it appears that more attention should be given to the changes produced in litmus milk and that hydrolysis of fat should be included as a differentiating character. The changes produced in litmus milk permit considerable differentiation of the organisms. The detection of fat hydrolysis is now relatively simple because of the development of methods that are easily applied.

The cultures that could not be identified by the descriptions of organisms listed by Bergey (3) were not given species names for several reasons. There has been a definite attempt, especially by Hucker (24), to reduce the number of recognized species of micrococci because the large number of species described has resulted in much confusion. It appears that new species should only be suggested when the evidence is complete and when a considerable number of cultures is available for study, particularly from the standpoint of the variations to be expected within the genus. If additional isolations of the cultures are made, species names must eventually be proposed.

SUMMARY AND CONCLUSIONS

By plating on milk agar and incubating at 37°C. for 48 to 72 hours, proteolytic micrococci were readily isolated from many samples of milk, butter, cheese and ice cream. They were obtained from 340 of 580 samples of milk, from 36 of 42 sample of butter, from 20 of 36 samples of cheese (cheddar) and from 30 of 46 samples of ice cream.

On the basis of chromogenesis, the proteolytic micrococci were readily divided into four groups--red, orange, yellow and white (non-pigmented). The red micrococci were not studied in detail because of the small numbers encountered.

Studies on the influence of various factors on chromogenesis indicated:

- (a) That when grown in an atmosphere of carbon dioxide, the general pigment production on agar was not significantly different than when grown in air, although with a few of the orange cultures the pigment was lighter in color.
- (b) That when grown on potato, some of the orange organisms produced a deeper color than when grown on agar.
- (c) That the pH value of the medium had an effect on the pigment production, a value of 7.0 being more favorable than values of 6.6 or 7.8, although pigment production was definite at all three values.
- (d) That holding organisms in sterile water for 6 days before

inoculating on agar did not influence pigment production.

- (e) That subjecting cultures to 55°C. for 10 minutes before inoculating on agar had little or no effect on pigment production.
- (f) That more highly colored pigment was produced at 21°C. than at 37°C. with the orange cultures but not with the yellow cultures.
- (g) That agar cultures held in darkness for 10 days were not as deeply pigmented as when held in a sun lighted room.

Milk cultures of the proteolytic micrococci were viable for 3 to 4 months and agar slant cultures for 10 to 12 months at room temperature.

When litmus milk was inoculated with the various organisms, four general types of changes were produced while a few cultures produced no noticeable change.

The titratable acidity produced in milk by 20 representative cultures varied from 0.12 to 0.66 per cent, calculated as lactic acid. Organisms of the orange and white groups generally gave higher values than those of the yellow group.

All of the cultures studied grew well on beef infusion agar slants and in beef broth.

A majority of the cultures studied reduced nitrates.

In general, the organisms did not assimilate amino

nitrogen, although ammonia nitrogen was assimilated by many cultures of the orange and yellow groups but by none of the white group. All cultures were capable of producing ammonia from peptone, whereas none produced ammonia from nitrates.

All cultures except a few of the white group liquefied gelatine.

Cultures of the yellow group generally produced larger amounts of carbon dioxide than cultures of the orange or white groups.

Hemolysis was shown by some cultures of the orange and yellow groups but by none of the white group.

Lipolysis was limited to organisms of the orange group. None of the yellow or white organisms studied were lipolytic, according to the methods used.

Diastatic action was exhibited by 1 culture of the orange group and by 96 of the yellow group. None of the white cultures produced diastase.

In general, most of the cultures studied fermented dextrose, galactose, lactose, levulose and maltose. Dextrine, glycerol, mannitol, salicin, sorbitol, sucrose and xylose were fermented by some cultures.

Arabinose and raffinose were not fermented.

In broth cultures all of the organisms were destroyed at temperatures ranging from 55° to 65°C. for 15 minutes.

Appreciable amounts of acetylmethylcarbinol plus diacetyl or volatile acid were not produced by the organisms.

Tests for hydrogen sulphide and indol were regularly negative.

Thirty representative cultures were found to be non-pathogenic when inoculated into guinea pigs intraperitoneally.

None of the organisms studied produced spores, and all were apparently non-motile.

Protein breakdown in skimmilk by four representative cultures of proteolytic micrococci was studied in detail. When grown in milk at 37° or 21°C. for 2 to 7 days, the organisms increased the amount of nitrogen in the serum, although the extent of proteolysis varied considerably with the cultures. In general, the fractions of nitrogen soluble in trichloroacetic acid, soluble and insoluble in ethyl alcohol or insoluble in phosphotungstic acid increased during the incubation, whereas the fractions insoluble in trichloroacetic acid or soluble in phosphotungstic acid remained about the same. The amounts of amino nitrogen increased in the serum during the holding.

The general action of 38 cultures of proteolytic micrococci on unsalted butter was studied by churning small portions of sterile cream inoculated with relatively large numbers of the organisms, and then examining the butter after various periods at 4° and 20°C. Definite defects were produced in many samples of butter after 4 to 8 weeks at 4°C. or 1 to 5 weeks at 20°C. Certain cultures of the orange group produced rancid, cheesy and pungent flavors, certain of the yellow group produced cheesy, musty or bitter flavors, while certain of the white group brought about oily or bitter flavors. In

general, members of the orange group changed the flavor of butter more rapidly and more extensively than members of the yellow or white groups.

An attempt was made to identify, on a species basis, the 426 cultures of proteolytic micrococci. The organisms were separated into 15 species of the genus Micrococcus on the basis of chromogenesis, action on nitrates, action on milk and the fermentation of various carbohydrates. Ten species were identified from descriptions given by other investigators, while 5 species were unidentified. A key to the identification of the 15 species of proteolytic micrococci is given.

ACKNOWLEDGMENTS

The author wishes to express his sincere appreciation to Dr. B. W. Hammer for his wise counsel in planning and directing the work herein reported and in the preparation of the manuscript; to Dr. C. B. Lane for his kind assistance on nitrogen work, and to Prof. F. H. Herzer for his assistance in scoring butter.

LITERATURE CITED

1. Allen, P. W. A simple method for the classification of bacteria as to diastase production. *Jour. Bact.* 3: 15-17. 1918.
2. Bailey, S. F. and Lacy, G. R. A modification of the Kligler lead acetate medium. *Jour. Bact.* 13: 183-189. 1927.
3. Bergey, D. H. Manual of determinative bacteriology. 4th Ed. Williams and Wilkins Co. Baltimore. 1954.
4. Breed, Alice F. Micrococci present in the normal cow's udder. N. Y. (Geneva) Agr. Exp. Sta., Tech. Bul. 152. 1928.
5. Buchanan, R. E. Studies on the nomenclature and classification of the bacteria IV. Subgroups and genera of the Coccaceae. *Jour. Bact.* 2: 605-617. 1917.
6. Burkey, L. A. Methods for determining the persistence of certain important bacteria in Swiss cheese. *Jour. Bact.* 21: 40-41. 1931.
7. Chester, F. D. A manual of determinative bacteriology. Macmillan Co. New York. 1901.
8. Cohn, F. Untersuchungen über Bakterien. *Beitr. z. Biol. d. Pflanzen*, 1 (Heft 2): 127-224. 1872. Original not seen; cited by G. J. Hucker, 22, p.7.
9. Cohn, F. Organismen in der Pockenlymphe. *Virchow's Arch. f. Path. Anat.*, 55: 229-258. 1872 Original not seen; cited by G. J. Hucker, 22, p.7.
10. Cohn, F. Untersuchungen über Bakterien. II. *Beitr. z. Biol. d. Pflanzen* 1: (Heft 3): 141-207. 1875. Original not seen; cited by G. J. Hucker, 22, p.7.
11. Copeland, Lynn, and Olson, T. M. The bacterial flora of normal cow's udder. S. D. Agr. Exp. Sta., Bul. 218. 1926.
12. Dudgeon, L. S. The differentiation of the Staphylococci. *Jour. Path. and Bact.* 12: 242-257. 1908.

13. Eldredge, E. E. and Rogers, L. A. The bacteriology of cheese of the Emmental type. *Centbl. Bakt.* 2. Abt. 40: 5-21. 1914.
14. Evans, Alice C. The bacteria of milk freshly drawn from normal udders. *Jour. Inf. Dis.* 18: 437-476. 1916.
15. Frazier, W. C. and Rupp, Phillip. Studies on the proteolytic bacteria of milk. II. Action of the proteolytic bacteria of milk on milk. *Jour. Bact.* 16: 65-78. 1936.
16. Freudenreich, E. von. Beitrag zur Kenntnis der Ursachen des bitteren Käses und der bitteren Milch. *Landw. Jahrb. d. Schweiz*, 8: 135-143. 1894. (Original not seen) Cited in Jordan, E. O. and Falk, I. S. A newer knowledge of bacteriology and immunology. p. 379; University Chicago Press. 1928.
17. Gorini, C. The acidoproteolytes in the pasteurization of milk for the manufacture of soft cheeses of "Bel Paese" type. *Proc. 10th World's Dairy Congress, Rome 10*: Sec. 3: 7-15. 1934.
18. Hallier, E. Bewis, dass der Micrococcus der Infektionskrankheiten keimfähig und von höheren Pilzformen abhängig ist, und Widerlegung der leichsinnigen Angriffe des Herrn Collegen Bary zu Halle. *Leitschrift für Parasitenkunde.* 2: 1-20. 1870.
19. Hamner, B. W. and Bailey, D. E. The volatile acid production of starters and of organisms isolated from them. *Iowa Agr. Exp. Sta., Res. Bul.* 55. 1919.
20. Hamner, B. W. and Collins, M. A. The numbers of lipolytic bacteria in various dairy products, as determined with Nileblue sulphate. *Iowa Agr. Exp. Sta., Res. Bul.* 169. 1934.
21. Harding, H. A. and Wilson, K. J. A study of the udder flora of cows. *N. Y. (Geneva) Agr. Exp. Sta., Tech. Bul.* 27. 1913.
22. Hucker, G. J. Studies on the Coccaceae. I. Previous taxonomic studies concerning the genera of the Coccaceae. *N. Y. (Geneva) Agr. Exp. Sta., Tech. Bul.* 99. 1924.

23. Hucker, G. J. The types of bacteria found in commercial cheddar cheese. N. Y. (Geneva) Agr. Exp. Sta., Tech. Bul. 90. 1922.
24. Hucker, G. J. Studies on the Coccaceae II. A study of the general characters of the micrococci. N. Y. (Geneva) Agr. Exp. Sta., Tech. Bul. 100. 1924.
25. Hucker, G. J. Studies on the Coccaceae IV. Classification of the genus Micrococcus Cohn. N. Y. (Geneva) Agr. Exp. Sta., Tech. Bul. 102. 1924.
26. Kligler, I. J. A systematic study of the Coccaceae in the collection of the Museum of Natural History. Jour. Inf. Dis. 12: 452-453. 1915.
27. Knowles, H. R. A study of the bacterial flora of fore-milk and of rennet extract with special reference to acid-proteolytic types. Jour. Dairy Res. 7: 63-74. 1936.
28. Lane, C. B. and Hammer, B. W. Bacteriology of cheese. I. Effect of pasteurizing the milk on the nitrogenous decomposition in cheddar cheese. Iowa Agr. Exp. Sta., Res. Bul. 183. 1935.
29. Long, H. F. A study of some lipolytic microorganisms isolated from dairy products. Unpublished Thesis, Iowa State College Library, Ames. 1936.
30. Michaelian, H. B. and Hammer, B. W. Studies on acetyl-methylcarbinol and diacetyl in dairy products. Iowa Agr. Exp. Sta., Res. Bul. 179. 1935.
31. Migula, W. System der Bakterien. G. Fischer, Jena. 1897.
32. Orla-Jensen, S. Dairy Bacteriology. P. Blakiston's Son and Co. Philadelphia. 1921.
33. Society of American Bacteriologists, Committee on Bacteriological Technique Manual of methods for pure culture study of bacteria. 2: Leaflet 5. 1934.
34. Van Slyke, D. D. A method for quantitative determination of aliphatic amino groups. Jour. Bio. Chem. 9: 185. 1911.

35. Winslow, C. E.-A. and Winslow, A. R. The systematic relationships of the Coccaceae. John Wiley and Sons, New York. 1908.
36. Winslow, C. E.-A., Rothberg, W., and Parsons, E. I. Notes on the classification of the white and orange staphylococci. Jour. Bact. 5: 145-167. 1920.
37. Winslow, C. E.-A., Broadhurst, Jean, Buchanan, R. E., Krumwiede, Charles, Rogers, L. A., and Smith, G. H. The families and genera of the bacteria. Jour. Bact. 5: 191-329. 1920.