



Arginine dephosphorylation propels spore germination in bacteria

Bing Zhou^{a,b}, Maja Semanjski^c, Natalie Orlovetskie^b, Saurabh Bhattacharya^b, Sima Alon^b, Liron Argaman^b, Nayef Jarrous^b, Yan Zhang^a, Boris Macek^c, Lior Sinai^{b,1}, and Sigal Ben-Yehuda^{b,1}

^aCollege of Food Science and Nutritional Engineering, National Engineering Research Center for Fruits and Vegetables Processing, China Agricultural University, 10083 Beijing, China; ^bDepartment of Microbiology and Molecular Genetics, Institute for Medical Research Israel-Canada, The Hebrew University-Hadassah Medical School, The Hebrew University of Jerusalem, 91120 Jerusalem, Israel; and ^cProteome Center Tuebingen, University of Tuebingen, 72076 Tuebingen, Germany

Edited by Caroline S. Harwood, University of Washington, Seattle, WA, and approved May 6, 2019 (received for review October 16, 2018)

Bacterial spores can remain dormant for years but possess the remarkable ability to germinate, within minutes, once nutrients become available. However, it still remains elusive how such instant awakening of cellular machineries is achieved. Utilizing *Bacillus subtilis* as a model, we show that YwE arginine (Arg) phosphatase is crucial for spore germination. Accordingly, the absence of the Arg kinase McsB accelerated the process. Arg phosphoproteome of dormant spores uncovered a unique set of Arg-phosphorylated proteins involved in key biological functions, including translation and transcription. Consequently, we demonstrate that during germination, YwE dephosphorylates an Arg site on the ribosome-associated chaperone Tig, enabling its association with the ribosome to reestablish translation. Moreover, we show that Arg dephosphorylation of the housekeeping σ factor A (SigA), mediated by YwE, facilitates germination by activating the transcriptional machinery. Subsequently, we reveal that transcription is reinitiated at the onset of germination and its recommencement precedes that of translation. Thus, Arg dephosphorylation elicits the most critical stages of spore molecular resumption, placing this unusual post-translational modification as a major regulator of a developmental process in bacteria.

Bacillus subtilis | spore dormancy | germination | arginine phosphorylation

In response to nutrient deprivation, the Gram-positive bacterium *Bacillus subtilis* (*B. subtilis*) and its relatives have the capacity to initiate a developmental process, termed sporulation, which culminates in the formation of a highly resilient durable spore (1–4). Sporulation commences with the formation of a polar septum, asymmetrically dividing the cell into a small forespore and a larger mother cell compartment. Subsequently, the forespore is engulfed by the mother cell, and through sequential activation of cell type specific σ factors, the spore matures within the mother cell (5, 6). Spore development is accompanied by the formation of a peptidoglycan layer called the cortex, surrounded by two proteinaceous coats, conferring spore robustness (7–12). The forespore core then undergoes dehydration, mainly by accumulating pyridine-2, 6-dicarboxylic acid [dipicolinic acid or (DPA)] in complex with Ca²⁺ (13). Eventually, the phase-bright spore is liberated by lysis of the mother cell but intriguingly can still undergo significant molecular changes, at least for a few days, influencing its emergence from quiescence (14–16).

Spores can maintain dormancy for years yet possess the remarkable ability to rapidly resume a vegetative life form once nutrients become available. This revival process includes three key phases: germination, ripening, and outgrowth (17–19). Germination, the earliest revival event, is characterized by transition from a phase-bright spore to a phase-dark cell (20), whereas no morphological changes are evident during the intermediate ripening period, exploited by the spore for molecular reorganization (18). The last phase of revival is outgrowth, in which the spore intensively synthesizes macromolecules to rebuild a rod shaped cell that emerges from the disintegrating spore shells (17, 21).

Germination is considered the most enigmatic event in the process of spore revival, lasting only for a few minutes (19). During germination, the spore undergoes rehydration, release of DPA, hydrolysis of the cortex, and coat disassembly (17, 21, 22). This phase is associated with the loss of heat resistance and a decrease in optical density (20). Recently, we demonstrated that translation is rapidly activated after DPA release and is required for the subsequent events of germination (23, 24), challenging the traditional view that germination occurs without the need for any macromolecule synthesis (13, 16, 17, 25, 26). Furthermore, we showed that protein synthesis during germination was facilitated by the bona fide translational factors RpmE, a ribosomal subunit, and Tig, a peptidyl-prolyl *cis-trans* isomerase, that is a ribosome-associated chaperone (23, 27–29).

Germination is triggered by binding of nutrients, termed germinants, to multiple germination receptors located in the spore membrane (30). Germinant factors include single amino acids, sugars, purine nucleosides, and cell wall muropeptides (17, 19, 31). In *B. subtilis*, the GerA receptor binds L-alanine (L-Ala) to induce germination, whereas GerB and GerK receptors respond jointly to a mixture of germinants consisting of asparagine (Asp), glucose, fructose, and potassium (AGFK) ions (17, 21, 22). This germinant sensing stimulates downstream effectors activating germination; however, relatively little is known about this signal transmission pathway. A potential mechanism for the fast resumption of multiple cellular processes in the germinating spore

Significance

Dormant bacterial spores can survive long periods of time, withstanding extreme conditions, but can rapidly resume a vegetative life form once nutrients become accessible. The key event of this revival process is termed germination, the earliest phase, lasting only for a few minutes, during which cellular awakening is established. Yet, the molecular events propelling this enigmatic phase remain unknown. Here, we revealed that dephosphorylation of Arg residues on key target proteins triggers germination in the model bacterium *B. subtilis*. We further demonstrate that Arg dephosphorylation of a ribosome-associated factor reactivates translation, whereas dephosphorylation of the major housekeeping σ factor reinitiates transcription. Our results provide a whole mechanism for spore awakening in bacteria.

Author contributions: B.Z., M.S., N.O., S.B., S.A., L.A., N.J., Y.Z., B.M., L.S., and S.B.-Y. designed research; B.Z., M.S., N.O., and L.S. performed research; B.Z., M.S., N.O., B.M., L.S., and S.B.-Y. analyzed data; and B.Z., L.S., and S.B.-Y. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Published under the PNAS license.

¹To whom correspondence may be addressed. Email: lior.sinai@mail.huji.ac.il or sigalb@ekmd.huji.ac.il.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1817742116/-DCSupplemental.

Published online June 20, 2019.

is protein phosphorylation, a ubiquitous means for mediating rapid cellular responses to various external stimuli in bacteria (32, 33). Evidence has been provided that peptidoglycan-derived muropeptides induce germination by binding to the Ser/Thr kinase PrkC, which appears to phosphorylate the translation elongation factor G (EF-G) (31, 34). However, the impact of this phosphorylation event on germination is still unexplored, and the absence of PrkC has no effect on the process of germination driven by nutrient germinants (31, 35). More recently, we described the dynamic of the Ser/Thr/Tyr phosphoproteome during germination, identifying a plethora of 155 phosphorylation sites, modified during germination, providing evidence that phosphorylation events facilitate spore revival. However, no role in promoting germination was assigned to these events (35).

Arg phosphorylation, an emerging protein modification, has been lately described to occur in *B. subtilis* and *Staphylococcus aureus* (36, 37). The kinase responsible for Arg phosphorylation in *B. subtilis* was shown to be McsB, which counteracts the protein Arg phosphatase YwIE (36–41). Here, we demonstrate that YwIE drives the progression of spore germination by dephosphorylating Arg phosphosites of target proteins involved in key cellular processes. Furthermore, we show that YwIE mediates the rapid reactivation of the translational machinery by dephosphorylating the translational component Tig, enabling its association with the ribosome. Surprisingly, we found that Arg dephosphorylation of the housekeeping factor SigA by YwIE is crucial for the execution of germination, and subsequently, we discovered that transcription is reestablished at the onset of the process.

Results

Spore Germination Is Driven by Arg Dephosphorylation. To identify genes required for spore germination, we designed a transposon-based genetic screen, searching for mutants that are able to form mature spores but deficient in the conversion from a phase-bright to a phase-dark state following germinant addition (*SI Appendix, Fig. S1A*). Using this approach, we identified a mutant disrupted in the *ywIE* gene, encoding an Arg phosphatase (39), exhibiting severe germination deficiency (*Fig. 1A*). Constructing a strain, fully deleted of *ywIE*, revealed that it was able to grow and sporulate normally but produced spores that were perturbed in germination as indicated by only a slight decrease in optical density achieved in response to the efficient L-Ala germinant (*SI Appendix, Figs. S1 B and C and S2A*). Monitoring germination efficiency by time lapse microscopy at a single spore resolution showed that 30 min following L-Ala addition only 20% of the $\Delta ywIE$ spores could convert from a phase-bright to a phase-dark state in comparison with 96% of the wild type (WT) spores (*Fig. 1 B and C*). Similar results were obtained when germination was induced by supplementing the spores with AGFK and even with LB (*SI Appendix, Fig. S2 B, C, and E*). In line with these findings, $\Delta ywIE$ mutant spores were slower to release DPA and lose their heat resistance relative to the WT spores (*Fig. 1D and SI Appendix, Fig. S2D*), signifying that YwIE is a key factor in prompting spore germination.

The role of YwIE as an Arg phosphatase brings about the possibility that some crucial factors required for germination are phosphorylated on their Arg sites during sporulation, a modification that should be removed to allow germination. Notably, only a single kinase, McsB, is known to be responsible for Arg phosphorylation in *B. subtilis* (38, 40). Hence, we hypothesized that, in $\Delta mcsB$ spores, these factors would be constantly in their dephosphorylated active form, and consequently, these spores would germinate rapidly. Indeed, $\Delta mcsB$ mutant spores germinated faster than WT spores as evidenced by their remarkable rapid transition into the phase-dark state in all tested conditions (*Fig. 1 B and C and SI Appendix, Fig. S2*). Furthermore, germinating $\Delta mcsB$ spores turned heat sensitive and released DPA faster than WT spores (*Fig. 1D and SI Appendix, Fig. S2D*). Examining

germination of $\Delta mcsB \Delta ywIE$ double mutant spores revealed kinetics similar to that of $\Delta mcsB$ spores (*SI Appendix, Fig. S3*). Taken together, we infer that Arg phosphorylation of target proteins is executed by McsB during sporulation and is removed by YwIE to propel spore germination.

Revealing the Arg Phosphoproteome of Dormant Spores. Having established that Arg dephosphorylation is required for spore germination, we next sought to identify Arg-phosphorylated sites in the dormant spore proteomes that could serve as substrates for YwIE during spore germination. As Arg phosphorylation was shown to be more stable and enriched in the *ywIE* mutant (37, 40), we carried out Arg-phosphoproteomic analysis of $\Delta ywIE$ dormant spores. In total, we identified 18 Arg-phosphorylation sites located in 18 proteins with very high confidence using stringent quality criteria for the validation of the phosphorylation sites (Table 1 and *SI Appendix, Table S1*). A comparison with the Arg phosphoproteome of vegetative cells (37, 40) revealed that Arg-phosphorylation events in spores are less frequent and largely differ from those of vegetative cells with 10 identified sites unique to spores (Table 1 and *SI Appendix, Table S1*). The detected Arg-phosphorylated proteins are involved in key biological functions, including carbon metabolism, sporulation, translation, and transcription (Table 1). Thus, we uncovered a unique set of protein Arg-phosphorylation sites existing in dormant spores that can be potentially dephosphorylated by YwIE to allow germination. Among the identified phosphorylated factors, the most intriguing were those involved in transcription and translation, raising the possibility that they provide the means for reestablishment of these basic cellular processes.

Dephosphorylation of an Arg Site on the Translational Factor Tig Powers Spore Germination. Previously, we revealed that protein synthesis takes place during germination and relies on the bona fide translational factor Tig (23, 27), identified in our phosphoproteomic analysis to harbor an Arg45-phosphorylated site (Table 1). Interestingly, the occurrence of this phosphorylation event was monitored in vegetative cells following exposure to heat stress (40). We reasoned that phosphorylation of Tig in spores blocks its activity, whereas dephosphorylation by YwIE restores the factor utility during germination. To investigate the impact of the Arg45 site on the activity of Tig, we introduced mutations that would either abolish the phosphorylation potential by replacing Arg with Ala or mimic a constitutive phosphorylation state by replacing Arg with Asp (42). Each mutation was inserted as a sole copy at the native *tig* chromosomal locus, replacing the original WT allele. Remarkably, *tig-R45D* mutant spores exhibited germination defects similar to that of Δtig spores, whereas *tig-R45A* mutant spores germinated similarly to WT spores as indicated by optical density and time lapse microscopy analyses (*Fig. 2 A and B and SI Appendix, Fig. S4 A–D*). Consistently, mimicking constitutive phosphorylation by replacing the Arg45 site with Glu (R45E), harboring a larger side chain group, led to a germination defect similar to that of R45D protein (*SI Appendix, Fig. S5*). Importantly, both *tig-R45A* and *tig-R45D* mutations had no effect on vegetative growth or sporulation (*SI Appendix, Fig. S6 A and B*). Further analysis of the *tig* mutant spores showed that, upon germination induction, *tig-R45D* spores were capable of releasing DPA and accordingly lost their heat resistance (*SI Appendix, Fig. S4 E and F*). Hence, similar to Δtig spores, *tig-R45D* spores initiated germination normally but were subsequently stalled in their phase-bright state.

To corroborate that the Tig Arg-phosphorylation state influences the function of Tig, we examined its impact on protein synthesis during germination. We employed the BioOrthogonal Non-Canonical Amino acid Tagging (BONCAT) protein tagging technique that enables specific labeling of newly synthesized proteins due to the incorporation of an artificial amino acid termed azidohomoalanine (AHA), which is a substitute for




Fig. 1. Spore germination is facilitated by Arg dephosphorylation. (A) Spores of PY79 [wild type (WT)] and *ywIE::Tn* were incubated with L-Ala (10 mM) and observed by light microscopy before ($t = 0$) and after ($t = 30$ min) L-Ala addition. Shown are phase contrast images from a representative experiment out of three independent biological repeats. (Scale bar: 1 μm .) (B) Spores of PY79 (WT), BZ16 ($\Delta ywIE$), and BZ129 ($\Delta mcsB$) strains were incubated on agarose supplemented with L-Ala (10 mM) and monitored by time lapse microscopy. Shown are phase contrast images from a representative experiment out of three independent biological repeats. (Scale bar: 1 μm .) (C) Spores of PY79 (WT), BZ16 ($\Delta ywIE$), and BZ129 ($\Delta mcsB$) strains were incubated on agarose supplemented with L-Ala (10 mM) and monitored by time lapse microscopy. Data are presented as percentages of the initial number of the phase-bright spores. Shown are average values and SDs obtained from three independent biological repeats ($n \geq 300$ for each strain). (D) Spores of PY79 (WT), BZ16 ($\Delta ywIE$), and BZ129 ($\Delta mcsB$) strains were incubated with L-Ala to trigger germination. DPA release to the medium was determined by Tb-DPA assay. Presented are relative fluorescence units (RFUs) measured at 545 nm with excitation at 270 nm. Shown is a representative experiment out of three independent biological repeats.

methionine (43). This assay was utilized to establish the requirement for Tig to facilitate translation during germination (23). To this end, spores of WT, *tig-R45D*, and *tig-R45A* strains were induced to germinate with L-Ala in the presence of AHA without supplementing any additional nutrients. Then, proteins were extracted from spores at $T = 0$ and 30 min postgermination induction, tagged with biotin, and subjected to dot blot analysis using antibiotin antibodies. Whereas a strong signal was readily detected from WT and *tig-R45A* germinating spores, only a faint signal was monitored from *tig-R45D* germinating spores, similar to that obtained from Δtig spores (Fig. 2C). In line with these findings, an apparent delay in protein synthesis during germination was detected in *tig-R45D* mutant spores by time lapse microscopy using MalS-GFP as a reporter for monitoring translation early in

germination (23) (SI Appendix, Fig. S7 A and B). Finally, Western blot analysis substantiated that *tig-R45D* mutation influences specifically the translation of MalS during germination but did not impact translation of RpsB, an essential ribosomal subunit, and purine nucleoside phosphorylase (PupG), which were not included in the germination proteome (23, 44, 45) (Fig. 2D). Taken together, spores harboring the phosphorylated-like *tig-R45D* allele are impaired in protein synthesis at the course of germination, similar to Δtig spores, highlighting the potential importance of this phosphosite for Tig activity.

Tig Association with Ribosomes Depends on Dephosphorylation of the Arg45 Site. During translation, Tig associates with the ribosome and interacts with newly synthesized nascent chains to chaperone the folding of cytosolic proteins (27). Since *tig-R45D*

Table 1. Proteins phosphorylated on Arg residues and their phosphosites in spores

Protein	Description	Phosphosite
Translation		
Tig	Trigger factor	pR45
Tuf	Elongation factor G	pR265
RplN	50S ribosomal protein L17	pR7
Transcription		
SigA	RNA polymerase σ factor	pR365
Protein quality control		
GroEL	Chaperonin	pR35
ClpC	Class III stress response-related ATPase	pR70
Carbon metabolism		
GapA	Glyceraldehyde-3-phosphate dehydrogenase	pR199
YtsJ	NADP-dependent malic enzyme	pR27
MenB	Dihydroxynaphthoic acid synthetase	pR269
Pgi	Glucose-6-phosphate isomerase	pR5
Sporulation		
SspA	Small acid-soluble spore protein A	pR55
SspB	Small acid-soluble spore protein B	pR65
SspE	Small acid-soluble spore protein E	pR16
SpolVA	Stage IV sporulation protein A	pR454
CotF	Spore coat protein	pR155
YtfJ	Germination protein	pR125
Unknown		
YeaD	Unknown function	pR250
YtxH	Unknown function	pR151

Shown is the phosphoproteome of BZ16 ($\Delta ywIE$) dormant spores. The names and descriptions of proteins are given according to the SubtiWiki (60). Shown in bold font are phosphosites, which were also detected in vegetative cells (37, 40).

mutant spores showed delayed protein synthesis during germination, we hypothesized that Arg45 phosphorylation interfered with Tig binding to ribosomes. To examine this premise, we investigated the localization of Tig and ribosomes in WT and *tig* mutants in vegetative growing cells whereby the typical ribosome localization to the cell circumference surrounding the nucleoid is profound (46, 47). Tig-R45D mutant protein was fused to GFP, and its localization was followed concomitantly with the RpsB-mCherry fusion, signifying ribosome position (18). Whereas Tig and RpsB perfectly colocalized, the signals from Tig-R45D and RpsB showed only marginal overlap (Fig. 3A). Furthermore, although it is challenging to covisualize RpsB-mCherry and Tig-GFP in the small sized spores, we could detect the specific localization of Tig-GFP to the spore periphery during germination, a pattern that was hardly visible in germinating *tig-R45D* spores (SI Appendix, Fig. S7C). Interestingly, examining the localization of Tig-R45A revealed partial association with RpsB (Fig. 3A and SI Appendix, Fig. S7C), highlighting the significance of the R45 position for proper association with the ribosome. Nevertheless, this partial association was sufficient to propel germination (Fig. 2). Of note, fusing GFP to WT Tig or to the phosphomimetic mutant proteins did not affect their germination capabilities (SI Appendix, Fig. S6C).

To further substantiate the notion that Tig-R45D could not bind to ribosomes, we utilized a direct biochemical approach, purifying ribosomes from vegetative cells and spores of *tig* and

tig-R45D strains. Subsequently, we carried out Western blot analysis to detect the levels of Tig in cell extracts (CEs) and in purified ribosomes (PRs). Although similar levels of Tig or Tig-R45D were detected in CEs, Tig-R45D levels were considerably lower in PR fractions from vegetative cells, dormant and germinating spores (Fig. 3B). Of note, the levels of the ribosomal component RpsB were relatively equal in PRs derived from *tig* and *tig-R45D* strains (SI Appendix, Fig. S7D). These results reinforce the view that phosphorylation on Arg45 of Tig causes its dissociation from ribosomes.

Germination Is Promoted by Arg Dephosphorylation of SigA. Intriguingly, Arg phosphorylation at position Arg365 of SigA, the house keeping σ factor of *B. subtilis* (48), was identified in our Arg-phosphoproteome analysis hinting that, similar to translation, transcription might be required to facilitate germination. Furthermore, the phosphorylation of this site appeared unique to spores and was not identified in vegetative cells (37, 40) (Table 1). To test whether the Arg365-phosphorylation state has an impact on spore germination, we constructed mutant strains in which Arg365 was replaced by either Ala or Asp and assessed their ability to germinate. The *sigA-R365A* mutant spores could germinate as efficiently as WT spores (Fig. 4 A and B and SI Appendix, Fig. S8). However, spores harboring the *sigA-R365D* mutation, mimicking the phosphorylated form of the protein, exhibited a profound germination defect with only 44% of the spores switching to phase-dark 30 min postgermination induction compared with 96% of the WT spores (Fig. 4 A and B and SI Appendix, Fig. S8). A similar germination defect was obtained in spores from a strain carrying the phosphomimetic R365E mutation (SI Appendix, Fig. S9). A slight decrease in vegetative growth could be assigned to R365D mutations but only when the cells were grown in minimal medium, although no apparent effect on sporulation could be detected (SI Appendix, Fig. S6). Additionally, monitoring the protein levels of germination receptors as well as of Tig and RpmE, required for germination (23), in *sigA-R365D* and WT dormant spores did not yield any significant difference (SI Appendix, Fig. S10). Examination of DPA release and heat resistance showed that germinating *sigA-R365D* mutant spores were halted before DPA release and remained heat resistant (Fig. 4 C and D), indicating a deficiency at an early stage of the process before protein translation (SI Appendix, Fig. S4 E and F). Thus, we surmise that the dephosphorylation of SigA promotes germination.

Notably, combining *sigA-R365D* with *tig-R45D* mutation failed to recapitulate the extent of the $\Delta ywIE$ germination defect (Fig. 4E and SI Appendix, Fig. S11A), indicating that, as inferred by our phosphoproteomic analysis (Table 1), additional substrates are dephosphorylated by YwIE during the process. In a similar manner, spores harboring both *sigA-R365A* and *tig-R45A* mutations did not display the accelerated germination phenotype exhibited by $\Delta mcsB$ spores (SI Appendix, Fig. S11 A and B).

Evidence That the Transcriptional Machinery Is Reactivated in Germination. The finding that SigA dephosphorylation is necessary for prompting germination suggested that transcription is reestablished during this phase. To explore this notion, we monitored the transcriptional activity of whole extracts derived from vegetative cells, dormant or germinating spores of WT, and *sigA-R365A* and *sigA-R365D* strains by the addition of radioactively labeled ribonucleotides. Transcriptional activity was evident in the extract from vegetative cells with a minor reduction observed for *sigA-R365D* cells (SI Appendix, Fig. S12A). This is in line with the slight decrease in growth seen for this mutant in minimal medium (SI Appendix, Fig. S6D). Intriguingly, no detectable transcription took place in extracts of dormant spores (Fig. 5A), indicating that the transcription machinery was halted at the course of dormancy. Nonetheless, triggering germination by L-Ala was sufficient to




Fig. 2. Dephosphorylation of the translational factor Tig is required for spore germination. (A) Spores of PY79 (WT), BZ98 (*tig-R45A*), BZ97 (*tig-R45D*), and LS38 (Δ *tig*) strains were incubated on agarose supplemented with L-Ala (10 mM) and monitored by time lapse microscopy. Shown are phase contrast images from a representative experiment out of three independent biological repeats. (Scale bar: 1 μ m.) (B) Spores of PY79 (WT), BZ98 (*tig-R45A*), BZ97 (*tig-R45D*), and LS38 (Δ *tig*) strains were incubated on agarose supplemented with L-Ala (10 mM) and monitored by time lapse microscopy. Data are presented as percentages of the initial number of the phase-bright spores. Shown are average values and SDs obtained from three independent biological repeats ($n \geq 300$ for each strain). (C) Spores of LS5 (Δ *metE*), BZ103 (Δ *metE*, *tig-R45A*), BZ102 (Δ *metE*, *tig-R45D*), and LS80 (Δ *metE*, Δ *tig*) strains were induced to germinate with L-Ala in the presence of AHA for 30 min. Shown is a dot blot analysis of protein samples (in duplicate) that were collected before ($t = 0$, dormant spores) and after ($t = 30$ min) germination induction. Samples (1:1 and 1:10 dilution) were spotted on a membrane that was subsequently probed with antibiotin antibodies. The obtained signal was compared with known amounts of biotinylated BSA. (D) Protein extracts from vegetative cells, dormant and germinating spores of strains: (i) AR71 (WT), BZ107 (*tig-R45A*), and BZ106 (*tig-R45D*) carrying *malS-gfp*; (ii) BZ118 (WT), BZ120 (*tig-R45A*), and BZ119 (*tig-R45D*) carrying *rpsB-mCherry*; and (iii) AR68 (WT), BZ144 (*tig-R45A*), and BZ145(*tig-R45D*) carrying *pupG-gfp* were subjected to Western blot analysis using an antibody against GFP or mCherry, respectively. Equal amounts of protein extracts were loaded. Germination was induced by suspending the spores in 10 mM L-Ala for 10 min.

release this inhibition and robustly reactivate the transcriptional machinery (Fig. 5A). Furthermore, although restoration of transcription was apparent in extracts from both WT and *sigA-R365A* germinating spores, it was nearly undetectable in extracts of *sigA-R365D* germinating spores (Fig. 5A), substantiating the idea that dephosphorylation of SigA mediated this molecular awakening.

Next, we wished to detect the transcription of specific genes *in vivo* during germination. To this end, qRT-PCR assays were performed on RNA extracted from vegetative cells, dormant and germinating spores of WT, and *sigA-R365A* and *sigA-R365D* strains, focusing on transcripts of Tig and RpmE, encoding for proteins synthesized during germination (23).

Remarkably, the mRNA levels of *tig* and *rpmE* profoundly increased after germination induction in WT and *sigA-R365A* mutant spores (Fig. 5B and *SI Appendix*, Fig. S12B). Nevertheless, no significant changes in mRNA levels were observed in germinating *sigA-R365D* mutant spores (Fig. 5B and *SI Appendix*, Fig. S12B). As a control, we measured the mRNA levels of *pupG* (a SigA controlled gene) and *yaaH* (a SigB controlled gene), which their corresponding proteins are excluded from the germination proteome (23). No difference in the mRNA levels was detected in any of the tested strains (Fig. 5B and *SI Appendix*, Fig. S12B). Furthermore, the transcript levels of *tig*, *rpmE*, or *pupG* were similar among the examined strains during




Fig. 3. Tig-ribosome association is dependent on Arg dephosphorylation. (A) BZ118 (*tig-gfp, rpsB-mCherry*), BZ119 (*tig-R45D-gfp, rpsB-mCherry*), and BZ120 (*tig-R45A-gfp, rpsB-mCherry*) strains were grown to a mid logarithmic phase in LB at 37 °C and visualized by fluorescence microscopy. Shown are images of phase contrast (Phase), fluorescence from mCherry (red), and GFP (green), and an overlay of red and green fluorescence from a representative experiment out of three independent biological repeats. (Scale bar: 1 μ m.) (B) Intact ribosomes were purified from CEs of the vegetative cells, dormant and germinating spores of BZ118 (*tig-gfp, rpsB-mCherry*) and BZ119 (*tig-R45D-gfp, rpsB-mCherry*) strains. Germination was induced by suspending the spores in 10 mM L-Ala for 10 min. Western blot analysis was carried out using an antibody against GFP, detecting the levels of Tig and TigR45D in CEs and in PRs (Left). The signal from GFP fusion proteins was quantified by MetaMorph software (version 7.7, Molecular Devices) (Right). The signal from Tig-GFP was used to normalize Tig expression levels in CEs and PRs, separately. Shown are average values and SDs obtained from two independent biological repeats.

vegetative growth (SI Appendix, Fig. S12C). In line with these findings, Northern blot analysis designated the *sigA-R365D* mutation to influence specifically the transcription of *rpmE* during germination, but to have no impact on the transcription of *pupG* (Fig. 5C). These results indicate that a subset of genes is being transcribed during germination and that spores harboring the phosphorylated-like *sigA-R365D* allele are impaired in executing this mRNA synthesis.

Discussion

Dormant bacterial spores undergo a fascinating awakening complex process to restore active metabolism and reinitiate growth (19). However, the molecular events underlining this dramatic cellular transition are still elusive. Here, we describe that the recently identified bacterial protein modification, Arg dephosphorylation (37, 38, 40), plays a key role in facilitating the

earliest molecular events, occurring during germination. We show that the Arg-phosphatase YwIE is required to prompt germination whereas the absence of the Arg kinase McsB results in accelerated germination. Based on our results, we propose the following model for spore germination (Fig. 6). The Arg-phosphatase YwIE interacts with germination receptors and is activated following the binding of a germinant to its cognate receptor, dephosphorylating its downstream substrate proteins. YwIE is a redox sensor, and thus, its activation could be achieved by switching from the disulfide-linked inactive form to the reduced active mode of the protein (49, 50). As a consequence of YwIE-phosphatase activity, transcription and translation are restored, at least in part, via Arg dephosphorylation of SigA and Tig, respectively.

Interestingly, additional factors were identified to be Arg phosphorylated in dormant spores, among them crucial enzymes




Fig. 4. The dephosphorylation of SigA facilitates spore germination. (A) Spores of PY79 (WT), BZ90 (*sigA-R365A*), and BZ91 (*sigA-R365D*) strains were incubated on agarose supplemented with L-Ala (10 mM) and monitored by time lapse microscopy. Shown are phase contrast images from a representative experiment out of three independent biological repeats. (Scale bar: 1 μ m.) (B) Spores of PY79 (WT), BZ90 (*sigA-R365A*), and BZ91 (*sigA-R365D*) strains were incubated on agarose supplemented with L-Ala (10 mM) and monitored by time lapse microscopy. Data are presented as percentages of the initial number of the phase-bright spores. Shown are average values and SDs obtained from three independent biological repeats ($n \geq 300$ for each strain). (C) Spores of PY79 (WT), BZ90 (*sigA-R365A*), and BZ91 (*sigA-R365D*) strains were incubated with L-Ala to trigger germination. DPA release to the medium was determined by Tb-DPA assay. Presented are RFUs measured at 545 nm with excitation at 270 nm. Shown is a representative experiment out of three independent biological repeats. (D) Spores of PY79 (WT), BZ90 (*sigA-R365A*), and BZ91 (*sigA-R365D*) strains were supplemented with L-Ala for 10 min and then incubated at 80 °C for 30 min. The percentage of nongerminating heat resistant spores was determined by the number of colonies after heat treatment/number of colonies before heat treatment. (E) Spores of PY79 (WT), BZ16 ($\Delta ywIE$), BZ91 (*sigA-R365D*), BZ97 (*tig-R45D*), and BZ104 (*tig-R45D*, *sigA-R365D*) strains were incubated with L-Ala (10 mM), and OD₆₀₀ was measured at the indicated time points. Data are presented as percentages of the initial OD₆₀₀ of spore suspension. Germination assays were carried out in triplicate, and representative data are presented. Decreasing OD₆₀₀ signifies spore germination.

involved in carbon metabolism, such as GapA (Glyceraldehyde-3-phosphate dehydrogenase) and Pgi (Glucose-6-phosphate isomerase), controlling central metabolic pathways (51, 52). The detection of Arg phosphorylation within spore specific proteins, including members of the Ssp (small, acid-soluble spore protein) family and spore coat proteins, suggests that Arg phosphomodification could mediate the degradation of spore proteins that need to be removed to enable revival (17, 53).

Arg phosphorylation was initially monitored in *B. subtilis* and was shown to function as part of the bacterial stress response and to mark proteins for degradation by the Clp protease complex (32, 37, 38, 40, 53). Phosphoproteomic analyses suggested that

Arg phosphorylation has a regulatory effect on many cellular pathways (32, 37, 40), although the *in vivo* impact of the Arg-phosphorylation pathway remained limited. Here, we show that Arg dephosphorylation elicits the most critical stage of spore awakening, placing it as a major regulator of a developmental process in bacteria.

Surprisingly, our analysis revealed that not only translation, but also transcription occurs during germination and are required for its progression. The transcription machinery restoration is likely to be mediated by the switching between the phosphorylation and the dephosphorylation state of the housekeeping SigA. The Arg-365 site seems to be located in the DNA




Fig. 5. Evidence that transcription occurs during germination. (A) An in vitro transcription reaction was carried out in whole extracts from vegetative cells of PY79 (WT) as well as from dormant and germinating spores of PY79 (WT), BZ90 (*sigA-R365A*), and BZ91 (*sigA-R365D*) strains and in a transcription buffer supplemented with NTPs (ATP, CTP, GTP, UTP, [α - 32 P]-UTP). After 40 min of incubation at 37 °C, the reaction was stopped, RNA was purified, radioactively labeled RNAs were analyzed in 8% polyacrylamide gels, and bands were visualized by autoradiography. The analysis was carried out in two biological repeats, and a representative experiment is presented. (B) Spores of PY79 (WT), BZ90 (*sigA-R365A*), and BZ91 (*sigA-R365D*) strains were incubated with 10 mM L-Ala to trigger germination, collected by centrifugation at the indicated time, and their RNAs were extracted. The mRNA levels of selective genes *tig*, *rpmE*, and *pupG* during germination were determined by quantitative RT-PCR. The result is presented as the fold change of target gene expression after germination relative to before germination. The assays were carried out in triplicate, and representative data are presented. (C) Total RNA was extracted from vegetative cells, dormant and germinating spores of PY79 (WT), BZ90 (*sigA-R365A*), and BZ91 (*sigA-R365D*) strains and subjected to Northern blot analysis using *rpmE* (201 nt) and *pupG* (2001 nt) specific biotinylated probes. knockout (KO), indicates control extracts derived from vegetative cells of the corresponding LS26 (Δ *rpmE*) and LS76 (Δ *pupG*) KO strains. The analysis was carried out in three independent biological repeats, and a representative experiment is presented. Numbers on the right correspond to RNA size marker.

binding domain of SigA (54), suggesting that its phosphorylation might interfere with the binding of the protein to its template. The transcription machinery of dormant spores appears to be silenced regardless of the Arg-phosphorylation state, indicating

that additional factors impose this arrest. It will be intriguing to elucidate at which stage of entering dormancy this halt is achieved as, according to our previous findings, this could take several days (14, 18). The activation of transcription precedes




Fig. 6. A model for propelling spore germination via Arg dephosphorylation. A model depicting the events occurring during germination based on the data obtained in our analysis. Upon nutrient sensing by germination receptors, the Arg phosphatase YwE is activated (A). Consequently, SigA is dephosphorylated by active YwE to reactivate the transcription machinery (B). Next, translation is reestablished by the YwE-mediated dephosphorylation of Tig (C). The source of the transcripts for protein synthesis during germination could be derived from the newly transcribed RNA (D).

that of translation as SigA-R365D mutant spores were stalled before DPA release, whereas Tig-R45D spores could liberate DPA normally. This observation raises the possibility that the source of the transcripts required for translation during germination could be derived from newly synthesized mRNA. The RNA profile of aging spores was found to be dynamic, mainly involving RNA degradation into nucleotides, which can serve as a source for de novo germination-induced RNA synthesis (14, 15, 18).

Dephosphorylation of Tig was found to mediate its association with ribosomes to reactivate the translational machinery, reinforcing the view that translation is required to execute germination. During sporulation, Tig may be phosphorylated by McsB thereby limiting its association with the ribosomes. However, when germination is induced, Tig activity could be restored by the action of YwE, propelling its rapid association with ribosomes and, consequently, allowing protein folding to resume. Interestingly, the Thr-phosphorylated form of the elongation factor Tu was shown to bind ribosomes during sporulation and inhibit translation elongation, hence, acting in an opposed manner to Tig (55). It is plausible that networks of co-occurring phosphorylation

events could serve as biological switches functioning to control translation when entering and exiting dormancy.

The observation that not all of the SigA-R365D and Tig-R45D mutant spores are stalled in germination could reflect the differences in the levels of RNA and proteins within spores shown to affect germination capacity (14, 15, 18, 56). This bet-hedging strategy could increase population fitness to maximize survival under changeable conditions. Yet, the connection between the spore molecular reservoir and its germination propensity remains puzzling, highlighting the gaps in our understanding of spore dormancy and awakening. This suggests that the spore molecular pool plays a significant role in germination capacity of spores residing in different natural niches.

Materials and Methods

Strains and General Methods. *B. subtilis* strains are derivatives of the WT PY79 and are listed in *SI Appendix, Table S2*. Plasmid construction is described in *SI Appendix, Experimental Procedures*, and primers are described in *SI Appendix, Table S3*. All general methods for *B. subtilis* were carried out as described previously (57). *B. subtilis* cultures were inoculated at OD₆₀₀ 0.05 from an overnight culture, and growth was carried out at 37 °C in LB medium (Difco). Antibiotics were used at the following concentrations: kanamycin 5 µg/mL, chloramphenicol 6 µg/mL, lincomycin 25 µg/mL, erythromycin 1 µg/mL, tetracycline 10 µg/mL, and spectinomycin 100 µg/mL. Sporulation was induced at 37 °C by suspending cells in Schaeffer's liquid medium (Difco Sporulation Medium) (58). Sporulation efficiency was evaluated by comparing the number of colony forming units before and after heat treatment. Spores were harvested, purified, and stored as described below according to (23). Purified spores were heat activated at 80 °C for 30 min before germination experiments. Spore germination is induced by L-Ala (10 mM), AGFK (2.5 mM L-Asp, 5 mg/mL D-glucose, 5 mg/mL D-fructose, and 50 mM KCl), or LB at 37 °C.

Light Microscopy. Light microscopy was carried out as described previously (23). Briefly, bacterial cells (0.5 mL) were collected by centrifugation and resuspended in 10 µL of PBS × 1. For time lapse experiments, spores were placed on 1% agarose pads supplemented with L-Ala (10 mM) or LB and incubated in a chamber where temperature was maintained at 37 °C with a temperature controller (Pecon-Zeiss). For fluorescence measurements, the intensity of a WT (PY79) strain, lacking *gfp*, *mCherry*, or *dronpa* genes, was subtracted from the net average fluorescence intensity. Samples were photographed using Axio Observer Z1 (Zeiss), equipped with a CoolSnap HQII camera (Photometrics, Roper Scientific). System control, image analysis, and processing were performed using MetaMorph software (version 7.7; Molecular Devices).

DPA Measurements. DPA release was measured as described previously (59). Spore germination was performed at 37 °C in 10 mM L-Ala suspended in 25 mM K-Hepes buffer (pH 7.4). At the indicated time, 200 µL aliquots were centrifuged, and TbCl₃ was added to the supernatant fluid in a final concentration of 50 µM. Tb-DPA fluorescence was measured at a fluorescence emission of 545 nm and with excitation of 270 nm.

Quantification and Statistical Analysis. Unless stated otherwise, bar charts display a mean ± SD from at least three independent biological experiments. MS Excel was used for all statistical analysis, data processing, and presentation.

ACKNOWLEDGMENTS. We thank Silke Wahl for preparing samples for MS measurements. We are grateful to the members of the S.B.-Y. laboratory for valuable discussions and comments. We thank P. Setlow (University of Connecticut) for generously providing antibodies against germination receptors. This work was supported by the European Research Council Advanced Grant (339984) and the Israel Science Foundation (774/16) awarded to S.B.-Y., and by The German-Israeli Foundation (GIF) (Grant I-1464-416.13/2018) awarded to S.B.-Y. and B.M. B.Z. was funded by the Key Project of National Natural Science Foundation of China (NSFC) (31530058) and China Scholarship Council (CSC) scholarship (201606350068).

1. D. Higgins, J. Dworkin, Recent progress in *Bacillus subtilis* sporulation. *FEMS Microbiol. Rev.* **36**, 131–148 (2012).
2. P. J. Piggot, Spore development in *Bacillus subtilis*. *Curr. Opin. Genet. Dev.* **6**, 531–537 (1996).
3. P. J. Piggot, D. W. Hilbert, Sporulation of *Bacillus subtilis*. *Curr. Opin. Microbiol.* **7**, 579–586 (2004).

4. P. Stragier, R. Losick, Molecular genetics of sporulation in *Bacillus subtilis*. *Annu. Rev. Genet.* **30**, 297–241 (1996).
5. P. Stragier, R. Losick, Cascades of sigma factors revisited. *Mol. Microbiol.* **4**, 1801–1806 (1990).
6. J. Dworkin, Protein targeting during *Bacillus subtilis* sporulation. *Microbiol. Spectr.* **2**, TBS-0006-2012 (2014).

7. A. Driks, *Bacillus subtilis* spore coat. *Microbiol. Mol. Biol. Rev.* **63**, 1–20 (1999).
8. D. L. Popham, Specialized peptidoglycan of the bacterial endospore: The inner wall of the lockbox. *Cell. Mol. Life Sci.* **59**, 426–433 (2002).
9. A. O. Henriques, C. P. Moran, Jr, Structure, assembly, and function of the spore surface layers. *Annu. Rev. Microbiol.* **61**, 555–588 (2007).
10. P. T. McKenney, A. Driks, P. Eichenberger, The *Bacillus subtilis* endospore: Assembly and functions of the multilayered coat. *Nat. Rev. Microbiol.* **11**, 33–44 (2013).
11. D. L. Popham, C. B. Bernhards, Spore peptidoglycan. *Microbiol. Spectr.* **3**, TBS-0005-2012 (2015).
12. A. Driks, P. Eichenberger, The spore coat. *Microbiol. Spectr.* **4**, TBS-0023-2016 (2016).
13. P. Setlow, Spores of *Bacillus subtilis*: Their resistance to and killing by radiation, heat and chemicals. *J. Appl. Microbiol.* **101**, 514–525 (2006).
14. E. Segev, Y. Smith, S. Ben-Yehuda, RNA dynamics in aging bacterial spores. *Cell* **148**, 139–149 (2012).
15. S. Ghosh, G. Korza, M. Maciejewski, P. Setlow, Analysis of metabolism in dormant spores of *Bacillus* species by ³¹P nuclear magnetic resonance analysis of low-molecular-weight compounds. *J. Bacteriol.* **197**, 992–1001 (2015).
16. G. Korza, B. Setlow, L. Rao, Q. Li, P. Setlow, Changes in *Bacillus* spore small molecules, rRNA, germination, and outgrowth after extended sublethal exposure to various temperatures: Evidence that protein synthesis is not essential for spore germination. *J. Bacteriol.* **198**, 3254–3264 (2016).
17. P. Setlow, Spore germination. *Curr. Opin. Microbiol.* **6**, 550–556 (2003).
18. E. Segev, A. Rosenberg, G. Mamou, L. Sinai, S. Ben-Yehuda, Molecular kinetics of reviving bacterial spores. *J. Bacteriol.* **195**, 1875–1882 (2013).
19. P. Setlow, Germination of spores of *Bacillus* species: What we know and do not know. *J. Bacteriol.* **196**, 1297–1305 (2014).
20. A. Moir, G. Cooper, Spore germination. *Microbiol. Spectr.* **3**, TBS-0014-2012 (2015).
21. A. Moir, How do spores germinate? *J. Appl. Microbiol.* **101**, 526–530 (2006).
22. P. Setlow, S. Wang, Y. Q. Li, Germination of spores of the orders *Bacillales* and *Clostridiales*. *Annu. Rev. Microbiol.* **71**, 459–477 (2017).
23. L. Sinai, A. Rosenberg, Y. Smith, E. Segev, S. Ben-Yehuda, The molecular timeline of a reviving bacterial spore. *Mol. Cell* **57**, 695–707 (2015).
24. L. Sinai, S. Ben-Yehuda, Commentary: Changes in *Bacillus* spore small molecules, rRNA, germination, and outgrowth after extended sublethal exposure to various temperatures: Evidence that protein synthesis is not essential for spore germination. *Front. Microbiol.* **7**, 2043 (2016).
25. W. Steinberg, H. O. Halvorson, A. Keynan, E. Weinberg, Timing of protein synthesis during germination and outgrowth of spores of *Bacillus cereus* strain T. *Nature* **208**, 710 (1965).
26. V. Vinter, Symposium on bacterial spores: V. Germination and outgrowth: Effect of inhibitors. *J. Appl. Bacteriol.* **33**, 50–59 (1970).
27. G. Kramer et al., L23 protein functions as a chaperone docking site on the ribosome. *Nature* **419**, 171–174 (2002).
28. M. A. Lauber, W. E. Running, J. P. Reilly, *B. subtilis* ribosomal proteins: Structural homology and post-translational modifications. *J. Proteome Res.* **8**, 4193–4206 (2009).
29. S. F. Göthel, C. Scholz, F. X. Schmid, M. A. Marahiel, Cyclophilin and trigger factor from *Bacillus subtilis* catalyze in vitro protein folding and are necessary for viability under starvation conditions. *Biochemistry* **37**, 13392–13399 (1998).
30. D. Paredes-Sabja, P. Setlow, M. R. Sarker, Germination of spores of *Bacillales* and *Clostridiales* species: Mechanisms and proteins involved. *Trends Microbiol.* **19**, 85–94 (2011).
31. I. M. Shah, M. H. Laaberki, D. L. Popham, J. Dworkin, A eukaryotic-like Ser/Thr kinase signals bacteria to exit dormancy in response to peptidoglycan fragments. *Cell* **135**, 486–496 (2008).
32. I. Mijakovic, C. Grangeasse, K. Turgay, Exploring the diversity of protein modifications: Special bacterial phosphorylation systems. *FEMS Microbiol. Rev.* **40**, 398–417 (2016).
33. J. Dworkin, Ser/Thr phosphorylation as a regulatory mechanism in bacteria. *Curr. Opin. Microbiol.* **24**, 47–52 (2015).
34. J. Dworkin, I. M. Shah, Exit from dormancy in microbial organisms. *Nat. Rev. Microbiol.* **8**, 890–896 (2010).
35. A. Rosenberg et al., Phosphoproteome dynamics mediate revival of bacterial spores. *BMC Biol.* **13**, 76 (2015).
36. K. Basell et al., The phosphoproteome and its physiological dynamics in *Staphylococcus aureus*. *Int. J. Med. Microbiol.* **304**, 121–132 (2014).
37. A. K. Elsholz et al., Global impact of protein arginine phosphorylation on the physiology of *Bacillus subtilis*. *Proc. Natl. Acad. Sci. U.S.A.* **109**, 7451–7456 (2012).
38. J. Fuhrmann et al., McsB is a protein arginine kinase that phosphorylates and inhibits the heat-shock regulator CtsR. *Science* **324**, 1323–1327 (2009).
39. J. Fuhrmann et al., Structural basis for recognizing phosphoarginine and evolving residue-specific protein phosphatases in gram-positive bacteria. *Cell Rep.* **3**, 1832–1839 (2013).
40. A. Schmidt et al., Quantitative phosphoproteomics reveals the role of protein arginine phosphorylation in the bacterial stress response. *Mol. Cell. Proteomics* **13**, 537–550 (2014).
41. S. Junker et al., Spectral library based analysis of arginine phosphorylations in *Staphylococcus aureus*. *Mol. Cell. Proteomics* **17**, 335–348 (2018).
42. N. Dephoure, K. L. Gould, S. P. Gygi, D. R. Kellogg, Mapping and analysis of phosphorylation sites: A quick guide for cell biologists. *Mol. Biol. Cell* **24**, 535–542 (2013).
43. D. C. Dieterich et al., Labeling, detection and identification of newly synthesized proteomes with bioorthogonal non-canonical amino-acid tagging. *Nat. Protoc.* **2**, 532–540 (2007).
44. R. Schuch, A. Garibian, H. H. Saxild, P. J. Piggot, P. Nygaard, Nucleosides as a carbon source in *Bacillus subtilis*: Characterization of the *drm-pupG* operon. *Microbiology* **145**, 2957–2966 (1999).
45. G. Akanuma et al., Inactivation of ribosomal protein genes in *Bacillus subtilis* reveals importance of each ribosomal protein for cell proliferation and cell differentiation. *J. Bacteriol.* **194**, 6282–6291 (2012).
46. P. J. Lewis, S. D. Thaker, J. Errington, Compartmentalization of transcription and translation in *Bacillus subtilis*. *EMBO J.* **19**, 710–718 (2000).
47. A. Rosenberg, L. Sinai, Y. Smith, S. Ben-Yehuda, Dynamic expression of the translational machinery during *Bacillus subtilis* life cycle at a single cell level. *PLoS One* **7**, e41921 (2012).
48. W. G. Haldenwang, The sigma factors of *Bacillus subtilis*. *Microbiol. Rev.* **59**, 1–30 (1995).
49. J. Fuhrmann, V. Subramanian, D. J. Kojetin, P. R. Thompson, Activity-based profiling reveals a regulatory link between oxidative stress and protein arginine phosphorylation. *Cell Chem. Biol.* **23**, 967–977 (2016).
50. J. Fuhrmann, V. Subramanian, P. R. Thompson, Targeting the arginine phosphatase YwIE with a catalytic redox-based inhibitor. *ACS Chem. Biol.* **8**, 2024–2032 (2013).
51. C. Prasad, E. Freese, Cell lysis of *Bacillus subtilis* caused by intracellular accumulation of glucose-1-phosphate. *J. Bacteriol.* **118**, 1111–1122 (1974).
52. A. DeLoughery, J. B. Lalanne, R. Losick, G. W. Li, Maturation of polycistronic mRNAs by the endoribonuclease RNase Y and its associated Y-complex in *Bacillus subtilis*. *Proc. Natl. Acad. Sci. U.S.A.* **115**, E5585–E5594 (2018).
53. D. B. Trentini et al., Arginine phosphorylation marks proteins for degradation by a Clp protease. *Nature* **539**, 48–53 (2016).
54. S. P. Haugen, W. Ross, R. L. Gourse, Advances in bacterial promoter recognition and its control by factors that do not bind DNA. *Nat. Rev. Microbiol.* **6**, 507–519 (2008).
55. S. F. Pereira, R. L. Gonzalez, Jr, J. Dworkin, Protein synthesis during cellular quiescence is inhibited by phosphorylation of a translational elongation factor. *Proc. Natl. Acad. Sci. U.S.A.* **112**, E3274–E3281 (2015).
56. A. Mutlu et al., Phenotypic memory in *Bacillus subtilis* links dormancy entry and exit by a spore quantity-quality tradeoff. *Nat. Commun.* **9**, 69 (2018).
57. C. R. Harwood, S. M. Cutting, *Molecular Biological Methods for Bacillus* (Wiley, Chichester, New York, 1990).
58. P. Schaeffer, [Problems of spore formation in bacteria]. *Zentralblatt für Bakteriologie, Parasitenkunde, Infektionskrankheiten und Hygiene. 1. Abt. Medizinisch-hygienische Bakteriologie, Virusforschung und Parasitologie. Originale* **198**, 72–75 (1965).
59. A. Perez-Valdespino et al., Isolation and characterization of *Bacillus subtilis* spores that are superdormant for germination with dodecylamine or Ca²⁺-dipicolinic acid. *J. Appl. Microbiol.* **114**, 1109–1119 (2013).
60. B. Zhu, J. Stülke, SubtiWiki in 2018: from genes and proteins to functional network annotation of the model organism *Bacillus subtilis*. *Nucleic Acids Res.* **46**, D743–D748 (2018).