

Integral and sustainable community self-management of the native fruit trees of Munhiba, Mozambique

Autogestión comunitaria integral y sostenible de los frutales nativos de Munhiba, Mozambique

doi: 10.15446/rfnam.v72n2.78980

Barbarita Mitjans Moreno^{1*}, Joel Pacheco Escobar¹ and Gervásia Musaico Lopes²

ABSTRACT

Keywords:

Community self-management
Economic importance
Forest deterioration
Fruit tree species
Mozambique
Working lines

With the aim of propose lines of work for the community's appropriate self-management of the native fruit trees in Munhiba, province of Zambézia (Mozambique), it was carried out a study of the perception of the local actors about the causes of deterioration of the fruit trees, as well as their economic, social, and environmental importance. It was used intentional sampling by selecting 118 local actors. In order to develop a Participatory Rapid Diagnosis, exchange workshops and field visits were also performed to identify the fruit species and their use. For the information gathering, semi-structured interviews were applied together with the use of simple and participant observation. Frequency and correlation of variables (Rho of Spearman) were developed for data processing. Among the most important results, it was found the botanical classification of the fruit species, the perception of the local actors about the economic, social, and environmental importance as well as causes of deterioration, identification of strengths and weaknesses for the management of the native fruit trees and a proposal of work lines for their self-management. Despite the potential of native fruit trees in the territory of study, it was not appreciated yet proper management due to the lack of knowledge of the production process that could lead to the progressive deterioration of the species and the ecosystem where they inhabit.

RESUMEN

Palabras clave:

Autogestión comunitaria
Importancia económica
Deterioro del bosque
Especies de frutales
Mozambique
Líneas de trabajo

Con el objetivo de proponer líneas de trabajo para la autogestión comunitaria y manejo adecuado de los frutales nativos en la comunidad de Munhiba, provincia de Zambézia (Mozambique), se realizó un estudio sobre la percepción de los actores locales sobre las causas de deterioro de los frutales, así como la importancia económica social y ambiental. Se realizó un muestreo intencional seleccionando 118 actores locales. En aras de desarrollar un diagnóstico rápido participativo, se realizaron talleres de intercambio y visitas al campo para identificar las especies frutales y usos que se le atribuyen. Para la recolección de información, se aplicó entrevista semi-estructurada junto con el uso de observación simple y participante. Para el tratamiento de datos se usó el análisis de frecuencia y de correlación de variables (Rho de Spearman). Entre los resultados más importantes destacan la clasificación botánica de las especies frutales, la percepción de los actores locales sobre la importancia económica, social, ambiental y las causas de deterioro, se identificaron las fortalezas y debilidades para el manejo de los frutales y se definieron líneas de trabajo para su autogestión. A pesar del potencial de los frutales en el territorio de estudio no se aprecia un manejo adecuado, principalmente por falta de conocimiento en el proceso de producción y recolección de las comunidades locales, lo que puede conducir al deterioro progresivo de las especies y el ecosistema donde habitan.

¹ Universidad Pinar del Río-Hermanos Saíz de Oca. CP 2700. Candelaria, Pinar del Río, Cuba.

² Direcção Provincial da Agricultura e Segurança Alimentar. Avenida Samora Machel Quelimane, Província de Zambézia, Mozambique.

* Corresponding author: <barbaritamozambique@gmail.com>

Native fruits (NF) in Mozambique are very diverse. Although many of them show broad perspectives of economic use, only a few of them have been studied at both the local and national level. At the same time, the current transfer of scientific and technical knowledge and business opportunities have an incipient value in new productive chains. Besides, a significant part of the indigenous knowledge for the protection of the forest ecosystem has not been considering (Simone, 2001).

The use and logging of the native fruit trees have become into an important practice for the subsistence of rural communities, not only during the seasonal periods of hunger but also in their day-to-day for the nutritious consumption. It must be kept in mind their good adaptability to arid areas without requiring the use of fertilizers, making them more accessible to the rural community, due to their low cost of production and acquisition (Goulão and Santo-Antonio, 2015).

Native fruit trees also play an important role in the traditional medicine and represent a high potential of economic exploitation, in the promotion of employment opportunities and the improvement of the family profit within the rural populations (Chiau, 2003).

Several native fruits, which can contribute to the solution of the alimentary crisis, have been identified within the forest ecosystems in Mozambique. However, the inadequate use and constantly logging are a concern due to their contribution to the decrease in the population of these fruit species, regardless of their importance for the nutritious diet in the Mozambican rural population.

Munhiba is a municipality of the district of Mocuba in which the nutrition security depends on a large part of the production and availability of food coming from a heterogeneous agricultural sector of smallholders, as well as a small incomplete and not well-distributed commercialization network. Agricultural production is characterized by low productivity due to the weak use of technology and incomes. Consequently, the production percentage that reaches the market is low (MAE, 2005).

The low levels of rainfall that have been registered in Munhiba during the last period have placed its population

under challenging situations to access to food supplies (nutrition insecurity). Under those conditions, the native fruits have been the fundamental mean of sustenance of local population.

Castro Paulino Comrade, the representative of Food and Agriculture Organization of the United Nations in Mozambique, underlined in the First National Workshop of Native Fruits, that the research in the native area of the fruits in Mozambique is still in its initial phase. He also asserted: "The research, domestication, and development processes of the valuable native fruits chains should deserve a special attention because 70% of the population live in rural areas and almost 80% depend on the natural resources, like fruits and medicinal plants to satisfy the nutritional and health necessities"(Goulão and Santo-Antonio, 2015).

These arguments evidence the necessity to find alternatives to commit the community in the self-management of their natural resources, a subject that has been debated in different trends and socio-political situations along the history of the practical social thinking (Reyes, 2013).

According to Brivio Borja (2001), the community self-management is an administration of the community that occurs due to the transformation of the descending spiral of poverty in an upward spiral of development. The source of the community self-management is the change from a fatalistic vision of the poverty, only as a summary of lacks, to a hopeful vision as a generator of the necessary impulse for development. The community self-management is the channel through which the inherent infinite potential of the human being is canalized toward the achievement of a worthy life, through improving the life quality of each one of the inhabitants. It also allows them to achieve their objectives and goals with solidary support of their fellow men; taking advantage of all the resources of the community, that in this case it is adjusted to the natural resources (native fruit trees).

A worldwide emphasis has been developed toward the managing processes at the community level. Several researchers converge in the topic based on the creation of participative processes that lead to the emergence of a critical popular conscience to understand the social practice as an inseparable unit of reflection and action.

Referring to the case some authors such as Valdés (2010), Mitjans (2012), Sabogal *et al.* (2012), Molero *et al.* (2012), Reyes (2013), and Marzin *et al.* (2014); which offer an updated and in-depth focus on the community actions from the participation and self-management with emphasis in an economic, social, and the environmental scope.

In this research, the self-management is represented as a suitable alternative to the local communities that inhabit the forest ecosystems of Mozambique. They need to be articulated among them to established functions in pro of the forest and their well-being. Therefore, the following hypothesis is presented: if it is considered the economic and environmental perception, as well as the knowledge, that the local actors of the community of Munhiba have about the native fruits of the region; it will be possible to elaborate working lines to contribute with an appropriate management of these native fruits. Regarding the former idea, this research centers its objective in proposing some lines of work for the community of Munhiba that contribute to an appropriate self-management of the native fruits.

MATERIALS AND METHODS

Localization of the study area

Munhiba municipality is located in the central part of Mozambique, in the county of Zambézia, district of Mocuba. It limits to the North with the Headquarters of the District of Mocuba, to the South with the district of Namacurra, to the East with the district of Maganja da Costa through the Licungo River, and to the West with the town of Namanjavira (MA, 2011).

Methodology used

To assess the economic, environmental, and social perception of the local actors about the native fruits, identify the causes of their deterioration and recognize the strengths and weaknesses for the community development; both quantitative and qualitative research methods were used. Considering the Ethnographic prevalence, the criteria of Rodríguez *et al.* (2008) were used; given that they declared that the study of the Ethnographic is a research method from which it is possible to know the way of life of a specific social unit. Besides, the Participatory Rapid Diagnosis (PRD) was used, following the criteria of Gomes *et al.* (2001) who identify the PRD as the approach most used to facilitate

the direct participation of the residents in the generation and the analysis of the obtained information.

For gathering the information, a semi-structured interview along with a simple and participant observation was used to increase the validity of the study and also to obtain a better understanding of the studied context, keeping in mind the approaches proposed by Dewalt and Dewalt (2002).

It was performed an intentional sampling to apply the interview. 118 residents were selected out of a universe of 326 to be interviewed. Three exchange workshops and six field visits were carried out to identify the native fruit species that were in the forest that limits the area where the community inhabits.

For the selection of interviewees, it was taking into consideration those who live up to 2 km of distance away from the forest. The sample size for the research was to determine following the procedures for social studies in finite populations during the estimation of a proportion, presented by Calero Vinelo (1978). A confidence level ($1-\alpha$) of 95% with a significance level (α) of 5% and a critical value (Z) of 1.96, as well as a positive variability (p) of 0.95 and a negative variability (q) of 0.05 were considered, assuming a maximum permissible error (E) of 0.05.

$$n = \frac{Z^2 pqN}{NE^2 + Z^2 pq}$$

Where:

n : sample size.

Z : level of confidence.

p : positive variability.

q : negative variability.

N : size of the population.

For the interpretation of the interview results, a variable codifier was used (Table 1). The information processing was carried out with the statistical package IBM-SPSS, version 20.0.

For the statistical analysis, non-parametric tests were carried out. It was used Bi-varied correlation using the coefficient of correlation, Rho of Spearman, to determine

Table 1. Variable codifier for the statistical analysis.

Dimensions	Group/Vision		
	1 (Low)	2 (Intermediate)	3 (Advanced)
Knowledge about the identified NF	They know less than 25% of the identified NF	They know between the 25 and 50% of the NF identified	They know 100 % of the identified NF
Perceptions of the NF socio-economic function	They do not consider economic and/or social NF contributions	They consider the NF contributions only to feed	They identify potentialities of the NF as economic and/or social benefits (edible, medicinal, wood, ornamental and other uses like sorcery and fibers)
Perceptions of the NF environmental function	They do not recognize any NF environmental function	They recognize the anti-erosive function of these species	They identify the NF function as a contribution to the soil erosion control and their utility for feeding the fauna
Perceptions about the NF degradation causation	They do not define the causes that have led to the NF degradation	They clearly recognize the responsibility of the NF degradation to the necessity of using these species in wood production	They consider multiple the NF degradation causes, among them: lack of knowledge about the management of these species, wooden use for coal and firewood and the annual fires

(NF=Native fruits)

the strength of the linear relation among the categorical variables. The evaluation of the perception of local actors was also made through a descriptive analysis, based on frequency distributions with the graphics generator.

In order to identify the fruit species taxonomically, there were used not only the criteria of Specialists in Botany and Dendrology from the Agronomy and Forestry Engineering Faculty of Zambeze University, but also the traditional knowledge of the more experienced local actors and the technical record on fruit species proposed by the FAO (1982). Furthermore, an ethnobotanical study was also carried out to assess the strength of the area for the use of the located species. They were classified in: groceries, medicinal, wood, ornamental, sorcery, water store, as well as soil and water protectors.

RESULTS AND DISCUSSION

Perception of local actors on the NF and their relation with the proximity to the forest

During the study of the perception of local actors on the environmental function, it is appreciated that 68 of the

interviewees (58%) possess a low vision, because they did not recognize any environmental function of the fruits; out of them, 49 individuals live at a distance between one and two kilometers from the forest. For an advanced vision, 38 individuals were identified, from which 34 live in the forest proximity. In general, it is observed that from the 61 interviewees that live inside the forest area and in its periphery, 40 somehow considered environmental importance of the native fruits, and 19 discarded all environmental value of these species (Figure 1).

Figure 2 shows the relationship between the causes of deterioration and the economic importance that is attributed to the fruit species. It is also appreciated that the largest number of individuals is in the low vision (1) who do not define the causes of deterioration (84/118) (71%), 12 of them do not attribute any benefit, while 72 of them consider its use only as food. So, it can be inferred that if these local actors were qualified about the economic importance of the fruits, they would have been motivated to diminish the causes of deterioration and to self-negotiate the management of the fruits.




Figure 1. Distribution of frequencies according to the perception of the local actors on the environmental function related to the distance to the forest.

About this concern, AIDER and FAO (2016) declared that “the training becomes the first step toward the self-help to take responsibilities on their destination, or what is known as self-management.”

It is necessary to train them, for them to be able to make use of the forest resources, including the native fruits.

These strategies of multiple uses would allow them to maintain a dual economy. On the one hand, they have production areas from which they sell some goods and services, and on the other hand, they satisfy their necessities of local consumption (García-Frapolli *et al.*, 2008; Infante-Ramírez, 2011; Infante-Ramírez and Arce Ibarra, 2013).




Figure 2. Distribution of frequencies according to the perception of the local actors about the causes of deterioration and economic importance.

Figure 3 shows the relationship between the proximity to the forest and the knowledge of the fruit species. In this case, the positions were favorable because all the local actors were in the advanced and intermediate positions, so they indeed know the native fruit species. It is observed that the 87% identify the 100% of the fruits (103/118) and that the largest representativeness of them (58) comes from those who live nearer the forest. The latter is an important question to be kept in mind according to the criteria of Mitjans (2012), due to the people know their region, the traditional use of the natural resources, the

location of the species and in some cases the way of plants propagation. These factors are important inquiries to be considered in the management plan of the different forest ecosystems.

According to Moreno (2013), during the last decades some geographers, anthropologists, sociologists, historians, naturalistic, jurists, among other specialists have continued the task of describing and building theories about the collective rural institutions and their knowledge and perceptions about the natural resources.




Figure 3. Distribution of frequencies according to the perception of the local actors on knowledge of the species and proximity to the forest.

Concerning the knowledge of the species in function of the school level of the local actors, it was verified that all of them know the native fruits since they were in the visions 2 and 3. 85% (100/118) of the interviewees know the 100% of native fruit species. Individuals in low positions were not found (Figure 4).

Traditional knowledge and the local actors participated in the administration of the natural resources should be considered during their management. Molina-Pelegrín *et al.* (2011) obtained important results concerning protected and conserved areas, with the intervention of the adjacent communities.

Likewise, Vargas Larreta (2013) asserted that it is necessary to use the local and traditional knowledge about the biodiversity to incorporate it in the forest planning and management to guarantee better results in their conservation.

On the other hand, Jiménez (2012) alluded to the relation between the anthropic action and distance from the forest, asserting that local players who are close to the forest know the species and their uses.

According to Packham (1993), although it exists a growing awareness of the importance of the wild fruits and other




Figure 4. Distribution of frequencies according to the perception of the local actors on knowledge of the species and the school level.

non-mature stand products from the forests, it is still a very little knowledge of their importance to keeping the families safe during difficult climatic, nutrition, and financial periods. This short literature review seems to show that the social and economic security should be attributed to the highly diverse environment that

sustains these forested areas of Miombo (Africa). The perception of the local actors on the environmental function was contrasted with the knowledge they had about the species, resulting in all local actors know the fruit species, while only the 46% recognized their environmental importance (Figure 5).




Figure 5. Distribution of frequencies according to the local actors' perception of the environmental function and the knowledge about the species.

The environmental perception is an important element to consider for the environmental administration since it reinforces the responsibility level and the local actors' right to the forest (Mitjans, 2012).

From the results of the correlation analysis (Table 2) by applying the coefficient of Rho of Spearman, it is observed a significant relationship between the variables causes of deterioration and economic importance. The relation among

these variables is inverse, which indicates that the lower the economic importance, the increasing in the causes of deterioration. It is conditioned by the lack of knowledge about the economic, environmental, and nutrition importance of the native fruits. It is an issue of great importance for the African countries since different studies in other parts of this continent have concluded that the wild fruits provide an important dietary component, mainly to the children (Packham, 1993).

Table 2. Correlation analysis between the studied variables according to the correlation coefficient Rho of Spearman.

	Environmental Function	Economic Importance	Knowledge of the species	Deterioration Causes
Environmental Function	1.000	-0.102	-0.072	0.05
Economic Importance	-0.102	1.000	0.137	-0.192*
Knowledge of the species	-0.072	0.137	1.000	-0.01
Deterioration Causes	0.05	-0.192*	-0.01	1.000

*: $P < 0.05$

Strengths and weaknesses found in the community of Munhiba

During the participatory diagnosis were also identified strengths and weaknesses that allowed to propose a group of working lines to facilitate the participatory NF self-management by the community of Munhiba.

Strengths

1. There is a diversity of NF species in the community.
2. Suitable edafo-climatic and ecological condition for NF production.
3. Conducted studies show that Mozambique's NF have potential uses (feeding, medicine, the industry of cosmetics, building supplies, wood, and extraction of essential oils).
4. Many of the NF are protected by beliefs and traditional myths.
5. Most of the local actors use the NF as the staple and daily food due to the economic situation and precarious conditions in which they inhabit.
6. There is an available budget for projects on feeding.
7. Simple technology and low-cost expenses will be required if local actors are included in the handling of NF.
8. A high percentage (82%) of the interviewees knew 100% of the NF of the study area.

Weaknesses

1. Limited community knowledge on NF nutritional security.
2. The absence of systematic and participatory approaches through which the community could negotiate its patrimony in an effective participatory way.
3. Insufficient availability of upgraded technical people in the community institutions to conduct NF research and production processes.
4. Inadequately qualified technicians in the necessary research areas.
5. Insufficient technical support for the producers.
6. Low applied investment in the research and implementation projects on NF.
7. There is not an intentioned production and processing of NF, so it responds to the necessities of the community.
8. The community does not value the NF importance to improve their economy and quality of life.
9. There is not an integral economic, environmental and social perception to facilitate the suitable NF self-management.
10. Faulty environmental management that leads to the NF exploitation from inadequate uses, as it is the felling for firewood and coal.

11. Insufficient training of the community on the participatory processes in NF administration and use.
12. Annual practices of forest fires in the herbaceous stratum where the NF are.

These elements related to the strengths and weaknesses show that it is necessary to promote community work. According to the approaches of Reyes *et al.* (2017), it becomes more and more opportune and indispensable for the economic and

social development of a country.

Just as it is reflected in Table 3, there were identified 17 native fruits species, distributed in 14 families and 16 genders, which are used for feeding, the control of the illnesses, and sorcery; as the local actors identify in a territory. The family with more representative species is Malvaceae with three species (*Adansonia digitata* L., *Anzanza garckeana* F. Hoffm, and *Ceiba pentandra* L.).

Table 3. Identified species in the study area.

No.	Common name	Scientific name	Family
1	Malambe- imbondeiro	<i>Adansonia digitata</i> L.	Malvaceae
2	Maebe	<i>Annona senegalensis</i> Pers.	Annonaceae
3	Maceraou umtanwala	<i>Ancylobotrys petersiana</i> Klotzsch.	Apocynaceae
4	Mukole	<i>Anzanza garckeana</i> F. Hoffm.	Malvaceae
5	Sumauma	<i>Ceiba pentandra</i> L.	Malvaceae
6	Tongoma	<i>Eugenia mosambicensis</i> Engels.	Myrtaceae
7	Nalu	<i>Ficus sycomorus</i> L.	Moraceae
8	Tubi	<i>Parinari curatellifolia</i> Planch.	Chrysobalonaceae
9	Chindo	<i>Phoenix reclinata</i> L.	Palmaceae
10	Matema	<i>Strychnos madagascariensis</i> Lam.	Loganiaceae
11	Massala, mutamba	<i>Strychnos spinosa</i> Lam.	Loganiaceae
12	Missica, uepa	<i>Tamarindus indica</i> L.	Fabaceae
13	Mafureira	<i>Trichilia emetica</i> Vahl subsp.	Meliaceae
14	Nyunkomazhanje.	<i>Uapaca kirkiana</i> Müll. Arg.	Phyllanthaceae
15	Mangiro, maphilo,	<i>Vangueria infausta</i> Burch.	Rubiaceae
16	Puro	<i>Vitex doniana</i> Sweet.	Lamiaceae
17	Pudho	<i>Zanha africana</i> (Radlk) Exell.	Sapindaceae

It is important to underline that some of the NF species found in the area were already studied by Cândido (2011) in the district of Quissanga, such as the case of *Annona senegalensis*, *Ancylobotrys petersiana*, *Phoenix reclinata*, *Adansonia digitata*, *Ceiba pentandra*, *Strychnos madagascariensis*, *Tamarindus indica*, *Trichilia emetica*, *Anzanza garckeana*, *Vangueria infausta*, *Zanha africana*, and *Vitex doniana*. This author explained that they are widely used as food and commercialized by the local actors.

Table 4 shows different uses of the species by category. Among the species three of them (*Phoenix reclinata* L.,

Tamarindus indica L., and *Trichilia emetica* Vahl subsp) have a wide scope of uses.

Out of the 118 interviewees, 77 (65%) declared that they eat the leaves of the *Ceiba peltandra* and its toasted seeds.

Among the identified species, it was found the *Strychnos madagascariensis*, which is used as a food and medicine, has been yet studied by Inguane *et al.* (2015). They found in it a considerable percentage of sugars, lipids, proteins, macronutrients (K, Na, Mg, and Ca), micronutrients (Cu, Ni, Zn, Mn, and F), as well as a content of vitamin A, alkaloids, and saponins (anti-cholesterol and anti-cancer).

A high nutritional and medicinal potential can be exploited for nutrition security during scarcity periods.

treatment. Other authors like Munyemana *et al.* (2015) found similar results in conducted studies.

Strychnos spinosa Lam. It is a fruit used for feeding, liquor production, and in the traditional medicine for illnesses

Among the interviewees, 79% (93/118) identified the NF as a necessary food and medicine source for their life development.

Table 4. Category by use declared of the species.

No	Scientific name	Category of use			
		Food	Medicine	Wood	Other uses
1	<i>Adansonia digitata</i> L.	x	x		
2	<i>Annona senegalensis</i> Pers.	x	x		
3	<i>Ancylobotrys petersiana</i> Klotzsch.	x			
4	<i>Anzanza garckeana</i> F. Hoffm.	x	x		x
5	<i>Ceiba pentandra</i> L.	x		x	x
6	<i>Eugenia mosambicensis</i> Engels.	x	x	x	
7	<i>Ficus sycomorus</i> L.	x	x		
8	<i>Parinari curatellifolia</i> Planch.	x	x	x	
9	<i>Phoenix reclinata</i> L.	x	x	x	x
10	<i>Strychnos madagascariensis</i> Lam.	x	x		
11	<i>Strychnos spinosa</i> Lam.	x	x	x	
12	<i>Tamarindus indica</i> L.	x	x	x	x
13	<i>Trichilia emetica</i> Vahl subsp.	x	x	x	x
14	<i>Uapaca kirkiana</i> Müll. Arg.	x	x		
15	<i>Vangueria infausta</i> Burch.	x	x		
16	<i>Vitex doniana</i> Sweet.	x		x	x
17	<i>Zanha africana</i> (Radlk) Exell.	x			

Working lines proposed for the NF self-management in the community of Munhiba

The inadequate NF management caused insufficient protection of them, leading to the damage of forest ecosystem, where they are. Besides, not taking advantage of their total benefit as nutritious potential and source of employments and earnings for the local actors in the community of Munhiba.

The proposed working lines aim to place the local actors as the immediate beneficiaries; they leaned on in their suggested ideas (from the groups' discussion) to help the promotion of a functionally participatory development process under a sociocultural vision. These work lines have been figured out as follows:

1. To carry out exchange participatory meetings with the community and the institutions, applying participatory

technics to the creation of democratic knots that allow, in a combined way, the decision-making about the NF management.

2. Upgrading the population's knowledge, both individual and collective, to assume in a participatory way the administration of the NF in their territory.
3. To elaborate and bring into the forest management, practical plans with an integral and holistic vision of the forest ecosystem, with specificities in the NF; contributing with intellectual instruments and supported on methodological tools that allow accessing and building up an environmental knowledge.
4. Designing and implementing integrated programs, to maximize the use and exploit the potential of each species.
5. To identify the more outstanding NF at the economic level and to adopt measures for their great scale domestication and production.

6. Implementing programs of secondary and high education students' formation, as well as technical education level in different application areas to the NF.
7. To document the local knowledge on NF (ethnobotanical studies) and to create mechanisms of popularization of the local know-how.
8. Promoting in the community the nutritive, medicinal, and commercial importance that have the NF.
9. To train the local actors in how to manage the NF.
10. To make a catalog that provides information regarding NF nutrition security and the collateral benefits for their management in a participative way.
11. To carry out protection practices against fires, pests, diseases, and anthropogenic activities.

CONCLUSIONS

The developed diagnosis and its analysis facilitated to verify the limited participation and local actors' knowledge (producers, processors, and merchants of fruits) from Munhiba town. The lack of work lines was also verified for a community self-management, so that contributes to its production and commercialization.

Despite the degradation by the man's influence in the study area, it is evident the high potential of NF mean for the community, mainly as edible, medicinal, and wood use, represented by 17-identified species; some of them of high commercial value as *Vangueria infausta* (Mangiro), *Strychnos spinosa* (Massala) and *Trichilia emetica* (Mafurra).

The working lines for NF self-management proposals in the community of Munhiba are based in the developed diagnosis and the theoretical conceptions, and they are conceived to be executed with a participatory social focus that contributes to their appropriate management.

REFERENCES

- AIDER and FAO. 2016. Prácticas de manejo para el uso múltiple sostenible en bosques comunitarios de la Amazonía peruana. Guía para el facilitador. Módulo Introductorio: Lineamientos metodológicos y pedagógicos para la capacitación en Manejo Forestal Comunitario. Lima, Perú.
- Brivio Borja A. 2001. La autogestión comunitaria. In: Gestipolis, <https://www.gestipolis.com/la-autogestion-comunitaria/>; accessed: March 2015
- Calero Vinelo A. 1978. Técnicas de muestreo. Editorial Pueblo y Educación, Ciudad de la Habana. 514 p.
- Cándido N. 2011. Testagem de diferentes métodos de

garfagem na Massaleira (*Strychnos spinosa* Lam.). Mater's Thesis in Rural Development. Faculty of Agronomy and Forest Engineering. Universidade Eduardo Mondlane. Maputo. 59 p.

Chiau E, Cruz Francisco JD, Bergenstahl B and Sjöholm I. 2003. Softening of dried *Vangueria infausta* (African medlar) using maltodextrin and sucrose. African Journal of Food Science 7(10): 382-391. doi: 10.5897/AJFS2013.1034

Dewalt KM and Dewalt BR. 2002. Participant observation: A guide for fieldworkers. Alta Mira Press, Walnut Creek. 285 p.

FAO. 1982. Especies frutales forestales. Fichas Técnicas. Roma, 157p.

García-Frapolli E, Toledo VM and Martínez-Alier J. 2008. Apropiación de la naturaleza por una comunidad maya Yucateca: Un análisis económico ecológico. Revista de la Red Iberoamericana de Economía Ecológica 7: 27-42.

Goulão LF and Santo-Antonio V. 2015. Avaliação do estado actual do conhecimento sobre fruteiras nativas em Moçambique. Instituto de Investigação Científica Tropical, Matola. 167 p.

Gomes M, Souza A and Carvalho R. 2001. Diagnóstico Rápido Participativo (DRP) como mitigador de impactos socioeconômicos negativos em empreendimentos agropecuários. pp. 63-78. In: Brose M. (ed.). Metodologia participativa: uma introdução a 29 instrumentos. First edition. Tomo Editorial. Porto Alegre. 328 p.

Infante-Ramírez KD. 2011. Valoración de unidades y paisaje en la zona maya de Quintana Roo. Master's Thesis in Science in Natural Resources and Rural Development. El Colegio de la Frontera Sur, México. 97 p.

Infante-Ramírez KD and Arce Ibarra A. 2013. Percepción local de los servicios ecológicos y de bienestar de la selva de la zona maya en Quintana Roo, México. Investigaciones Geográficas, Boletín del Instituto de Geografía, UNAM (86): 67-81. doi: 10.14350/ig.36593

Inguane S, Maida A and Pagula F. 2015. Avaliação nutricional da Macuacua (*Strychnos madagascariensis*) e dos seus subprodutos. In: 1º Workshop Nacional de Fruteiras Nativas. Matola, Moçambique. 10 p.

Jiménez A. 2012. Contribución a la ecología del bosque semideciduo mesófilo en el sector oeste de la Reserva de la Biosfera "Sierra del Rosario", orientada a su conservación. Ph.D's Thesis in Forestry Science. Universidad de Pinar del Río, Cuba. 100 p.

MA-Ministério da Agricultura. 2011. Plano Estratégico de Desenvolvimento do Sector Agrário (PEDSA 2011-2020). In: Land Portal, <https://landportal.org/library/resources/plano-estrat%C3%A9gico-para-o-desenvolvimento-do-sector-agr%C3%A1rio-pedsa-2011-2020> 76 p.; accessed: January, 2016.

MAE-Ministerio da Administração Estatal. 2005. Perfil do distrito de Mocuba, provincia da Zambézia. In: Portal do Governo de Moçambique, <http://www.portaldogoverno.gov.mz/por/content/download/2890/23502/version/1/file/Mocuba.pdf> 87 p.; accessed: March 2015.

Marzin J, Benoit S, López TV, Cid G, Peláez O, Almaguer N, Herrera JA and Rose M. 2014. Herramientas metodológicas para una extensión agraria generalista, sistémica y participativa. First edition. Editora Agroecológica, La Habana. 150 p.

Molero M, Cubas C, Calderón L, Aguirre C and Fassbender D. 2012. Manejo forestal comunitario. Lecciones aprendidas. In: X Congreso Forestal Nacional "Bosque Sostenible con Inclusión Social". Pucallpa. 13 p.

Mitjans B. 2012. Líneas estratégicas participativas para la rehabilitación del bosque de ribera del río Cuyaguaje, municipio Guane. Ph.D's Thesis in Forestry Science. Universidad Pinar del Río. Pinar del Río. 90 p.

Molina-Peegrín Y, Santos-Chacón W, Sosa-López A, Hechavarría-Kindelán O and Cruza-La Paz M. 2011. Percepción ambiental por los actores sociales de la reserva ecológica El Gigante. *Revista Forestal Baracoa* 30(1): 79-86.

Moreno J. 2013. La gestión comunitaria de recursos naturales, agrosilvopastoriles y pesqueros en la Sierra de Santa Marta, Veracruz, México: ¿una alternativa posible al discurso desarrollista y a la globalización capitalista? *Universitas Humanística* 75(75): 189-217.

Munyemana F and Nhaca IAA. 2015. Avaliação comparativa da composição fitoquímica e actividade antioxidante da polpa, casca e sementes do fruto de *Strychnos spinosa* (Massala). In: Goulão LF and Santo-Antonio V. 2015. Avaliação do estado actual do conhecimento sobre fruteiras nativas em Moçambique. Instituto de Investigação Científica Tropical, Matola. 167 p.

Packham J. 1993. The value of indigenous fruit-bearing trees in miombo woodland areas of South-central Africa. *Social Forestry Network*: 9-15.

Reyes E. 2013. Estrategia de autogestión comunitaria en el sector

campesino, Circunscripción 16 del municipio Guane. Alternativa para el cambio social. Master's Thesis in Social Development. Faculty of Social Sciences and Humanities. Universidad de Pinar del Río. Pinar del Río. 80 p.

Reyes E, Mitjans B and Camallea O. 2017. La autogestión comunitaria como alternativa al cambio social. Diagnostico comunitario participativo. *Revista Caribeña de Ciencias*. 11 p.

Rodríguez G, Gil J and García E. 2008. Metodología de la investigación cualitativa. Félix Varela, La Habana. 300 p.

Sabogal C, de Jong W and Louman B. 2008. Manejo forestal comunitario en América Latina. Experiencias, lecciones aprendidas y retos para el futuro. CIFOR-CATIE, Belém. 274 p.

Simone M. 2001. Estudo de plantas medicinais em uso pelas comunidades locais no posto administrativo de Mahel e sua propagação. In: Links project-FAO (eds.). Síntese dos Trabalhos de Teses e de Investigação em LinKs existentes nas Instituições de Ensino Superior e de Pesquisa, 1990 - 2003. 88 p.

Valdés J. 2010. Los procesos de organización agraria en Cuba 1959-2006. Fundación Antonio Núñez Jiménez, La Habana.

Vargas Larreta B. 2013. Manual de mejores prácticas de manejo forestal para la conservación de la biodiversidad, en ecosistemas templados de la región de México. First edition. Comisión Nacional Forestal, Zapopan. 90 p.

© 2019. This work is published under

<https://creativecommons.org/licenses/by-nc-sa/4.0/>(the “License”).

Notwithstanding the ProQuest Terms and Conditions, you may use this content
in accordance with the terms of the License.