

## Baseline Study of Mercury in Fish of Shatt Al-Arab river by Cold Vapor Atomic Absorption Spectrometry

N. A.N.Awad and H.T. Abdulsahib

Chemistry Department, Collage of science, University of Basrah

### Abstract

Cold vapor atomic absorption spectrometric determination of mercury in six fish species from Shatt Al-Arab river, provide values that are lower than the most literature data. We found significant species differences, with *Cyprins Carpio* having the highest levels ( $0.160 \pm 0.045 \mu\text{g/gm}$ ) and *Barbus sharpeyi* the lowest levels ( $0.053 \pm 0.026 \mu\text{g/gm}$ ). Statistically significant differences of total mercury concentration were also found among the individually tested tissues (muscle, kidney, liver, skin and gills), since the muscle of the tested fish accumulate the highest amount of mercury ( $0.067- 0.160 \mu\text{g/gm}$ ). The data illustrates that Shatt Al-Arab river is still a non mercury– polluted area at the time of analysis.

,  
*Cyprins Carpio* , *Barbus luteus* , *Liza abu* , *Tilapia zillii*, *Barbus*  
 . % 50 *sharpeyi* , *Tenaulosa ilisha*  
*CyprinsCarpio* / ( $0.160 \pm 0.045$ )  
 / ( $0.137 \pm 0.053$ )  
 . / ( $0.056 \pm 0.014$ )

### ***Introduction***

It is well know that fish can be very nutritious and are packed with great nutrients such as omega-3's, the B vitamins and lean protein that are essential for growth and development [Daviglius et al 2002, Patterson 2002]. But unfortunately, fish can also have some unhealthy contaminants.

Mercury and its compounds are included among the most toxic substances found in the environment [Renner, 1997; Wheatley and Wyzga, 1997]and it is exists in a large number of different physical and chemical forms with a wide range of physical, chemical, and ecotoxicological properties that are of fundamental importance to its environmental behavior [Lambertsson et al., 2001; Queuvallier et al.2000] . Mercury in fish can affect brain development and damaging the central nervous system. The three most important inorganic chemical forms of mercury (elemental mercury –  $Hg^0$ , mercurous –  $Hg_2^{2+}$  and mercuric –  $Hg^{2+}$  ions) are far more soluble and have a strong affinity for many inorganic and organic legends, especially those containing sulfur. Those forms are strongly accumulated in aquatic organisms and also concentrated in aquatic food chains i.e., fish, birds etc. [polyi et al 2004; Ebinghauser et al., 2000]Fishes in the surface water serve as bioindicators of pollutant loads [Teh et al., 1997],it is therefore a product for which suitable measures should be taken to provide chemical monitoring of the risks deriving from its consumption.

The major health impacts caused by mercury affect mostly individuals who have a regular fish diet. Human exposure to mercury intoxication through the ingestion of mercury contaminated fish has been well review[Harada 1995].

The FDA has released guidelines for children, women who are pregnant and women who are trying to become pregnant. These guidelines state that no more than 12 oz of low mercury fish should be consumed weekly. "Highest" mercury fish should be avoided and "high" mercury fish should be kept to only three 6-oz servings per month. Most concern has centred on the presence of mercury in fish since seafood is a major source of

mercury in non-occupationally exposed populations [Faro and Seidler 1960].

Fish and fish products account for most of the organic mercury in food. The average daily intake of mercury from food is in the range 2–20 µg/day, but may be much higher in regions where ambient waters have become contaminated with mercury and where fish constitute a high proportion of the diet [Gorchev 1991]. Mercury concentration in fish are influenced by fish age , mercury concentration in water ecosystem , food contamination , chemical , biological and physical process in aquatic environment and seasonal variations [Svobodova et al 2004, Sarica et al 2005 ].

The people are faced with conflicting information about the risks and benefits of consuming fish. The aim of this study is to determine the mercury concentration levels in the six fish species from Shatt Al-arab river to providing data on the most commonly eaten fish that will help people make informed decisions about risks from fish consumption.

### *Experimental*

#### **Apparatus:**

The determination of mercury in the fish samples achieved by means of cold mercury vapor atomic absorption spectrometry (CVAAS) using shimadzu atomic absorption spectrophotometer model (AA-630-12). The cold vapor system used has been described in detail elsewhere [Ali 2006].

#### **Reagents:**

Chemicals of analytical – reagent grade and distilled – deionized water were used for the preparation of all solutions:

- 1- 1000 ppm Hg<sup>2+</sup> solution, 1.3535 gm of HgCl<sub>2</sub> was dissolved in 50 ml of concentrated HCl and diluted to 1 Litre with deionized water .
- 2- 25% (w/v) SnCl<sub>2</sub> ,25 gm of SnCl<sub>2</sub> was dissolved in 100 ml of 20% (v/v) HCl .
- 3- 0.05 % (w/v) K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> .0.05 gm of K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> was dissolved in 2% (v/v) HNO<sub>3</sub> solution .

#### 4- Crystal potassium persulphate

##### **Sample preparation:**

The freshwater Shatt Al-Arab is the longer river in Basrah (180 km). Fish from this river serve as the main source of protein for the population of Basrah city Six species of fish commonly consumed in Basrah were collected in September 2006 at various sites on the shore of Shatt Al-Arab as shown in Fig.1. Some fish samples were bought directly from local fish markets. Total length (TL) of fish was ranged among 22.5 to 44.6 cm. Ages of the fish were between 3–7 years. Muscle flesh from the mid-dorsal region of the five species was removed. The samples were first sun dried on the beach sand until nearly dry (about 10% of the water remained) .Dried sample (0.5 gm) and 15 ml HNO<sub>3</sub> were placed in a digestion tube fitted with a 40 cm air-cooled condenser .A temperature and time controlled heated digestion unit was used as follows: 25°C (4 hrs.) ;70°C (3hrs.) and 140°C(6 hrs.). The condenser was removed to reduce the volume to 2 ml .After cooling ,5 ml HClO<sub>4</sub> was added, the air condenser replaced and the following temperature-time schedule implemented : 140°C (10 min.), 160°C (10 min.), 180°C(10 min.)and finally 210°C for 1 hr . This gradual increase in temperature is necessary to prevent charring of the material. The digestion block was cooled to 170 °C and the air condenser removed to reduce the volume to 1 ml . After cooling to room temperature , the solution was quantitatively transferred to a standard flask and diluted to 20 ml with deionized water.

##### **Procedure:**

Two ml of digested fish samples was introduced in the reduction vessel, 2 ml of SnCl<sub>2</sub> solution was added and mixed using magnetic stirrer for 2 min. then the mercury vapor forced by nitrogen gas with rate of 0.25 L/min in range (1.0 – 25 ng/ml) (Fig.2). A calibration graph prepared from aqueous standard mercury solution to which SnCl<sub>2</sub> solution was added and measured by the above procedure. The optimum parameters of atomic absorption spectrometer for the determination of mercury were listed in

Table (1). Detection limit calculated following the recommendations of IUPAC was 0.73 ng Hg /gm dry weight.

#### Statistical analysis:

The statistical analysis were carried out using two-way analysis of variance with unbalanced repeated measurements .Statistical between individual time points was made by using Revised Least Significant Difference (RLSD) test. The probability level for significance was 5% or less .

#### Result and discussion

Six fish species were selected for monitoring bioaccumulation of total mercury in river ecosystems. There are four possible routes for substances such as mercury to enter fish body. One way is by means of the food ingested [ Baudin 1987] and another is by means of absorption of the mercury in its ionic form by the fish through the gills [ Heath 1987]. This can occur probably by simple diffusion and possibly through the pores in the gills [ Bryan 1979] . Indications are that metal uptake in the gill tissues may be correlated with mass specific rates, with small fish accumulating metals more rapidly than larger ones [Anderson and Spear 1980]. A third possible route is through drinking water [Eddy 1981] while a fourth is by absorption through the skin.

The bioaccumulation of mercury was evaluated in selected fish tissues of *Tenaulosa ilisha* ,the mercury concentrations in the tested tissues are presented in Table (2) .The results showed that the highest concentrations of the total mercury were determined in muscle tissues ( $0.137 \pm 0.053\mu\text{g/gm}$ ) and the lowest concentrations of total mercury were found in gills ( $0.056 \pm 0.014\mu\text{g/gm}$ ) and skin ( $0.042 \pm 0.017\mu\text{g/gm}$ ) of tested fish .Statistically significant differences of mercury concentrations were found among the individually tested tissues with the following exceptions: the skin tissues have statistically insignificant differences compared to the gill ( $P < 0.05$ ) also statistically insignificant differences of the mercury concentrations were found between liver and kidney(  $P < 0.01$ ) The mercury content in tested tissues of *Tenaulosa ilisha* decreased in order: muscle >>

kidney  $\approx$  liver  $>$  skin  $\approx$  gills .; this can attributed to the influences of the living environment on the total mercury content in muscle tissues of fish where mercury is bound to cysteine rich proteins [Boening 2000,Anonymous 2002] . Also the other authors [Anonymous 1999,2002 ,Dusek et al 2005] observed high concentrations of the total mercury in muscle tissues of various fish species .Vigh et al 1996 found the highest mercury concentrations in the kidney of grass carp .Only the flesh of freshwater fish in England revealed higher values than those in the other tissues [Barak and Mason 1990 ) .

Mercury concentrations between fish species *Cyprins Carpio* , *Barbus lutaus*, *Liza abu*, *Tilapia zillii*,*Barbus sharpeyi* and *Tenaulosa ilisha* were significantly different ( $p < 0.0001$ ). Table (3) showed that *Cyprins Carpio* had the highest mercury concentrations ( $0.160 \pm 0.045\mu\text{g/gm}$ ), followed by *Barbus lutaus* ( $0.149\pm 0.075\mu\text{g/gm}$ )The lowest mercury concentrations were found in the two species *Liza abu* and *Barbus sharpeyi* with mean concentrations of ( $0.067 \pm 0.027\mu\text{g/gm}$ ) and ( $0.053 \pm 0.026\mu\text{g/gm}$ ) respectively. The high mean mercury concentrations observed in *Cyprins Carpio* can be attributed to the widest food web (algae, aquatic plants and terrestrial seeds ,larvae and molluse animals). Because of the low range fish age (3-7 years) their age did not influence the content of total mercury in the tested tissues.

In Table(4) literature data for levels of total mercury in fish are presented. Even for the same type of water interspecies comparison has limited value. Consequently our data can only be compared with the scarce values for the same or related *Tilapia* species as shown in Table (5). The total mercury content in the muscle of *Tilapia zillii* from Egypt (Maryut Lake) [ El Nabawi et al 1987 was 4-times higher compared to the muscle samples from Egypt (Niger Delta)[ Kakulu and Osibanjo 1986] and 65-times lower compared to the muscle samples from Egypt(Wadi El-Raiyan Lake) [Saleh et al 1988]. Levels of mercury for the Egypt (Nile River ) fish samples were identical for those found in the fish samples from Egypt (Maryut Lake) [ El Nabawi et al 1987] . In the fish samples from Kenya

(Lake Nakuru) the total mercury levels obtained by Koeman et al [1972] were significantly lower than those found by Greichus et al [1978b] .

The FAO limit for total mercury in fish products , which is also adopted by various countries , is  $0.5\mu\text{g/gm}$  ,(FAO 1983). The WHO (1987) recommendeds a permitted value of  $0.31\mu\text{g/gm}$  .In all fish samples analyzed a concentration level of at least thirty times lower was observed

Mercury in fish has been featured in the media frequently ,and people are faced with conflicting information about the risks and benefits of consuming fish.We suggest that the state agencies responsible for protecting the health of their citizens should obtain information on fish availability in markets and fish preferences of diverse groups of citizens and use this information to select fish for analysis of contaminant levels , providing data on the most commonly eaten fish that will help people make informed decisions about risks from fish consumption.

### **Conclusions**

The levels of mercury determined are higher in *Cyprins Carpio* than in the other fish species, all of which are consumed by humans The muscle tissue had the highest mercury concentrations in comparison with other organism. The levels of mercury in the tested fish species are , as far as could be compared , below published values for other fish and well below the acceptable limits for mercury in foods and thus unlikely to cause adverse effects to aquatic organisms. Our results help us to understand better the accumulation of the individual mercury in the selected tissues of the fish.

**Table (1) :Instromental Parameters**

|   |        |
|---|--------|
| Wave length (nm)                        | 253.65 |
| Band width (nm)                         | 0.3    |
| Lamp current (mA)                       | 6.0    |
| Back ground correction                  | None   |
| Time of mixing (sec.)                   | 60     |
| Flow rate of N <sub>2</sub> gas (L/min) | 0.25   |
| Conc. Of SnCl <sub>2</sub> (%)          | 25     |
| Volume of SnCl <sub>2</sub> (ml)        | 2.0    |
| Temprature (°C )                        | 80     |
| Volume of sample (ml)                   | 2.0    |

**Table (2):Concentrations of mercury (µg/gm ,wet weight) in different organs in *Tenaulosa ilisha* fish from Shatt Al-Arab river.**

| Organs | No. of sample | Hg Conc.(Mean±S.D)<br>For 5 replicates |
|--------|---------------|--|
| Kidney | 8             | 0.076±0.030                            |
| Liver  | 8             | 0.079±0.025                            |
| Muscle | 10            | 0.137±0.053                            |
| Skin   | 9             | 0.042±0.017                            |
| Gills  | 8             | 0.056±0.014                            |



**Table (3): Concentrations of mercury ( $\mu\text{g/gm}$ , wet weight) in the muscle of different species of fish from Shatt Al-Arab river.**

| Species                | No. of sample | Hg Conc For 5 replicates.<br>(Mean $\pm$ S.D) |
|------------------------|---------------|---|
| <i>Liza abu</i>        | 9             | 0.067 $\pm$ 0.027                             |
| <i>Barbus luteus</i>   | 11            | 0.149 $\pm$ 0.075                             |
| <i>Barbus sharpeyi</i> | 10            | 0.053 $\pm$ 0.026                             |
| <i>Tenauosa ilisha</i> | 10            | 0.137 $\pm$ 0.053                             |
| <i>Cyprins Carpio</i>  | 10            | 0.160 $\pm$ 0.045                             |
| <i>Tilapia zillii</i>  | 8             | 0.077 $\pm$ 0.036                             |

**Table(4): Concentrations of mercury in freshwater fish .**

| species                     | Location                           | No. of samples | Hg contents ( $\mu\text{g/gm}$ ) | References                |
|-----------------------------|------------------------------------|----------------|----------------------------------|---------------------------|
| <i>Godus morhua</i>         | Norway (fjords)                    | -              | 0.1-0.4                          | Stenner and Nickless 1974 |
| Commercial species          | SW Spain Portugal (Atlantic coast) | -              | 0.79                             | Stenner and Nickless 1975 |
| Composite                   | South Africa (Transvaal)           | -              | 0.52                             | Greichus et al 1977       |
| Composite                   | Zimbabwe (Mcllwaine Lake)          | 3              | 0.23                             | Greichus et al 1978a      |
| Twelve species              | Finland (Lake Pajanne)             | 1774           | 0.05-4.68                        | Hattula et al 1978        |
| <i>Lates niloticus</i>      | Nigeria (Niger River)              | -              | 0.67                             | Ndiokwere 1983            |
| All fish                    | Nigeria (Niger Delta)              | -              | 0.10                             | Kakulu and Osibanjo 1986  |
| <i>Salmon sole</i>          | Pakistan (coastal waters)          | 16             | 0.05 $\pm$ 0.02                  | Jaffar and Ashraf 1988    |
| <i>Forimio niger</i>        | Pakistan (coastal waters)          | 16             | 0.07 $\pm$ 0.02                  | Jaffar and Ashraf 1988    |
| Paralichthys species (sole) | Argentina (Bahia Blanca estuary)   | 188            | 0.85 $\pm$ 0.18                  | Marcovecchio et al 1988   |
| Finfish                     | USA (Louisiana)                    | 52             | 0.4                              | Ramelow et al 1989        |
| Four species                | Spain (Castellan Coast)            | -              | 0.16-0.84                        | Hernandes et al 1990      |
| <i>Rutilus rutilus</i>      | Eastern England(Rivers)            | 338            | 0.01-0.4                         | Barak and Mason 1990a     |
| Nine fish species           | USA (Savannah River)               | -              | <0.5                             | Winger et al 1990         |
| <i>Serranus species</i>     | Italy (Coastal waters)             | -              | 0.09-0.63                        | Giadano et al 1991        |
| Commercial species          | Czechoslovakia                     | 30             | 0.01-0.12                        | Palusova et al 1991       |
| Common fish                 | Spain (Granada Coast)              | 32             | 0.03-1.21                        | Navarro et al 1992        |
| Two species                 | Burundi (Lake Tanganyika)          | 100            | <0.05                            | Benemariya et al 1991     |
| Six species                 | Iraq (Shatt Al-Arab River)         | 58             | <0.3                             | Present study             |

Table(5): Concentrations of mercury in some *Tilapia* species.

| Species                     | Location                       | No. of samples | Hg contents ( $\mu\text{g/gm}$ ) | References               |
|-----------------------------|--------------------------------|----------------|----------------------------------|--------------------------|
| <i>Tilapia grahami</i>      | Kenya<br>(Lake Nakuru)         | 10             | 0.016                            | Koeman et al 1972        |
| <i>Tilapia</i>              | Kenya<br>(Lake Nakuru)         | 10             | 0,22                             | Greichus et al 1978b     |
| <i>Tilapia melanopleura</i> | Egypt<br>(Niger Delta)         | -              | 1.33                             | Ndiokwere 1983           |
| <i>Tilapia zillii</i>       | Egypt<br>(Niger Delta)         | -              | 0.013                            | Kakulu and Osibanjo 1986 |
| <i>Tilapia niloticus</i>    | Egypt<br>(Nile River)          | 5              | 0.05                             | El Nabawi et al 1987     |
| <i>Tilapia zillii</i>       | Egypt<br>(Maryut Lake)         | 4              | 0.06                             |                          |
| <i>Tilapia niloticus</i>    | Egypt<br>(Maryut Lake)         | 5              | 0.04                             |                          |
| <i>Tilapia zillii</i>       | Egypt<br>(Wadi El-Raiyan Lake) | -              | 4.00                             | Saleh et al 1988         |
| <i>Tilapia zillii</i>       | Iraq<br>(Shatt Al-ArabRiver)   | 10             | 0.077                            | Present study            |





Fig. 1. Shatt Al-Arab River map showing sampling stations



**References**

- Ali, M.T, 2006."Long term exposure to mercury and its effect on B1 vitamin in human " M.Sc. Thesis, University of Basrah, Basrah, Iraq.
- Anderson, P.D. and Spear , P.A., (1980) : Copper pharmacokinetics in fish gills, 1 : Kinetics of pumpkinseed sunfish, *Lepomis gibbosus* of different body sizes. Water Res., 14, 1101-1105.
- Anonymous (1999): Toxicological Profile for Mercury-U.S. Department of Health and Human Services. Public Health Service Agency for Toxic Substances and Disease Registry .
- Anonymous (2002): Ecosystem Health-Science Based Solution. Canadian Tissue Residue Guildelines for the Protection of Consumers of Aquatic Life: Methylmercury. National Guidelines and Standard Office Environmental Quality Branch Environmental Canada ,Ottawa.
- Barak N.A.E. and Mason C.F., (1990a) : Mercury ,Cadmium and lead in five species of freshwater fish from eastern England. Sci. Total Environ., 92, 257-263 .
- Barak N.A.E. and Mason C.F., (1990b): Mercury ,Cadmium and lead in eels and roach: the effects of size, season and locality on metal concentrations in flesh and liver . Sci. Total Environ., 92,249-256.
- Baudin,J.P.,(1987): Investigation into the retention of <sup>65</sup>Zn absorbed by the trophic pathway in *Cyprinus carpio* L.:influence of the ingestion frequency and the radiozink content of food.Water Res., 21,285-294.
- Benemariya H., Robberecht H. and Deelstra H.(1991)."Atomic absorption spectrometric determination of zinc,copper,and selenium in fish from Lake Tanganyika ,Burundi ,Africa" The Science of the Total Environment,105,73-85.
- Boening D.W. (2000): Ecological effects ,transport. And fate of mercury :a general review. Chemosphere ,40, 1335-1351.
- Bryan , G.W., (1979) : Bioaccumulation of marine pollutants. Philos . Trans. Res. Soc. (London) B,286,483.
- Daviglus M., Sheeshka J., Murkin E.. (2002), Health benefits from eating fish. Comments Toxicol 8(4-6):345-374

- Dusek L., Svobodova Z., Janouskova B., Jarkovsky J., Smid R., Pavlis P., (2005) :Bioaccumulation of mercury in muscle tissue of fish in the Elbe River (Czech Republic ) : multispecies monitoring study 1991-1996. *Ecotoxicology and Environmental Safety*. 61,256-267.
- Ebinghauser R., Kock R.R., Schmolke S.R. (2000): Measurements of atmospheric mercury with high time resolution. Recent applications in environmental research and monitoring. *Fresenius Journal of Analytical Chemistry*, 371, 806–815.
- Eddy , F.B., (1981) : Effects of stress on osmotic and ionic regulation in fish, In: A.D.Packering (Ed.) , *Stress and Fish* , Academic Press , New York.
- El-Nabawi,A., Heinzow,B. and Kruse,H., (1987). As ,Cd ,Cu, Pb , Hg and Zn in fish from Alexandria region, Egypt . *Bull. Environ. Contam. Toxicol.*, 39, 889-897 .
- Food and Agriculture Organisation ,(1983), *Compilation of Legal Limits of Hazardous Substances in Fish and Fishery Products (FAO Fisheries Circular no.764)*. FAO ,Rome.
- Giardano, R., Arata, P., Ciaralli,L., Rinaldi, S., Giani, M.,Cicero, A.M. and Costantini,S.,( 1991) . Heavy metals in mussels and fish from Italian coastal waters . *Mar. Pollut. Bull.*, 22,10-14 .
- Gorchev G.H. (1991), Dietary intake of pesticide residues, cadmium, mercury, and lead. *Food additives and contaminants*, 8:793-806.
- Greichus,Y.A., Greichus,A., Draager, H.A. and Marshall,B.,( 1978a), Insecticides polychlorinated biphenyls and metals in African lake ecosystems . II : Lake McIlwaine, Rhodesia .*Bull. Environ. Contam.*, 19, 444-453 .
- Greichus,Y.A., Greichus,A., Amman ,B.D., and Hopcraft ,J., (1978b) Insecticides , polychlorinated biphenyls and metals in African lake ecosystems.III :Lake Nakuru,Kenya. *Bull . Environm. Contam. Toxicol.*, 19, 454-461.
- Greichus,Y.A., Greichus,A., Amman ,B.A, Call,D.J., Hamman,C.D. and Potts. R.M., (1977). Insecticides polychlorinated biphenyls and metals in

- African lake ecosystems . I:Hartbeenspoort Dam, Transvaal and Voelvlei Dam ,Cape Province , Republic of south Africa . Arch .Contam. Toxicol., 6,371-383 .
- Harada M.(1995). Minamata disease: methylmercury. Poisoning in Japan caused by environmental pollution.CritRevToxicol25:1-24.
- Hattula, M.L., Sarkka,J.,Janatuinen,J., Paasivirta ,J. and Ross,A., (1978). Total mercury and methyl mercury contents in fish from lake Pijanne. Environ. Pollut., 17,19-29 .
- Heath , A.G.,(1987). Water Pollution and Fish Physiology . CRC Press,Boca Raton (Florida).
- Hernandez, F.H., Medina,J., Ansuategui,J. and Lopez, F.J.,(1990) . Application of simple procedure of digestion for the determination of trace metals in marine organisms. Analysis ,18,327-330.
- Ipolyi I., Massanisso P.,Sposato S., Fodor P., Morabito R.,(2004):Concentration levels of total and methylmercury in mussel samples collected along the coasts of Sardinia Island(Italy). Analytica Chimica Acta, 505,145-151.
- Jaffar,M. and Ashraf ,M., (1988). Selected trace metal concentrations in different tissues of fish from coastel waters in Pakistan (Arabian Sea ) . Indian J. Marine Sci., 17,231-234 .
- Kakulu,S.E. and Osibanjo, O.,(1986). A baseline study of mercury in fish and sediments in the Niger Delta area of Nigeria .Environ. Pollut., (Series B) 11,322 .
- Koeman, J.H.,Pennings,J.H.,de Goeij,J.J.M., Tsjoe ,P.S., Olindo,P.M., and Hopcraft ,J., (1972). A preliminary survey of the possible contamination of lake Nakuru in Kenya with some metals and chlorinated hydrocarbon pesticides.J. Appl. Ecol., 9, 411-416.]
- Kurland, L.T., Faro ,S.N. and Seidler ,H., (1960), Minamata Disease. World Neurology **1**, 370-90.

- Lambertsson L., Lundberg E., Nilsson M., Frech W. (2001): Applications of enriched stable isotope tracers in combination with isotope dilution C-ICP-MS to study mercury species transformation in sea sediments during in situ ethylation and determination. *Journal of Analytical Atomic Spectroscopy*, 16, 296–301.
- Marcovecchio , J.E., Moreno,V.J. and Perez, (1988b). The sole, *Paralichthys* sp., as an indicator species for heavy metal pollution in the Bahia Blanca estuary, Argentina. *Sci. Total Environ.* , 75,191-199 .
- Navarro , M., Lopez , M.C. and Lopez,H.,(1992). Microwave dissolution for the determination of mercury in fish by cold vapour atomic absorption spectrometry. *Anal. Chim . Acta*, 257, 155-158 .
- Ndiokwere, C.L.,(1983). Arsenic ,gold and mercury concentration levels in freshwater by neutron activation analysis.*Environ. Pollut. (Series B)*, 6,263-269.
- Palusova , O., Ursinyova, M. and Uhnak,J.,(1991). Mercury levels in the components of the environment and diets. *Sci. Total Environ.*, 101, 79-82.
- Patterson J., (2002), Introduction--comparative dietary risk: balance the risks and benefits of fish consumption. *Comments Toxicol* 8(4-6):337-344.
- Queuvallier P., Filippelli M., Horvat M. (2000): Method performance evaluation for methylmercury determination in fish and sediment. *TrAC Trends in Analytical Chemistry*, 19, 157–166.
- Ramelow, G.J., Webre,C.L.,Mueller,C.S., Beck, J.N.,Young,J.C. and Langley,M.P., (1989). Variations of heavy metals and arsenic in fish and other organisms from Calsasien River and Lake , Louisiana. *Arch. Environ. Contam . Toxicol.*, 18,804-818 .
- Renner R. (1997): Rethinking water quality standards for metal toxicity. *Environmental Science and Technology*, 31, 466A–468A.
- Saleh M.A.,Fouda M.M., Saleh M.A., Lattif M.S.A. and Walson B.L., (1988). "Inorganic pollution of the man-made lakes of Wadi-Raiyan and

- its impact on aquaculture and wildlife of the surrounding Egyptian desert". *Arch. Environ. Contam. Toxicol.*, 17,391-403.
- Sarica J., Amyot M., Hare L., Blanchfield P., Bodaly R.A., Hintelmann H., Lucotte M. (2005) Mercury transfer from fish carcasses to scavengers in boreal lakes: the use of stable isotopes of mercury . *Environmental Pollution* ,134,13-22.
- Stenner, R.D. and Nickless, G., (1974). Distribution of some heavy metals in organisms in Hardangerfjord and Skjerstadvjord , Norway. *Water Air Soil Pollut.*, 3,279-291.
- Stenner, R.D. and Nickless, G., (1975). Heavy metals in organism of the Atlantic coast of S.W. Spain and Portugal. *Mar . Pollut . Bull .* , 6, 89-92.
- Svobodova Z., Celechovska O., Kolarova J., Randak T., Zlabek V., (2004): Assessment of metal contamination IN the upper reaches of the Ticha Orlice River . *Czech Journal of Animal Science*, 49,458-464.
- Teh S.J., Adams S.M., Hinton D.E. (1997): Histopathologic biomarkers in feral freshwater fish populations exposed to different types of contaminant stress. *Aquatic Toxicology*, 37, 51–70.
- Vigh P., Mastala Z., Balogh K.V. (1996): Comparison of heavy metal concentration of grass carp (*Ctenopharyngodon idella* ) in shallow eutrophic lake and a fish pond ( possible effects of food contamination). *Chemosphere* ,32,691-701.
- Wheatley B., Wyzga R. (1997): Mercury as a global pollutant– human health issues. *Water, Air and Soil Pollution*,97, 5–7.
- World Health Organization, (1989), *Mercury—environmental aspects*. Geneva, (Environmental Health Criteria, No. 86).
- World Health Organisation ,(1987). *Global Pollution and Health: Global Environmental Monitoring System Report* . WHO, Geneva .
- Winger, P.V., Schultz ,D.P. and Johnson, W.W., (1990). Environmental contaminant concentrations in biota from the lower Savannah River , Georgia and South Carolina . *Arch. Environ. Contam. Toxicol.*, 19, 101-117 .