


December 2020

## Does Invasion Science Encompass the Invaded Range? A Comparison of the Geographies of Invasion Science Versus Management in the U.S.

Lara Munro  
*University of Massachusetts Amherst*

Follow this and additional works at: [https://scholarworks.umass.edu/masters\\_theses\\_2](https://scholarworks.umass.edu/masters_theses_2)

 Part of the [Biodiversity Commons](#), [Natural Resources Management and Policy Commons](#), [Other Earth Sciences Commons](#), and the [Other Plant Sciences Commons](#)

---

### Recommended Citation

Munro, Lara, "Does Invasion Science Encompass the Invaded Range? A Comparison of the Geographies of Invasion Science Versus Management in the U.S." (2020). *Masters Theses*. 978.  
<https://doi.org/10.7275/18685032> [https://scholarworks.umass.edu/masters\\_theses\\_2/978](https://scholarworks.umass.edu/masters_theses_2/978)

This Open Access Thesis is brought to you for free and open access by the Dissertations and Theses at ScholarWorks@UMass Amherst. It has been accepted for inclusion in Masters Theses by an authorized administrator of ScholarWorks@UMass Amherst. For more information, please contact [scholarworks@library.umass.edu](mailto:scholarworks@library.umass.edu).

Does invasion science encompass the invaded range? A comparison of the  
geographies of invasion science versus management in the U.S.

A Thesis Presented

By

LARA MUNRO

Submitted to the Graduate School of the  
University of Massachusetts Amherst in partial fulfillment  
of the requirements for the degree of

MASTER OF SCIENCE

September 2020

GEOSCIENCES

Does invasion science encompass the invaded range? A comparison of the  
geographies of invasion science versus management in the U.S.

A Thesis Presented

By

LARA MUNRO

Approved as to style and content by:

---

Bethany Bradley, Chair

---

Forrest J. Bowlick, Member

---

John T. Finn, Member

---

Qian Yu, Member

---

Stephen J. Burns, Department Head, Geosciences

*It's here they got the range  
And the machinery for change  
And it's here they got the spiritual thirst  
- Leonard Cohen, Democracy*

## ACKNOWLEDGEMENTS

I would like to start by thanking Bethany Bradley, for being an amazing, patient, and empathetic mentor and role model. Thank you so much for taking me on, giving me the tools to do this, and including me in all the other lab projects. The past two years have been tumultuous, but ultimately my time at UMass was positive and that is largely thanks to the spaces that you created, thank you again! I would also like to give an enormous thank you to Brittany Laginhas for the work that she put into developing the data extraction protocol and building the Global Invaders database – this project would probably not exist without them. Another big thank you goes to Bridget Griffin and Evelyn Beaury for the data extraction and GIS and aesthetic consults. More thanks go out to Courtney O’Connell who triaged articles from 2016-2018. Thank you to everyone in the lab and extended group that I have not mentioned yet, Emily, Audrey, Jack, Mike, and Will, for the discussions and coffee breaks that helped me work through and develop this project. I also wanted to thank the other members of my committee, Forrest and Qian, who have been incredibly supportive throughout this process.

I wanted to give a shout out to everyone in the RISCC group, it has been a blast learning and working with all 10-15 (or is it more?) of you! I am also grateful to everyone not mentioned yet who helped keep up my morale/listened to me rant/taught me how to DnD with me over the past 2 years: Corey, Camila, Jeimy, Andréanne, Alexe, Liz, Florence, Ben, Jonah, Jonah, Maurice, and the CCPH. Many thanks to my parents for all the support (including driving over to take me out for supper when I broke my ankle!) these past two years (but also the ones before). Thank you to Akim, for being a great brother, and giving me a place to work in the middle of a pandemic in exchange for “artwork”. Finally, thank you to Michele Cooke, who has been a stellar GPD and helped navigate awkward administrative tasks, I was lucky to have had you around!

## ABSTRACT

# DOES INVASION SCIENCE ENCOMPASS THE INVADED RANGE? A COMPARISON OF THE GEOGRAPHIES OF INVASION SCIENCE VERSUS MANAGEMENT IN THE U.S.

SEPTEMBER 2020

LARA MUNRO,

B.SC., UNIVERSITÉ DE MONTRÉAL

M.SC., UNIVERSITÉ DE MONTRÉAL

M.S., UNIVERSITY OF MASSACHUSETTS AMHERST

Directed by: Professor Bethany Bradley

Biases in invasion science lead to a taxonomic focus on plants, particularly a subset of well-studied plants, and a geographic focus on invasions in Europe and North America. Geographic biases could also cause some branches of invasion science to focus on a subset of environmental conditions in the invaded range, potentially leading to an incomplete understanding of the ecology and management of plant invasions. While broader, country-level geographic biases are well known, it is unclear whether these biases extend to a finer scale and thus affect research within the invaded range. This study assessed whether research sites for ten well-studied invasive plants in the U.S. are geographically biased relative to each species' invaded range. We compared the distribution, climate, and land uses of research sites for 735 scientific articles to manager records from EDDMapS and iMap Invasives representing the invaded range. We attributed each study to one of five types: impact, invasive trait, mapping, management, and recipient community traits. While the number of research sites was much smaller than the number of manager records, they generally encompassed similar geographies. However, research sites tended to skew towards species' warm range

margins, indicating that researchers have knowledge on how these plants might behave in a warming climate. For all but one species, at least one study type encompassed a significantly different climate space from manager records, suggesting that some level of climatic bias is common. Impact and management studies occurred within the same climate space for all species, suggesting that these studies focus on similar areas – likely those with the greatest impacts and management needs. Manager records were more likely to be found near roads, which are both habitats and vectors for invasive plants, and on public land. Research sites were more likely to be found near a college or university. Studies on these plants largely occur across their invaded range, however, different study types occur within a narrower climate range. This clustering can create gaps in our general understanding of how these plants interact with different environments, which can have important policy and management consequences.

Keywords: Biological invasions; Geographic bias; EDDMapS; iMap Invasives; Invasive plant; Spatial bias; Disturbance

## TABLE OF CONTENTS

ACKNOWLEDGEMENTS .....	iv
ABSTRACT.....	v
LIST OF TABLES .....	viii
LIST OF FIGURES .....	ix
CHAPTER	
1. DOES INVASION SCIENCE ENCOMPASS THE INVADED RANGE? A COMPARISON OF THE GEOGRAPHIES OF INVASION SCIENCE VERSUS MANAGEMENT IN THE U.S.....	1
1.1 Introduction.....	1
1.2 Methods.....	4
1.3 Results.....	10
1.4 Discussion .....	17
1.5 Conclusion .....	22
APPENDICES .....	24
A. SUPPLEMENTAL TABLES .....	24
B. SUPPLEMENTAL FIGURES .....	26
BIBLIOGRAPHY .....	28



## LIST OF TABLES

Table	Page
Table 1: Species analyzed in this study .....	5
Table 2: Descriptions and examples of the categories used to classify articles.....	7
Table 3: Mean annual temperature and precipitation of literature and manager record... ..	24
Table 4: Proportion of records within 100 m of a road, on public land, and in a state with a listing for that species.....	24
Table 5: Precision of locations reported in the literature for each species in the lower 48.....	25

## LIST OF FIGURES

Figure	Page
Figure 1: Distribution of literature and manager records in the United States .....	11
Figure 2: Proportion of each study type attributed to each species (A) and distinct study locations (B) in the lower 48. ....	12
Figure 3: Mean annual temperature and precipitation of study sites and manager records for each species .....	14
Figure 4: Combined distribution of distance to roads for manager and literature records in the Lower 48 .....	15
Figure 6: Proportion of records found within a given distance to a road.....	26
Figure 7: Proportion of records found within a given distance to a college or university.....	27

## CHAPTER 1.

### DOES INVASION SCIENCE ENCOMPASS THE INVADED RANGE? A COMPARISON OF THE GEOGRAPHIES OF INVASION SCIENCE VERSUS MANAGEMENT IN THE U.S.

#### 1.1 Introduction

It is well known that spatial and taxonomic biases exist in the invasive plant literature (Pyšek et al. 2008, Hulme et al. 2013). Geographic regions such as the U.S. and invasive plants such as *Phragmites australis* have an oversized footprint in invasion ecology research (Laginhas and Bradley in prep., Hulme et al. 2013). Geographic biases are a problem because they lead to an incomplete view of which species are potentially invasive, their likely impacts, and the efficacy of management options. Even well studied species in well studied regions like the U.S. could be biased in terms of the type and location of scientific analyses. If some types of scientific studies only occur in a portion of the range, for example, treatment methods at a species' cool range margin, this could lead to ineffective management in other parts of the species' range. Thus, an important next step in understanding biases in invasion ecology involves delving deeper into biases associated with particular types of studies.

Studies on taxonomic biases in invasion literature show that plants make up a significant majority of studies on invasive species, among which grasses, forbs, and herbs are overrepresented (Pyšek et al. 2006, 2008, Jeschke et al. 2012, Hulme et al. 2013, Lowry et al. 2013, Stricker et al. 2015, Tekiel and Barney 2017). Of these, a select few species are exceptionally well studied. Hulme et al. (2013) found that a third of all impact studies focus on only nine species, including *Bromus tectorum* and *P. australis*. Large scale geographic biases have also led to an overrepresentation of

Europe and North America and the underrepresentation of Asia, Africa, and South and Central America (Pyšek et al. 2008, Hulme et al. 2013, Bellard and Jeschke 2016). These biases indicate that plants that are invasive in Europe and North America are the most well-studied invasive species and have, in turn, played an important role in the development of central invasion hypotheses (Colautti and Barrett 2013).

On a finer scale, geographic biases could include easily accessible sites, notably sites near roads and research institutions (e.g. herbaria, universities) (Graham et al. 2004, Boakes et al. 2010, Stolar and Nielsen 2015, Daru et al. 2018). A bias in ecological sampling towards roads could be problematic because invasive plants are often linked to landscape scale disturbances associated with road corridors (Vilà and Ibáñez 2011, Menuz and Kettenring 2013, Bhattarai and Cronin 2014). Roadsides can be considered as distinct micro-environments, with distinct soil and climate conditions (Kadmon et al. 2004, Rotholz and Mandelik 2013), that are both habitats and vectors for invasive plants (Jodoin et al. 2008, Christen and Matlack 2009). Thus, biased sampling adjacent to roads could inflate the reported impacts of invasive plants if they are instead a result of disturbance (MacDougall and Turkington 2005). Additionally, plant specimen and samples are often found near research institutions, where they are kept and analyzed (Daru et al. 2018). However, these are not evenly distributed throughout the U.S. Coastal and Great Lake states are home to a higher density of these institutions than the rest of the country (Oak Ridge National Laboratory 2010), which could be the source of a spatial bias towards certain parts of the country. A third potential source of regional bias could result from species being prioritized through state-level noxious weed lists. These lists are mainly used to prevent the sale and import of invasive plants; however, they can also be used to set management priorities (Skinner et al. 2000, Quinn et al. 2013). The identity of state listed species varies considerably between states (Beaury &

Fusco et al. in prep., Buerger et al. 2016) and could create biases in research and management priorities. Collectively, landscape-scale geographic biases could produce a false portrait of the impacts of and vulnerability to plant invasions.

Larger-scale spatial biases in invasion ecology studies could also lead to an overrepresentation of a portion of the climatic range in our scientific understanding of invasions. A bias in sampling towards one margin of the range could produce imprecise or ineffective recommendations for management or understanding of impacts. For example, herbicide efficacy has been found to vary at different temperatures, higher temperatures can notably reduce their effect through reduced stomatal openings and thus uptake, increased plant growth and metabolism, which leads to dilution, and increased soil temperature and volatilization (Bailey 2004, Matzrafi et al. 2016, Ziska 2016). Mechanical removal can also be affected by temperature; for example, *Eichhornia* spp., an aquatic invasive, can be managed by pulling, but only in environments that experience winter freezing (Hellmann et al. 2008, U.S. EPA 2008). Invasive plant traits, notably their phenology, also likely vary across climatic conditions (Hou et al. 2014). For example, *Lythrum salicaria* plants from different North American populations flowered at different times and had different growth rates when grown under a single climatic regime (Colautti and Barrett 2013). Thus, spatial biases towards one climatic range margin could lead to an inaccurate understanding of invasive plant competitiveness throughout its range.

Given the extensive documentation of spatial biases in invasion ecology globally (Pyšek et al. 2008, Hulme et al. 2013, Lowry et al. 2013) combined with the need to use relevant science to guide management and policy actions, it is important to understand how well scientific studies encompass the invaded range. Here, we analyze ten widespread and commonly studied invasive plants in the conterminous U.S. We

compare the spatial distributions of management records to locations of scientific studies to determine 1) whether researchers study these plants in the same range in which managers record them, and 2) whether ecological studies are biased with regards to land use or climate. By measuring landscape- and regional-scale biases in the literature, this study highlights areas that might be overlooked by researchers, which can, in turn, influence management and policy priorities.

## 1.2 Methods

### 1.2.1 Study species

We chose ten invasive plant species that are well studied in the scientific literature and, also, widespread within the lower 48 states (Table 1). We identified well-studied species using the Global Invaders database (Laginhas and Bradley in prep.), which provides an inventory of scientific articles on invasive plants from 1999 to 2018. We used this database to identify well-studied species with scientific articles that also included geographic information (coordinates or a map). We also identified widespread species using spatial records contributed by managers and the public and compiled by the Early Detection and Distribution Mapping System (EDDMapS; Barger and Moorhead 2007) or iMap Invasives (iMap; NatureServe 2019). Although they were slightly less well studied than some other species, we included *Tamarix ramossissima* and *Ailanthus altissima*, a shrub and a tree, respectively, to encompass multiple growth forms. Thus, our ten study species are sufficiently reported in the scientific literature and in management databases to enable a comparison of their spatial overlap.

Table 1: Species analyzed in this study. Species are sorted by the total number of scientific articles with geographic location data published between 1999-2018. Manager records include data from EDDMapS and iMap Invasives. All species except *Lonicera maackii* were among the top 50 most recorded plants in EDDMapS.

Common name	Scientific name	USDA code	Growth Form	Articles (n)	Articles lower 48 (n)	EDDMapS records (n)	iMap Invasives records (n)
Common reed	<i>Phragmites australis</i>	PHAU7	Graminoid	247	148	28763	216
Cheatgrass	<i>Bromus tectorum</i>	BRTE	Graminoid	170	162	28136	6329
Japanese knotweed	<i>Fallopia japonica</i>	POCU6	Forb/herb	93	18	32484	19847
Japanese stiltgrass	<i>Microstegium vimineum</i>	MIVI	Graminoid	86	84	29661	4496
Reed canarygrass	<i>Phalaris arundinacea</i>	PHAR3	Graminoid	83	69	36082	3337
Garlic mustard	<i>Alliaria petiolata</i>	ALPE4	Forb/herb	78	69	51600	13163
Amur honeysuckle*	<i>Lonicera maackii</i>	LOMA6	Shrub	74	74	5657	512
Purple loosestrife	<i>Lythrum salicaria</i>	LYSA2	Forb/herb	68	46	41200	24042
Tree-of-heaven	<i>Ailanthus altissima</i>	AIAL	Tree	56	24	28416	6341
Saltcedar	<i>Tamarix ramosissima</i>	TARA	Shrub	47	41	29637	7675

\**L. maackii* was less widespread in management records, ranking #107 in number of occurrences in EDDMapS

## 1.2.2 Data collection

### 1.2.2.1 Spatial data from the scientific literature

We extracted spatial data for the target species from all articles identified in the Global Invaders database (Laginhas and Bradley in prep.) as having geographic information. This database includes species from all papers from 1999-2016 returned using the search term “INVASI\* PLANT” in Web of Science (Web of Science 2020). To gather consistent information for 2017-2018 for our ten target species, we conducted a Web of Science search (Web of Science 2020) for “INVASI\* PLANT” AND “genus species”, as well as all reported synonyms (ITIS 2020). To be included, an article needed to have recorded the occurrence of the invasive species at a given location and have geographic coordinates with a minimal precision equivalent to 0.1 decimal

degrees (~11 km), or include a map, or an aerial photograph of the study locations. For occurrences reported on maps or aerial photographs, we estimated the location based on toponomy or landmarks using Google maps and recorded these locations to a 0.1 decimal degree precision. For maps with many clustered locations, level of precision that was given or recorded in a map often led to multiple locations being identically recorded, we therefore estimated the centroid of the cluster and reported that location.

To assess whether some subfields of invasion ecology were spatially biased, we classified articles into one of five study types (Table 2). These categories represent research topics that focus on invasion risk factors (invader traits and recipient community traits), ways in which scientists and stakeholders can monitor or respond to plant invasion (management and mapping), or the impact of invasion. A small number of studies, such as reviews, did not fit into any of these categories and were grouped as “other”. We excluded them from comparative analyses as their subfield was ambiguous.



Table 2: Descriptions and examples of the categories used to classify articles.

Study type	Article Focus	Examples
Impact	Impact of the invasive plant on the abiotic environment or biotic communities	Impact of invasion on hydrology (Martinez 2017), native plants (McGlynn 2009) or native fauna (Wiesenborn 2005)
Invasive Trait	Traits of the invasive plant	Germination (McCaughey and Stephenson 2000); population differences (Shi et al. 2018); genetics (Pyšek et al. 2018); allelopathy (Gómez-Aparicio and Canham 2008); plant growth (Collins et al. 2010); seed dispersal (Kaproth and McGraw 2008)
Mapping	Occurrence of the invasive plant and/or its spread	Remote sensing (Narumalani et al. 2009); predictive modelling (occurrence data only; Jarnevich et al. 2014); historical reconstruction of invasion (Lavoie et al. 2005)
Management	Management strategies for the invasive plant.	Efficacy of herbicides (Adams and Galatowitsch 2006) or biocontrol agents (Craine et al. 2016); effect of treatments on native species (Hovick and Carson 2015)
Recipient Community Traits	Traits of the invaded ecosystem prior to invasion or ecosystem traits that facilitate plant invasion.	Abiotic properties of invaded areas (Uddin and Robinson 2018); disturbance (Hager 2004); invaded plant communities (Peter and Burdick 2010); effects of soil fungi (Shearin et al. 2018); effect of herbivory (Williams and Sahli 2016)

#### 1.2.2.2 Spatial data from managers

We compiled occurrence data reported by invasive species managers from EDDMapS (Bargeron and Moorhead 2007) and the iMap databases (NatureServe 2019). EDDMapS is the most used database by managers and citizens to record and track invasive species. However, some states use iMap as their primary repository for invasive species occurrences. Therefore, we also compiled iMap data from Arizona,

Kentucky, Maine, New York, Oregon, and Pennsylvania. Data were downloaded from EDDMapS on March 2, 2020 and from iMap on October 2, 2019. We removed duplicate points from the combined manager database to avoid double-counting sites that were visited multiple times or were reported in both datasets.

### 1.2.3 Data Analysis

Because the majority of EDDMapS and iMap records are in the lower 48 states, we focused our spatial comparison on this region. Records located outside of these states were excluded. Articles sometimes provided measurements at the same plot recorded over time or gave a single latitude and longitude or map location to represent multiple nearby plots with differing occurrence or abundance values. We extracted all abundance or occurrence data for these plots. However, including replicates of the same location could bias our analysis of the spatial characteristics of invasion ecology studies. Thus, we retained only one data point for each individual location, defined by their reported latitude and longitude, in each article. We retained spatial information at the level of precision that each author gave, if only one set of coordinates was given in an article, we reported only one location, but if multiple coordinates were given, we reported each one individually.

In order to visualize the distribution of the two datasets, we created a grid of equal area hexagons with a 50 km cell size height (1623.8 km<sup>2</sup>) encompassing the lower 48. Within each hexagon, we recorded the presence of an occurrence from the literature, from a manager record, or both. To determine whether studies on these species focused on any particular aspect of invasions, we calculated the proportion of papers for a given species that was associated with each study type. We also assessed the number of distinct locations reported within each study type to identify study types or species with more spatial data.

### 1.2.3.1 Climate comparisons

In order to test for differences in climate space between literature and management records, we compared spatial occurrences to 30-year (1981-2010) average annual precipitation and temperature created by the PRISM climate group (PRISM Climate Group 2004). To avoid skewing the comparison with locations that have been studied or sampled multiple times, we performed this analysis using a 4 x 4 km grid size, matching the resolution of the PRISM data. Thus, only one point within each grid cell was retained for analysis. We used a Student t-test to compare the manager data to the literature data as a whole. To determine if any differences exist between the different study types and manager records, we used a Kruskal-Wallis and Dunn post-hoc tests to compare the mean climate conditions.

### 1.2.3.2 Disturbance and other biases

To assess whether either dataset is biased towards more disturbed areas, we calculated the proximity of each independent location, within a given article, to a road using US census data for road locations (U.S. Census Bureau 2016). We used these data to create and compare proportional histograms for the literature and management datasets. The same approach was used to compare the distances to colleges and universities (Oak Ridge National Laboratory 2010), which could influence the sampling strategies of studies reported in the literature. Lastly, we compared the proportion of literature vs. management records found on private vs. public land (USGS Gap Analysis Project 2018) as well as within vs. outside states where the species was regulated (i.e. prohibited from sales or planting; Beaury & Fusco et al. in prep.).

## 1.3 Results

### 1.3.1 Distribution of literature and manager records

The distribution of literature records and manager records is presented in Figure 1. Manager records occur in more grid cells than literature records for all species, which is consistent with the larger number of manager records for all species (Table 1). The mean number of grid cells for manager records across all species is 749 +/- 192 (95% CI), whereas the mean number of grid cells for literature records is 200 +/- 143 (95% CI). In general, literature records appear to encompass the invaded range described by manager records with no appearance of strong spatial biases.

Almost all species have at least one spatially explicit study in each of the five study type categories (Figure 2), except for *F. japonica*, which had no spatial studies on management techniques in the lower 48. On average, impact studies are the most common (29% overall). However, study types vary between species. For example, impact studies represent 45%, 39% and 35% of studies on *L. maackii*, *F. japonica*, and *M. vimineum*, respectively, but less than 15% of studies on *A. altissima* (Figure 2A). In contrast, when comparing numbers of individual study locations, invasive trait studies are the most common (46% overall). This pattern is driven by the large number of sample locations from studies focusing on comparing genetics or plant traits from different populations (Figure 2B).




Figure 1: Distribution of literature and manager records in the United States. Each color represents a 1624 km<sup>2</sup> hexagon in which one or more manager record (light blue), literature record (red), or both (purple) were present.




Figure 2: Proportion of each study type attributed to each species (A) and distinct study locations (B) in the lower 48.

### 1.3.2 Climate comparisons

For most species, we found that mean temperature and mean precipitation differ significantly ( $p < 0.05$ ) between the manager and literature datasets (Table S1). Differences in absolute mean temperature average  $1.0^{\circ}\text{C} \pm 0.8^{\circ}\text{C}$  (95% CI; median  $0.6^{\circ}\text{C}$ ), with *P. arundinacea* and *P. australis* showing the largest difference. For all but two species (*A. altissima* and *T. ramosissima*), the literature records skew towards warmer climate conditions. Precipitation ranges are more variable between the two datasets: six species have less than 50 mm difference between mean annual precipitation, while four (*B. tectorum*, *F. japonica*, *L. maackii*, and *P. australis*) have precipitation differences as high as 180 mm (mean absolute precipitation difference:  $53.0\text{ mm} \pm 45.1\text{ mm}$  (95% CI; median 41 mm)). Literature records for six species tend towards drier conditions, while records for the remaining four species (*B. tectorum*, *F. japonica*, *L. salicaria*, and *T. ramosissima*) tend towards wetter conditions.

For most species, at least one study type has a significantly different average climate (precipitation or temperature) from manager records (Figure 3). One species, *A. altissima* has no significant climatic differences between study types and manager records; however, this species also had low sample sizes. Impact studies are the most likely to occur in a significantly different climate space than manager records (35%; 7 out of 20 possible differences). These differences are significant for both temperature and precipitation in the cases of *L. salicaria* and *P. australis*. Management studies were most climatically similar to manager records (22% significantly different; 4 out of 18 possible cases due to a lack of spatial management studies for *F. japonica*). The other three categories have a significantly different climate for 6 out of 20 possible values (30%). There is no consistent directionality (hotter vs. colder or wetter vs. drier) for the differences between study types and manager records.

Most species also show at least one significant difference in mean climate between study types. For 14 of 20 possible species and climate variable combinations, there is at least one significant study type difference. Impact and management studies are the only study types with no significant differences in mean climate across all species. We found three instances of differences between recipient community trait and impact studies (*L. salicaria*, *P. australis*, and *T. ramosissima*) and three instances of differences between recipient community trait and management studies (*B. tectorum*, *P. australis*, and *T. ramosissima*). As a whole, ecological studies on plant invasions (impact, management, and recipient community trait studies) tend to occur in similar climate spaces.




Figure 3: Mean annual temperature and precipitation of study sites and manager records for each species. Letters indicate significant differences ( $p < 0.05$ ) between study types and/or manager records. *F. japonica* and *P. arundinacea* have a maximum mean annual precipitation of 4314 mm and 5244 mm, respectively.



### 1.3.3 Disturbances and other biases

Both literature and manager datasets tend to be located close to roads, with manager records more likely to be near roads. 54% of manager records are found within 100 m of a road versus 45% of literature records (Figure 4; Figure S1). These values vary between species. Over 70% of *F. japonica* records are found within 100 m of a road (74% manager and 71% literature), whereas less than 25% of *T. ramosissima* records are next to roads (22% manager and 13% literature). The pattern of manager records located closer to roads is consistent across all species (Figure S1).




Figure 4: Combined distribution of distance to roads for manager and literature records in the Lower 48. Values represent the proportion of records found in each distance class for each dataset, all species combined.

For both datasets, most records, regardless of species, are found within 50 km of a college or university, although records for western species (*B. tectorum* and *T. ramosissima*) tend to be farther away. As a whole, 70% of literature records are found within a 25 km radius of a college or university versus 60% of manager records. The proportion of records found within this radius varies between species. *T. ramosissima* represents the low end of records near higher education institutions (25% literature and 16% manager records) whereas *L. maackii* represents the high end (94% literature and 83% manager records) (Figure S2). Literature records are consistently closer to a college or university than manager records.

A majority of manager records are on public land (67%), whereas only half of literature records are on public land (50%). The proportions of points on public land varied between species. *F. japonica* is least commonly recorded on public land (38% literature and 37% manager records) whereas *T. ramosissima* is mostly recorded on public land (72% literature and 89% manager records) (Table S2). Only two species, *F. japonica* and *L. salicaria*, had a slightly higher proportion of literature records on public land compared to manager records (<2% difference between datasets). All other species had a higher proportion of manager records on public land than literature records.

Finally, the presence of a species listing does not relate to increased reporting in the literature or by managers. An average of 34% of literature and 30% of manager records are found in states where that species is listed as a noxious weed. *L. maackii* is listed in four states (2% land area in the lower 48) and a low proportion of records are in states with a listing (0.5% literature and 8% manager records). On the other hand, *L. salicaria* is listed in 34 states (70.2% land area in the lower 48) and most of the records are found in states with a listing (90% literature and 92% manager records) (Table S2). Species

that are listed in a larger number of states and over a larger area have more records in areas with a listing.

#### 1.4 Discussion

Geographic biases in invasion science are common (Pyšek et al. 2008, Hulme et al. 2013, Lowry et al. 2013, Bellard and Jeschke 2016) and have the potential to skew our understanding of invasive plant impacts, efficacy of management, and native community susceptibility if these studies focus on a portion of the invaded range. Our results suggest that there is often a significant geographic bias in one or more study type, though it is more common for scientific studies to match manager records. The distribution of scientific studies is not as extensive as the distribution of manager records, but the geographies are similar (Figure 1). Overall, we do not find strong evidence of consistent geographic or climatic biases in scientific studies (Figure 1, Figure 3).

##### 1.4.1 Distribution of records

With an order of magnitude more occurrence records, manager records typically described a larger invaded range than scientific studies (Table 1, Figure 1). Two exceptions to this trend were *B. tectorum* and *P. australis*. In both cases, the broader geography is due to genetic studies seeking to understand the introduction, spread, or hybridization of these species (Meyerson et al. 2016, Arnesen et al. 2017). However, genetic studies likely include locations where the species are naturalized, but not necessarily invasive (spreading or having impact). For example, *B. tectorum* is recorded throughout the lower 48, but is most problematic in western states (Knapp 1996, Bradley et al. 2018), which aligns with manager records. Because EDDMapS and iMap records tend to focus on areas of high priority for monitoring and management, it is likely that manager records provide an effective description of the invaded range.

Although the invaded range is encompassed by scientific studies in general, different study types are not as evenly distributed or pursued. Between each species, the relative proportion of each study type is variable, indicating that the research priorities for these species are not the same. These uneven prioritizations of study types affect the total number of study locations for each species, as certain study types, like genetic studies (recorded under invasive trait), retrieve data from more sites than others, like impact studies (Figure 2B). As a whole, management and mapping studies, which relate to ways in which stakeholders can track and respond to plant invasions, consistently represent a low proportion of the total number of studies (Figure 2A). Considering their more applied nature, questions related to management and mapping might be more extensively addressed in grey literature instead of scientific literature. The variable research priorities between species suggests that scientists are not consistently gaining ecological or evolutionary knowledge on these plants. It also produces an uneven distribution of studies because they do not all collect data from the species' entire invaded or naturalized range.

On average, literature records tend to occur in warmer environments than manager records, although, for most species, the temperature differences between datasets are small ( $< 1^{\circ}\text{C}$ ). With climate change, temperatures are likely to increase in the near term (Allen et al. 2018). When ecological studies varied climatically from manager records, they tended to occur towards the warm range margin (Figure 3, Table S1). This focus on the warm range margin suggests that ecological studies could provide an effective illustration of the future ecology of invasions as temperatures warm. However, a focus on the warm range margin could also suggest that invasions are of greatest concern or highest impact in these areas. In this case, climate warming could make plant invasions worse than anticipated (Bradley et al. 2010) as more of the invaded range becomes

climatically similar to the more problematic warm range margin. For example, experimental studies have shown that management with herbicide can be less effective in warmer climates due to herbicide dilution following increased growth, and increased herbicide volatilization (Bailey 2004, Matzrafi et al. 2016, Ziska 2016). It is unclear why researchers have focused on the warm margin of these species' range, but it implies that we may know how these plants will behave and interact with their environment in a warming climate.

In contrast, differences in mean annual precipitation had up to 180 mm difference between literature and manager records but showed inconsistent directionality towards wetter or drier climates. Similarly, for all but one species, at least one study type occurred on a climate range margin, compared to manager records, but there is no consistent trend towards warmer/colder or wetter/drier climates across species (Figure 3). This suggests that the focus on a more limited climate range is driven by other factors, such as impact, land use, or access, which varies between species. Two study types that never differed significantly in climatic space for any species were impact and management studies. The similarity between studies on management and studies on impact suggests that they focus on areas with the largest impacts that are also the most important to control. Impact studies were also most likely to be found at a climatic range margin (Figure 3), which also suggests that these studies tend to focus on areas where invasions are more pronounced, and the impacts are highest. As a result, reported invasive plant impacts may not apply to the entire invaded range. The varied biases with respect to precipitation and inconsistent biases by study type suggest that invasion science does not always encompass the climate of the invaded range, leading to higher uncertainty in ecological forecasting.

#### 1.4.2 Disturbances and other biases

Invasive species are known to preferentially colonize disturbed areas, especially roadsides, which are both a habitat and a dispersal corridor for invasive plants (Christen and Matlack 2009, Menuz and Kettenring 2013). However, a major question in invasion ecology is whether invasive species are drivers of ecological impacts or passengers taking advantage of disturbance, but not the main drivers of impact (MacDougall and Turkington 2005). If invasion science tends to occur in more disturbed areas such as adjacent to roads, it could suggest that reported impacts of invasive plants are inflated. Our results show that scientific studies across all species are clearly skewed away from roads relative to manager records (Figure 4, Figure S1). While managers report invasions near roads, scientists focus on less disturbed areas, further from roads.

Scientific studies are, however, biased towards proximity to colleges and universities. This finding is not surprising given a desire for easy access to field sites and is consistent with past research on biases in herbarium records showing high proportions of specimen from locations near the herbarium itself (Daru et al. 2018). Nonetheless, a bias towards universities can lead to larger-scale biases because higher learning institutions are not evenly distributed throughout the country. This may be particularly problematic for species located in the less dense western U.S., such as *B. tectorum* and *T. ramosissima*. While our results do not suggest that this bias affects the geography of ecological research on invasive plants, the bias towards a more populated and university-dense eastern U.S. may contribute to the overrepresentation of these ten plants specifically in invasion literature.

The large proportion of manager records on public lands in comparison to researchers, is likely due to their focus on public land management. Nonetheless, public land is overrepresented in both datasets, it only represents 7.8% of the total land area in the

lower 48 (Jenkins et al. 2015) but 67% manager records and 50% literature records. Public lands tend to have more natural areas and be accessible to federal researchers, so this finding is consistent with larger trends in ecology that focus on natural areas (Martin et al. 2012). Cities are disturbed landscapes that are experiencing faster warming than their neighboring environments, they have also been found to have different plant succession than surrounding environments (George et al. 2009). If research is mostly focused on natural environments, we may not have an accurate portrait of invasion processes that affect most of the land area in the country. This bias could also affect predictive models, which often use manager records for calibration, by overrepresenting these natural sites.

Finally, the inclusion of a plant on a noxious weed list did not seem to impact the reporting of that plant. These lists are mostly made to regulate the sale and distribution of these plants and do not often include plants that are found in unmanaged areas (Quinn et al. 2013). It is therefore unsurprising that they do not affect where these plants are reported or studied.

#### 1.4.3 Data limitations

The results from these two datasets highlight differences between where studies occur and where invasive plants are found, however, they are both imperfect records. Considering the volume of literature records analyzed, it was not possible to confirm site locations with more precision than was given in an article. This means that some sites could be up to 11 km (~0.1 decimal degrees) away from their recorded locations, this uncertainty is even greater for studies that present locations as broad, regional scale maps. Nonetheless, 69% of study locations were recorded with greater than 0.1 decimal degree precision, so these errors are a small portion of the overall dataset (Table S3).

We also find consistent trends with regards to distance to roads and to colleges and

universities, the two most sensitive variables to these imprecisions, which suggests that our data reasonably capture where studies occur. Not all states record the presence or distribution of invasive species in accessible databases, which can create blind spots in manager datasets. It is also possible that certain counties and states have specific priorities with regards to invasive plants that are not reflected in any legislation but affect reports and create overreporting in certain areas. For example, *A. petiolata* has been reported throughout Wisconsin in EDDMapS, but is unreported in neighboring states. Ultimately this work reflects where researchers and managers report the presence of these plants, even if it diverges from where the plants are actually found.

### **1.5 Conclusion**

On a global scale, North America is overrepresented in invasion literature (Pyšek et al. 2008, Hulme et al. 2013, Lowry et al. 2013), whereas, at a finer scale, land uses such as roadsides and sites near research institutions are favored (Graham et al. 2004, Daru et al. 2018). Environmental variables such as climate affect invasive plant impacts, management techniques and native community vulnerability (Bailey 2004, Hou et al. 2014, Matzrafi et al. 2016, Ziska 2016). Landscape level biases in the literature could therefore have important consequences for our understanding and management of invasions. The distribution of studies reveals that research encompasses these plants' invaded ranges, but different study types tend to occur in a subset of that range. This produces an uneven distribution of knowledge on these plants that may be linked to the invasion intensity. These biases towards a narrower climate range are compounded with general biases associated with different human features, like roads, colleges and universities, and public lands. Land use biases contribute to the distribution patterns found across the lower 48 because these features are not evenly distributed across the territory. The uneven geography of invasive plant research, either with regards to



climate or land use, implies that our understanding of plant invasions is limited, even for well-studied plants. In turn, these limitations can under- or over-inflate the threat posed by these plants by misrepresenting their local invasion potential, impact, or management feasibility.

APPENDIX A

SUPPLEMENTAL TABLES

Table 3: Mean annual temperature and precipitation of literature and manager records.  
\*p < 0.05 \*\* p<0.01

Species	Mean annual temperature (°C)			Mean annual precipitation (mm)		
	Literature records	Manager records	Difference	Literature records	Manager records	Difference
<i>A. altissima</i>	11.0	12.2	1.2**	1102.4	1122.0	19.6
<i>A. petiolata</i>	10.0	9.5	-0.6**	1062.1	1091.8	29.7
<i>B. tectorum</i>	9.5	9.1	-0.4**	465.2	398.9	-66.3**
<i>F. japonica</i>	9.5	9.3	-0.3	1342.6	1162.6	-180.0**
<i>L. maackii</i>	11.9	11.9	0.0	1064.3	1123.6	59.3**
<i>L. salicaria</i>	9.6	8.1	-1.5**	960.3	921.1	-39.1**
<i>M. vimineum</i>	12.8	12.2	-0.6**	1206.8	1222.9	16.2**
<i>P. arundinacea</i>	9.2	6.9	-2.3**	897.2	939.3	42.0*
<i>P. australis</i>	12.1	9.7	-2.4**	897.3	947.6	50.3**
<i>T. ramosissima</i>	11.4	11.9	0.5	371.2	344.2	-27.0

Table 4: Proportion of records within 100 m of a road, on public land, and in a state with a listing for that species.

Species	Literature records within 100 m of a road (%)	Manager records within 100 m of a road (%)	Literature records on public land (%)	Manager records on public land (%)	Literature records in state with listing (%)	Manager records in state with listing (%)	States with listing (n)
<i>A. altissima</i>	44.8	53.4	46.0	69.1	4.6	1.8	4
<i>A. petiolata</i>	46.7	54.4	47.5	64.4	25.3	40.5	9
<i>B. tectorum</i>	42.3	50.8	66.9	82.9	37.4	26.3	6
<i>F. japonica</i>	71.5	74.4	38.1	36.6	73.8	77.2	10
<i>L. maackii</i>	39.1	54.2	49.1	82.0	0.5	7.9	4
<i>L. salicaria</i>	46.2	64.9	48.7	48.7	89.5	91.9	34
<i>M. vimineum</i>	54.6	44.8	31.4	84.5	11.7	12.3	4
<i>P. arundinacea</i>	45.4	65.0	65.5	83.8	3.5	4.2	3
<i>P. australis</i>	44.1	55.5	47.1	60.3	16.4	11.9	7
<i>T. ramosissima</i>	13.3	22.3	72.2	88.7	80.6	26.1	11

Table 5: Precision of locations reported in the literature for each species in the lower 48.

	Tenths		Hundredths		Thousandths		Map estimate		Total (n)
	(n)	(%)	(n)	(%)	(n)	(%)	(n)	(%)	
<i>A. altissima</i>	9	10.3	1	1.1	48	55.2	29	33.3	87
<i>A. petiolata</i>	50	19.2	39	14.9	129	49.4	43	16.5	261
<i>B. tectorum</i>	70	9.3	9	1.2	404	53.9	266	35.5	749
<i>F. japonica</i>	13	8.1	11	6.9	116	72.5	20	12.5	160
<i>L. maackii</i>	26	12.4	6	2.9	131	62.4	47	22.4	210
<i>L. salicaria</i>	10	3.6	53	19.3	134	48.7	78	28.4	275
<i>M. vimineum</i>	30	3.4	7	0.8	784	89.2	58	6.6	879
<i>P. arundinacea</i>	47	15.0	38	12.1	171	54.6	57	18.2	313
<i>P. australis</i>	60	5.4	245	21.9	539	48.2	275	24.6	1119
<i>T. ramosissima</i>	13	7.2	12	6.7	24	13.3	131	72.8	180
TOTAL	328	7.7	421	9.9	2480	58.6	1004	23.7	4233

APPENDIX B

SUPLMENTAL FIGURES




Figure 5: Proportion of records found within a given distance to a road.




Figure 6: Proportion of records found within a given distance to a college or university.

## BIBLIOGRAPHY

- Adams CR, Galatowitsch SM (2006) Increasing the Effectiveness of Reed canary grass (*Phalaris arundinacea* L.) Control in Wet Meadow Restorations. *Restoration Ecology* 14: 441–451. <https://doi.org/10.1111/j.1526-100X.2006.00152.x>
- Allen MR, Dube OP, Solecki W, Aragón-Durand F, Cramer W, Humphreys S, Kainuma M, Kala J, Mahowald N, Mulugetta Y (2018) Framing and context. In: Masson-Delmotte V, P. Zhai, H.-O. Pörtner, Roberts D, Skea J, Shukla PR, Pirani A, Moufouma-Okia W, Péan C, Pidcock R, Connors S, Matthews JBR, Chen Y, Zhou X, Gomis MI, Lonnoy E, Maycock T, Tignor M, Waterfield T (Eds), *Global Warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change.*, 49–91.
- Arnesen S, Coleman CE, Meyer SE (2017) Population genetic structure of *Bromus tectorum* in the mountains of western North America. *American Journal of Botany* 104: 879–890. <https://doi.org/10.3732/ajb.1700038>
- Bailey SW (2004) Climate change and decreasing herbicide persistence. *Pest Management Science* 60: 158–162. <https://doi.org/10.1002/ps.785>
- Barger CT, Moorhead DJ (2007) EDDMapS—early detection and distribution mapping system for the southeast exotic pest plant council. *Wildland weeds* 10: 4–8.
- Beaury EM.\*, Fusco EJ.\*, Allen JA, Bradley BA ( Reactive and inconsistent state policies do little to prevent plant invasions.
- Bellard C, Jeschke JM (2016) A spatial mismatch between invader impacts and research publications. *Conservation Biology* 30: 230–232. <https://doi.org/10.1111/cobi.12611>
- Bhattarai GP, Cronin JT (2014) Hurricane Activity and the Large-Scale Pattern of Spread of an Invasive Plant Species. Adam P (Ed). *PLoS ONE* 9: e98478. <https://doi.org/10.1371/journal.pone.0098478>
- Boakes EH, McGowan PJK, Fuller RA, Chang-Qing D, Clark NE, O'Connor K, Mace GM (2010) Distorted views of biodiversity: Spatial and temporal bias in species occurrence data. *PLoS Biology* 8: e1000385. <https://doi.org/10.1371/journal.pbio.1000385>
- Bradley BA, Wilcove DS, Oppenheimer M (2010) Climate change increases risk of plant invasion in the Eastern United States. *Biological Invasions* 12: 1855–1872. <https://doi.org/10.1007/s10530-009-9597-y>
- Bradley BA, Curtis CA, Fusco EJ, Abatzoglou JT, Balch JK, Dadashi S, Tuanmu M-N (2018) Cheatgrass (*Bromus tectorum*) distribution in the intermountain Western United States and its relationship to fire frequency, seasonality, and ignitions. *Biological Invasions* 20: 1493–1506. <https://doi.org/10.1007/s10530-017-1641-8>

- Buerger A, Howe K, Jacquart E, Chandler M, Culley T, Evans C, Kearns K, Schutzki R, Riper L Van (2016) Risk Assessments for Invasive Plants: A Midwestern U.S. Comparison. *Invasive Plant Science and Management* 9: 41–54. <https://doi.org/DOI: 10.1614/IPSM-D-15-00018.1>
- Christen DC, Matlack GR (2009) The habitat and conduit functions of roads in the spread of three invasive plant species. *Biological Invasions* 11: 453–465. <https://doi.org/10.1007/s10530-008-9262-x>
- Colautti RI, Barrett SCH (2013) Rapid adaptation to climate facilitates range expansion of an invasive plant. *Science* 342: 364–366. <https://doi.org/10.1126/science.1242121>
- Collins A, Hart EM, Molofsky J (2010) Differential response to frequency-dependent interactions: an experimental test using genotypes of an invasive grass. *Oecologia* 164: 959–969. <https://doi.org/10.1007/s00442-010-1719-9>
- Craine EB, Evankow A, Wolfson KB, Dalton K, Swedlund H, Bowen C, Heschel MS (2016) Physiological Response of *Tamarix ramosissima* (Tamaricaceae) to a Biological Control Agent. *Western North American Naturalist* 76: 339–351. <https://doi.org/10.3398/064.076.0310>
- Daru BH, Park DS, Primack RB, Willis CG, Barrington DS, Whitfeld TJS, Seidler TG, Sweeney PW, Foster DR, Ellison AM, Davis CC (2018) Widespread sampling biases in herbaria revealed from large-scale digitization. *New Phytologist* 217: 939–955. <https://doi.org/10.1111/nph.14855>
- George K, Ziska LH, Bunce JA, Quebedeaux B, Hom JL, Wolf J, Teasdale JR (2009) Macroclimate associated with urbanization increases the rate of secondary succession from fallow soil. *Oecologia* 159: 637–647.
- Gómez-Aparicio L, Canham CD (2008) Neighbourhood analyses of the allelopathic effects of the invasive tree *Ailanthus altissima* in temperate forests. *Journal of Ecology* 96: 447–458. <https://doi.org/10.1111/j.1365-2745.2007.01352.x>
- Graham CH, Ferrier S, Huettman F, Moritz C, Peterson AT (2004) New developments in museum-based informatics and applications in biodiversity analysis. *Trends in Ecology & Evolution* 19: 497–503. <https://doi.org/https://doi.org/10.1016/j.tree.2004.07.006>
- Hager HA (2004) Differential Effects of *Typha* Litter and Plants on Invasive *Lythrum Salicaria* Seedling Survival and Growth. *Biological Invasions* 6: 433–444. <https://doi.org/10.1023/B:BINV.0000041558.22300.3a>
- Hellmann JJ, Byers JE, Bierwagen BG, Dukes JS (2008) Five Potential Consequences of Climate Change for Invasive Species. *Conservation Biology* 22: 534–543. <https://doi.org/10.1111/j.1523-1739.2008.00951.x>
- Hou Q-Q, Chen B-M, Peng S-L, Chen L-Y (2014) Effects of extreme temperature on seedling establishment of nonnative invasive plants. *Biological Invasions* 16: 2049–2061. <https://doi.org/10.1007/s10530-014-0647-8>
- Hovick SM, Carson WP (2015) Tailoring biocontrol to maximize top-down effects: on the importance of underlying site fertility. *Ecological Applications* 25: 125–139. <https://doi.org/10.1890/13-2050.1>

- Hulme PE, Pyšek P, Jarošík V, Pergl J, Schaffner U, Vilà M (2013) Bias and error in understanding plant invasion impacts. *Trends in Ecology and Evolution* 28: 212–218. <https://doi.org/10.1016/j.tree.2012.10.010>
- ITIS (2020) Integrated Taxonomic Information System (ITIS). : Accessed 07/01/2020. Available from: <http://www.itis.gov>.
- Jarnevich CS, Holcombe TR, Bella EM, Carlson ML, Graziano G, Lamb M, Seefeldt SS, Morissette J (2014) Cross-Scale Assessment of Potential Habitat Shifts in a Rapidly Changing Climate. *Invasive Plant Science and Management* 7: 491–502. <https://doi.org/DOI: 10.1614/IPSM-D-13-00071.1>
- Jenkins CN, Van Houtan KS, Pimm SL, Sexton JO (2015) US protected lands mismatch biodiversity priorities. *Proceedings of the National Academy of Sciences* 112: 5081 LP – 5086. <https://doi.org/10.1073/pnas.1418034112>
- Jeschke JM, Aparicio LG, Haider S, Heger T, Lortie CJ, Pyšek P, Strayer DL (2012) Taxonomic bias and lack of cross-taxonomic studies in invasion biology. *Frontiers in Ecology and the Environment* 10: 349–350. <https://doi.org/10.1890/12.WB.016>
- Jodoin Y, Lavoie C, Villeneuve P, Theriault M, Beaulieu J, Belzile F (2008) Highways as corridors and habitats for the invasive common reed *Phragmites australis* in Quebec, Canada. *Journal of Applied Ecology* 45: 459–466. <https://doi.org/10.1111/j.1365-2664.2007.01362.x>
- Kadmon R, Farber O, Danin A (2004) Effect of Roadside Bias on the Accuracy of Predictive Maps Produced by Bioclimatic Models. *Ecological Applications* 14: 401–413. <https://doi.org/10.1890/02-5364>
- Kaproth MA, McGraw JB (2008) Seed Viability and Dispersal of the Wind-Dispersed Invasive *Ailanthus altissima* in Aqueous Environments. *Forest Science* 54: 490–496. <https://doi.org/10.1093/forestscience/54.5.490>
- Knapp PA (1996) Cheatgrass (*Bromus tectorum* L) dominance in the Great Basin Desert: History, persistence, and influences to human activities. *Global Environmental Change* 6: 37–52. [https://doi.org/https://doi.org/10.1016/0959-3780\(95\)00112-3](https://doi.org/https://doi.org/10.1016/0959-3780(95)00112-3)
- Laginhas BB, Bradley BA ( Global Invaders (version 1.0): a compendium of invasive plant species documented by the peer-reviewed literature.
- Lavoie C, Dufresne C, Delisle F (2005) The spread of reed canarygrass (*Phalaris arundinacea*) in Québec: A spatio-temporal perspective. *Écoscience* 12: 366–375. <https://doi.org/10.2980/i1195-6860-12-3-366.1>
- Lowry E, Rollinson EJ, Laybourn AJ, Scott TE, Aiello-Lammens ME, Gray SM, Mickley J, Gurevitch J (2013) Biological invasions: a field synopsis, systematic review, and database of the literature. *Ecology and Evolution* 3: 182–196. <https://doi.org/10.1002/ece3.431>
- MacDougall AS, Turkington R (2005) Are Invasive Species the Drivers or Passengers of Change in Degraded Ecosystems? *Ecology* 86: 42–55. <https://doi.org/10.1890/04-0669>
- Martin LJ, Blossey B, Ellis E (2012) Mapping where ecologists work: biases in the



- global distribution of terrestrial ecological observations. *Frontiers in Ecology and the Environment* 10: 195–201. <https://doi.org/10.1890/110154>
- Martinez AE (2017) Sensitivity of Modeled Channel Hydraulic Variables to Invasive and Native Riparian Vegetation. *International Journal of Applied Geospatial Research (IJAGR)* 8: 47–61.
- Matzrafi M, Seiwert B, Reemtsma T, Rubin B, Peleg Z (2016) Climate change increases the risk of herbicide-resistant weeds due to enhanced detoxification. *Planta* 244: 1217–1227. <https://doi.org/10.1007/s00425-016-2577-4>
- McCaughey TL, Stephenson GR (2000) Time from flowering to seed viability in purple loosestrife (*Lythrum salicaria*). *Aquatic Botany* 66: 57–68. [https://doi.org/10.1016/S0304-3770\(99\)00018-2](https://doi.org/10.1016/S0304-3770(99)00018-2)
- McGlynn CA (2009) Native and invasive plant interactions in wetlands and the minimal role of invasiveness. *Biological Invasions* 11: 1929–1939. <https://doi.org/10.1007/s10530-008-9370-7>
- Menuz DR, Kettenring KM (2013) The importance of roads, nutrients, and climate for invasive plant establishment in riparian areas in the northwestern United States. *Biological Invasions* 15: 1601–1612. <https://doi.org/10.1007/s10530-012-0395-6>
- Meyerson LA, Cronin JT, Bhattarai GP, Brix H, Lambertini C, Lučanová M, Rinehart S, Suda J, Pyšek P (2016) Do ploidy level and nuclear genome size and latitude of origin modify the expression of *Phragmites australis* traits and interactions with herbivores? *Biological Invasions* 18: 2531–2549. <https://doi.org/10.1007/s10530-016-1200-8>
- Narumalani S, Mishra DR, Wilson R, Reece P, Kohler A (2009) Detecting and Mapping Four Invasive Species along the Floodplain of North Platte River, Nebraska. *Weed Technology* 23: 99–107. <https://doi.org/DOI: 10.1614/WT-08-007.1>
- NatureServe (2019) iMapInvasives: an online data system supporting strategic invasive species management. : Accessed 10/2/2019. Available from: <http://imapinvasives.org>.
- Oak Ridge National Laboratory GIS and TG (2010) Colleges and Universities. : Accessed 01/16/2020. Available from: <https://www.sciencebase.gov/catalog/item/4f4e4acee4b07f02db67fb39>.
- Peter CR, Burdick DM (2010) Can Plant Competition and Diversity Reduce the Growth and Survival of Exotic *Phragmites australis* Invading a Tidal Marsh? *Estuaries and Coasts* 33: 1225–1236. <https://doi.org/10.1007/s12237-010-9328-8>
- PRISM Climate Group (2004) 30-Year Normals. Oregon State University: Accessed 07/01/2020. Available from: <http://prism.oregonstate.edu>.
- Pyšek P, Skálová H, Čuda J, Guo W-Y, Suda J, Doležal J, Kautzál O, Lambertini C, Lučanová M, Mandáková T, Moravcová L, Pyšková K, Brix H, Meyerson LA (2018) Small genome separates native and invasive populations in an ecologically important cosmopolitan grass. *Ecology* 99: 79–90. <https://doi.org/10.1002/ecy.2068>
- Pyšek P, Richardson DM, Jarošík V (2006) Who cites who in the invasion zoo:

- insights from an analysis of the most highly cited papers in invasion ecology. *Preslia* 78: 437–468.
- Pyšek P, Richardson DM, Pergl J, Jarošík V, Sixtová Z, Weber E (2008) Geographical and taxonomic biases in invasion ecology. *Trends in Ecology & Evolution* 23: 237–244. <https://doi.org/https://doi.org/10.1016/j.tree.2008.02.002>
- Quinn LD, Barney JN, McCubbins JSN, Endres AB (2013) Navigating the “Noxious” and “Invasive” Regulatory Landscape: Suggestions for Improved Regulation. *BioScience* 63: 124–131. <https://doi.org/10.1525/bio.2013.63.2.8>
- Rotholz E, Mandelik Y (2013) Roadside habitats: effects on diversity and composition of plant, arthropod, and small mammal communities. *Biodiversity and Conservation* 22: 1017–1031. <https://doi.org/10.1007/s10531-013-0465-9>
- Shearin ZRC, Filipek M, Desai R, Bickford WA, Kowalski KP, Clay K (2018) Fungal endophytes from seeds of invasive, non-native *Phragmites australis* and their potential role in germination and seedling growth. *Plant and Soil* 422: 183–194. <https://doi.org/10.1007/s11104-017-3241-x>
- Shi J, Macel M, Tielbörger K, Verhoeven KJF (2018) Effects of admixture in native and invasive populations of *Lythrum salicaria*. *Biological Invasions* 20: 2381–2393. <https://doi.org/10.1007/s10530-018-1707-2>
- Skinner K, Smith L, Rice P (2000) Using noxious weed lists to prioritize targets for developing weed management strategies. *Weed Science* 48: 640–644. [https://doi.org/10.1614/0043-1745\(2000\)048\[0640:UNWLTP\]2.0.CO;2](https://doi.org/10.1614/0043-1745(2000)048[0640:UNWLTP]2.0.CO;2)
- Stolar J, Nielsen SE (2015) Accounting for spatially biased sampling effort in presence-only species distribution modelling. Franklin J (Ed). *Diversity and Distributions* 21: 595–608. <https://doi.org/10.1111/ddi.12279>
- Stricker KB, Hagan D, Flory SL (2015) Improving methods to evaluate the impacts of plant invasions: lessons from 40 years of research. *AoB PLANTS* 7. <https://doi.org/10.1093/aobpla/plv028>
- Tekiela DR, Barney JN (2017) Not All Roads Lead to Rome: A Meta-analysis of Invasive Plant Impact Methodology. *Invasive Plant Science and Management* 10: 304–312. <https://doi.org/DOI: 10.1017/inp.2017.39>
- U.S. Census Bureau (2016) TIGER/Line Shapefiles. : Accessed 01/2016. Available from: <https://www.census.gov/geographies/mapping-files/time-series/geo/tiger-line-file.html>.
- U.S. EPA (2008) Effects of Climate Change on Aquatic Invasive Species and Implications for Management and Research. Washington, DC
- Uddin MN, Robinson RW (2018) Can nutrient enrichment influence the invasion of *Phragmites australis*? *Science of The Total Environment* 613–614: 1449–1459. <https://doi.org/10.1016/J.SCITOTENV.2017.06.131>
- USGS Gap Analysis Project (2018) Protected Areas Database of the United States (PAD-US). U.S. Geological Survey (USGS): Accessed. Available from: <https://usgs.gov/gapanalysis/PAD-US/>.
- Vilà M, Ibáñez I (2011) Plant invasions in the landscape. *Landscape Ecology* 26:

461–472. <https://doi.org/10.1007/s10980-011-9585-3>

Web of Science (2020) Web of Science. : Accessed 07/09/2019. Available from: [www.webofknowledge.com](http://www.webofknowledge.com).

Wiesenborn WD (2005) Biomass of Arthropod Trophic Levels on *Tamarix ramosissima* (Tamaricaceae) Branches. *Environmental Entomology* 34: 656–663. <https://doi.org/10.1603/0046-225X-34.3.656>

Williams V-RJ, Sahli HF (2016) A Comparison of Herbivore Damage on Three Invasive Plants and Their Native Congeners: Implications for the Enemy Release Hypothesis. *Castanea* 81: 128–137. Available from: <http://www.jstor.org/stable/26353883>.

Ziska LH (2016) The role of climate change and increasing atmospheric carbon dioxide on weed management: Herbicide efficacy. *Agriculture, Ecosystems & Environment* 231: 304–309. <https://doi.org/https://doi.org/10.1016/j.agee.2016.07.014>