

1-1-2018

## Effect of Curcumin Supplementation on Exercise-Induced Oxidative Stress, Inflammation, and Muscle Damage

Steven Allen Basham

Follow this and additional works at: <https://scholarsjunction.msstate.edu/td>

---

### Recommended Citation

Basham, Steven Allen, "Effect of Curcumin Supplementation on Exercise-Induced Oxidative Stress, Inflammation, and Muscle Damage" (2018). *Theses and Dissertations*. 1650.  
<https://scholarsjunction.msstate.edu/td/1650>

This Graduate Thesis - Open Access is brought to you for free and open access by the Theses and Dissertations at Scholars Junction. It has been accepted for inclusion in Theses and Dissertations by an authorized administrator of Scholars Junction. For more information, please contact [scholcomm@msstate.libanswers.com](mailto:scholcomm@msstate.libanswers.com).

Effect of curcumin supplementation on exercise-induced oxidative stress, inflammation,  
and muscle damage

By

Steven Allen Basham

A Master's Thesis  
Submitted to the Faculty of  
Mississippi State University  
in Partial Fulfillment of the Requirements  
for the Degree of Master of Science  
in Kinesiology  
in the Department of Kinesiology

Mississippi State, Mississippi

May 2018

Copyright by  
Steven Allen Basham  
2018

Effect of curcumin supplementation on exercise-induced oxidative stress, inflammation,  
and muscle damage

By

Steven Allen Basham

Approved:

---

Matthew J. McAllister  
(Major Professor)

---

JohnEric William Smith  
(Committee Member)

---

John Lamberth  
(Committee Member)

---

Adam C. Knight  
(Graduate Coordinator)

---

Richard L. Blackburn  
Dean  
College of Education

Name: Steven Allen Basham

Date of Degree: May 3, 2018

Institution: Mississippi State University

Major Field: Kinesiology

Major Professor: Matthew McAllister

Title of Study: Effect of curcumin supplementation on exercise-induced oxidative stress, inflammation, and muscle damage

Pages in Study 66

Candidate for Degree of Master of Science

Oxidative stress (OS) and inflammation can be detrimental to exercise performance. Curcumin has shown to reduce OS, inflammation, muscle damage, and soreness. The purpose of this study was to examine the effect of curcumin on biomarker markers of OS (MDA, TAC), inflammation (TNF- $\alpha$ ), muscle damage (CK) and soreness. Participants performed a muscle damaging protocol, before and after supplementation. Subjects were randomly assigned to curcumin (1.5 g/day) or placebo for 28 days. Blood was sampled pre and immediately post exercise (additional post: 60 min, 24, and 48 h). No significant differences were observed for OS or inflammation. There was a treatment X condition interaction for CK, where CK were significantly lower post supplementation in the curcumin group ( $p < 0.0001$ ). Curcumin resulted in significantly lower muscle soreness compared to the placebo ( $p = 0.0120$ ). In conclusion, curcumin may reduce muscle damage, and soreness without affecting OS and inflammatory after exercise.

## DEDICATION

This project is dedicated to my family and their unconditional support throughout this entire process. Although this project has been very rewarding, it has been very challenging. Without unwavering love and motivation given from my family, this project would not have been successful.

## ACKNOWLEDGEMENTS

First, I would like to thank my advisor and mentor, Dr. McAllister. Without his guidance and support, I would not have been able to complete this project. Secondly, I would like to acknowledge and thank my committee members, Dr. Smith and Dr. Lamberth for their support and guidance. Lastly, without the help of other graduate students involved, specifically, Hunter Waldman and Ben Krings, this project would not have been as successful.

## TABLE OF CONTENTS

DEDICATION .....	ii
ACKNOWLEDGEMENTS .....	iii
LIST OF TABLES .....	vi
LIST OF FIGURES .....	vii
CHAPTER	
I. INTRODUCTION .....	1
BACKGROUND .....	1
II. LITERATURE REVIEW .....	4
INTRODUCTION .....	4
Defining Oxidative Stress, Reactive Oxygen Species.....	4
Exercise-Induced Oxidative Stress .....	6
Eccentric Muscle Contraction .....	8
Inflammation/Cytokine Production and Muscle Damage/Recovery.....	9
Antioxidants .....	11
Endogenous .....	11
Exogenous .....	13
Curcumin .....	15
Description/Mechanism.....	15
Supporting Evidence .....	17
Summary/Conclusion .....	18
III. MATERIALS AND METHODS.....	19
IRB Approval .....	19
Design and Subjects .....	19
Testing Procedures .....	20
Supplementation .....	21
Eccentric Muscle Damage Protocol .....	21
Instrumentation.....	22
Blood Sampling and Analysis .....	22
Statistical Analysis .....	23



IV.	RESULTS .....	24
	Inflammation .....	24
	Total Antioxidant and Oxidative Stress .....	24
	Muscle Damage and Soreness .....	25
V.	DISCUSSION .....	32
	Summary.....	32
	Total Antioxidant and Oxidative Stress .....	33
	Inflammation .....	34
	Muscle damage and Soreness .....	36
	Limitations.....	37
	Conclusion.....	38
VI.	REFERENCES .....	39

## LIST OF TABLES

1.	Subject Characteristics.....	26
----	------------------------------	----

## LIST OF FIGURES

1.	Timeline of testing procedures.....	26
2.	TNF- $\alpha$ concentrations, overall.....	27
3.	Total Antioxidant Capacity concentrations, overall. ....	28
4.	Malondialdehyde concentrations, overall. ....	29
5.	Creatine Kinase concentrations, overall. ....	30
6.	Post Supplementation Perceived Muscle Soreness.....	31

## CHAPTER I

### INTRODUCTION

#### **Background**

Oxidative stress (OS) and inflammation may be detrimental to exercise performance (Cesari et al., 2004) and health (Huang et al., 2015). OS is the result of an imbalance between reactive oxygen species (ROS), reactive nitrogen species (RNS) and antioxidants (Sies, 1997). Antioxidants serve as a protectant mechanism from the damaging effects of ROS (Kunwar & Priyadarsini, 2011). Excessive ROS production has been shown to impair metabolic function (Costa, Amorim, Quintanilha, & Moradas-Ferreira, 2002; Grant, Quinn, & Dawes, 1999), which may cause further risk for cardiometabolic and cardiovascular diseases (Montezano & Touyz, 2012). Although exercise is a protectant against many diseases (Kullo, Khaleghi, & Hensrud, 2007), it elicits acute OS and inflammation (Park & Kwak, 2016; Tiidus, 1998), with single bouts of exercise shown to increase OS (Koska et al., 2000; Maughan et al., 1989). In addition to aerobic exercise, unaccustomed resistance training (RT) (eccentric muscle actions) also elicits an OS response (Proske & Morgan, 2001), along with muscle damage (Newham, McPhail, Mills, & Edwards, 1983) and muscle soreness (Maughan et al., 1989). Following the onset of muscle damage, an initial inflammatory response initiates for muscle regeneration (Philippou, Maridaki, Theos, & Koutsilieris, 2012). Under normal conditions, the inflammatory process is regulated shortly after by anti-

inflammatory processes (Philippou et al., 2012). However, if chronic inflammation persists, it may hinder exercise performance (Cesari et al., 2004).

Although exercise may reduce OS (Hollander et al., 1999), it may not be sufficient; thus, antioxidant supplementation may aid in reducing OS and inflammation (Kodama et al., 2009). Curcumin, the active ingredient in the spice turmeric, supplementation reduces OS (Kim et al., 2007; Ruby, Kuttan, Dinesh Babu, Rajasekharan, & Kuttan, 1995) and inflammation (Gupta et al., 2011; Kim et al., 2007; Kondamudi, Kovelamudi, Nayak, Rao, & Shenoy, 2015), while enhancing antioxidant capacity (Hamidie, Yamada, Ishizawa, Saito, & Masuda, 2015; Sarker et al., 2015; Takahashi et al., 2014). Curcumin may act on several different cytokine/ROS mediated inflammatory pathways. Curcumin has shown to specifically reduce expression of nuclear factor kappa-light-chain-enhancer of activated  $\beta$  cells (NF- $\kappa$ B)(Kang, Lee, Price, & Kim, 2009)/cyclooxygenase-2 (COX-2) (Kang et al., 2004), while increasing antioxidants via activation of nuclear factor-like 2 (Nrf2) (Shehzad & Lee, 2013). While there is a consensus on the potent anti-inflammatory (Davis et al., 2007; McFarlin et al., 2016) and antioxidant (Kawanishi et al., 2013; Takahashi et al., 2014) effects of curcumin, there are inconsistent reports for reduced muscle damage and soreness (Drobnic et al., 2014; Jäger et al., 2017; McFarlin et al., 2016). Most recent studies have demonstrated reduced creatine kinase concentration (i.e. biomarker of muscle damage) (Jäger et al., 2017; McFarlin et al., 2016) following exercise-induced muscle damage. However, both reported confounding results for inflammation and muscle soreness. McFarlin et al. (2016) reported a reduction in tumor necrosis factor alpha (TNF- $\alpha$ ) up to 25%, whereas, Jäger et al. (2017) demonstrated no changes. In contrast, Jäger et al.

(2017) reported a significant reduction in perceived muscle soreness, while McFarlin et al. (2016) did not. Differences may lie in the assessment technique for subjective measurements, subject training status, and exercise protocol. In an animal model, Davis et al. (2007) utilized a similar protocol of downhill running (Jäger et al., 2017), and reported a reduction in both inflammation and muscle damage using more invasive techniques (plasma vs. tissue sample), thus, suggesting differences may also lie in species/tissue sampling.

Further, rather than acute supplementation (e.g. hours-days) used in previous studies (Drobic et al., 2014; McFarlin et al., 2016; Nicol, Rowlands, Fazakerly, & Kellett, 2015), this study consisted of 28 days of curcumin supplementation. The primary aim of this study was to examine the relationship between a commercially available curcumin (CurcuFresh™) on markers of inflammation, total antioxidants, muscle damage, and muscle soreness. We hypothesized curcumin supplementation would reduce inflammation and increase antioxidant status at rest. Additionally, we hypothesized curcumin supplementation would reduce inflammation, muscle damage, and muscle soreness following exercise-induced muscle damage (EIMD).

## CHAPTER II

### LITERATURE REVIEW

#### **Introduction**

Oxidative Stress (OS) may negatively affect exercise performance and health, thus, controlling OS may be critical for optimizing both. This review will define OS and the different mechanisms that can perpetuate OS. Moreover, this review will specifically examine exercise-induced OS, inflammation, and muscle damage. Aerobic exercise and eccentric actions have been shown to induce OS (Fisher-Wellman & Bloomer, 2009; Powers & Jackson, 2008; Proske & Morgan, 2001). Further, endogenous antioxidant status and exogenous supplementation with regard to exercise will be discussed. Lastly, this review will focus on the effects of curcumin on inflammation and antioxidants regarding exercise-induced responses.

#### **Defining Oxidative Stress, Reactive Oxygen Species**

Although the mitochondria are the energy producing organelle in the cell, they are the source of potentially harmful reactive oxygen species (ROS) and/or reactive nitrogen species (RNS) (Balaban, Nemoto, & Finkel, 2005). Aerobic metabolism is responsible for the greatest production of ROS (Balaban, Newmoto, & Finkel, 2005). The electron transport chain may lead to an incomplete reduction in oxygen which gives rise to the

superoxide anion ( $O_2^-$ ) (Jastroch, Divakaruni, Mookerjee, Treberg, & Brand, 2010). Superoxide is relatively unreactive; however, in the presence of other radicals, such as nitric oxide, may react very quickly to form peroxynitrate, which is a relatively strong oxidizing agent (Halliwell & Gutteridge, 2015). Within the cell membrane of the mitochondria, “Complex I” and “Complex III” is reported to generate the majority of superoxide under resting conditions (Chen, Vazquez, Moghaddas, Hoppel, & Lesnefsky, 2003; Liu, Fiskum, & Schubert, 2002). Further, superoxide may be reduced to a less active molecule, hydrogen peroxide ( $H_2O_2$ ) (Loschen, Azzi, Richter, & Flohe, 1974), where it can readily cross cell-membranes with a relatively longer half-life compared to superoxide. Although hydrogen peroxide is relatively stable, in the presence of iron, it may be converted to a more highly reactive radical, hydroxyl radical ( $\cdot OH$ ), in the Fenton reaction (Powers & Jackson, 2008). Due to the dependency of aerobic metabolism, the generation of ROS may be inevitable. However, the production of ROS may be a trigger for endogenous antioxidants (Brieger, Schiavone, Miller, & Krause, 2012) and improved mitochondrial redox environment (Hollander et al., 1999; Lander, 1997). The imbalance among antioxidants and the generation of oxidants or ROS has been termed OS (Sies, 1985, 1986, 1991). OS signifies a disruption/challenged redox environment most notably observed through molecular changes to redox sensitive molecules and increased production of ROS (Powers & Jackson, 2008). There is an unwavering consensus that OS is the underlying pathological cause for impaired metabolic function (Costa et al., 2002; Grant et al., 1999), thus, leading to increased risk for cardiovascular disease (Huang et al., 2015).



## Exercise-Induced Oxidative Stress

Most research has confirmed as mitochondria the main site for ROS production (Davies, Maguire, Brooks, Dallman, & Packer, 1982), however, some researchers suggest that it is only at rest when the mitochondria is the main culprit, and there is less ROS generation in the mitochondria during exercise in contrast to previous theories (Kozlov et al., 2005). Other, more prominent sites of ROS production during exercise, may be to blame, which include the sarcoplasmic reticulum, transverse tubules, plasma membranes and cytoplasmic spaces within the muscle cell (Powers & Jackson, 2008). In these areas, other ROS generating enzymes such as NADPH oxidases (Ushio-Fukai, 2006) and xanthine oxidases (McNally et al., 2003) have been identified as viable sources of ROS production. ROS production from these enzymes have been shown to alter the function of calcium pumps, and contractile structures in the muscle fiber, thus, possibility leading to ROS-induced muscle fatigue (Powers & Jackson, 2008). Nonetheless, exercise can increase ROS generation resulting in an OS response (Park & Kwak, 2016); however, several exercise training elements dictate the extent of ROS production and damage.

The extent of oxidative damage is related to both the duration (Bloomer, Davis, Consitt, & Wideman, 2007) and the intensity (Goto et al., 2003) of exercise. Most literature investigating acute aerobic exercise consisting of longer durations have examined elite runners following marathons and triathlons (McAnulty et al., 2005; Nieman et al., 2004; Radák et al., 2003), as well as other modalities such as cycling (McAnulty et al., 2007). OS is prevalent via increased lipid peroxidation measurements such as thiobarbituric acid reactive substances (TBARS) which is typically measured as malondialdehyde (MDA) equivalents (Fisher-Wellman & Bloomer, 2009). Other modes

of exercise, such as short-term anaerobic exercise has also shown to increase ROS/RONS resulting in OS (Groussard et al., 2003). Additionally, other anaerobic exercise modalities such as RT elicits similar responses of increased ROS/OS as seen from aerobic exercise (Fisher-Wellman & Bloomer, 2009). Studies that have examined similar outcomes in increased biomarkers of lipid peroxidation have used full-body RT paradigms with multiple sets (McBride, Kraemer, Triplett-McBride, & Sebastianelli, 1998; Ramel, Wagner, & Elmadfa, 2004; Vina et al., 2000; Vincent, Morgan, & Vincent, 2004), while some have demonstrated elevated markers of OS only utilizing one exercise (Bloomer, Falvo, Schilling, & Smith, 2007; Subudhi, Davis, Kipp, & Askew, 2001; Volek et al., 2002). In contrast, Bloomer et al. (2005) investigated differences in blood markers of OS in ten cross-trained men during 30-minutes of cycling at 70%  $\dot{V}O_{2max}$  and dumbbell squats at 70% of 1-RM (repetition maximum). When compared to cycling, blood markers of OS were significantly higher in the RT protocol post exercise (Bloomer, Goldfarb, Wideman, McKenzie, & Consitt, 2005). These results suggest there may be differences between aerobic and anaerobic exercises (i.e. resistance exercise) and their respective roles for perpetuating OS. A later study by the same authors demonstrated training status may modulate the OS response to exercise (Bloomer et al., 2006). In this study (Bloomer et al., 2006), 12 anaerobically trained males performed six, 10-s sprints, on a separate occasion participants performed barbell squats matching the same volume (total work [kJ]). There were no significant changes in MDA following either exercise. The authors note, repeated exposure to OS may potentiate protective mechanisms, resulting in greater resistance to OS. Given the association between OS and RT, exact mechanisms responsible for excessive ROS during RT should be identified.

## **Eccentric Muscle Contraction**

During dynamic exercise, muscles undergo concentric and eccentric actions (D. B. Hollander et al., 2015). Morgan and Allen (1999), and Warren, Ingalls, Lowe, and Armstrong (2001) identified two major components that contribute to muscle damage, damaged sarcomeres and an impaired “excitation-contraction coupling” system. Ultimately, eccentric muscle actions elicit the most significant muscle damage (Newham et al., 1983; Proske & Morgan, 2001) and are linked to muscle soreness (Maughan et al., 1989; Radak, Pucsok, Mecseki, Csont, & Ferdinandy, 1999). A universally accepted method to promote muscle damage is through downhill running, allowing for controlled lengthening, while stimulating damage (Davis et al., 2007; Drobic et al., 2014; Powers & Jackson, 2008). Maughan et al. (1989) showed increases in MDA after downhill-running, while also reporting evidence indicative of muscle damage via increased creatine kinase (CK) levels. Another study observed increased CK levels of approximately 150 fold, compared to resting levels after a one-legged eccentric exercise protocol (Hellsten, Frandsen, Orthenblad, Sjødin, & Richter, 1997). Other studies have used other methods including timed eccentric contractions during dynamic resistance exercises resulting in similar muscle damage (Nicol et al., 2015). However, other studies reported no change in MDA after eccentric muscle contractions (Saxton, Donnelly, & Roper, 1994). Following exercise-induced muscle damage, an inflammatory response results lead to more ROS production (Azlina, Kamisah, Chua, Ibrahim, & Qodriyah, 2015).

## **Inflammation/Cytokine Production and Muscle Damage/Recovery**

Muscle damage involves an inflammatory response which is initiated to modulate the repair process or degradation of muscle tissue before the final repair stages (Philippou et al., 2012). ROS generation may be responsible for the initial inflammatory response in damaged muscle tissue (Tiidus, 1998). Specific cells such as neutrophils, macrophages, and lymphocytes engorge the damaged tissue (Pedersen & Hoffman-Goetz, 2000; Smith, Kruger, Smith, & Myburgh, 2008), which secrete cytokines that modulate repair of damaged muscle tissue (Pedersen, Ostrowski, Rohde, & Bruunsgaard, 1998a; Tidball & Villalta, 2010). The main cytokine/ROS mediated inflammatory cascades consist of the COX-2 and nuclear factor kappa-light-chain-enhancer of activated  $\beta$  cells (Nf-kB) (Peterson, Bakkar, & Guttridge, 2011). The COX-2 pathway is responsible for the production of prostaglandins, which are responsible for the regeneration of damage muscle tissue and muscle soreness (Bondesen, Mills, Kegley, & Pavlath, 2004). Stimulation of cytokines are found within the damaged muscle tissue, in addition to other tissues and circulation, and can promote localized inflammation (Pedersen & Hoffman-Goetz, 2000; Pedersen, Ostrowski, Rohde, & Bruunsgaard, 1998b). The responses to damaged muscle tissue consists of break-down, repair, and regeneration of muscle tissues (Charge & Rudnicki, 2004; Pedersen et al., 1998a; Smith et al., 2008), whereas muscle repair after damage can be explained declination, inflammation, and growth (Mourkioti & Rosenthal, 2005; Prisk & Huard, 2003; Tidball, 2005; Tidball & Villalta, 2010). However, if inflammatory response is prolonged, additional damage will occur, disrupting muscle growth (Philippou et al., 2012).

One major cytokine involved in the initial inflammatory response is tumor necrosis factor alpha (TNF- $\alpha$ ) (Figarella-Branger, Civatte, Bartoli, & Pellissier, 2003; Peake, Nosaka, & Suzuki, 2005). Increased secretion of this pro-inflammatory cytokine perpetuates muscle damage (Philippou et al., 2012; Tidball & Villalta, 2010) which, may inhibit performance (Cassatella, 1995). Hardin et al. (2008) reported decreased contractile force in mice diaphragms after TNF- $\alpha$  administrations, which supports the relationship between cytokines and performance. Further, TNF- $\alpha$  activates the transcriptional factor Nf-kB which initiates the inflammatory cascade, ultimately increasing adhesion molecules in the endothelial lining of vessels; thus TNF- $\alpha$  is strongly associated with the progression and initiation of atherogenesis/atherosclerosis (Lastra, Manrique, & Hayden, 2006). Unlike the pro-inflammatory cytokine TNF- $\alpha$ , interleukin 6 (IL-6), also known as a myokine due to direct secretion from muscle fibers (Pedersen, Akerstrom, Nielsen, & Fischer, 2007), is an anti-inflammatory cytokine (Philippou et al., 2012). IL-6 plays pivotal roles in mobilizing energy stores during exercise (Febbraio & Pedersen, 2002) while also regulating several neurotransmitters and other signaling cascades (Gleeson, 2000). The reported increased production of IL-6 during exercise increases the synthesis of other anti-inflammatory cytokines, which can inhibit TNF- $\alpha$  production (Petersen & Pedersen, 2005). Although IL-6 is classified as anti-inflammatory (Tilg, Dinarello, & Mier, 1997), if chronically elevated, it can act as a pro-inflammatory cytokine (Pedersen et al., 2007; Pedersen & Hoffman-Goetz, 2000). IL-6 is related to exercised-induced muscle damage (Bruunsgaard et al., 1997; Hellsten et al., 1997; Pedersen et al., 1998a; Steensberg et al., 2000). Hellsten et al. (1997) and Bruunsgaard et al. (1997) both reported a substantial increase in IL-6, 90 min and 120

min after eccentric protocols, respectively. Many studies have also demonstrated a relationship between increased IL-6 and strenuous aerobic exercise (Castell et al., 1996; Nieman et al., 2003; Ostrowski et al., 1998; Ostrowski, Rohde, Asp, Schjerling, & Pedersen, 1999; Sprenger et al., 1992; Ullum et al., 1994). Pedersen and Hoffman-Goetz (2000) and Pedersen et al. (2007) further discuss the possible relationship between increased IL-6 secretion and the type of exercise modality, intensity, and duration. Ultimately, increased levels of inflammatory cytokines are associated with deferments in performance (Cesari et al., 2004; Taaffe, Harris, Ferrucci, Rowe, & Seeman, 2000; Visser et al., 2002). If inflammation persists, it may inhibit muscle regeneration and possibly reduce exercise performance.

## **Antioxidants**

### **Endogenous**

Improved endogenous antioxidant status may be a result from exercise training (Wiecek et al., 2016). Although some studies have demonstrated increased endogenous antioxidants with RT (Çakir-Atabek, Demir, PinarbaSili, & Gündüz, 2010; Ramel et al., 2004), research is scarce, and the relationship between RT and endogenous antioxidants remains unclear. Although excessive ROS production can be detrimental to health (Ji, Gomez-Cabrera, & Vina, 2006) and performance (Donges, Duffield, & Drinkwater, 2010), acute ROS production may improve the mitochondrial redox environment (Dada et al., 2003; Hollander et al., 1999). Cellular processes depend heavily on the mitochondria aerobic capacity (Jastroch et al., 2010). In response to ROS production, endogenous antioxidants such as superoxide dismutase (SOD), catalase (CAT), and

glutathione peroxidase (GPX) are up-regulated (Brieger et al., 2012). These antioxidants help scavenge ROS and reduce oxidative strain. First discovered in 1969, SOD is the first line of protection against the anion superoxide (McCord & Fridovich, 1969), which catalyzes the reaction to create hydrogen peroxide.

However, chronic aerobic training is shown to increase endogenous antioxidants (Brites et al., 1999; Mena et al., 1991; Ortenblad, Madsen, & Djurhuus, 1997). Mena et al. (1991) reported an increase in SOD in professional cyclist after a 2800-kilometer ride in 20 days. Brites et al. (1999) and Ortenblad et al. (1997) both observed increased SOD activity in soccer and volleyball players, respectively. Similarly, Miyazaki et al. (2001) reported increased SOD levels in 9 untrained males after a 12-week endurance program consisting of running on a treadmill at 80% maximal heart rate. Increased antioxidants such as SOD are strongly related to decreased OS (Fielding & Meydani, 1997). Increased fitness levels appear to positively influence OS during exercise (Kocabas et al., 2016). This is due to the activation of peroxisome proliferator activated receptor gamma coactivator 1 alpha (PGC1- $\alpha$ ), a major metabolic adaptation to exercise, which increases endogenous antioxidants through binding to its' downstream targets (St-Pierre et al., 2006). Other possible mechanisms such as direct activation of Nrf2 improves the mitochondria redox environment (Cantó & Auwerx, 2009; Kobayashi & Yamamoto, 2005). Although these studies reported increased antioxidants with trained individuals, literature is inconsistent with these findings. One study reported no changes in SOD regarding untrained individuals after an 8 week cycling program (Tiidus, Pushkarenko, & Houston, 1996). Likewise, Tauler, Gimeno, Aguilo, Guix, and Pons (1999) reported no increases in SOD, but did find increased CAT activity in aerobically trained athletes. It is

pertinent to mention that observable changes in SOD concentrations differ among exercise intensity and duration (Powers et al., 1994), which may give rise to possible explanations for confounding results.

Once superoxide is catalyzed to hydrogen peroxide, secondary enzymes, GPX and CAT, may reduce the radical to water. Similar to SOD, many studies have identified increased concentrations of GPX contingent upon both exercise intensity and duration (Powers et al., 1994). In one study Powers et al. (1994) examined significant increases GPX activity and expression following high-intensity/long-duration exercises when compared to low-intensity/short-duration exercises. Further, it was reported that elevated concentration/activity of GPX was increased in Type I/IIa muscle fibers. In the same study, there were no observable differences found for CAT concentration/activity, suggesting CAT activity and expression is independent of changes seen in GPX and SOD. Along with exercise, dietary interventions consisting of increased fruit and vegetable consumption are attributed to increase endogenous antioxidant status (Teixeira, Mill, Pereira, & Molina Mdel, 2016). Exercise as well as dietary interventions are well established factors that can enhance endogenous antioxidants.

### **Exogenous**

In conjunction with exercise, supplemental or exogenous antioxidants may be used to decrease OS and/or improve antioxidant status. Some supplemental antioxidants that have demonstrated reduced OS include (but are not limited to): quercetin (Duarte et al., 2002; Duarte et al., 2001; Garcia-Saura et al., 2005), resveratrol (Ates et al., 2007; Rege et al., 2013; Venturini et al., 2010), and coenzyme Q10 (CoQ10) (Pala et al., 2016; Zhang et al., 2013). Pala et al., (2016) reported anti-inflammatory potential through



supplementation with CoQ10. Other dietary supplements such as fish oil or flax seed oil are shown to lower cytokines (IL-6) and upregulate endogenous antioxidants such as SOD, GSH, and CAT (Jangale, Devarshi, Bansode, Kulkarni, & Harsulkar, 2016). However, these studies reported decreased markers of OS and inflammatory cytokines in animals. One study examined quercetin supplementation in humans and found no evidence of decreased OS, but demonstrated decreased blood pressure, which may reduce the risk of cardiovascular disease (Edwards et al., 2007). Edwards et al., (2007) hypothesized possible metabolic differences between species, which may result in contraindicating findings in human models. Further, Edwards et al., (2007) suggested that quercetin may have a localized antioxidant effect that was undetectable.

In addition, two of the most popular antioxidants are vitamin C and E, where they work collectively to provide protection against lipid peroxidation (Knez, Coombes, & Jenkins, 2006). Multiple studies have reported decreased biomarkers of exercise-induced OS following vitamin C (Alessio, Goldfarb, & Cao, 1997; Goldfarb, Patrick, Bryer, & You, 2005; Sastre et al., 1992) and vitamin E supplementation (Kawai et al., 2000; Satoshi, Kiyoji, Hiroyo, & Fumio, 1989). Despite advantageous findings supporting vitamin C and vitamin E supplementation, some studies have indicated null findings (Bryant, Ryder, Martino, Kim, & Craig, 2003; Gaeini, Rahnama, & Hamedinia, 2006). Bryant et al. (2003) reported possible differences in vitamin C and vitamin E supplementation following independent dosage phases, where vitamin E supplementation was superior for reducing MDA concentrations after 60 min and 30 min cycling tests. In contrast, one study demonstrated vitamin E supplementation was insufficient for reducing muscle damage, protein, and lipid oxidation (Gaeini et al., 2006). Moreover, both

antioxidants in high dosages may be associated with adverse metabolic adaptations to exercise (Gomez-Cabrera et al., 2008; Picklo & Thyfault, 2015). Paulsen et al. (2014) examined vitamin C and vitamin E ingestion in 54 young men and women. This study utilized a double-blind, randomized, controlled trial, where 1000 mg of vitamin C and 235 mg of vitamin E or placebo were randomly assigned during 11 weeks of an endurance training program. Paulsen and colleagues reported decreased metabolic adaptations via decreased PGC- $\alpha$  activation, which is known to stimulate mitochondrial biogenesis. However, the treatment group exhibited a greater  $\dot{V}O_{2max}$  than the control group (Paulsen et al., 2014). Conversely, previous literature indicated no adverse effect on exercise-induced metabolic adaptations through high doses of vitamin C (Wadley & McConell, 2010). In all, there are mixed findings that both support and dismiss beneficial reductions for exercise-induced OS along with enhanced exercise performance following exogenous antioxidants. Future research should investigate these antioxidant effects in humans due to the unclear effects and other dependent variables such as training status and sex.

## Curcumin

### Description/Mechanism

Curcumin, the active ingredient of the spice turmeric, is found within the *Curcuma Longa* plant (Hamidie et al., 2015). Curcumin is a lipophilic polyphenol, suggesting it can penetrate all cell membranes (i.e. blood brain barrier, cell wall, and cell membrane) (Goel, Kunnumakkara, & Aggarwal, 2008). Evidence suggests curcumin may enhance the redox environment, and induce mitochondrial biogenesis (Hamidie et al., 2015), along with inhibiting inflammatory cytokines (Gupta et al., 2011; Kim et al., 2007;

Kondamudi et al., 2015). Several mechanisms are thought to explain curcumin's effect of inflammation and OS. One mechanism is indirect inhibition of Nf-kB (Singh & Aggarwal, 1995), which is responsible for the production of many inflammatory cytokine such as TNF- $\alpha$  and IL-6 in muscle. Curcumin may also decrease other inflammatory cascades such as COX-2 (Chun et al., 2003). Curcumin, may up-regulate endogenous antioxidants through direct activation of the transcriptional factor, Nrf2, which then activates and binds to the antioxidant response element (ARE) (Shehzad & Lee, 2013). Additionally, the chemical structures of polyphenols, such as curucmin, provide a direct mechanism for scavenging ROS. The disassociation of the hydroxyl group may potentially decrease the scavenging of superoxide radicals through donation of electrons in vitro (Singh, Barik, Singh, & Priyadarsini, 2011). Further, due to the chelation properties, curucmin may inhibit the Fenton reaction where hydrogen peroxide is oxidized to a more unstable radical, hydroxyl radical, in the presence of iron (Perron & Brumaghim, 2009). The decreased ROS production may also lead to indirect inhibition of Nf-kB (Biswas, McClure, Jimenez, Megson, & Rahman, 2005) in the aforementioned mechanisms. Many studies have examined the effects of curcumin on the modulation of inflammation and ROS production in animals, with promising results regarding human trials. In one study, Hamidie et al. (2015) reported increased expression of AMP-activated kinase (AMPK) from curcumin supplementation in rats. AMPK in turns, phosphorylates and primes PGC-1 $\alpha$  for deacetylation via sirtuin 1 (SIRT1). The increased activity of PGC-1 $\alpha$  results in greater levels of mitochondria biogenesis. These adaptations improve the redox environment within the cell and may lead to reduced oxidative strain.

## Supporting Evidence

Along with curcumin's ability to enhance mitochondrial biogenesis in conjunction with endurance training, curcumin may improve recovery, thus, potentially improving exercise performance indirectly (Davis et al., 2007). Davis et al. (2007) reported reductions in markers of muscle damage with down-hill running (predominately eccentric muscle contractions). Furthermore, curcumin blunted the inflammatory cytokines, TNF- $\alpha$  and IL-6. Another study demonstrated reductions in blood markers of inflammation such as C-reactive proteins (CRP) through curcumin supplementation (Sarker et al., 2015). Sarker et al. (2015) also reported an increase in GSH and improvements in the redox environment. Earlier studies support improved antioxidant status (preventing reduction of SOD and CAT) and reduction of inflammatory cytokines in diabetic rats with curcumin supplementation (Gupta et al., 2011). Likewise, another study reported decreases in CRP and increased nitric oxide with chronic inflammation during curcumin ingestion (Banerjee, Tripathi, Srivastava, Puri, & Shukla, 2003). However, Banerjee et al., (2003) found no change in the inflammatory cytokine TNF- $\alpha$ . Given the association of improved antioxidant capacity and reduced OS, curcumin may attenuate factors accompanied with cardiovascular disease. Most importantly, the aforementioned studies investigated these effects of curcumin supplementation in mice. Literature that investigates curcumin supplementation in human subjects is limited and inconsistent. Although a few studies reported slightly lower levels of CK, they demonstrated no significant decreases in markers of OS, inflammation, and endogenous antioxidants (Drobnic et al., 2014; Nicol et al., 2015; Sciberras et al., 2015). Some limitations to the above studies are inadequate dosage (i.e. reduced bioavailability) (Nicol et al., 2015) and absence of an eccentric

muscle damage protocol (Sciberras et al., 2015). In contrast, other studies showed significant decreases in OS, muscle damage, and inflammation, as well as significant increases in antioxidants (McFarlin et al., 2016; Soni & Kuttan, 1992; Takahashi et al., 2014; Usharani, Mateen, Naidu, Raju, & Chandra, 2008). Most recently, McFarlin et al. (2016) indicated minimal supplementation of curcumin (i.e., 400 mg/day for 5 days) effectively lowered levels of TNF- $\alpha$  and CK following an eccentric muscle damage protocol. As little as 90 mg of curcumin two hours before and after exercise, is shown to effectively reduce OS and increase antioxidants (Takahashi et al., 2014). Nonetheless, curcumin may lower OS, inflammation, and increase antioxidant status.

### **Summary/Conclusion**

Although OS is essential to induce antioxidant and mitochondrial adaptations, excessive and/or chronic OS is detrimental to exercise performance and health; moreover, elevated ROS and inflammatory cytokines may adversely affect recovery and performance. Despite increases in endogenous antioxidants due to exercise, the demand for dietary or supplemental alterations including exogenous antioxidants may be inevitable. Specifically, McBride and Kraemer (1999) suggested supplemental antioxidants may be beneficial at the start of a new training program. Ultimately, curcumin may provide benefits, which include, but not limited to, increased exercise performance, enhanced recovery post exercise, and overall better health. More research is necessary to further solidify curcumin supplementation and its effect on oxidative stress, inflammation, and muscle damage.

## CHAPTER III

### MATERIALS AND METHODS

#### **IRB Approval**

This investigation was approved by the Institute of Biosafety Committee (IBC) at Mississippi State University, as well as the Institutional Review Board (IRB) at Mississippi State University. All participants reported to the J.A. Chromiak Applied Research Laboratory in the Department of Kinesiology at Mississippi State University between 5-9 a.m. for testing. Subjects were recruited via word of mouth, as well as email. Recruitment flyers were also posted throughout the Department of Kinesiology at Mississippi State University.

#### **Design and Subjects**

The study design was a randomized, double-blinded, placebo-controlled trial to compare curcumin supplementation and a placebo on markers of OS, inflammation, antioxidant capacity, muscle damage, and muscle soreness. Twenty apparently healthy non-smoking males aged 18-39 years were selected for this study, with 19 subjects completing all phases. One subject was dropped due to compliance. Subject characteristics can be found in Table 1. Participants were randomly assigned to either the treatment or placebo group. Subjects must be participating in light or moderate exercise activity ( $\geq 30$  min of moderate intensity of physical activity at least three days/week for at least three months or  $\geq 150$  minutes) (American College of Sports, 2000). Exclusion

criterion included: regular ingestion of any anti-inflammatory drugs/foods (NSAIDS) (at least 3 of 7 days) (McFarlin et al., 2016), pre-existing medical diagnoses of cardio metabolic disorders (e.g., hypertension, diabetes), musculoskeletal disorders (e.g. arthritis, osteoporosis), known allergy to curcumin, taking blood thinners (e.g., coumadin), and any medical condition which could hinder completion of the exercise protocol (e.g., lower body extremity injury). Participants underwent an 8 hour fast, and refrained from any alcohol or caffeine consumption 48 h before each session. Subjects were also instructed not to perform any exercise 72 h before testing and remain inactive until the last blood analysis was taken, 48 h post exercise. A priori, sample size was chosen based on previous studies that analyzed similar blood markers (McFarlin et al., 2016; Nicol et al., 2015; Soni & Kuttan, 1992) and G\*Power (Faul, Erdfelder, Lang, & Buchner, 2007). An effect size of  $f^2 = 0.25$  and an  $\alpha$  error probability of 0.05 were set for a power of 0.95.

### **Testing Procedures**

Participants completed two testing sessions along with follow up blood draws (BD) (Figure 1). During the initial visit, the participants provided written informed consent and completed a health history report. Additionally, the general testing procedures were explained. During each testing session, anthropometric measures were taken before the exercise protocol. An initial BD for baseline measurements was taken immediately before the exercise protocol. Participants performed an exercise-induced muscle damaging (EIMD) protocol before and after 28 days of supplementation. Blood was sampled pre exercise, immediately post exercise, as well as 60 min, 24 h and 48 h

post exercise. Upon each BD, perceived muscle soreness was taken using a visual analog scale (VAS).

### **Supplementation**

Participants were randomly assigned to three, 500 mg capsules of curcumin once a day for a total of 1.5 g of curcumin/69 mg of curcuminoids or a cellulose placebo capsule daily for 28 days. This study implemented a newly-enhanced absorption and pharmacokinetics of fresh turmeric derived from curcuminoids in comparison with the standard curcumin from dried rhizomers (Krishnakumar, Kumar, Ninan, Kuttan, & Maliakel, 2015). Subjects were instructed to consume two capsules of either the placebo or curcumin upon waking in the morning with breakfast and one capsule in the afternoon with dinner. To ensure compliance, subjects were reminded weekly to ingest the suggested amount of their provided supplement. Further, each subject was required to maintain a 90% supplement usage rate, which was calculated using the following equation:

$$\text{Amount of Capsules (End of 28 days)} \div \text{Total Amount of Capsules} \times 100$$

### **Eccentric Muscle Damage Protocol**

Participants completed 15 minutes of continuous sitting with one leg (i.e., while the opposite leg is used for balance) at a bench height of 16.5 inches. A metronome was set at a pace of 30 beats/min, and the participants were instructed to follow a cadence of 15 sit-stand repetitions/minute for a total of 225 repetitions. The participants were allowed to stand with both legs. The lower phase of the sitting protocol consists primarily of eccentric muscle actions. Previous literature has utilized various modalities to initiate



muscle damage via eccentric muscle actions (Davis et al., 2007; McFarlin et al., 2016; Nicol et al., 2015). A single session of eccentric muscle actions can elicit OS and muscle damage (Mair et al., 1995). The participant's body weight was used for resistance making the load relative for each participant. The exercise leg was randomized for both pre and post supplementation.

### **Instrumentation**

An aerobic step bench (Step Fitness & Recreation Inc, Marietta, GA) was used for the EIMD protocol. An appropriate bench height was selected based on each participant's height. A heightronic digital stadiometer (Quick Medical 235-D Digital Stadiometer; Issaquah, WA) and digital scale (DEFENDERT 5000; Parsippany, NJ) was used to assess anthropometric measures for height and weight. A VAS was also used to measure muscle soreness and weakness (Boonstra, Preuper, Reneman, Posthumus, & Stewart, 2008). The VAS consisted of a horizontal line, 20 cm in length, with endpoints of "no soreness" to "unbearable soreness".

### **Blood Sampling and Analysis**

A 14 mL sample of blood was collected at each BD. On the two exercise testing sessions, an intravenous catheter was inserted into the antecubital vein, where an initial 3 mL of blood was drawn into a syringe and discarded before the sample blood volume was transferred into the appropriate vacutainer. On non-testing days, venous blood samples were harvested from the antecubital veins for each sample using a butterfly needle, a while subjects were in a supine position. Subsequently, blood was transferred into non-coated vacutainers for serum and sodium heparin vacutainers for plasma and centrifuged

for 15 minutes at 2500 rpm at 4°C. Plasma samples were stored at -80° C. Blood samples were drawn: pre exercise, post exercise, 60 min, 24 h, and 48 h post exercise for both conditions (i.e., before and after supplementation) for a total of 10 BDs. Blood samples were used to analyze markers of OS (thiobarbituric acid reactive substances, (TBARS), muscle damage (creatin kinase, (CK)), inflammation (tumor necrosis factor alpha, (TNF- $\alpha$ )), and antioxidant capacity (total antioxidant capacity, (TAC)) expressed in units of Trolox equivalents. All blood analysis except for CK concentrations, which was analyzed using a spectrophotometer (Pointe Scientific; Canton, MI), was analyzed in duplicates and absorbance was measured using an iMark Bio-Rad Microplate Absorbance Reader (Life Science Research, Hercules, CA).

Plasma was used to measure TBARS, expressed in units of malondialdehyde (MDA) and TAC using a commercial available assay kit (Cayman Chemical, Ann Arbor, MI). Serum was used to analyze TNF- $\alpha$  via end-point method (R & D systems, Minneapolis, MN). Additionally, CK concentrations were measured via kinetic assay (Pointe Scientific; Canton, MI).

### **Statistical Analysis**

Data are reported as mean  $\pm$  SD. All statistical procedures were conducted using SAS version 9.4 (SAS Institute, Cary, NC). A 2 X 2 X 5 RMANOVA (Treatment X Condition X Time) was used to measure the changes in TBARS, CK, TNF- $\alpha$ , and TAC. A 2 X 2 X 4 RMANOVA was used to measure perceived muscle soreness between time points and treatment. A Fisher's least significance different (LSD) post-hoc analysis was used in the instances of significant main effect ( $p < 0.05$ ).

## CHAPTER IV

### RESULTS

#### **Inflammation**

Mean TNF- $\alpha$  levels are shown in Figure 2. There were no significant interactions (treatment X condition [F = 0.00, p = 0.975]; time X condition [F = 0.58, p = 0.675]; or treatment X time X condition [F = 0.36, p = 0.938]) regarding plasma TNF- $\alpha$  concentrations. However, there was an overall significant main effect for time (F = 3.89, p < 0.005) at 24 h post exercise where plasma levels were significantly lower than all other time points.

#### **Total Antioxidant and Oxidative Stress**

As for TAC plasma concentrations, there were no interactions (treatment X condition [F = 0.92, p = 0.339]; time X condition [F = 0.81, p = 0.52]; treatment X time X condition [F = 0.40, p = 0.919]) or main effects for treatment (F = 1.92, p = 0.1679), time (F = 1.43, p = 0.228), or condition (F = 1.33, p = 0.252) (Figure 3).

Moreover, for plasma MDA concentrations, there were no interactions (treatment X condition [F = 1.73, p = 0.191]; time X condition [F = 0.96, p = 0.431]; treatment X time X condition [F = 1.08, p = 0.384]). Additionally, there were no significant main effects for treatment (F = 1.29, p = 0.278), time (F = 0.04, p = 0.834), or condition (F = 1.22, p = 0.271) (Figure 4).

## Muscle Damage and Soreness

Mean CK plasma levels can be found in Figure 5. There were no interactions for treatment X time X condition ( $F = 0.54$ ,  $p = 0.826$ ) or time X condition ( $F = 0.923$ ,  $p = 0.23$ ). However, there was a treatment X condition interaction ( $F = 4.69$ ,  $p = 0.032$ ) for CK plasma concentration. CK plasma concentrations for both the curcumin (211.75 U/L) and placebo (234.32 U/L) group were not significantly different before supplementation ( $p = 0.318$ ). In contrast, CK concentration were significantly lower, overall, post supplementation in the curcumin group (199.62 U/L) compared to the placebo group (287.03 U/L) ( $p < 0.0001$ ). In addition, CK concentrations significantly increased in the placebo pre (234.31 U/L) to post (287.03 U/L) exercise ( $p = 0.0181$ ). There was a significant main effect for treatment ( $F = 13.09$ ,  $p = 0.0004$ ) where CK levels were significantly lower in the curcumin group (205.69 U/L) compared to the placebo group (260.67 U/L), overall.

Lastly, post supplementation VAS muscle soreness means are demonstrated in Figure 6. There were no interactions for: treatment X condition ( $F = 0.89$ ,  $p = 0.514$ ), location X condition ( $F = 1.25$ ,  $p = 0.271$ ), treatment X condition ( $F = 2.26$ ,  $p = 0.133$ ), time X condition ( $F = 1.75$ ,  $p = 0.137$ ); or treatment X time X condition ( $F = 1.44$ ,  $p = 0.178$ ). However, there was a time X location interaction ( $F = 4.20$ ,  $p < 0.0001$ ). Additionally, there was a significant main effect for treatment ( $F = 6.34$ ,  $p = 0.012$ ). Curcumin resulted in significantly lower soreness (2.88) compared to the placebo (3.36) ( $p = 0.0120$ ), overall. Further, there was a significant main effect for both condition ( $F = 20.85$ ,  $p < .0001$ ) and location ( $F = 48.95$ ,  $p < .0001$ ), where soreness was significantly

lower post supplementation (2.68) compared to pre supplementation (3.56) ( $p < .0001$ ), and the exercises leg was reported more sore compared to the non-exercised trained leg.

Table 1

Subject Characteristics

Descriptive Data	Mean $\pm$ SD
Age (years)	21.7 $\pm$ 2.9
Height (cm)	177.7 $\pm$ 7.4
Mass (kg)	83.7 $\pm$ 12.4




Figure 1. Timeline of testing procedures.

BD = blood draw; EIMD = Exercise Induced Muscle Damage




Figure 2. TNF- $\alpha$  concentrations, overall.

Concentrations reported as Mean  $\pm$  SE

\*Indicates significant difference in TNF- $\alpha$  concentrations at 24 hrs post exercise compared to all other time points ( $p < 0.05$ ). CUR = Curcumin; PLA = Placebo; Pre = Before exercise; Post = Immediately After exercise; min = minute; hr = hour.




Figure 3. Total Antioxidant Capacity concentrations, overall.

Concentrations reported as Mean  $\pm$  SE

Pre versus Post supplementation. CUR = Curcumin; PLA = Placebo; PRE = Before Supplementation; Post = After Supplementation.




Figure 4. Malondialdehyde concentrations, overall.

Concentrations reported as Mean  $\pm$  SE

Pre versus Post supplementation. CUR = Curcumin; PLA = Placebo; PRE = before supplementation; Post = after supplementation.






Figure 5. Creatine Kinase concentrations, overall.

Concentrations reported as Mean  $\pm$  SE

Pre versus Post supplementation. \*Indicates significant difference between treatments ( $p < 0.0001$ ); †Indicates significant difference from pre to post supplementation ( $p = 0.0181$ ). CUR = Curcumin; PLA = Placebo; PRE = Before Supplementation; Post = After Supplementation.




Figure 6. Perceived Muscle Soreness, Overall.

Units reported as Mean  $\pm$  SE

\*Indicated significant difference in perceived soreness from pre to post supplementation, overall ( $p < 0.0001$ ). †Indicates significant difference between treatments, overall ( $p = 0.0120$ ). CUR = Curcumin; PLA = Placebo; Pre = Before exercise; Post = Immediately After exercise; min = minute; hr = hour.

## CHAPTER V

### DISCUSSION

#### Summary

This study examined the effect of a newly formulated curcumin (CurcuFresh™) in humans on biomarkers of inflammation, OS, muscle damage, and muscle soreness. Secondly, this is one of only two studies to implement a 28-day supplemental period or greater (Jäger et al., 2017; Oliver et al., 2017), while others have only examined acute ingestion (Davis et al., 2007; Drobnic et al., 2014; Kawanishi et al., 2013; McFarlin et al., 2016; Nicol et al., 2015; Sciberras et al., 2015; Takahashi et al., 2014). The main findings from our study were that curcumin ingestion resulted in significantly lower plasma concentrations of CK and perceived muscle soreness following the eccentric muscle damage protocol, despite any changes in biomarkers of inflammation and OS. Decreased muscle damage and muscle soreness support along with null findings both support and disagree with our initial hypothesis. Further, these finding agree (decreased muscle damage and muscle soreness) with and disagree (unchanged inflammation and OS) with earlier studies examining similar biomarkers (Davis et al., 2007; Drobnic et al., 2014; Kawanishi et al., 2013; McFarlin et al., 2016; Nicol et al., 2015; Sciberras et al., 2015; Takahashi et al., 2014).

## Total Antioxidant and Oxidative Stress

Exercise-induced OS has been associated with fatigue along with muscle damage resulting in decreased performance (Powers & Jackson, 2008). Hypothetically, if OS is reduced along with inflammation, performance may be improved. Therefore, given the strong anti-inflammatory and antioxidant effect of curcumin (Shehzad & Lee, 2013), we sought to examine markers of inflammation (TNF- $\alpha$ ), antioxidant capacity (TAC) and OS (MDA). Previous mechanistic work has shown that curcumin works directly to increase endogenous antioxidants via activation of Nrf2, which subsequently binds to downstream targets and up-regulates antioxidants (Goel et al., 2008; Shehzad & Lee, 2013). Other direct mechanisms have been shown as well in that curcumin may potentiate its' ROS scavenging abilities via dissociation of the OH group (Singh et al., 2011). Given the aforementioned, it is difficult to explain the lack of change in MDA and overall antioxidant status in the present study. Takahashi et al. (2014) recruited participants with similar training statuses as the present study, and instructed subjects to run at 65%  $\dot{V}O_{2max}$  for 60 min. Subsequently, Takahashi et al. (2014) reported increased biological antioxidant potential and reduced glutathione, along with reduced derivatives of reactive oxygen metabolites following curcumin supplementation. From the same study, MDA was unaffected via curcumin treatment, which agrees with our data. One possible reason for these inconsistent findings may be the difference between exercise modality/intensity. Further possible explanations in confounding results concerning antioxidant status may lie in the measurements used in our study, where Takahashi et al. (2014) analyzed specific antioxidant enzymes (GPX, SOD, CAT, and glutathione reductase) (Takahashi et

al., 2014). Lastly, differences have been found in individual antioxidant enzymes (i.e. SOD, CAT) rather than overall antioxidant status following curcumin supplementation in mice in tissue specific analysis (Avci et al., 2012; Kalpana, Sudheer, Rajasekharan, & Menon, 2007), suggesting that curcumins' anti-inflammatory affects may potentially be tissue and/or enzyme specific.

### **Inflammation**

Additionally, curcumin also modulates inflammatory pathways/cascades of both NF-kB (Kang et al., 2009) and COX-2 (Kang et al., 2004), which are both activated following muscle damage. TNF- $\alpha$  was chosen due to its' role as an activator of these pathways, along with its role in the degenerative processes of muscle tissue (Townsend et al., 2015). While one study demonstrated similar findings in TNF- $\alpha$  (Nicol et al., 2015), other studies have reported dissimilar results (Davis et al., 2007; McFarlin et al., 2016). Two studies using similar protocols to elicit muscle damage, Nicol et al. (2015) and McFarlin et al. (2016), recruited non-resistance trained males, but reported differences in TNF- $\alpha$  levels. However, an underlying factor between the two and in the present study, is the disparity of aerobic training status found in our participants (< 30 min of moderate intensity of physical activity at least three days/week for at least three months or  $\leq$  150 minutes) and those in Nicol et al. (2015) (2-4 hours of endurance training/week; 1-2 hours of team training/week), thus potentially explaining the lack of observed differences in inflammation. Further, it is pertinent to note that unlike the present study, Nicol et al. (2015) reported increased IL-6 concentrations immediately following exercise. Although we did not measure IL-6, the rise IL-6 may mediate TNF- $\alpha$  levels following exercise

(Pedersen et al., 2001), which could also explain the significant reduction found in the present study at 24 hours post exercise. In contrast, other potential dissimilarities found in inflammation in the present study may lie in the sample selected, where active individuals were examined in the present study, whereas, untrained participants were recruited by McFarlin et al. (2016). This can be seen when comparing the pre-exercise values of TNF- $\alpha$  concentrations in the present study (1.2 pg/mL) to McFarlin et al. (2016) (4.0 pg/mL). One might speculate that the aerobic training status seen in our subjects provided enough stimulus for adaptations resulting in lower resting levels of TNF- $\alpha$ , thus, negating any potential anti-inflammatory benefits of curcumin supplementation. Another possible reason for discrepancies in inflammatory responses could be the intensity of the protocol. The increased musculature may have exemplified a greater metabolic demand causing greater oxidative damage, thus, leading to greater inflammatory response (McFarlin et al., 2016).

It is pertinent to mention that for proper muscle regeneration, there must be a balance of both an inflammatory and anti-inflammatory environment following muscle damage (Philippou et al., 2012). Previous works have identified TNF- $\alpha$  as a main pro-inflammatory cytokine secreted following muscle damage (Figarella-Branger et al., 2003). However, if concentrations remain elevated for extended periods, TNF- $\alpha$  may inhibit muscle regeneration (Townsend et al., 2015). Independent of curcumin supplementation, our data show a slight (non-significant) rise in plasma TNF- $\alpha$  concentrations, along with a significant decrease 24 h post exercise ( $p = 0.005$ ). Therefore, our findings suggest that curcumin may aid in decreased muscle damage (i.e.

decreased CK) without hindering the natural inflammatory response needed for adequate muscle growth.

### **Muscle damage and Soreness**

Our results are consistent in regards to muscle damage (CK) with previous data in humans (Jäger et al., 2017; McFarlin et al., 2016; Nicol et al., 2015) and animal models (Davis et al., 2007; Huang et al., 2015). Independent of reduced inflammation, curcumin significantly blunted a rise in CK post supplementation (199.62 U/L) compared to the placebo (287.03 U/L) ( $p < 0.0001$ ), overall. When compared to a variety of eccentric muscle damage protocols, our mean peak levels of plasma CK post exercise (211.7 U/L) resembles those reported by Nicol et al. (2015) ( $\approx 200$  IU/L), both of which are substantially less than other studies. These differences could be attributed to substantially less musculature involvement during the exercise modality compared to other studies (Drobic et al., 2014; McFarlin et al., 2016) due to the nature of the uni-lateral, sit/stand protocol which involved only one leg. Our reports of reduced CK concentrations following supplementation in the absence of noticeable changes in inflammation both agree (Jäger et al., 2017; Nicol et al., 2015) and conflict with previous studies (Davis et al., 2007; McFarlin et al., 2016). As mentioned previously, curcumin may act upon multiple inflammatory signal transduction pathways (i.e. Nf-kB, COX-2). However, curcumin may act upon other unidentified pathways to support muscle cell integrity other than these inflammatory/ROS mediated pathways (Nicol et al., 2015).

In respect to VAS muscle soreness, there was a significant reduction in perceived muscle soreness following curcumin supplementation. Additionally, there were no

differences in location of muscle soreness in the lower extremity following the exercise. These findings support our initial hypothesis, while agreeing with several human studies (Drobnic et al., 2014; Jäger et al., 2017; Nicol et al., 2015) and disagreeing with others (McFarlin et al., 2016). Previous works have shown curcumin may reduce prostaglandins via the COX-2 pathway (Kang et al., 2004), which in turn should decrease muscle soreness (Ferreira, 1972). Curcumin may act through other pathways such as inhibition of transient receptor potential ion channels (Yeon et al., 2010), which are responsible for the acute reduction in pain. The possible differences between the present study and that of McFarlin et al. (2016) may lie assessment techniques of muscle soreness. In retrospect, decreased perceived muscle soreness from curcumin supplementation may have substantial effects on mediating greater subsequent exercise performance along with increasing exercise training volume.

### **Limitations**

Some of the limitations are presented in the techniques used for subsequent analysis of biomarkers of antioxidants and OS. Previous research has shown that specific enzymes related to both endogenous antioxidant status and OS are more subject to noticeable changes with antioxidant supplementation (Sarker et al., 2015; Takahashi et al., 2014). Other limitation can be found in the exercise modality chosen. The lack of overall musculature used in our protocol may have resulted in insufficient muscle damage and inflammation. Further, the purpose of this study was to examine the effects of chronic curcumin supplementation. However, the lack of a cross-over design along with



the absence of allocating training status among participants in each group (placebo and treatment), may pose potential limitations.

### **Conclusion**

In conclusion, our data suggests curcumin supplementation of 28 days may result in decreased muscle damage and muscle soreness following exercise. Although there were no differences inflammatory/OS markers, a consensus is still evident in support of curcumin's anti-inflammatory and antioxidant capacities (Kawanishi et al., 2013; McFarlin et al., 2016; Takahashi et al., 2014). Inevitably, curcumin supplementation may be a viable source for enhancing exercise performance in trained individuals. Future research should investigate long-term (> 28 days) curcumin supplementation and its effect on the redox environment and muscle growth as it pertains to aerobic exercise and RT.

## REFERENCES

- Alessio, H. M., Goldfarb, A. H., & Cao, G. (1997). Exercise-induced oxidative stress before and after vitamin C supplementation. *International journal of sport nutrition*, 7(1), 1-9.
- American College of Sports, M. (2000). *ACSM's guidelines for exercise testing and prescription*: Sixth edition. Philadelphia : Lippincott Williams & Wilkins, [2000] ©2000.
- Ates, O., Cayli, S. R., Yucel, N., Altinoz, E., Kocak, A., Durak, M. A., . . . Yologlu, S. (2007). Central nervous system protection by resveratrol in streptozotocin-induced diabetic rats. *J Clin Neurosci*, 14(3), 256-260.  
doi:10.1016/j.jocn.2005.12.010
- Avci, G., Kadioglu, H., Sehirli, A. O., Bozkurt, S., Guclu, O., Arslan, E., & Muratli, S. K. (2012). Curcumin protects against ischemia/reperfusion injury in rat skeletal muscle. *Journal of Surgical Research*, 172(1), e39-e46.
- Azlina, M. F., Kamisah, Y., Chua, K. H., Ibrahim, I. A. A., & Qodriyah, H. M. S. (2015). Preventive Effects of Tocotrienol on Stress-Induced Gastric Mucosal Lesions and Its Relation to Oxidative and Inflammatory Biomarkers. *PLoS ONE*, 10(10), e0139348. doi:10.1371/journal.pone.0139348

Balaban, R. S., Nemoto, S., & Finkel, T. (2005). Mitochondria, oxidants, and aging. *Cell*, 120(4), 483-495. doi:10.1016/j.cell.2005.02.001

Banerjee, M., Tripathi, L., Srivastava, V., Puri, A., & Shukla, R. (2003). Modulation Of Inflammatory Mediators By Ibuprofen And Curcumin Treatment During Chronic Inflammation In Rat#. *Immunopharmacology & Immunotoxicology*, 25(2), 213.

Biswas, S. K., McClure, D., Jimenez, L. A., Megson, I. L., & Rahman, I. (2005). Curcumin induces glutathione biosynthesis and inhibits NF- $\kappa$ B activation and interleukin-8 release in alveolar epithelial cells: mechanism of free radical scavenging activity. *Antioxidants & redox signaling*, 7(1-2), 32-41.

Bloomer, Davis, P. G., Consitt, L., & Wideman, L. (2007). Plasma protein carbonyl response to increasing exercise duration in aerobically trained men and women. *International journal of sports medicine*, 28(01), 21-25.

Bloomer, Falvo, M. J., Fry, A. C., Schilling, B. K., Smith, W. A., & Moore, C. A. (2006). Oxidative stress response in trained men following repeated squats or sprints. *Medicine & Science in Sports & Exercise*, 38(8), 1436-1442.

Bloomer, Falvo, M. J., Schilling, B. K., & Smith, W. A. (2007). Prior exercise and antioxidant supplementation: effect on oxidative stress and muscle injury. *Journal of the International Society of Sports Nutrition*, 4(1), 9.

- Bloomer, Goldfarb, A. H., Wideman, L., McKenzie, M. J., & Consitt, L. A. (2005). Effects of acute aerobic and anaerobic exercise on blood markers of oxidative stress. *J Strength Cond Res*, 19(2), 276-285. doi:10.1519/14823.1
- Bondesen, B. A., Mills, S. T., Kegley, K. M., & Pavlath, G. K. (2004). The COX-2 pathway is essential during early stages of skeletal muscle regeneration. *American Journal of Physiology-Cell Physiology*, 287(2), C475-C483.
- Boonstra, A. M., Preuper, H. R. S., Reneman, M. F., Posthumus, J. B., & Stewart, R. E. (2008). Reliability and validity of the visual analogue scale for disability in patients with chronic musculoskeletal pain. *International Journal of Rehabilitation Research*, 31(2), 165-169.
- Brieger, K., Schiavone, S., Miller, F. J., Jr., & Krause, K. H. (2012). Reactive oxygen species: from health to disease. *Swiss Med Wkly*, 142, w13659. doi:10.4414/smw.2012.13659
- Brites, F. D., Evelson, P. A., Christiansen, M. G., Nicol, M. F., Basilico, M. J., Wikinski, R. W., & Llesuy, S. F. (1999). Soccer players under regular training show oxidative stress but an improved plasma antioxidant status. *Clin Sci (Lond)*, 96(4), 381-385.
- Bruunsgaard, H., Galbo, H., Halkjaer-Kristensen, J., Johansen, T., MacLean, D., & Pedersen, B. (1997). Exercise-induced increase in serum interleukin-6 in humans is related to muscle damage. *The Journal of physiology*, 499(Pt 3), 833.

Bryant, R. J., Ryder, J., Martino, P., Kim, J., & Craig, B. W. (2003). Effects of vitamin E and C supplementation either alone or in combination on exercise-induced lipid peroxidation in trained cyclists. *The Journal of Strength & Conditioning Research*, 17(4), 792-800.

Çakir-Atabek, H., Demir, S., PinarbaSili, R. D., & Gündüz, N. (2010). Effects of different resistance training intensity on indices of oxidative stress. *The Journal of Strength & Conditioning Research*, 24(9), 2491-2497.

Cantó, C., & Auwerx, J. (2009). PGC-1alpha, SIRT1 and AMPK, an energy sensing network that controls energy expenditure. *Curr Opin Lipidol*, 20(2), 98-105.  
doi:10.1097/MOL.0b013e328328d0a4

Cassatella, M. A. (1995). The production of cytokines by polymorphonuclear neutrophils. *Immunology today*, 16(1), 21-26.

Castell, L., Poortmans, J., Leclercq, R., Brasseur, M., Duchateau, J., & Newsholme, E. (1996). Some aspects of the acute phase response after a marathon race, and the effects of glutamine supplementation. *Eur J Appl Physiol Occup Physiol*, 75(1), 47-53.

Cesari, M., Penninx, B. W., Pahor, M., Lauretani, F., Corsi, A. M., Williams, G. R., . . . Ferrucci, L. (2004). Inflammatory markers and physical performance in older persons: the InCHIANTI study. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, 59(3), M242-M248.

- Charge, S. B., & Rudnicki, M. A. (2004). Cellular and molecular regulation of muscle regeneration. *Physiol Rev*, 84(1), 209-238. doi:10.1152/physrev.00019.2003
- Chen, Q., Vazquez, E. J., Moghaddas, S., Hoppel, C. L., & Lesnefsky, E. J. (2003). Production of reactive oxygen species by mitochondria central role of complex III. *Journal of Biological Chemistry*, 278(38), 36027-36031.
- Chun, K.-S., Keum, Y.-S., Han, S. S., Song, Y.-S., Kim, S.-H., & Surh, Y.-J. (2003). Curcumin inhibits phorbol ester-induced expression of cyclooxygenase-2 in mouse skin through suppression of extracellular signal-regulated kinase activity and NF- $\kappa$ B activation. *Carcinogenesis*, 24(9), 1515-1524.
- Costa, V. M., Amorim, M. A., Quintanilha, A., & Moradas-Ferreira, P. (2002). Hydrogen peroxide-induced carbonylation of key metabolic enzymes in *Saccharomyces cerevisiae*: the involvement of the oxidative stress response regulators Yap1 and Skn7. *Free Radical Biology and Medicine*, 33(11), 1507-1515.
- Dada, L. A., Chandel, N. S., Ridge, K. M., Pedemonte, C., Bertorello, A. M., & Sznajder, J. I. (2003). Hypoxia-induced endocytosis of Na,K-ATPase in alveolar epithelial cells is mediated by mitochondrial reactive oxygen species and PKC-zeta. *J Clin Invest*, 111(7), 1057-1064. doi:10.1172/jci16826
- Davies, K., Maguire, J., Brooks, G., Dallman, P., & Packer, L. (1982). Muscle mitochondrial bioenergetics, oxygen supply, and work capacity during dietary iron deficiency and repletion. *American Journal of Physiology-Endocrinology And Metabolism*, 242(6), E418-E427.

Davis, J. M., Murphy, E. A., Carmichael, M. D., Zielinski, M. R., Groschwitz, C. M., Brown, A. S., . . . Mayer, E. P. (2007). Curcumin effects on inflammation and performance recovery following eccentric exercise-induced muscle damage. *Am J Physiol Regul Integr Comp Physiol*, 292(6), R2168-2173.  
doi:10.1152/ajpregu.00858.2006

Donges, C. E., Duffield, R., & Drinkwater, E. J. (2010). Effects of resistance or aerobic exercise training on interleukin-6, C-reactive protein, and body composition. *Medicine and Science in Sports and Exercise*, 42(2), 304-313.

Drobic, F., Riera, J., Appendino, G., Togni, S., Franceschi, F., Valle, X., . . . Tur, J. (2014). Reduction of delayed onset muscle soreness by a novel curcumin delivery system (Meriva®): a randomised, placebo-controlled trial. *Journal of the International Society of Sports Nutrition*, 11(1), 1.

Duarte, Jiménez, R., O'Valle, F., Galisteo, M., Pérez-Palencia, R., Vargas, F., . . . Tamargo, J. (2002). Protective effects of the flavonoid quercetin in chronic nitric oxide deficient rats. *Journal of hypertension*, 20(9), 1843-1854.

Duarte, Perez-Palencia, R., Vargas, F., Ocete, M. A., Perez-Vizcaino, F., Zarzuelo, A., & Tamargo, J. (2001). Antihypertensive effects of the flavonoid quercetin in spontaneously hypertensive rats. *Br J Pharmacol*, 133(1), 117-124.  
doi:10.1038/sj.bjp.0704064

- Edwards, R. L., Lyon, T., Litwin, S. E., Rabovsky, A., Symons, J. D., & Jalili, T. (2007). Quercetin reduces blood pressure in hypertensive subjects. *J Nutr*, *137*(11), 2405-2411.
- Faul, F., Erdfelder, E., Lang, A.-G., & Buchner, A. (2007). G\* Power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behavior research methods*, *39*(2), 175-191.
- Febbraio, M. A., & Pedersen, B. K. (2002). Muscle-derived interleukin-6: mechanisms for activation and possible biological roles. *The FASEB Journal*, *16*(11), 1335-1347.
- Ferreira, S. (1972). Prostaglandins, aspirin-like drugs and analgesia. *Nature*, *240*(102), 200-203.
- Fielding, R., & Meydani, M. (1997). Exercise, free radical generation, and aging. *Aging Clinical and Experimental Research*, *9*(1-2), 12-18.
- Figarella-Branger, D., Civatte, M., Bartoli, C., & Pellissier, J. F. (2003). Cytokines, chemokines, and cell adhesion molecules in inflammatory myopathies. *Muscle & nerve*, *28*(6), 659-682.
- Fisher-Wellman, K., & Bloomer, R. J. (2009). Acute exercise and oxidative stress: a 30 year history. *Dynamic medicine*, *8*(1), 1.



- Gaeini, A., Rahnama, N., & Hamedinia, M. (2006). Effects of vitamin E supplementation on oxidative stress at rest and after exercise to exhaustion in athletic students. *Journal of sports medicine and physical fitness*, 46(3), 458.
- Garcia-Saura, M. F., Galisteo, M., Villar, I. C., Bermejo, A., Zarzuelo, A., Vargas, F., & Duarte, J. (2005). Effects of chronic quercetin treatment in experimental renovascular hypertension. *Mol Cell Biochem*, 270(1-2), 147-155.
- Gleeson, M. (2000). Interleukins and exercise. *The Journal of physiology*, 529(1), 1-1.
- Goel, A., Kunnumakkara, A. B., & Aggarwal, B. B. (2008). Curcumin as “Curecumin”: from kitchen to clinic. *Biochemical pharmacology*, 75(4), 787-809.
- Goldfarb, A. H., Patrick, S. W., Bryer, S., & You, T. (2005). Vitamin C supplementation affects oxidative-stress blood markers in response to a 30-minute run at 75% VO<sub>2</sub>max. *International Journal of Sport Nutrition and Exercise Metabolism*, 15(3), 279-290.
- Gomez-Cabrera, M. C., Domenech, E., Romagnoli, M., Arduini, A., Borrás, C., Pallardo, F. V., . . . Vina, J. (2008). Oral administration of vitamin C decreases muscle mitochondrial biogenesis and hampers training-induced adaptations in endurance performance. *Am J Clin Nutr*, 87(1), 142-149.
- Goto, C., Higashi, Y., Kimura, M., Noma, K., Hara, K., Nakagawa, K., . . . Nara, I. (2003). Effect of different intensities of exercise on endothelium-dependent vasodilation in humans. *Circulation*, 108(5), 530-535.

- Grant, C. M., Quinn, K. A., & Dawes, I. W. (1999). Differential protein S-thiolation of glyceraldehyde-3-phosphate dehydrogenase isoenzymes influences sensitivity to oxidative stress. *Molecular and cellular biology*, *19*(4), 2650-2656.
- Groussard, C., Rannou-Bekono, F., Machefer, G., Chevanne, M., Vincent, S., Sergent, O., . . . Gratas-Delamarche, A. (2003). Changes in blood lipid peroxidation markers and antioxidants after a single sprint anaerobic exercise. *European Journal of Applied Physiology*, *89*(1), 14-20.
- Gupta, S. K., Kumar, B., Nag, T. C., Agrawal, S. S., Agrawal, R., Agrawal, P., . . . Srivastava, S. (2011). Curcumin prevents experimental diabetic retinopathy in rats through its hypoglycemic, antioxidant, and anti-inflammatory mechanisms. *J Ocul Pharmacol Ther*, *27*(2), 123-130. doi:10.1089/jop.2010.0123
- Halliwell, B., & Gutteridge, J. M. (2015). *Free radicals in biology and medicine*: Oxford University Press, USA.
- Hamidie, R. D., Yamada, T., Ishizawa, R., Saito, Y., & Masuda, K. (2015). Curcumin treatment enhances the effect of exercise on mitochondrial biogenesis in skeletal muscle by increasing cAMP levels. *Metabolism*, *64*(10), 1334-1347. doi:10.1016/j.metabol.2015.07.010
- Hardin, B. J., Campbell, K. S., Smith, J. D., Arbogast, S., Smith, J., Moylan, J. S., & Reid, M. B. (2008). TNF- $\alpha$  acts via TNFR1 and muscle-derived oxidants to depress myofibrillar force in murine skeletal muscle. *Journal of applied physiology*, *104*(3), 694-699. doi:10.1152/jappphysiol.00898.2007

- Hellsten, Y., Frandsen, U., Orthenblad, N., Sjødin, B., & Richter, E. A. (1997). Xanthine oxidase in human skeletal muscle following eccentric exercise: a role in inflammation. *The Journal of physiology*, 498(Pt 1), 239.
- Hollander, Fiebig, R., Gore, M., Bejma, J., Ookawara, T., Ohno, H., & Ji, L. L. (1999). Superoxide dismutase gene expression in skeletal muscle: fiber-specific adaptation to endurance training. *Am J Physiol*, 277(3 Pt 2), R856-862.
- Hollander, D. B., Worley, J. R., Asoodeh, M., Wakesa, D., Magnuson, M., Dantzler, D. K., . . . Kraemer, R. R. (2015). A comparison of resistance exercise perceived exertion and muscle activation at varied submaximal durations, loads, and muscle actions. *J Strength Cond Res*. doi:10.1519/jsc.0000000000001290
- Huang, Chiu, W.-C., Chuang, H.-L., Tang, D.-W., Lee, Z.-M., Wei, L., . . . Huang, C.-C. (2015). Effect of curcumin supplementation on physiological fatigue and physical performance in mice. *Nutrients*, 7(2), 905-921.
- Jäger, R., Caldwell, A. R., Sanders, E., Mitchell, J. B., Rogers, J., Purpura, M., & Oliver, J. M. (2017). Curcumin Reduces Muscle Damage and Soreness Following Muscle-Damaging Exercise. *The FASEB Journal*, 31(1 Supplement), lb766-lb766.
- Jangale, N. M., Devarshi, P. P., Bansode, S. B., Kulkarni, M. J., & Harsulkar, A. M. (2016). Dietary flaxseed oil and fish oil ameliorates renal oxidative stress, protein glycation, and inflammation in streptozotocin-nicotinamide-induced diabetic rats. *J Physiol Biochem*, 72(2), 327-336. doi:10.1007/s13105-016-0482-8

- Jastroch, M., Divakaruni, A. S., Mookerjee, S., Treberg, J. R., & Brand, M. D. (2010). Mitochondrial proton and electron leaks. *Essays Biochem*, 47, 53-67.  
doi:10.1042/bse0470053
- Ji, L. L., Gomez-Cabrera, M. C., & Vina, J. (2006). Exercise and hormesis: activation of cellular antioxidant signaling pathway. *Ann N Y Acad Sci*, 1067, 425-435.  
doi:10.1196/annals.1354.061
- Kalpana, C., Sudheer, A., Rajasekharan, K., & Menon, V. (2007). Comparative effects of curcumin and its synthetic analogue on tissue lipid peroxidation and antioxidant status during nicotine-induced toxicity. *Singapore medical journal*, 48(2), 124.
- Kang, Kong, P.-J., Yuh, Y.-J., Lim, S.-Y., Yim, S.-V., Chun, W., & Kim, S.-S. (2004). Curcumin suppresses lipopolysaccharide-induced cyclooxygenase-2 expression by inhibiting activator protein 1 and nuclear factor  $\kappa$ B bindings in BV2 microglial cells. *Journal of pharmacological sciences*, 94(3), 325-328.
- Kang, Lee, S. H., Price, J. E., & Kim, L. S. (2009). Curcumin Suppresses the Paclitaxel-Induced Nuclear Factor- $\kappa$ B in Breast Cancer Cells and Potentiates the Growth Inhibitory Effect of Paclitaxel in a Breast Cancer Nude Mice Model. *The breast journal*, 15(3), 223-229.
- Kawai, Y., Shimomitsu, T., Takanami, Y., MURASE, N., KATSUMURA, T., & MARUYAMA, C. (2000). Vitamin E level changes in serum and red blood cells due to acute exhaustive exercise in collegiate women. *Journal of nutritional science and vitaminology*, 46(3), 119-124.

- Kawanishi, N., Kato, K., Takahashi, M., Mizokami, T., Otsuka, Y., Imaizumi, A., . . . Suzuki, K. (2013). Curcumin attenuates oxidative stress following downhill running-induced muscle damage. *Biochemical and biophysical research communications*, 441(3), 573-578.
- Kim, Y. S., Ahn, Y., Hong, M. H., Joo, S. Y., Kim, K. H., Sohn, I. S., . . . Kim, W. (2007). Curcumin attenuates inflammatory responses of TNF- $\alpha$ -stimulated human endothelial cells. *Journal of cardiovascular pharmacology*, 50(1), 41-49.
- Knez, W. L., Coombes, J. S., & Jenkins, D. G. (2006). Ultra-endurance exercise and oxidative damage. *Sports Medicine*, 36(5), 429-441.
- Kobayashi, M., & Yamamoto, M. (2005). Molecular mechanisms activating the Nrf2-Keap1 pathway of antioxidant gene regulation. *Antioxidants & redox signaling*, 7(3-4), 385-394.
- Kocabas, R., Namiduru, E. S., Bagceci, A. M., Erenler, A. K., Karakoc, O., Orkmez, M., . . . Tarakcioglu, M. (2016). The acute effects of interval exercise on oxidative stress and antioxidant status in volleyball players. *J Sports Med Phys Fitness*.
- Kodama, S., Saito, K., Tanaka, S., Maki, M., Yachi, Y., Asumi, M., . . . Sone, H. (2009). Cardiorespiratory fitness as a quantitative predictor of all-cause mortality and cardiovascular events in healthy men and women: a meta-analysis. *Jama*, 301(19), 2024-2035. doi:10.1001/jama.2009.681

Kondamudi, P. K., Kovelamudi, H., Nayak, P. G., Rao, M. C., & Shenoy, R. R. (2015).

Curcumin half analog modulates interleukin-6 and tumor necrosis factor-alpha in inflammatory bowel disease. *Pharmacogn Mag*, *11*(Suppl 2), S296-302.

doi:10.4103/0973-1296.165991

Koska, J., Blazicek, P., Marko, M., Grna, J. D., Kvetnansky, R., & Vidas, M. (2000).

Insulin, catecholamines, glucose and antioxidant enzymes in oxidative damage during different loads in healthy humans. *Physiol Res*, *49 Suppl 1*, S95-100.

Kozlov, A. V., Szalay, L., Umar, F., Kropik, K., Staniek, K., Niedermüller, H., . . . Nohl,

H. (2005). Skeletal muscles, heart, and lung are the main sources of oxygen radicals in old rats. *Biochimica et Biophysica Acta (BBA)-Molecular Basis of Disease*, *1740*(3), 382-389.

Krishnakumar, I. M., Kumar, D., Ninan, E., Kuttan, R., & Maliakel, B. (2015). Enhanced

absorption and pharmacokinetics of fresh turmeric (*Curcuma Longa* L) derived curcuminoids in comparison with the standard curcumin from dried rhizomes.

*Journal of Functional Foods*, *17*, 55-65.

doi:<http://dx.doi.org/10.1016/j.jff.2015.04.026>

Kullo, I. J., Khaleghi, M., & Hensrud, D. D. (2007). Markers of inflammation are

inversely associated with VO<sub>2</sub> max in asymptomatic men. *J Appl Physiol* (1985), *102*(4), 1374-1379. doi:10.1152/jappphysiol.01028.2006

Kunwar, A., & Priyadarsini, K. (2011). Free radicals, oxidative stress and importance of

antioxidants in human health. *Journal of Medical & Allied Sciences*, *1*(2), 53.

- Lander, H. M. (1997). An essential role for free radicals and derived species in signal transduction. *The FASEB Journal*, 11(2), 118-124.
- Lastra, G., Manrique, C. M., & Hayden, M. R. (2006). The Role of  $\beta$ -Cell Dysfunction in the Cardiometabolic Syndrome. *Journal of the cardiometabolic syndrome*, 1(1), 41-46.
- Liu, Y., Fiskum, G., & Schubert, D. (2002). Generation of reactive oxygen species by the mitochondrial electron transport chain. *J Neurochem*, 80(5), 780-787.
- Loschen, G., Azzi, A., Richter, C., & Flohe, L. (1974). Superoxide radicals as precursors of mitochondrial hydrogen peroxide. *FEBS Lett*, 42(1), 68-72.
- Mair, J., Mayr, M., Mullet, E., Koller, A., Haid, C., Artner-Dworzak, E., . . . Puschendorf, B. (1995). Rapid adaptation to eccentric exercise-induced muscle damage. *International journal of sports medicine*, 16(06), 352-356.
- Maughan, R. J., Donnelly, A. E., Gleeson, M., Whiting, P. H., Walker, K. A., & Clough, P. J. (1989). Delayed-onset muscle damage and lipid peroxidation in man after a downhill run. *Muscle & nerve*, 12(4), 332-336.
- McAnulty, McAnulty, L., Nieman, D., Morrow, J., Dumke, C., & Utter, A. (2007). Carbohydrate effect: hormone and oxidative changes. *International journal of sports medicine*, 28(11), 921-927.

- McAnulty, McAnulty, L. S., Nieman, D. C., Morrow, J. D., Shooter, L. A., Holmes, S., . . . Henson, D. A. (2005). Effect of alpha-tocopherol supplementation on plasma homocysteine and oxidative stress in highly trained athletes before and after exhaustive exercise. *The Journal of nutritional biochemistry*, 16(9), 530-537.
- McBride, J. M., Kraemer, W. J., Triplett-McBride, T., & Sebastianelli, W. (1998). Effect of resistance exercise on free radical production. *Medicine and Science in Sports and Exercise*, 30(1), 67-72.
- McCord, J. M., & Fridovich, I. (1969). Superoxide dismutase an enzymic function for erythrocyte hemocuprein (hemocuprein). *Journal of Biological Chemistry*, 244(22), 6049-6055.
- McFarlin, B. K., Venable, A. S., Henning, A. L., Sampson, J. N. B., Pennel, K., Vingren, J. L., & Hill, D. W. (2016). Reduced inflammatory and muscle damage biomarkers following oral supplementation with bioavailable curcumin. *BBA clinical*, 5, 72-78.
- McNally, J. S., Davis, M. E., Giddens, D. P., Saha, A., Hwang, J., Dikalov, S., . . . Harrison, D. G. (2003). Role of xanthine oxidoreductase and NAD (P) H oxidase in endothelial superoxide production in response to oscillatory shear stress. *American Journal of Physiology-Heart and Circulatory Physiology*, 285(6), H2290-H2297.



- Mena, P., Maynar, M., Gutierrez, J. M., Maynar, J., Timon, J., & Campillo, J. E. (1991). Erythrocyte free radical scavenger enzymes in bicycle professional racers. Adaptation to training. *Int J Sports Med*, 12(6), 563-566. doi:10.1055/s-2007-1024734
- Miyazaki, H., Oh-ishi, S., Ookawara, T., Kizaki, T., Toshinai, K., Ha, S., . . . Ohno, H. (2001). Strenuous endurance training in humans reduces oxidative stress following exhausting exercise. *Eur J Appl Physiol*, 84(1-2), 1-6. doi:10.1007/s004210000342
- Montezano, A. C., & Touyz, R. M. (2012). Reactive oxygen species and endothelial function—role of nitric oxide synthase uncoupling and Nox family nicotinamide adenine dinucleotide phosphate oxidases. *Basic & clinical pharmacology & toxicology*, 110(1), 87-94.
- Morgan, D. L., & Allen, D. G. (1999). Early events in stretch-induced muscle damage. *J Appl Physiol (1985)*, 87(6), 2007-2015.
- Mourkioti, F., & Rosenthal, N. (2005). IGF-1, inflammation and stem cells: interactions during muscle regeneration. *Trends Immunol*, 26(10), 535-542. doi:10.1016/j.it.2005.08.002
- Newham, D. J., McPhail, G., Mills, K. R., & Edwards, R. H. (1983). Ultrastructural changes after concentric and eccentric contractions of human muscle. *J Neurol Sci*, 61(1), 109-122.

- Nicol, L. M., Rowlands, D. S., Fazakerly, R., & Kellett, J. (2015). Curcumin supplementation likely attenuates delayed onset muscle soreness (DOMS). *European Journal of Applied Physiology*, *115*(8), 1769-1777.
- Nieman, D. C., Davis, J. M., Henson, D. A., Walberg-Rankin, J., Shute, M., Dumke, C. L., . . . Brown, A. (2003). Carbohydrate ingestion influences skeletal muscle cytokine mRNA and plasma cytokine levels after a 3-h run. *Journal of applied physiology*, *94*(5), 1917-1925.
- Nieman, D. C., Henson, D. A., Mcanulty, S. R., Mcanulty, L. S., Morrow, J. D., Ahmed, A., & Heward, C. B. (2004). Vitamin E and immunity after the Kona triathlon world championship. *Medicine and Science in Sports and Exercise*, *36*, 1328-1335.
- Oliver, J. M., Caldwell, A., Sanders, E., Mitchell, J., Rowlands, D., Purpura, M., & Jager, R. (2017). Novel form of curcumin attenuates performance decrements following muscle damaging exercise. *The FASEB Journal*, *31*(1 Supplement), lb415-lb415.
- Ortenblad, N., Madsen, K., & Djurhuus, M. S. (1997). Antioxidant status and lipid peroxidation after short-term maximal exercise in trained and untrained humans. *Am J Physiol*, *272*(4 Pt 2), R1258-1263.
- Ostrowski, K., Hermann, C., Bangash, A., Schjerling, P., Nielsen, J. N., & Pedersen, B. K. (1998). A trauma-like elevation of plasma cytokines in humans in response to treadmill running. *The Journal of physiology*, *513*(3), 889-894.

- Ostrowski, K., Rohde, T., Asp, S., Schjerling, P., & Pedersen, B. K. (1999). Pro-and anti-inflammatory cytokine balance in strenuous exercise in humans. *The Journal of physiology*, *515*(1), 287-291.
- Pala, R., Orhan, C., Tuzcu, M., Sahin, N., Ali, S., Cinar, V., . . . Sahin, K. (2016). Coenzyme Q10 Supplementation Modulates NFkappaB and Nrf2 Pathways in Exercise Training. *J Sports Sci Med*, *15*(1), 196-203.
- Park, S. Y., & Kwak, Y. S. (2016). Impact of aerobic and anaerobic exercise training on oxidative stress and antioxidant defense in athletes. *J Exerc Rehabil*, *12*(2), 113-117. doi:10.12965/jer.1632598.299
- Paulsen, G., Cumming, K. T., Holden, G., Hallen, J., Ronnestad, B. R., Sveen, O., . . . Raastad, T. (2014). Vitamin C and E supplementation hampers cellular adaptation to endurance training in humans: a double-blind, randomised, controlled trial. *J Physiol*, *592*(8), 1887-1901. doi:10.1113/jphysiol.2013.267419
- Peake, J., Nosaka, K., & Suzuki, K. (2005). Characterization of inflammatory responses to eccentric exercise in humans. *Exerc Immunol Rev*, *11*, 64-85.
- Pedersen, Akerstrom, Nielsen, & Fischer. (2007). Role of myokines in exercise and metabolism. *J Appl Physiol (1985)*, *103*(3), 1093-1098.  
doi:10.1152/jappphysiol.00080.2007
- Pedersen, & Hoffman-Goetz. (2000). Exercise and the Immune System: Regulation, Integration, and Adaptation. *Physiological reviews*, *80*(3), 1055-1081.

- Pedersen, Ostrowski, Rohde, & Bruunsgaard. (1998a). The cytokine response to strenuous exercise. *Canadian journal of physiology and pharmacology*, 76(5), 505-511.
- Pedersen, Ostrowski, Rohde, & Bruunsgaard. (1998b). The cytokine response to strenuous exercise. *Can J Physiol Pharmacol*, 76(5), 505-511.
- Pedersen, Steensberg, A., Fischer, C., Keller, C., Ostrowski, K., & Schjerling, P. (2001). Exercise and cytokines with particular focus on muscle-derived IL-6. *Exerc Immunol Rev*, 7, 18-31.
- Perron, N. R., & Brumaghim, J. L. (2009). A review of the antioxidant mechanisms of polyphenol compounds related to iron binding. *Cell Biochem Biophys*, 53(2), 75-100. doi:10.1007/s12013-009-9043-x
- Petersen, A. M. W., & Pedersen, B. K. (2005). The anti-inflammatory effect of exercise. *Journal of applied physiology*, 98(4), 1154-1162.
- Peterson, J. M., Bakkar, N., & Guttridge, D. C. (2011). 4 NF-kB Signaling in Skeletal Muscle Health and Disease. *Current topics in developmental biology*, 96, 85.
- Philippou, A., Maridaki, M., Theos, A., & Koutsilieris, M. (2012). Cytokines in muscle damage. *Adv Clin Chem*, 58, 49-87.
- Picklo, M. J., & Thyfault, J. P. (2015). Vitamin E and vitamin C do not reduce insulin sensitivity but inhibit mitochondrial protein expression in exercising obese rats. *Appl Physiol Nutr Metab*, 40(4), 343-352. doi:10.1139/apnm-2014-0302

- Powers, S. K., Criswell, D., Lawler, J., Ji, L. L., Martin, D., Herb, R. A., & Dudley, G. (1994). Influence of exercise and fiber type on antioxidant enzyme activity in rat skeletal muscle. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, 266(2), R375-R380.
- Powers, S. K., & Jackson, M. J. (2008). Exercise-induced oxidative stress: cellular mechanisms and impact on muscle force production. *Physiological reviews*, 88(4), 1243-1276.
- Prisk, V., & Huard, J. (2003). Muscle injuries and repair: the role of prostaglandins and inflammation. *Histol Histopathol*, 18(4), 1243-1256.
- Proske, U., & Morgan, D. L. (2001). Muscle damage from eccentric exercise: mechanism, mechanical signs, adaptation and clinical applications. *J Physiol*, 537(Pt 2), 333-345.
- Radák, Z., Apor, P., Pucsok, J., Berkes, I., Ogonovszky, H., Pavlik, G., . . . Goto, S. (2003). Marathon running alters the DNA base excision repair in human skeletal muscle. *Life sciences*, 72(14), 1627-1633.
- Radak, Z., Pucsok, J., Mecseki, S., Csont, T., & Ferdinandy, P. (1999). Muscle soreness-induced reduction in force generation is accompanied by increased nitric oxide content and DNA damage in human skeletal muscle. *Free Radic Biol Med*, 26(7-8), 1059-1063.

- Ramel, A., Wagner, K.-H., & Elmadfa, I. (2004). Plasma antioxidants and lipid oxidation after submaximal resistance exercise in men. *European journal of nutrition*, 43(1), 2-6.
- Rege, S. D., Kumar, S., Wilson, D. N., Tamura, L., Geetha, T., Mathews, S. T., . . . Babu, J. R. (2013). Resveratrol Protects the Brain of Obese Mice from Oxidative Damage. *Oxid Med Cell Longev*, 2013. doi:10.1155/2013/419092
- Ruby, A. J., Kuttan, G., Dinesh Babu, K., Rajasekharan, K. N., & Kuttan, R. (1995). Anti-tumour and antioxidant activity of natural curcuminoids. *Cancer Letters*, 94(1), 79-83. doi:[http://dx.doi.org/10.1016/0304-3835\(95\)03827-J](http://dx.doi.org/10.1016/0304-3835(95)03827-J)
- Sarker, M. R., Franks, S., Sumien, N., Thangthaeng, N., Filipetto, F., & Forster, M. (2015). Curcumin Mimics the Neurocognitive and Anti-Inflammatory Effects of Caloric Restriction in a Mouse Model of Midlife Obesity. *PLoS One*, 10(10). doi:10.1371/journal.pone.0140431
- Sastre, J., Asensi, M., Gasco, E., Pallardo, F. V., Ferrero, J., Furukawa, T., & Vina, J. (1992). Exhaustive physical exercise causes oxidation of glutathione status in blood: prevention by antioxidant administration. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, 263(5), R992-R995.
- Satoshi, S., Kiyoji, T., Hiroyo, K., & Fumio, N. (1989). Exercise-induced lipid peroxidation and leakage of enzymes before and after vitamin E supplementation. *International Journal of Biochemistry*, 21(8), 835-838.

- Saxton, J. M., Donnelly, A. E., & Roper, H. P. (1994). Indices of free-radical-mediated damage following maximum voluntary eccentric and concentric muscular work. *Eur J Appl Physiol Occup Physiol*, 68(3), 189-193.
- Sciberras, J. N., Galloway, S. D., Fenech, A., Grech, G., Farrugia, C., Duca, D., & Mifsud, J. (2015). The effect of turmeric (Curcumin) supplementation on cytokine and inflammatory marker responses following 2 hours of endurance cycling. *Journal of the International Society of Sports Nutrition*, 12(1), 1.
- Shehzad, A., & Lee, Y. S. (2013). Molecular mechanisms of curcumin action: signal transduction. *Biofactors*, 39(1), 27-36.
- Sies, H. (1985). Oxidative stress: introductory remarks. *Oxidative stress*, 1-8.
- Sies, H. (1986). Biochemistry of oxidative stress. *Angewandte Chemie International Edition in English*, 25(12), 1058-1071.
- Sies, H. (1991). Oxidative stress: from basic research to clinical application. *The American journal of medicine*, 91(3), S31-S38.
- Sies, H. (1997). Oxidative stress: oxidants and antioxidants. *Experimental Physiology*, 82(2), 291-295.
- Singh, & Aggarwal. (1995). Activation of transcription factor NF-B is suppressed by curcumin (diferuloylmethane). *Journal of Biological Chemistry*, 270(50), 30235b-30235b.

- Singh, Barik, A., Singh, B. G., & Priyadarsini, K. I. (2011). Reactions of reactive oxygen species (ROS) with curcumin analogues: Structure–activity relationship. *Free radical research*, 45(3), 317-325.
- Smith, Kruger, Smith, & Myburgh. (2008). The inflammatory response to skeletal muscle injury. *Sports Medicine*, 38(11), 947-969.
- Soni, K., & Kuttan, R. (1992). Effect of oral curcumin administration on serum peroxides and cholesterol levels in human volunteers. *Indian journal of physiology and pharmacology*, 36, 273-273.
- Sprenger, H., Jacobs, C., Nain, M., Gressner, A., Prinz, H., Wesemann, W., & Gemsa, D. (1992). Enhanced release of cytokines, interleukin-2 receptors, and neopterin after long-distance running. *Clinical immunology and immunopathology*, 63(2), 188-195.
- St-Pierre, J., Drori, S., Uldry, M., Silvaggi, J. M., Rhee, J., Jäger, S., . . . Yang, W. (2006). Suppression of reactive oxygen species and neurodegeneration by the PGC-1 transcriptional coactivators. *Cell*, 127(2), 397-408.
- Steensberg, A., Hall, G., Osada, T., Sacchetti, M., Saltin, B., & Pedersen, B. K. (2000). Production of interleukin-6 in contracting human skeletal muscles can account for the exercise-induced increase in plasma interleukin-6. *The Journal of physiology*, 529(1), 237-242.



- Subudhi, A. W., Davis, S. L., Kipp, R. W., & Askew, E. W. (2001). Antioxidant status and oxidative stress in elite alpine ski racers. *International Journal of Sport Nutrition and Exercise Metabolism*, 11(1), 32-41.
- Taaffe, D. R., Harris, T. B., Ferrucci, L., Rowe, J., & Seeman, T. E. (2000). Cross-sectional and prospective relationships of interleukin-6 and C-reactive protein with physical performance in elderly persons MacArthur Studies of Successful Aging. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, 55(12), M709-M715.
- Takahashi, M., Suzuki, K., Kim, H., Otsuka, Y., Imaizumi, A., Miyashita, M., & Sakamoto, S. (2014). Effects of curcumin supplementation on exercise-induced oxidative stress in humans. *International journal of sports medicine*, 35(06), 469-475.
- Tauler, P., Gimeno, I., Aguilo, A., Guix, M. P., & Pons, A. (1999). Regulation of erythrocyte antioxidant enzyme activities in athletes during competition and short-term recovery. *Pflugers Arch*, 438(6), 782-787.
- Teixeira, M. G., Mill, J. G., Pereira, A. C., & Molina Mdel, C. (2016). Dietary intake of antioxidant in ELSA-Brasil population: baseline results. *Rev Bras Epidemiol*, 19(1), 149-159. doi:10.1590/1980-5497201600010013
- Tidball. (2005). Inflammatory processes in muscle injury and repair. *Am J Physiol Regul Integr Comp Physiol*, 288(2), R345-353. doi:10.1152/ajpregu.00454.2004

- Tidball, & Villalta. (2010). Regulatory interactions between muscle and the immune system during muscle regeneration. *American Journal of Physiology - Regulatory, Integrative and Comparative Physiology*, 298(5), R1173-R1187.  
doi:10.1152/ajpregu.00735.2009
- Tiidus, P. M. (1998). Radical species in inflammation and overtraining. *Can J Physiol Pharmacol*, 76(5), 533-538.
- Tiidus, P. M., Pushkarenko, J., & Houston, M. E. (1996). Lack of antioxidant adaptation to short-term aerobic training in human muscle. *Am J Physiol*, 271(4 Pt 2), R832-836.
- Tilg, H., Dinarello, C. A., & Mier, J. W. (1997). IL-6 and APPs: anti-inflammatory and immunosuppressive mediators. *Immunology Today*, 18(9), 428-432.  
doi:[http://dx.doi.org/10.1016/S0167-5699\(97\)01103-1](http://dx.doi.org/10.1016/S0167-5699(97)01103-1)
- Townsend, J. R., Hoffman, J. R., Fragala, M. S., Jajtner, A. R., Gonzalez, A. M., Wells, A. J., . . . Stout, J. R. (2015). TNF- $\alpha$  and TNFR1 responses to recovery therapies following acute resistance exercise. *Front Physiol*, 6.
- Ullum, H., Haahr, P. M., Diamant, M., Palmo, J., Halkjaer-Kristensen, J., & Pedersen, B. K. (1994). Bicycle exercise enhances plasma IL-6 but does not change IL-1 alpha, IL-1 beta, IL-6, or TNF-alpha pre-mRNA in BMNC. *Journal of applied physiology*, 77(1), 93-97.

- Usharani, P., Mateen, A. A., Naidu, M. U., Raju, Y. S., & Chandra, N. (2008). Effect of NCB-02, atorvastatin and placebo on endothelial function, oxidative stress and inflammatory markers in patients with type 2 diabetes mellitus: a randomized, parallel-group, placebo-controlled, 8-week study. *Drugs R D*, 9(4), 243-250.
- Ushio-Fukai, M. (2006). Redox signaling in angiogenesis: role of NADPH oxidase. *Cardiovascular research*, 71(2), 226-235.
- Venturini, C. D., Merlo, S., Souto, A. A., Fernandes Mda, C., Gomez, R., & Rhoden, C. R. (2010). Resveratrol and red wine function as antioxidants in the nervous system without cellular proliferative effects during experimental diabetes. *Oxid Med Cell Longev*, 3(6), 434-441.
- Vina, J., Gimeno, A., Sastre, J., Desco, C., Asensi, M., Pallardó, F. V., . . . Repine, J. E. (2000). Mechanism of free radical production in exhaustive exercise in humans and rats; role of xanthine oxidase and protection by allopurinol. *IUBMB life*, 49(6), 539-544.
- Vincent, H. K., Morgan, J. W., & Vincent, K. R. (2004). Obesity exacerbates oxidative stress levels after acute exercise. *Medicine and Science in Sports and Exercise*, 36(5), 772-779.

Visser, M., Pahor, M., Taaffe, D. R., Goodpaster, B. H., Simonsick, E. M., Newman, A. B., . . . Harris, T. B. (2002). Relationship of interleukin-6 and tumor necrosis factor- $\alpha$  with muscle mass and muscle strength in elderly men and women The Health ABC Study. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, 57(5), M326-M332.

Volek, J. S., Kraemer, W. J., Rubin, M. R., Gómez, A. L., Ratamess, N. A., & Gaynor, P. (2002). L-Carnitine L-tartrate supplementation favorably affects markers of recovery from exercise stress. *American Journal of Physiology-Endocrinology And Metabolism*, 282(2), E474-E482.

Wadley, G. D., & McConell, G. K. (2010). High-dose antioxidant vitamin C supplementation does not prevent acute exercise-induced increases in markers of skeletal muscle mitochondrial biogenesis in rats. *J Appl Physiol (1985)*, 108(6), 1719-1726. doi:10.1152/jappphysiol.00127.2010

Warren, G. L., Ingalls, C. P., Lowe, D. A., & Armstrong, R. B. (2001). Excitation-contraction uncoupling: major role in contraction-induced muscle injury. *Exerc Sport Sci Rev*, 29(2), 82-87.

Wiecek, M., Maciejczyk, M., Szymura, J., Wiecha, S., Kantorowicz, M., & Szygula, Z. (2016). Effect of body composition, aerobic performance and physical activity on exercise-induced oxidative stress in healthy subjects. *J Sports Med Phys Fitness*.

Yeon, K., Kim, S., Kim, Y., Lee, M., Ahn, D., Kim, H., . . . Oh, S. (2010). Curcumin produces an antihyperalgesic effect via antagonism of TRPV1. *Journal of dental research*, 89(2), 170-174.

Zhang, Y. P., Song, C. Y., Yuan, Y., Eber, A., Rodriguez, Y., Levitt, R. C., . . . Candiotti, K. A. (2013). Diabetic neuropathic pain development in type 2 diabetic mouse model and the prophylactic and therapeutic effects of coenzyme Q10. *Neurobiol Dis*, 58, 169-178. doi:10.1016/j.nbd.2013.05.003