



# European Society for Clinical Nutrition and Metabolism (ESPEN) Malnutrition Criteria for Predicting Major Complications After Hepatectomy and Pancreatectomy

Yasuyuki Fukami<sup>1</sup> · Takuya Saito<sup>1</sup> · Takashi Arikawa<sup>1</sup> · Takaaki Osawa<sup>1</sup> · Shunichiro Komatsu<sup>1</sup> · Kenitiro Kaneko<sup>1</sup> · Yuria Ishida<sup>2</sup> · Keisuke Maeda<sup>2,3</sup> · Naoharu Mori<sup>2,3</sup> · Tsuyoshi Sano<sup>1</sup>

Accepted: 16 August 2020 / Published online: 3 September 2020  
© Société Internationale de Chirurgie 2020

## Abstract

**Background** Recently, diagnostic criteria for malnutrition have been proposed by the European Society for Clinical Nutrition and Metabolism (ESPEN). This study aimed to investigate the utility of the ESPEN malnutrition criteria as a predictor for major complications following hepatectomy and pancreatectomy.

**Methods** Data were reviewed from 176 consecutive patients who underwent hepatectomy ( $n = 103$ ) or pancreatectomy ( $n = 73$ ) between November 2017 and December 2019. Patients were divided into two groups according to the ESPEN malnutrition criteria using a prospectively collected database. The clinical data and the surgical outcomes of patients in the malnourished and normal groups were retrospectively analyzed.

**Results** Thirty-five (20%) patients were diagnosed with malnourishment according to ESPEN criteria. The malnourished group had a significantly low preoperative albumin concentration ( $p = 0.001$ ). After hepatectomy, major complications (Clavien grade  $\geq 3a$ ) occurred significantly more frequently in the malnourished group than in the normal group ( $p = 0.013$ ). Multivariate analysis indicated that operative duration  $\geq 300$  min (hazard ratio: 22.47, 95% CI: 2.17 to 232.73,  $p = 0.009$ ) and malnourishment (hazard ratio: 14.56, 95% CI: 2.58 to 82.17,  $p = 0.002$ ) were independently associated with major complications after hepatectomy. On the other hand, malnutrition was not associated with major complications after pancreatectomy.

**Conclusions** The ESPEN malnutrition criteria are a valuable predictor for major complications following hepatectomy.

## Introduction

Surgical resection is a standard treatment for hepatopancreatobiliary (HPB) malignancies [1–4]. Although hepatectomy and pancreatectomy have become safe surgeries with low mortality rates due to improvements in surgical techniques and perioperative management, postoperative morbidity rates remain high [5, 6]. Postoperative complications are risk factors for prolonged hospital stays and increased costs of care in surgical patients [7, 8]. Therefore, minimizing postoperative complications can potentially reduce medical costs and improve patient quality of life.

✉ Yasuyuki Fukami  
yasuyuki490225@yahoo.co.jp

<sup>1</sup> Division of Gastroenterological Surgery, Department of Surgery, Aichi Medical University, 1-1 Yazakokarimata, Nagakute 480-1195, Aichi, Japan

<sup>2</sup> Department of Nutrition, Aichi Medical University, Nagakute, Aichi, Japan

<sup>3</sup> Department of Palliative and Supportive Medicine, Aichi Medical University, Nagakute, Aichi, Japan

Malnutrition is a well-known risk factor for postoperative complications after abdominal surgery [9, 10]. However, for HPB surgery, the data on the influence of malnutrition are limited by retrospective study designs and insufficient sample sizes. Another issue is that it is difficult to determine which criteria of malnutrition to adopt because numerous different clinical scores have been developed to detect malnutrition [11–13]. Therefore, identifying criteria that are practical and easy to use for assessing malnutritional status in HPB surgery is necessary.

Recently, diagnostic criteria for malnutrition have been proposed by the European Society for Clinical Nutrition and Metabolism (ESPEN), and these criteria have been validated by some studies [14–16]. However, the value of the ESPEN malnutrition criteria as a predictor for postoperative complications after HPB surgery is unclear. This study aimed to investigate the utility of the ESPEN malnutrition criteria as a predictor for major complications following hepatectomy and pancreatectomy.

## Patients and methods


Between November 2017 and December 2019, 176 consecutive patients who underwent hepatectomy ( $n = 103$ ) or pancreatectomy ( $n = 73$ ) at our institution were enrolled in the study. Patients were divided into two groups according to the ESPEN malnutrition criteria using a prospectively collected database (Fig. 1). The clinical data and the surgical outcomes of patients in the malnourished and normal

groups were retrospectively analyzed. Major hepatectomy was defined as the resection of three or more Couinaud's segments. All patients provided signed informed consent before undergoing therapy. This study was approved by the Institutional Review Board of our institution (No. 2019-136) and was in accordance with the 1964 Helsinki Declaration and its later amendments or comparable ethical standards.

## The new ESPEN diagnostic criteria for malnutrition

Multidisciplinary nutrition support teams provided nutritional assessment, as well as nutritional screening, as part of standard medical care in our hospital [17, 18]. The diagnosis of malnutrition was made by nutritional support teams based on the 2015 ESPEN criteria [14]. ESPEN - defined malnutrition is diagnosed in two steps. The first step is screening patients at risk of malnutrition using a validated nutritional screening tool. In the present study, we applied the Nutritional Assessment-Short Form as the screening tool. Consequently, screened patients were assessed if they satisfied at least one of the following criteria: (i) BMI  $<18.5 \text{ kg/m}^2$ ; (ii) unintentional weight loss  $>5\%$  over the past 3–6 months combined with BMI  $<20 \text{ kg/m}^2$  if age  $<70$  years or  $<22.0 \text{ kg/m}^2$  if age  $\geq 70$  years; and (iii) unintentional weight loss combined with fat-free mass index  $<15.0 \text{ kg/m}^2$  for women or  $<17.0 \text{ kg/m}^2$  for men. The fat-free mass index was calculated by dividing the patient's estimated fat-free mass by the patient's height in meters squared ( $\text{m}^2$ ) [19, 20]. The fat-free mass was obtained using the following formula that

**Fig. 1** Patient malnutrition assessment flow chart.  
\*unintentional weight loss  $>5\%$  over the past 3–6 months combined with BMI  $<20 \text{ kg/m}^2$  if age  $<70$  years or  $<22.0 \text{ kg/m}^2$  if age  $\geq 70$  years and/or unintentional weight loss combined with fat-free mass index  $<15.0 \text{ kg/m}^2$  for women or  $<17.0 \text{ kg/m}^2$  for men



**Table 1** Patient characteristics

	ESPEN criteria		<i>P</i>
	Normal ( <i>n</i> = 141)	Malnourished ( <i>n</i> = 35)	
Age (year)	69 (31–89)	72 (35–85)	0.204
Sex (male/female)	96/45	13/22	0.002
BMI (kg/m <sup>2</sup> )	23.2 (18.7–40.8)	17.8 (15.4–21.2)	< 0.001
Albumin (g/dL)	3.8 (1.7–5.0)	3.4 (2.4–4.1)	0.001
ASA status (I/II/III)	57/78/6	17/14/4	0.120
Comorbidity			
Cardiovascular disease	25 (18%)	6 (17%)	1.000
Cerebrovascular disease	7 (5%)	3 (9%)	0.419
Pulmonary disease	11 (8%)	3 (9%)	1.000
Chronic kidney disease	10 (7%)	2 (6%)	1.000
Diabetes mellitus	31 (22%)	6 (17%)	0.646
Surgery			0.851
Hepatectomy	83 (59%)	20 (57%)	
Pancreatectomy	58 (41%)	15 (43%)	

Expressed as *N* (%) or median (range)

ESPEN European Society for Clinical Nutrition and Metabolism; BMI Body mass index; ASA American Society of Anesthesiologists

includes the estimated 24-h urine creatinine excretion rate (eCER): fat-free mass =  $13.0 + 0.03 \times \text{eCER}$ ; eCER (mg/day) =  $879.89 + 12.51 \times \text{weight (kg)} - 6.19 \times \text{age}$  ( $- 379.42$  for female patients).

### Postoperative complications

Postoperative complications were assessed according to the Clavien–Dindo classification system, and complications of grade 3a or worse were defined as major [21]. Any complications that developed within 90 days after the operation were included. No patients were lost to follow-up. Postoperative pancreatic fistula ( $\geq$  grade B) and bile leakage were defined according to the definitions of an international study group (i.e., ISGPF or ISGLS) [22, 23].

### Outcome measurements

The first outcome measurement was performed by comparing the variables of patient demographic and clinical data as well as postoperative parameters between the malnourished and normal groups. The second outcome measurement was performed by analyzing the risk factors affecting the postoperative major complications in patients who underwent hepatectomy or pancreatectomy.

### Statistical analysis

The continuous data are expressed as the medians (ranges). The statistical analyses were performed using chi square tests, Mann–Whitney *U* tests, or Fisher’s exact probability tests as appropriate. The variables identified as potentially significant by univariate analysis (*p* value < 0.10) were selected for multivariate analysis with a logistic regression model to identify the independent predictors of postoperative major complications. All *P* values were 2-sided, and *P* < 0.05 was considered to indicate a statistically significant difference. All statistical calculations were performed using the IBM SPSS Statistics 21 software package (IBM Japan Inc., Tokyo, Japan).

### Results

The patient characteristics related to hepatectomy and pancreatectomy are shown in Table 1. Thirty-five (20%) patients were diagnosed as malnourished according to the ESPEN criteria. The malnourished group had a significantly lower preoperative albumin concentration than the normal group (*p* = 0.001). The median age, ASA status, and incidence of comorbidities did not differ between the two groups. The incidence of chemotherapy prior to hepatectomy was 19% in the normal group and 25% in the malnourished group (*p* = 0.549). The incidence of chemotherapy prior to pancreatectomy was 9% in the

**Table 2** Surgical outcome after hepatectomy

	ESPEN criteria		P
	Normal (n = 83)	Malnourished (n = 20)	
Major hepatectomy ( $\geq 3$ segments*)	14 (17%)	9 (45%)	0.014
Multiple sites hepatectomy	15 (18%)	3 (15%)	1.000
Laparoscopic hepatectomy	29 (35%)	3 (15%)	0.109
Operative duration (min)	295 (123–877)	247.5 (113–569)	0.379
Total blood loss (mL)	224 (5–2856)	265.5 (1–2734)	0.802
Disease			0.340
Hepatocellular carcinoma	43 (52%)	8 (40%)	
Metastatic liver cancer	29 (35%)	6 (30%)	
Bile duct cancer	6 (7%)	3 (15%)	
Others	5 (6%)	3 (15%)	
Morbidity (overall)	18 (22%)	8 (40%)	0.149
Intraabdominal abscess	6 (7%)	2 (10%)	0.651
Bile leakage	4 (5%)	2 (10%)	0.330
Wound infection	3 (4%)	0	1.000
Refractory ascites	2 (2%)	0	1.000
Pleural effusion <sup>a</sup>	1 (1%)	1 (5%)	0.352
Liver failure	0	0	-
Other	4 (5%)	4 (20%)	0.044
Morbidity ( $\geq$ Clavien grade 3a)	4 (5%)	5 (25%)	0.013
Mortality	0	1 (5%)	0.194

Expressed as N (%) or median (range)

ESPEN European Society for Clinical Nutrition and Metabolism

\*Couinaud's segments

<sup>a</sup>Required thoracentesis

normal group and 0% in the malnourished group ( $p = 0.576$ ).

Table 2 shows the surgical outcomes after hepatectomy: 51 patients had hepatocellular carcinoma (HCC), 35 patients had metastatic liver cancer, 9 patients had bile duct cancer, and 8 patients had other diseases. Hepatectomy at multiple sites, use of a laparoscopic approach, operative duration, total blood loss volume, disease type, and overall morbidity did not differ between the two groups. A list of the observed morbidities is shown in Table 2, and the details of other morbidities are as follows: delayed gastric emptying ( $n = 1$  in the normal group;  $n = 2$  in the malnourished group), portal vein thrombosis ( $n = 1$  in the normal group), anastomotic leakage of the colon after simultaneous colectomy ( $n = 1$  in the normal group), pneumonia ( $n = 1$  in the normal group), septicemia ( $n = 1$  in the malnourished group), and cerebral infarction ( $n = 1$  in the malnourished group). Although the mortality rate was not significantly different between the two groups, one patient in the malnourished group who underwent minor hepatectomy (i.e., left lateral sectionectomy) for HCC died

from cerebral infarction postoperatively. There were significant differences between the two groups in the major hepatectomy rate (17% in the normal group vs. 45% in the malnourished group) and the major postoperative complication rate (5% in the normal group vs. 25% in the malnourished group). Among the nine malnourished patients who underwent major hepatectomy, three patients (33%) had major complications.

Table 3 shows the surgical outcomes after pancreatectomy: 41 patients had pancreatic cancer, 9 patients had bile duct cancer, and 23 patients had other diseases. Type of pancreatectomy, use of a laparoscopic approach, operative duration, total blood loss volume, disease type, pancreatic texture, and morbidity did not differ between the two groups. A list of observed morbidities is shown in Table 3, and the details of other morbidities are as follows: ascites ( $n = 3$  in the normal group;  $n = 2$  in the malnourished group), pneumonia ( $n = 2$  in the normal group;  $n = 1$  in the malnourished group), cholangitis ( $n = 1$  in the normal group;  $n = 1$  in the malnourished group), pleural effusion requiring thoracentesis ( $n = 1$  in the malnourished group),

**Table 3** Surgical outcome after pancreatectomy

	ESPEN criteria		P
	Normal (n = 58)	Malnourished (n = 15)	
Type of pancreatectomy			0.633
Pancreatoduodenectomy	31 (53%)	10 (67%)	
Distal pancreatectomy	23 (40%)	4 (27%)	
Total pancreatectomy	4 (7%)	1 (7%)	
Laparoscopic pancreatectomy	6 (10%)	2 (13%)	0.664
Operative duration (min)	353 (157–668)	343 (190–448)	0.633
Total blood loss (mL)	302 (10–4250)	200 (3–1234)	0.170
Disease			0.756
Pancreatic cancer	32 (55%)	9 (60%)	
Bile duct cancer	8 (14%)	1 (7%)	
Others	18 (31%)	5 (33%)	
Pancreatic texture			0.229
Soft	39 (67%)	7 (47%)	
Hard	19 (33%)	8 (53%)	
Morbidity (overall)	35 (60%)	10 (67%)	0.770
Pancreatic fistula ( $\geq$ grade B)	24 (41%)	3 (20%)	0.147
Bile leakage	2 (3%)	0	1.000
Intraabdominal abscess	7 (12%)	2 (13%)	0.664
Intraabdominal bleeding	3 (5%)	0	1.000
Delayed gastric emptying	1 (2%)	2 (13%)	0.105
Septicemia	0	2 (13%)	0.040
Other	6 (10%)	6 (40%)	0.013
Morbidity ( $\geq$ Clavien grade 3a)	24 (41%)	3 (20%)	0.147
Mortality	0	0	-

Expressed as *N* (%) or median (range)

ESPEN European Society for Clinical Nutrition and Metabolism

and portal vein thrombosis ( $n = 1$  in the malnourished group). The in-hospital and 90-day postoperative mortality rates were zero in both groups.

The multivariate analysis indicated that operative duration  $\geq 300$  min (hazard ratio: 22.47, 95% CI: 2.17 to 232.73,  $p = 0.009$ ) and malnourishment (hazard ratio: 14.56, 95% CI: 2.58 to 82.17,  $p = 0.002$ ) were independently associated with major complications after hepatectomy (Table 4). On the other hand, malnutrition was not associated with major complications after pancreatectomy (Table 5). The multivariate analysis indicated that soft pancreas (hazard ratio: 4.50, 95% CI: 1.25 to 16.19,  $p = 0.021$ ) and total blood loss  $\geq 400$  mL (hazard ratio: 4.91, 95% CI: 1.57 to 15.34,  $p = 0.006$ ) were independently associated with major complications after pancreatectomy.

## Discussion

This is the first nutritional study in the literature to note the new ESPEN malnutrition criteria in patients who underwent HPB surgery. In this study, 20% of all patients were diagnosed as malnourished according to the ESPEN criteria. The most interesting finding of this study was the difference between the results of the two surgeries (i.e., hepatectomy and pancreatectomy). Malnourishment was independently associated with major complications after hepatectomy. On the other hand, malnutrition was not associated with major complications after pancreatectomy.

In 2015, ESPEN presented new consensus criteria for the diagnosis of malnutrition with the aim of reaching uniformity between countries and/or studies [14]. According to a previous study on the prevalence of malnutrition diagnosed by the new ESPEN criteria, 0.5% of healthy

**Table 4** Uni- and multivariate predictors of major complications after hepatectomy

Variable	Univariate			Multivariate		
	HR	95% CI	P	HR	95% CI	P
<b>Age</b>						
> 5 (years) (n = 34)	2.46	0.32–19.0	0.390			
<75 (years) (n = 69)	1					
<b>Sex</b>						
Female (n = 31)	1.32	0.17–10.01	0.788			
Male (n = 72)	1					
<b>Albumin</b>						
<3.0 (g/dL) (n = 5)	1.20	0.02–78.52	0.755			
>3.0 (g/dL) (n = 98)	1					
<b>Diabetes mellitus</b>						
Present (n = 22)	0.39	0.03–4.96	0.470			
Absent (n = 81)	1					
<b>ASA status</b>						
III (n = 9)	1.23	0.04–40.47	0.907			
I/II (n = 94)	1					
<b>Extent of hepatectomy</b>						
Major (n = 23)	1.38	0.18–10.39	0.755			
Minor (n = 80)	1					
<b>Operative duration</b>						
>300 (min) (n = 48)	16.59	1.01–271.67	0.049	22.47	2.17–232.73	0.009
< 300 (min) (n = 55)	1			1		
<b>Total blood loss</b>						
>400 (mL) (n = 27)	3.60	0.58–22.23	0.169			
<400 (mL) (n = 76)	1					
<b>ESPEN criteria</b>						
Malnourished (n = 20)	10.39	0.98–109.90	0.052	14.56	2.58–82.17	0.002
Normal (n = 83)	1			1		

ASA American Society of Anesthesiologists; *ESPEN* European Society for Clinical Nutrition and Metabolism

elderly individuals and 6% of geriatric outpatients were identified as malnourished [24]. These values are very low compared with those of the present study, which indicates that patients with HPB disease might have a higher prevalence of malnutrition than the healthy and geriatric populations. Similarly, in a recent study of geriatric gastrointestinal cancer, 20% of the patients were classified as malnourished according to the new ESPEN criteria [16]. In another report, 16.9% of the patients admitted with a proximal femoral fracture were diagnosed as malnourished by the new ESPEN criteria [25]. Therefore, it should be noted that the number of patients requiring surgery that have malnutrition is higher than expected.

Hepatectomy is one of the common procedures in HPB surgery that can lead to postoperative complications. Patients with HCC have underlying chronic liver disease, and patients with metastatic liver cancer may have a

history of chemotherapy prior to hepatectomy; thus, these patients are at risk of developing postoperative complications. Preoperative nutritional status is one of the key factors for the success of a hepatectomy. Currently, various nutritional assessment tools are widely used for patients with HPB disease. Schiesser et al. [26] reported that 27.3% of the patients who underwent elective hepatobiliary surgery were malnourished, as determined by the Nutrition Risk Screening 2002 (NRS 2002) score. The Patient-Generated Subjective Global Assessment (PG-SGA) score is another nutritional assessment method [27, 28]. Huang et al. [29] showed that 33.4% of the patients who underwent hepatectomy for HCC were malnourished by using the PG-SGA, and malnourishment was associated with more postoperative complications than a healthy nutritional status. Therefore, the authors concluded that the PG-SGA is an effective tool for predicting postoperative

**Table 5** Uni- and multivariate predictors of major complications after pancreatectomy

Variable	Univariate			Multivariate		
	HR	95% CI	P	HR	95% CI	P
<b>Age</b>						
>75 (years) (n = 17)	1.39	0.30–6.42	0.675			
<75 (years) (n = 56)	1					
<b>Sex</b>						
Female (n = 36)	0.48	0.14–1.62	0.236			
Male (n = 37)	1					
<b>Albumin</b>						
<3.0 (g/dL) (n = 10)	4.97	0.80–31.02	0.086	3.80	0.79–18.40	0.097
>3.0 (g/dL) (n = 63)	1			1		
<b>Diabetes mellitus</b>						
Present (n = 15)	0.73	0.18–2.92	0.661			
Absent (n = 58)	1					
<b>Preoperative chemotherapy</b>						
Present (n = 5)	1.54	0.17–14.21	0.703			
Absent (n = 68)	1					
<b>Pancreatic texture</b>						
Soft (n = 46)	4.71	1.12–19.77	0.034	4.50	1.25–16.19	0.021
Hard (n = 27)	1			1		
<b>Operative duration</b>						
>300 (min) (n = 52)	0.48	0.12–1.99	0.313			
<300 (min) (n = 21)	1					
<b>Total blood loss</b>						
>400 (mL) (n = 29)	4.76	1.20–18.91	0.027	4.91	1.57–15.34	0.006
<400 (mL) (n = 44)	1			1		
<b>ESPEN criteria</b>						
Malnourished (n = 15)	0.55	0.10–2.97	0.487			
Normal (n = 58)	1					

ESPEN European Society for Clinical Nutrition and Metabolism

complications in patients with HCC following hepatectomy. However, the PG-SGA is a questionnaire completed by the individual, and thus the nutritional status can be underestimated or overestimated. In addition, the NRS 2002 or PG-SGA are just nutritional screening tools, not diagnostic tools. Conversely, the new ESPEN criteria are diagnostic criteria for malnutrition that are simple and practical to use.

When comparing the short-term results of hepatectomy and pancreatectomy, although the mortality rates are generally similar, pancreatectomy has a higher complication rate [5, 6]. Pancreatectomy is a highly invasive procedure with an inherent risk of complications. Postoperative pancreatic fistula (POPF) was the most common complication and subsequently triggered other infectious complications. Many studies have described risk factors for POPF and reported that pancreatic texture and duct size were

significant factors [30, 31]. In our series of patients receiving pancreatectomy, the POPF rate was as high as 37% in the total patients, and POPF was responsible for 50% of the overall morbidity. Taking this into consideration, nutritional status may be only a secondary factor in patients who undergo pancreatectomy. The beneficial effect of nutrition may not be sufficient to overcome these major factors affecting the outcome of pancreatectomy.

Many studies have addressed different screening tools and their ability to identify malnourished patients and patients who are at nutritional risk. In contrast, only a few studies have compared different screening tools to investigate their correlation and reliability [11]. Probst et al. [12] reported a prospective trial to evaluate the prognostic value of different nutritional assessment scores in pancreatectomy. They concluded that none of the nutritional assessment scores that assessed malnutrition were relevant to



complications after pancreatectomy and that these scores may thus be abandoned. McKenna et al. [13] investigated the optimal definition of malnutrition before major oncologic resection for 6 cancer types (i.e., colorectal, gastric, esophageal, liver, lung, and pancreatic cancer). They concluded that postoperative risk due to malnutrition varies based on both the definition of malnutrition used and the specific cancer type. Even in our study, a clinically relevant definition of malnutrition for patients undergoing pancreatectomy remains to be established.

This study has some limitations that need to be addressed. First, this was a retrospectively designed study, and a propensity score matching analysis was not possible because the number of patients, who were identified with the new malnutrition criteria at a single center, was too small. Second, there were no comparisons of different diagnostic criteria for malnutrition to investigate their correlation and reliability. Biochemical nutritional marker analysis is sometimes performed to evaluate the immunonutritional status, but this is sometimes unreliable and should only be used as a subsidiary tool to supplement nutrition risk screening assessment. In the future, multi-center large-series studies evaluating the clinical impact of malnutrition criteria, including the ESPEN malnutrition criteria after HPB surgery, will compensate for the limitations of this study.

In conclusion, our study indicates that the ESPEN malnutrition criteria are a valuable predictor for major complications following hepatectomy. Routine preoperative assessment of the ESPEN malnutrition criteria could help manage meticulous follow-up after hepatectomy.

**Acknowledgments** Not applicable.

**Author contributions** YF designed the study and wrote the initial draft of the manuscript. TS contributed to interpretation of the data and the critical revision of the manuscript for important intellectual content. All the other authors (TS, TA, TO, SK, KK, YI, KM, and NM) contributed to the data collection and interpretation and critically reviewed the manuscript. All the authors have read and approved the final version of the manuscript and have agreed to be accountable for all aspects of the study, ensuring that any questions related to the accuracy or integrity of any part of the work are answerable.

**Funding** The author(s) received no financial support for the research, authorship, and/or publication of this article.

**Compliance with ethical standards**

**Conflict of interest** The authors have no conflicts of interest to disclose.

**Ethical approval** This study was approved by the Institutional Review Board of Aichi Medical University (No. 2019-136) and was in accordance with the 1964 Helsinki Declaration and its later amendments or comparable ethical standards.

**Informed consent** All patients provided signed informed consent before undergoing therapy.

## References

- Sadot E, Groot Koerkamp B, Leal JN, Shia J, Gonen M, Allen PJ et al (2015) Resection margin and survival in 2368 patients undergoing hepatic resection for metastatic colorectal cancer: surgical technique or biologic surrogate? *Ann Surg* 262:476–485 **Discussion 483–485**
- Gemenetzis G, Groot VP, Blair AB, Laheru DA, Zheng L, Narang AK et al (2019) Survival in locally advanced pancreatic cancer after neoadjuvant therapy and surgical resection. *Ann Surg* 270:340–347
- Sano T, Shimada K, Sakamoto Y, Yamamoto J, Yamasaki S, Kosuge T (2006) One hundred two consecutive hepatobiliary resections for perihilar cholangiocarcinoma with zero mortality. *Ann Surg* 244:240–247
- Fukami Y, Kaneoka Y, Maeda A, Kumada T, Tanaka J, Akita T et al (2019) Liver resection for multiple hepatocellular carcinomas: a Japanese nationwide survey. *Ann Surg*. <https://doi.org/10.1097/SLA.0000000000003192> [Epub ahead of print]
- Spolverato G, Ejaz A, Hyder O, Kim Y, Pawlik TM (2014) Failure to rescue as a source of variation in hospital mortality after hepatic surgery. *Br J Surg* 101:836–846
- Hartwig W, Gluth A, Hinz U, Koliogiannis D, Strobel O, Hackert T et al (2016) Outcomes after extended pancreatectomy in patients with borderline resectable and locally advanced pancreatic cancer. *Br J Surg* 103:1683–1694
- Gani F, Hundt J, Makary MA, Haider AH, Zogg CK, Pawlik TM (2016) Financial impact of postoperative complication following hepato-pancreatico-biliary surgery for cancer. *Ann Surg Oncol* 23:1064–1070
- Wang J, Ma R, Churilov L, Eleftheriou P, Nikfarjam M, Christophi C et al (2018) The cost of perioperative complications following pancreaticoduodenectomy: a systematic review. *Pancreatol* 18:208–220
- Kuzu MA, Terzioğlu H, Genç V, Erkek AB, Ozban M, Sonyürek P et al (2006) Preoperative nutritional risk assessment in predicting postoperative outcome in patients undergoing major surgery. *World J Surg* 30:378–390. <https://doi.org/10.1007/s00268-005-0163-1>
- Braga M, Gianotti L, Nespoli L, Radaelli G, Di Carlo V (2002) Nutritional approach in malnourished surgical patients: a prospective randomized study. *Arch Surg* 137:174–180
- Schiesser M, Kirchoff P, Müller MK, Schäfer M, Clavien PA (2009) The correlation of nutrition risk index, nutrition risk score, and bioimpedance analysis with postoperative complications in patients undergoing gastrointestinal surgery. *Surgery* 145:519–526
- Probst P, Haller S, Bruckner T, Ulrich A, Strobel O, Hackert T et al (2017) Prospective trial to evaluate the prognostic value of different nutritional assessment scores in pancreatic surgery (NURIMAS Pancreas). *Br J Surg* 104:1053–1062
- McKenna NP, Bews KA, Al-Refaie WB, Colibaseanu DT, Pemberton JH, Cima RR et al (2020) Assessing malnutrition before major oncologic surgery: one size does not fit all. *J Am Coll Surg* 230:451–460
- Cederholm T, Bosaeus I, Barazzoni R, Bauer J, Van Gossum A, Klek S et al (2015) Diagnostic criteria for malnutrition—An ESPEN consensus statement. *Clin Nutr* 34:335–340
- Rondel ALMA, Langius JAE, de van der Schueren MAE, Kruijzena HM (2018) The new ESPEN diagnostic criteria for



- malnutrition predict overall survival in hospitalised patients. *Clin Nutr* 37:163–168
16. Ye XJ, Ji YB, Ma BW, Huang DD, Chen WZ, Pan ZY et al (2018) Comparison of three common nutritional screening tools with the new European society for clinical nutrition and metabolism (ESPEN) criteria for malnutrition among patients with geriatric gastrointestinal cancer: a prospective study in China. *BMJ Opens* 8(4):e019750
  17. Maeda K, Ishida Y, Nonogaki T, Mori N (2020) Reference body mass index values and the prevalence of malnutrition according to the Global leadership Initiative on malnutrition criteria. *Clin Nutr* 39:180–184
  18. Ishida Y, Maeda K, Nonogaki T, Shimizu A, Yamanaka Y, Matsuyama R et al (2020) Malnutrition at admission predicts in-hospital falls in hospitalized older adults. *Nutrients* 12(2):E541
  19. Ix JH, Wassel CL, Stevens LA, Beck GJ, Froissart M, Navis G et al (2011) Equations to estimate creatinine excretion rate: the CKD epidemiology collaboration. *Clin J Am Soc Nephrol* 6:184–191
  20. Jassal SK, Wassel CL, Laughlin GA, Barrett-Connor E, Rifkin DE, Ix JH (2015) Urine creatinine-based estimates of fat-free mass in community-dwelling older persons: the Rancho Bernardo study. *J Ren Nutr* 25:97–102
  21. Dindo D, Demartines N, Clavien PA (2004) Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg* 240:205–213
  22. Bassi C, Dervenis C, Butturini G, Fingerhut A, Yeo C, Izbicki J et al (2005) Postoperative pancreatic fistula: an international study group (ISGPF) definition. *Surgery* 138:8–13
  23. Koch M, Garden OJ, Padbury R, Rahbari NN, Adam R, Capussotti L et al (2011) Bile leakage after hepatobiliary and pancreatic surgery: a definition and grading of severity by the International Study Group of Liver Surgery. *Surgery* 149:680–688
  24. Rojer AG, Kruijzena HM, Trappenburg MC, Reijnierse EM, Sipilä S, Narici MV et al (2016) The prevalence of malnutrition according to the new ESPEN definition in four diverse populations. *Clin Nutr* 35:758–762
  25. van der Sijp MPL, van Eijk M, Krijnen P, Schipper IB, Achterberg WP, Niggebrugge AHP (2018) Screening for malnutrition in patients admitted to the hospital with a proximal femoral fracture. *Injury* 49:2239–2243
  26. Schiesser M, Müller S, Kirchhoff P, Breitenstein S, Schäfer M, Clavien PA (2008) Assessment of a novel screening score for nutritional risk in predicting complications in gastro-intestinal surgery. *Clin Nutr* 27:565–570
  27. Laky B, Janda M, Cleghorn G, Obermair A (2008) Comparison of different nutritional assessments and body-composition measurements in detecting malnutrition among gynecologic cancer patients. *Am J Clin Nutr* 87:1678–1685
  28. Gabrielson DK, Scaffidi D, Leung E, Stoyanoff L, Robinson J, Nisenbaum R et al (2013) Use of an abridged scored patient-generated subjective global assessment (abPG-SGA) as a nutritional screening tool for cancer patients in an outpatient setting. *Nutr Cancer* 65:234–239
  29. Huang TH, Hsieh CC, Kuo LM, Chang CC, Chen CH, Chi CC et al (2019) Malnutrition associated with an increased risk of postoperative complications following hepatectomy in patients with hepatocellular carcinoma. *HPB (Oxford)* 21:1150–1155
  30. Gaujoux S, Cortes A, Couvelard A, Noullet S, Clavel L, Rebours V et al (2010) Fatty pancreas and increased body mass index are risk factors of pancreatic fistula after pancreaticoduodenectomy. *Surgery* 148:15–23
  31. Mathur A, Pitt HA, Marine M, Saxena R, Schmidt CM, Howard TJ et al (2007) Fatty pancreas: a factor in postoperative pancreatic fistula. *Ann Surg* 246:1058–1064

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Reproduced with permission of copyright owner. Further reproduction prohibited without permission.