

# Fission yeast *orb6*, a ser/thr protein kinase related to mammalian rho kinase and myotonic dystrophy kinase, is required for maintenance of cell polarity and coordinates cell morphogenesis with the cell cycle

FULVIA VERDE\*†, DAVID J. WILEY\*, AND PAUL NURSE‡

\*Department of Biochemistry and Molecular Biology, University of Miami School of Medicine, P.O. Box 016129, Miami, FL, 33101-6129; and †Imperial Cancer Research Fund, 44 Lincoln's Inn Fields, WC2A 3PX London, United Kingdom

Edited by Kai Simons, European Molecular Biology Laboratory, Heidelberg, Germany, and approved April 6, 1998 (received for review November 13, 1997)

**ABSTRACT** The molecular mechanisms that coordinate cell morphogenesis with the cell cycle remain largely unknown. We have investigated this process in fission yeast where changes in polarized cell growth are coupled with cell cycle progression. The *orb6* gene is required during interphase to maintain cell polarity and encodes a serine/threonine protein kinase, belonging to the myotonic dystrophy kinase/*cot1/warts* family. A decrease in Orb6 protein levels leads to loss of polarized cell shape and to mitotic advance, whereas an increase in Orb6 levels maintains polarized growth and delays mitosis by affecting the p34<sup>cdc2</sup> mitotic kinase. Thus the Orb6 protein kinase coordinates maintenance of cell polarity during interphase with the onset of mitosis. *orb6* interacts genetically with *orb2*, which encodes the Pak1/Shk1 protein kinase, a component of the Ras1 and Cdc42-dependent signaling pathway. Our results suggest that Orb6 may act downstream of Pak1/Shk1, forming part of a pathway coordinating cell morphogenesis with progression through the cell cycle.

Eukaryotic cells establish and maintain polarized cellular domains that are essential for intracellular transport, cell differentiation, cell shape, and cell locomotion (1). Changes in cell shape and cell polarity occur during the cell cycle and in response to exogenous signals, but little is known about how cell morphogenesis is regulated or how polarized cell growth is integrated with the cell cycle. Fission yeast *Schizosaccharomyces pombe* cells grow in a polarized fashion that undergoes distinct changes during the cell cycle. Cells grow from one tip during G<sub>1</sub> and S-phase, and activate the second tip during early G<sub>2</sub>, becoming bipolar in growth (2). Actin dots are located at one end of the cell when cells are growing with one tip and are found at both ends when cells grow in a bipolar manner (3). At the onset of mitosis, polarized cell growth ceases, and actin relocates from the cell tips to the middle of the cell where an actin ring forms. Both activation of bipolar growth and onset of mitosis require the attainment of a minimal cell size (2, 4).

In fission yeast, cell polarity and morphology is thought to be regulated by the product of the *ras* homologue, *ras1*, acting upstream of the Cdc42 GTPase (5, 6). Ras1 and Cdc42 are part of a complex including Scd1, the putative Cdc42 guanine nucleotide exchange factor, and Scd2, a Src homology 3 (SH3) domain-containing protein (6). Cdc42 also has been shown to be associated with the serine/threonine kinase Pak1/Shk1, a homologue of mammalian PAKs kinases and the budding yeast Ste20 protein kinase (7). In fission yeast, *pak1/shk1* has been shown to be essential for cell viability and to participate in the

Ras1-dependent mating response pathway (8, 9). Although overexpression studies suggest a role for the Pak1/Shk1 kinase in the control of cell morphology (9), the molecular details of Pak1/Shk1-dependent regulation of the cytoskeleton have yet to be fully clarified.

We previously identified 19 genes that are required for various aspects of cell morphogenesis (10). The *orb6* gene is required to maintain cell polarity throughout the interphase period of the cell cycle. Mutants in *orb6* lose growth polarity and become spherical with disorganized microtubule arrays and delocalized actin dots (10). In this paper, we report the cloning of Orb6 and its identification as a fission yeast kinase closely related to a number of higher eukaryotic kinases, including *Drosophila warts* kinase, mammalian Rho-associated kinase, and the human myotonic dystrophy kinase (DMPK). We show that Orb6 is required for maintenance of cell polarity during interphase and to promote actin reorganization during morphological transitions. Moreover, we demonstrate that Orb6 also has another role during the cell cycle, specifically to delay onset of mitosis, suggesting that it functions to coordinate cell morphogenesis with the cell cycle. Finally, we show that *orb6* interacts genetically with Pak1/Shk1 protein kinase and that *pak1/shk1* is required for proper Orb6 intracellular localization. We propose that the Orb6 kinase acts downstream of a morphogenetic control pathway involving Cdc42 and Pak1/Shk1, which maintains the cell in a polarized state during interphase while simultaneously delaying the onset of mitosis.

## MATERIALS AND METHODS

**Strains and Growth Conditions.** The strains used in this paper were *orb6-25 ade6-M210 leu1-32h<sup>-</sup>*; *orb2-34 ade6-M210 leu1-32h<sup>-</sup>*; *wee1-50 leu1-32h<sup>-</sup>*; and *leu1-32h<sup>-</sup>*. Cells were cultivated in rich medium (yeast extract liquid or agar), or Edinburgh minimal medium liquid with the appropriate supplements (11), as indicated. Cells in liquid cultures had been growing exponentially for at least eight generations, at densities below 10<sup>7</sup> cells/ml, before observation.

**Libraries and Transformation Procedures.** The libraries used for cloning *orb6* and *orb2* were a cDNA library cloned in the plasmid Rep3X (29) and a genomic library in plasmid pDB248' (30). Cells were transformed by the lithium acetate procedure (11).

This paper was submitted directly (Track II) to the *Proceedings* office. Abbreviations: HA, hemagglutinin; DMPK, myotonic dystrophy kinase.

Data deposition: The sequence reported in this paper has been deposited in the GenBank database (accession no. AF009512).

†To whom reprint requests should be addressed. e-mail: fverde@chroma.med.miami.edu.

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. §1734 solely to indicate this fact.

© 1998 by The National Academy of Sciences 0027-8424/98/957526-6\$2.00/0  
PNAS is available online at <http://www.pnas.org>.

**Cloning of *orb6*<sup>+</sup>.** *orb6* mutants were transformed with a cDNA library controlled by the thiamine-repressible *nmt1* promoter (29); 145,000 colonies were screened for transformants showing reversion of lethality and morphological phenotype. The screen yielded 14 rescuing plasmids. Nine plasmids (pFV 10, 44, 39, 6, 75, 42, 23, 271, and 622) rescued both lethality and shape defects under low level of expression (in thiamine-containing medium). They all identified the same gene sequence. One plasmid, pFV10, was chosen and used for all successive experiments. The coding sequence was cloned in a REP6X plasmid (containing the *sup3-5* sequence) and integrated in an *orb6-25 ade6-704* strain. Thirty-four integrants were isolated and crossed to a wild-type strain: on average one mutant colony appeared for 500 colonies screened, indicating integration at the *orb6* locus and consistent with a 0.5–1 cM distance between the mutant gene and its wild-type integrated copy. As a control, one of these integrants (number 27) was crossed to an *orb6-25 ade6-704* strain: *orb6*<sup>+</sup> colonies were found to be adenine prototrophs, indicating that the rescuing sequence was linked to the *sup3-5* marker and excluding the possibility of reversion. Plasmid pFV10 contained a single continuous ORF of 1,410 bp.

**Deletion of *orb6*<sup>+</sup>.** We identified a genomic clone containing the *orb6* sequence again by complementation of *orb6* mutants. The genomic sequence, ≈10 kilobases long, was shown by PCR and Southern blotting analysis to contain the *orb6*<sup>+</sup> gene, with a 267-bp intron. The deletion construct, substituting the whole catalytic domain of *orb6* with the *ura4*<sup>+</sup> gene sequence, was built leaving a 2.9-kilobase 5' flanking sequence and 0.530-kilobase 3' flanking sequence.

**Cloning of *orb2*<sup>+</sup>.** To clone the *orb2* gene, we identified a 12-kilobase suppressing genomic fragment that was shown to integrate very closely to the *orb2-34* mutation. We identified the suppressing gene by transposon knockout (12) and sequenced outwardly from the transposon insertion site. The *orb2-34* mutation also was suppressed by transformation with *pak1/shk1* cDNA, which was a kind gift of J. Chernoff (Fox Chase Cancer Center, Philadelphia) (9).

**Cytology.** Cells were grown exponentially at 25°C for at least eight generations, then incubated at the indicated temperature for the indicated time. Immunofluorescence was performed as described (13). Cells were fixed in methanol and stained with mouse anti-actin antibody (Sigma) or anti-hemagglutinin (HA) primary antibody (Babco, Richmond, CA) and a CY3-conjugated anti-mouse secondary antibody. Cells then were immobilized on coverslips and observed using a Zeiss Axioskop microscope. Cells were photographed using a Bio-Rad 600 confocal microscope, with a 0.2-μm interval between successive pictures and analyzed using Bio-Rad software. For quantification of actin delocalization, the two-dimensional projections of the confocal microscope pictures were analyzed by using National Institutes of Health IMAGE software, measuring the intensity of immunofluorescence signal in different areas of the cell (% of intensity throughout the cell and at the cell tips). Ten cells were measured for each time point. Finally, the penetrance of the corresponding phenotype was quantified in the original sample, by counting 200 cells for each time point.

## RESULTS

***orb6* Encodes a Serine Threonine Protein Kinase.** Mutant *orb6* cells are round at restrictive temperature (Fig. 1*b*). Moreover, the number of cells undergoing the process of cell division increases from 14% to 60% after 5 hr at 36°C: 15% of these septa are aberrant (see Fig. 5*b*). The *orb6* gene was cloned by complementation. *orb6* mutants were transformed with a cDNA library controlled by the thiamine-repressible *nmt1* promoter (29); 145,000 colonies were screened for transformants showing a rescue of the morphological phenotype. The screen yielded 14 rescuing plasmids. Nine plasmids contained the same gene sequence that was shown by integra-




FIG. 1. Cloning of *orb6*<sup>+</sup>. (a) *leu1-32*, pRep3X. (b) *orb6-25 leu1-32*, pRep3X. (c) *orb6-25 leu1-32*, pRep3X *orb6*<sup>+</sup>. Cells were exponentially grown at 25°C for at least eight generations, then shifted at 36°C for 5 hr.

tion to encode the *orb6*<sup>+</sup> gene (see *Materials and Methods*). These plasmids complemented the shape defects of the *orb6-25* mutant with low level of expression (in thiamine-containing medium). They contained a single continuous ORF predicted to encode a protein of 469 amino acids; its amino acid sequence was found to encode a member of a serine/threonine protein kinase family, including *Neurospora cot1* that affects hyphal morphology (14), *Drosophila warts* that controls cell shape and cell proliferation (15, 16), Rho-associated kinase (17–19), and the human DMPK (20) (Fig. 2*A*).

**The Function of the *orb6* Gene Is Essential.** A genomic clone containing the *orb6*<sup>+</sup> gene was isolated from a fission yeast genomic library (30), again by complementation of the *orb6* mutation. A deletion mutation was built, which removed most of the *orb6*<sup>+</sup> gene and replaced it with the *ura4*<sup>+</sup> gene. This




FIG. 2. The *orb6*<sup>+</sup> gene. (A) Sequence comparison of Orb6 and homologous proteins. An homology search using EMBL and GenBank databases identified *S. cerevisiae* 1050791, *Neurospora crassa cot1* (14), human Ndr (23), *Drosophila warts* (15, 16), human PK428, Rho-associated kinase (17–19), and human DMPK (20). (B) *orb6*<sup>+</sup> gene deletion. Spores were plated on yeast extract plates and allowed to germinate for 16 hr at 32°C. (a) *ade- leu1-32 ura4-D18* microcolony. (b) *orb6::ura4+ ade- leu1-32 ura4-D18* microcolony. (c) *orb6::ura4+ ade- leu1-32 ura4-D18* microcolony after 24 hr at 32°C. (Bar = 5 μm.)




Fig. 3. *Orb6*<sup>+</sup> gene dosage effects. *orb6::ura4<sup>+</sup> ade- leu1-32 ura4-D18* cells containing integrated HA-tagged *orb6*<sup>+</sup> were grown exponentially for at least eight generations at 32°C, then thiamine was added to the culture. (a and d) Cells grown in the absence of thiamine. (b and e) 7.5 hr after thiamine addition. (c and f) 12 hr after thiamine addition. Cells were photographed immediately (a–c) or fixed and stained for actin (d–f). (Bar, a–c = 5 μm.; d–f = 10 μm.)

construct deleted all 11 kinase domains in the *orb6* gene, leaving 106 bp at the 5' end and 224 bp at the 3' end. The *ura4<sup>+</sup>* gene was substituted for the whole *orb6* kinase domain by the one-step gene replacement method in diploids cells using this deletion construct (11). Deletion of the *orb6* gene led to cell lethality, establishing that *orb6* is essential. Spores deleted for the *orb6* gene germinate to form spherical cells that undergo 3–4 cell divisions before lysing (Fig. 2B). Lysis is not prevented by addition of 1 M sorbitol as an osmotic stabilizer.

***orb6* Is Necessary for Actin Reorganization After Mitosis and During Activation of Bipolar Growth.** To investigate *Orb6* function more fully we constructed a triple HA-tagged *orb6* gene under control of the thiamine repressible *nmt1* promoter. This construct was integrated into the genome of a strain in which the endogenous *orb6*<sup>+</sup> gene was replaced by the *ura4<sup>+</sup>* marker (*orb6::ura4<sup>+</sup>*). When *orb6-HA* was expressed in the absence of thiamine, this strain was perfectly viable, showing a generation time of 2 hr and 50 min (at 32°C) as compared with the generation time of 2 hr and 40 min of the *ade6-M210* control (determined by monitoring the optical density and the cell number increase of the cell culture). Cells displayed a normal cell morphology with a correctly localized actin cytoskeleton (Fig. 3 a and d). In contrast, when thiamine was added to repress *orb6-HA* expression, we observed changes in cell morphology and a reorganization of the actin cytoskeleton. Early in the time course, between 4 and 8 hr after thiamine addition, cell length shortened (Fig. 3b) and actin became less




Fig. 4. Effects of *orb6* overexpression. (a–c) Overexpression of HA-tagged *orb6*<sup>+</sup> in a *leu1-32* strain (experiment 1 in Table 1). (a) *leu1-23* pRep3X. (b) *leu1-23* cells, carrying integrated HA-tagged *orb6*<sup>+</sup>, were grown exponentially for at least eight generations at 32°C in the absence of thiamine. (c) *leu1-32* cells, carrying HA-tagged *orb6*<sup>+</sup> on the multicopy plasmid pRep3X, were grown in the absence of thiamine for 18 hr at 32°C. (d–f) Overexpression of HA-tagged *orb6*<sup>+</sup> in a *wee1-50* mutant strain. Cells were grown exponentially at 25°C in the presence of thiamine for at least eight generations; thiamine was washed away and cells were grown in the absence of thiamine at 25°C for 16 hr, then shifted at 36°C for 5 hr. (d) *leu1-32*, pRep3X *orb6HA*. (e) *wee1-50 leu1-32* pRep3X. (f) *wee1-50 leu1-32* pRep3X *orb6HA*. (g) Western blot comparing the levels of expression of *orb6-HA* in the experiments described above, as compared with the control tubulin: lane 1, *leu1-32* with an integrated copy of *orb6-HA*; lane 2, *leu1-23* pRep3X; lane 3, *leu1-32* pRep3X *orb6 HA*; lane 4, *leu1-32* pRep3X *orb6 K122A HA*; lane 5, *wee1-50 leu1-32* pRep3X; lane 6, *wee1-50 leu1-32* pRep3X *orb6-HA*. (Bar = 5 μm.)

localized at the cell tips. Delocalization of 61.5% of actin was found in 67% of cells, as compared with time zero where 91% of actin was found localized at the cell tips (see *Materials and Methods*) (Fig. 3e). Most cells showed a monopolar pattern of growth (see below). Actin dots also were found in a ring pattern at the new end after septation in 10% of cells, suggesting a delay in actin reorganization after mitosis (Fig. 3e, arrow). Later in the time course, 12 hr after thiamine addition, cells became spherical and actin was found dispersed throughout the cell cortex in 81% of cells (Fig. 3 c and f), with the exception of actin clusters, which persisted in the area where

Table 1. *orb6* gene dosage effects

Strain	Plasmid	Cell length at division*	% wt length
(1) <i>leu1-32</i>	Multicopy plasmid (Rep3X)	14.6 ± 1.0	100%
(1) <i>leu1-32</i>	Integrand <i>nmt1 orb6HA</i>	17.8 ± 1.9	121%
(1) <i>leu1-32</i>	Multicopy <i>nmt1 orb6HA</i>	26.6 ± 4.9	181%
(1) <i>leu1-32</i>	Multicopy <i>nmt1 orb6HA ala122</i>	13.7 ± 2.7	93%
(2) <i>leu1-32</i>	Multicopy plasmid (Rep3X)	14.2 ± 1.0	100%
(2) <i>leu1-32</i>	Multicopy <i>nmt1 orb6HA</i>	28.9 ± 5.0	203%
$\Delta orb6$	Integrand <i>nmt1 orb6HA</i>	18.0 ± 1.3	122%
$\Delta orb6$	Integrand <i>nmt1 orb6HA (+th., 5h.)</i>	12.4 ± 0.7	84%
$\Delta orb6$	Integrand <i>nmt1 orb6HA (+th., 7.5h.)</i>	9.5 ± 0.6	65%
<i>leu1-32</i>	Multicopy <i>nmt1 orb6HA</i>	20.0 ± 5.6	137%
<i>wee1-50 leu1-32</i>	Multicopy plasmid (Rep3X)	7.4 ± 0.9	50%
<i>wee1-50 leu1-32</i>	Multicopy <i>nmt1 orb6HA</i>	7.5 ± 0.9	51%

\*Thirty cells at division were measured for each sample. The mean values ± SD are shown.




FIG. 5. Overexpression of Wee1 in *orb6* mutants and actin localization in *orb6-25* and *wee1-50* mutants. (a–d) Overexpression of Wee1 in *orb6* mutants. (a) *leu1-32h<sup>-</sup>* pRep3x. (b) *orb6-25 leu1-32h<sup>-</sup>* pRep3x. (c) *leu1-32h<sup>-</sup>* pRep3x *wee1<sup>+</sup>*. (d) *orb6-25 leu1-32h<sup>-</sup>* pRep3x *wee1<sup>+</sup>*. (e) *wee1-50h<sup>-</sup>*. Cells were grown exponentially at 25°C in the presence of thiamine for at least eight generations; thiamine was washed away, and cells were grown in the absence of thiamine at 25°C for 16 hr, then shifted at 36°C for 5 hr. (f–h) Actin localization in *orb6-25* and *wee1-50* mutants. (f) *leu1-32h<sup>-</sup>*. (g) *orb6-25 leu1-32h<sup>-</sup>*. (h) *wee1-50h<sup>-</sup>*. (Bar = 5  $\mu$ m.)

the actin ring had formed during the previous cell division (in 12% of cells) (Fig. 3f, arrow). These observations indicate that when Orb6 levels decrease, the actin cytoskeleton becomes disorganized. Moreover, early in the time course, the reorganization of actin during the cell cycle transitions between monopolar and bipolar growth and between mitosis and interphase is defective, as indicated by an increase in the number of cells growing with one tip only (68% compared with 28% in *orb6<sup>+</sup>* cells) or undergoing the process of cell division (24% compared with 14% in *orb6<sup>+</sup>* cells) (Fig. 3b). We conclude that the Orb6 protein kinase is required to maintain polarity of the actin cytoskeleton during interphase and to promote actin reorganization both after mitosis and during activation of bipolar growth.

This experiment also revealed that Orb6 delays the onset of mitosis (Table 1). Five hours after switching off the *orb6* gene, cell length was reduced to 84% of *orb6<sup>+</sup>* cells, and at 7.5 hr cell length was reduced to 65%. This reduction in average cell length was found to be highly significant ( $P < 0.05$ ). Cell width was unchanged, indicating that these cells were advanced into mitosis at a reduced cell size. Shortening of cell length was not

caused by a reduction in cell growth rate, which was unchanged until 8.5 hr after thiamine addition (data not shown).

**Orb6 Acts as a Dose-Dependent Inhibitor of Mitosis.** To confirm the function of Orb6 in the regulation of mitosis, we overexpressed *orb6-HA* in wild-type cells (Fig. 4 and Table 1). Expressing *orb6-HA* from a single-copy integrant caused cells to divide at a larger cell size (Fig. 4b). Expression of *orb6-HA* on a multicopy plasmid using the *nmt1* promoter led to an increase of cell length at division, indicating that onset of mitosis was substantially delayed (Fig. 4c). FACS analysis established that these cells were delayed in the G<sub>2</sub> phase of the cell cycle (data not shown). This delay of mitotic onset required Orb6 protein kinase activity because an *orb6* kinase inactive mutant (K122A) did not induce cell elongation when overexpressed (Table 1). *orb6-HA* overexpression in a *wee1-50* strain, which is defective in the Wee1 tyrosine protein kinase that phosphorylates and inhibits p34<sup>cdc2</sup>, failed to increase cell length at division (Fig. 4d–f and Table 1). This experiment demonstrated that the Orb6-dependent delay over mitosis acts through the p34<sup>cdc2</sup> mitotic kinase. Overexpression of *wee1<sup>+</sup>* in *orb6-25* mutant cells did not suppress their morphological defects (Fig. 5 a–d). Moreover, *wee1-50* and *orb6-25* mutants show a different morphological phenotype at restrictive temperature: *orb6-25* cells are round and their actin cytoskeleton is dispersed (Fig. 5 b and g) whereas *wee1-50* cells are very short, but still grow in a polarized fashion (Fig. 5e) and show a polarized actin cytoskeleton (Fig. 5h). These experiments indicate that Orb6 kinase has a specific function in the control of cell morphology, maintaining a polarized actin cytoskeleton and promoting polarized cell growth, while delaying the onset of mitosis.

**Orb6 Localizes to Sites of Cell Growth.** The intracellular location of Orb6-HA protein was determined by using the Orb6-HA integrant strain (Fig. 6). Immunofluorescence showed that Orb6-HA protein is enriched at the cell tips during interphase (Fig. 6 a and e). During mitosis and cytokinesis Orb6-HA protein disappears from the tips and is found in the region of the developing septum (Fig. 6 b and f), indicating that Orb6-HA protein localizes to areas of cellular growth. This observation was confirmed by staining Orb6-HA in *cdc25<sup>ts</sup>* and *orb2<sup>ts</sup>* mutant cells. Both tips were stained in the *cdc25-22<sup>ts</sup>* mutant cells, which were arrested in G<sub>2</sub> with both ends growing (Fig. 6c), whereas in the *orb2-34* mutant, which cannot activate bipolar growth and only grows at one end (10) (Fig. 6h), the Orb6-HA protein was found only at one growing tip (Fig. 6d).




FIG. 6. *orb6<sup>+</sup>* gene product intracellular localization in the cell cycle. (a, b, e, and f) *orb6::ura4<sup>+</sup> ade-leu1-32 ura4-D18* cells containing integrated HA-tagged *orb6<sup>+</sup>* were grown exponentially at 32°C in the absence of thiamine, then fixed and stained with an antibody recognizing the HA tag (a and b) and with 4',6-diamidino-2-phenylindole (e and f). (g) Control staining of *orb6::ura4<sup>+</sup> ade-leu1-32 ura4-D18* cells containing integrated untagged *orb6<sup>+</sup>*. (c) *cdc25-22 leu1-32 Rep3X orb6 HA* cells were grown exponentially at 25°C in the presence of thiamine for at least eight generations; thiamine was washed away and cells were grown in the absence of thiamine at 25°C for 16 hr, then shifted at 36°C for 4 hr. (d) *orb2-34 leu1-32 pRep3X orb6 HA* at 32°C; cells were stained 14 hr after thiamine depletion, which is about 2 hr after derepression of the *nmt1* promoter. (h) *orb2-34 leu1-32* stained for actin at 32°C. (Bar = 10  $\mu$ m.)




FIG. 7. Cloning of the *orb2*<sup>+</sup> gene. (a) top, *ade6-M210 leu1-32h*<sup>-</sup>; left, *orb6-25 ade6-M210 leu1-32h*<sup>-</sup>; right, *orb2-34 orb6-25 ade6-M210 leu1-32h*<sup>-</sup>. Cells were grown at 32°C. (b) *ade6-M210 leu1-32h*<sup>-</sup>, Rep3X. (c) *orb2-34 ade6-M210 leu1-32h*<sup>-</sup>, pRep3X. (d) *orb2-34 ade6-M210 leu1-32h*<sup>-</sup> pDB248'-8 (genomic fragment). (e) *orb2-34 ade6-M210 leu1-32h*<sup>-</sup> pRep3X *pak1/shk1*<sup>+</sup>. (f) *orb2-34 ade6-M210 leu1-32h*<sup>-</sup> pRep3X *orb6*<sup>+</sup>. Cells were grown exponentially at 25°C in the presence of thiamine for at least eight generations; thiamine was washed away and cells were grown in the absence of thiamine at 25°C for 16 hr, then shifted at 36°C for 6 hr. (Bar = 5 μm.)

***orb6* Interacts Genetically with *orb2*.** *orb6* mutants were found to be synthetically lethal with *orb2* mutants (10). Synthetic lethality interactions were determined by tetrad analysis, and phenotypes were observed at 25°C, 32°C, and 36°C. At 32°C mutants in either *orb6* or *orb2* are viable, but the double-mutant *orb6-25 orb2-34* is lethal, with the cells rapidly undergoing lysis (Fig. 7a). *orb2* mutants are unable to activate bipolar growth and grow with only one tip (10). Analysis of the actin cytoskeleton in *orb2* mutants revealed that actin is localized only at one pole of growth (Fig. 6h) and that its reorganization after mitosis is delayed (data not shown). This phenotype is reminiscent of the one observed when expression levels of Orb6 decrease, suggesting that Orb2 and Orb6 might fulfill a similar function.

***orb2* Encodes the Protein Kinase Pak1/Shk1.** Given this interaction with *orb6*, *orb2* was cloned by complementation (Fig. 7c and d), and its sequence confirmed that *orb2* is identical to the previously identified *pak1/shk1* gene (8, 9). The *orb2-34* mutant was also suppressed by *pak1/shk1* cDNA (Fig. 7e). The *pak1/shk1* gene encodes a protein kinase that interacts with Cdc42 and is similar to the CDC42 and RAC1-interacting PAK1 protein kinase identified in mammalian cells and STE20 found in budding yeast cells (7). Overexpression of *orb6*<sup>+</sup> partially suppressed the *pak1/shk1/orb2*<sup>ts</sup> mutant shape defect (Fig. 7f). Overexpression of *pak1/shk1*<sup>+</sup> did not rescue the *orb6*<sup>ts</sup> mutant phenotype (data not shown), but did induce cell elongation in wild-type cells (data not shown). These results suggest that the Orb6 kinase might function downstream of the Pak1/Shk1 protein kinase. This finding also is supported by the observation that in *pak1/shk1/orb2*<sup>ts</sup> mutants, which are defective in the activation of the second tip, the Orb6-HA protein was found only at the one growing tip, suggesting that *pak1/shk1* is required to bring about bipolar growth and to localize Orb6 protein correctly (Fig. 6d).

## DISCUSSION

**Orb6 Belongs to a Family of Kinases Essential for Morphological Control.** In the present study, we report the cloning and characterization of a fission yeast gene, *orb6*, which is required for maintenance of cell polarity during interphase and to promote actin reorganization during morphological transi-




FIG. 8. Orb6 kinase is proposed to be part of a regulatory cascade contributing to the mitotic control and responding to the Ras1 and Cdc42-dependent morphological control pathway via the Pak1/Shk1 kinase. Orb6 kinase regulates independently cell morphogenesis and the cell cycle, mediating its effect on p34<sup>cdc2</sup> kinase directly through the Wee1 protein kinase or through some other mechanism dependent on p34<sup>cdc2</sup> tyrosine phosphorylation.

tions. Orb6 encodes a serine/threonine kinase and belongs to a kinase family that includes several serine/threonine kinases whose catalytic domains are highly conserved. These kinases are more closely related to each other than to their closest relatives, the cyclic nucleotide-dependent protein kinases, and, at least in the case of the DMPK and Rho-associated kinase, show distinct biochemical properties (17–19, 21). *Warts*, *orb6*, and *cot1* have been identified by mutagenesis screens and are required for maintenance of cell shape and polarity. Rho-associated kinase is regulated and translocated to the plasma membrane by rho-GTPase (17–19), which is involved in the control of various aspects of the actin cytoskeleton. Rho-associated kinase is the only member of this kinase family with known substrates and has been shown to have a role in the regulation of actin-myosin interaction by modulating myosin phosphorylation (22). Only one related kinase has been identified in the budding yeast, *Saccharomyces cerevisiae*, suggesting that this kinase family might have diverged in higher Eukaryotes. We presently are investigating to determine whether *S. pombe orb6* and *S. cerevisiae* 1050791 are functionally homologous to mammalian Rho-associated kinase and whether the modality of regulation by rho-GTPase is conserved.

**Orb6 Acts as a Dose-Dependent Inhibitor of Mitosis.** Orb6 also acts as a dose-dependent inhibitor of mitosis, a function requiring the activity of *wee1* kinase, an inhibitor of p34<sup>cdc2</sup> kinase activation. Thus Orb6 kinase has two roles, to control cell shape and to regulate G<sub>2</sub>/M transition, possibly through the *wee1* pathway, or some other mechanism dependent on Cdc2 tyrosine phosphorylation (Fig. 8). Although immunofluorescence microscopy showed that Orb6 is predominantly localized in the cytoplasm, Orb6 has a conserved sequence (KDKMATWKKNRR) that has been shown to contain the nuclear localization signal in Ndr kinase (23), so possibly it may directly interact with Wee1. These findings also suggest an interesting functional similarity with *Drosophila warts*. The *warts* gene is one of more than 50 genes in which mutations give an overgrowth phenotype, and which therefore are considered tumor suppressor genes (24). Homozygous loss of *warts* (*wts*) in somatic clones results in cell-autonomous formation of epithelial tumors (15, 16), whose cells also show an hypertro-

phy of the apical ends. Thus, the *warts* gene product is important for controlling cellular morphogenesis as well as proliferation. Myotonic dystrophy is also sometimes associated with calcifying epitheliomas, neurofibromas, and parathyroid adenomas, suggesting that the DMPK also may have a role in the control of cell proliferation (25).

What is the role of these kinases in the regulation of the cell cycle and what is its functional significance? One possibility is that the deregulation of cell proliferation is an indirect consequence of cell shape alteration. Our results, though, suggest that a decrease of Orb6 levels leads to mitotic advance before an alteration of cell shape and that Orb6 overexpression can delay onset of mitosis without altering cell morphology. This finding might point to a novel pathway in the control of mitotic onset and in the coordination of cell growth and cell proliferation. Future work will clarify the extent of functional similarity between *warts* and *orb6*. If this pathway is conserved, the identification of *orb6* in fission yeast, which allows the dissection of the cell cycle machinery by the use of cell cycle (*cdc*) mutants, may provide the opportunity to easily study the mechanism of cell cycle regulation by these kinases.

**Orb6 Protein Kinase Interacts Genetically with *pak1/shk1*.** To identify other partners in Orb6 function, we cloned *orb2*, because *orb2* mutants showed a strong synthetic lethality interaction with *orb6* mutants and their phenotype displayed similarities to the effects of *orb6* loss of function. Cloning of *orb2* showed that *orb2* is identical to the previously identified *pak1/shk1* gene (8, 9). The *pak1/shk1* gene encodes a protein kinase that interacts with Cdc42, and is similar to the CDC42 and RAC1-interacting PAK1 protein kinase identified in mammalian cells and STE20 found in budding yeast cells (7). Ras1, Cdc42, and Pak1/Shk1 have a regulatory role in fission yeast cell morphogenesis and are thought to function in an hierarchical order whereby ras1 GTPase interacts functionally with Cdc42 GTPase (6) and Pak1/Shk1 kinase acts as an effector of Cdc42 (8, 9).

We identified temperature-sensitive mutants of *pak1/shk1*, which confirm the morphological role of *pak1/shk1* previously indicated by overexpression studies (9). Moreover, our study suggests that Pak1/Shk1 also might have a function in cell cycle control because *pak1/shk1/orb2<sup>ts</sup>* cells are short in cell length, and overexpression of *pak1/shk1* induces cell elongation in wild type (data not shown). These results are consistent with the observation that Skb1, which interacts with Pak1/Shk1, also causes cell elongation when overexpressed (26). The fact that *orb6* overexpression partially suppresses the *orb2<sup>ts</sup>* phenotype suggests that the Orb6 kinase might function downstream of the Pak1/Shk1 protein kinase, in a pathway that controls and coordinates maintenance of cell polarity with progression through the cell cycle (Fig. 8). Alternatively, *pak1/shk1* and *orb6* might represent parallel pathway with similar and partially overlapping functions. The former model, though, is supported also by the observation that in *pak1/shk1/orb2<sup>ts</sup>* mutants, which are defective in the activation of the second tip, the Orb6-HA protein is found only at the one growing tip, whereas another marker of cell polarity, Tea1, is found localized at both ends (27). This finding indicates that *pak1/shk1* is required to bring about bipolar growth and to localize Orb6 protein correctly. Future work will address the interaction between Pak1/Shk1 and Orb6 and will clarify whether the Orb6 kinase activity or cortical localization is dependent on Pak1/Shk1.

## CONCLUSION

We conclude that Orb6 has two roles during the cell cycle: to maintain polarized cell growth and to delay onset of mitosis.

We propose that the Orb6 kinase acts downstream of a morphogenetic control pathway involving Cdc42 and Pak1/Shk1, which maintains the cell in a polarized state during interphase while delaying the onset of mitosis (Fig. 8). When the cell has reached the appropriate size and is ready to undergo mitosis, a reduction in Orb6 kinase activity could remove an inhibitory signal over the p34<sup>cdc2</sup> mitotic kinase and thus contribute to bringing about the onset of mitosis. The similarity of *orb6* with *Drosophila warts*, which is thought to function as a tumor suppressor (15, 16) and other genes in mammalian cells that affect cell shape (22, 28), suggests that its function in coordinating cell morphogenesis with the cell cycle may be conserved in higher eukaryotes. Future work will compare the functional properties of this kinase family in different organisms and elucidate their role in the mechanisms integrating cell morphogenesis and proliferation.

We thank Dr. J. Chernoff for kindly providing the *pak1/shk1* cDNA. We thank Dr. R. Myers, Dr. R. Assoian, and Z. Chen for critically reading the manuscript, and Dr. P. Salas for help with the confocal microscope. This work has been supported by an European Economic Community and an Imperial Cancer Research Fund fellowship and by the Sylvester Comprehensive Cancer Center (University of Miami).

1. Drubin, D. G. & Nelson, W. J. (1996) *Cell* **84**, 335–344.
2. Mitchison, J. M. & Nurse, P. (1985) *J. Cell Sci.* **75**, 357–376.
3. Marks, J. & Hyams, J. S. (1985) *Eur. J. Cell Biol.* **110**, 417–425.
4. Nurse, P. (1975) *Nature (London)* **256**, 547–551.
5. Miller, P. J. & Johnson, D. I. (1994) *Mol. Cell. Biol.* **14**, 1075–1083.
6. Chang, E. C., Barr, M., Wang, Y., Jung, V., Xu, H. P. & Wigler, M. H. (1994) *Cell* **79**, 131–141.
7. Sells, M. A. & Chernoff, J. (1997) *Trends Cell Biol.* **7**, 162–167.
8. Marcus, S., Polverino, A., Chang, E., Robbins, D., Cobb, M. H. & Wigler, M. H. (1995) *Proc. Natl. Acad. Sci. USA* **92**, 6180–6184.
9. Otilite, S., Miller, P. J., Johnson, D. I., Creasy, C. L., Sells, M. A., Bagrodia, S., Forsburg, S. L. & Chernoff, J. (1995) *EMBO J.* **14**, 5908–5919.
10. Verde, F., Mata, J. & Nurse, P. (1995) *J. Cell Biol.* **131**, 1–10.
11. Moreno, S., Klar, A. & Nurse, P. (1991) *J. Gen. Microbiol.* **194**, 795–823.
12. Morgan, B. A., Conlon, F. L., Manzanares, M., Millar, J. B., Kanuga, N., Sharpe, J., Krumlauf, R., Smith, J. C., & Sedgwick, S. G. (1996) *Proc. Natl. Acad. Sci. USA* **93**, 2801–2806.
13. Hagan, I. M. & Hyams, J. S. (1988) *J. Cell Sci.* **89**, 343–357.
14. Yarden, O., Plamann, M., Ebbole, D. J. & Yanofsky, C. (1992) *EMBO J.* **11**, 2159–2166.
15. Justice, R. W., Zilian, O., Woods, D. F., Noll, M. & Bryant, P. J. (1995) *Genes Dev.* **9**, 534–546.
16. Xu, T., Wang, W., Zhang, S., Stewart, R. A. & Yu, W. (1995) *Development (Cambridge, U.K.)* **121**, 1053–1063.
17. Leung, T., Manser, E., Tan, L. & Lim, L. (1995) *J. Biol. Chem.* **270**, 29051–29054.
18. Matsui, T., Amano, M., Yamamoto, T., Chihara, K., Nakafuku, M., Ito, M., Nakano, T., Okawa, K., Iwamatsu, A. & Kaibuchi, K. (1996) *EMBO J.* **15**, 2208–2216.
19. Ishizaki, T. M. M., Fujisawa, K., Okawa, K., Iwamatsu, A., Fujita, A., Watanabe, N. S. Y., Kakizuka, A., Morii, N. & Narumiya, S. (1996) *EMBO J.* **15**, 1885–1893.
20. Brook, J. D., McCurrach, M. E., Harley, H. G., Buckler, A. J., Church, D., Aburatani, H., Hunter, K., Stanton, V. P., Thirion, J. P., Hudson, T., *et al.* (1992) *Cell* **68**, 799–808.
21. Dunne, P. W., Walch, E. T. & Epstein, H. F. (1994) *Biochemistry* **6**, 10809–10814.
22. Kimura, K. I. M., Amano, M., Chihara, K., Fukata, Y., Nakafuku, M., Yamamori, B., Feng, J., Nakano, T., Okawa, K., Iwamatsu, A. & Kaibuchi, K. (1996) *Science* **273**, 245–248.
23. Millward, T. C. P. & Hemmings, B. A. (1995) *Proc. Natl. Acad. Sci. USA* **92**, 5022–5026.
24. Watson, K. L., Justice, R. W. & Bryant, P. J. (1994) *J. Cell. Sci.* **18**, Suppl., 19–33.
25. Hamshere, M. G. B. (1996) *Trends Genet.* **12**, 332–334.
26. Gilbreth, M., Yang, P., Wang, D., Frost, J., Polverino, A., Cobb, M. H. & Marcus, S. (1996) *Proc. Natl. Acad. Sci. USA* **93**, 13802–13807.
27. Mata, J. & Nurse, P. (1997) *Cell* **89**, 939–949.
28. Ishizaki, T. N. M., Fujisawa, K., Maekawa, M., Watanabe, N., Saito, Y. & Narumiya, S. (1997) *FEBS Lett.* **404**, 118–124.
29. Kelly, T. J., Martin, G. S., Forsburg, S. L., Stephen, R. J., Russo, A. & Nurse, P. (1993) *Cell* **74**, 371–382.
30. Hirano, T., Hiraoka, Y. & Yanagida, M. (1988) *J. Cell Biol.* **106**, 1171–1183.