



Flowering plant composition shapes pathogen infection intensity and reproduction in bumble bee colonies

Lynn S. Adler^{a,1}, Nicholas A. Barber^b, Olivia M. Biller^c, and Rebecca E. Irwin^d

^aDepartment of Biology, University of Massachusetts, Amherst, MA 01003; ^bEcology Program Area, Department of Biology, San Diego State University, San Diego, CA 92182; ^cDepartment of Occupational Therapy, Thomas Jefferson University, Philadelphia, PA 19107; and ^dDepartment of Applied Ecology, North Carolina State University, Raleigh, NC 27695

Edited by Nils Chr. Stenseth, University of Oslo, Oslo, Norway, and approved April 10, 2020 (received for review January 3, 2020)

Pathogens pose significant threats to pollinator health and food security. Pollinators can transmit diseases during foraging, but the consequences of plant species composition for infection is unknown. In agroecosystems, flowering strips or hedgerows are often used to augment pollinator habitat. We used canola as a focal crop in tents and manipulated flowering strip composition using plant species we had previously shown to result in higher or lower bee infection in short-term trials. We also manipulated initial colony infection to assess impacts on foraging behavior. Flowering strips using high-infection plant species nearly doubled bumble bee colony infection intensity compared to low-infection plant species, with intermediate infection in canola-only tents. Both infection treatment and flowering strips reduced visits to canola, but we saw no evidence that infection treatment shifted foraging preferences. Although high-infection flowering strips increased colony infection intensity, colony reproduction was improved with any flowering strips compared to canola alone. Effects of flowering strips on colony reproduction were explained by nectar availability, but effects of flowering strips on infection intensity were not. Thus, flowering strips benefited colony reproduction by adding floral resources, but certain plant species also come with a risk of increased pathogen infection intensity.

hedgerows | pathogen transmission | pollinator decline | pollinator habitat | wildflower strips

Community composition changes species dynamics, including the probability of disease transmission between hosts. For pollinators, understanding how plants shape disease transmission is important because pollination services contribute an estimated US\$235 to 577 billion to our global economy (1) and pathogens have been implicated as one of the factors underlying pollinator declines (2). Governments, industry, and private individuals worldwide are investing in planting pollinator-friendly habitat (called “flowering strips” hereafter to encompass any supplemental floral resources) to mitigate pollinator declines. These habitats can increase pollinator species richness and abundance (e.g., refs. 3–6) and sometimes pollination and crop yield (7–9). However, floral cover by particular plant groups, such as high-quality forage, can be more important than overall floral cover for bumble bee family lineage survival across years (10). Plant species vary in the amount and quality of resources for pollinators (e.g., refs. 11 and 12), and particular plant groups can play larger roles than overall plant diversity for bee colony growth (13). Thus, some plant species or groups may be more effective than others for managing pollinator health.

In addition to providing resources, plants can be sites of disease transmission between pollinators (e.g., refs. 14–16). However, the role of plant species composition in shaping parasite or pathogen infection in pollinators is largely unknown. Sown wildflower fields increased prevalence of several bee pathogens as well as bee abundance in landscapes with few seminatural elements (17). In an observational study across 19 urban gardens,

the number of trees and shrubs was positively correlated with phorid fly parasitism in both honey and bumble bees (18). In another study, the prevalence of deformed wing virus and black queen cell virus was higher in bumble bees and on flowers near honey bee apiaries, suggesting that flowers are the site of virus transmission from commercial honey bees to wild bumble bees (19). All of these studies suggest that floral resources can increase both bee abundance and risks of pathogen or parasite infection, but we do not yet know whether plant species composition plays significant roles in shaping bee pathogen infection.

Variation in floral traits within and among plant species can change the likelihood of vectoring or transmitting pathogens or parasitic mites (14, 15, 20, 21), and such variation can have consequences for disease transmission dynamics (22). In particular, a recent study found fourfold variation across 14 plant species in transmission of the gut pathogen *Criethidia bombi* to foraging bumble bees (*Bombus impatiens*) (20), and defecation on flowers by infected bees varied with plant species (23). However, we do not know whether these individual dynamics scale up to plant community consequences for bumble bee colony-level pathogen infection and reproduction.

The role of plant species in shaping infection intensity could be influenced by bee behavior. If infected bees increase visitation

Significance

Pollinator decline affects food security, and pollinators are threatened by stressors including pathogens and insufficient food resources. Flowering strips are increasingly planted to increase pollinator abundance and diversity in agricultural settings, but flowers can also be disease transmission sites between pollinators. However, the effect of plant species composition on bee disease is unknown. We compared the effect of flowering strips with high- or low-infection plant species, or no flowering strips, on bee infection and reproduction in tents. Using high-infection flowering strips doubled bee infection intensity compared to low-infection flowering strips. However, bee reproduction was higher with any flowering strips. Thus, floral resources in flowering strips benefited bees, but certain plants also come with a risk of increased pathogen infection intensity.

Author contributions: L.S.A. and R.E.I. designed research; L.S.A. and O.M.B. performed research; N.A.B. analyzed data; and L.S.A. wrote the paper.

The authors declare no competing interest.

This article is a PNAS Direct Submission.

Published under the PNAS license.

Data deposition: All data and R scripts are deposited in Dryad (<https://doi.org/10.5061/dryad.p2ngf1vn1>).

¹To whom correspondence may be addressed. Email: lsadler@umass.edu.

This article contains supporting information online at <https://www.pnas.org/lookup/suppl/doi:10.1073/pnas.2000074117/-DCSupplemental>.

First published May 11, 2020.

to antimicrobial plant species as a form of self-medication (24), such plant species could play a larger role than predicted in disease dynamics. Alternatively, antimicrobial plant species may be less effective than expected if pathogens manipulate host behavior to avoid such plants (25). Sunflower (*Helianthus annuus*) has pollen that dramatically reduces *C. bombi* in *B. impatiens* (26, 27) and several plant species produce nectar with secondary compounds that can reduce pathogens (28–30), although such effects are not always consistent (31, 32). Only a few studies have assessed whether infection alters bee preference. In the field, infected *B. impatiens* and *Bombus vagans* had greater preference than uninfected bees for inflorescences with high nectar iridoid glycosides that can reduce pathogen infection (33). However, a laboratory study with *Bombus terrestris* found only weak evidence that infected bees had increased preference for nectar nicotine compared to uninfected bees (28). Thus, there are conflicting results across species and compounds, and very few data overall to assess whether infection changes foraging preferences.

We assessed whether flowering strip species composition would affect pathogen infection intensity and bumble bee colony reproduction, and whether flowering strips and infection affected foraging behavior and pollination services. We designed our experiment in an agricultural context because flowering strips are increasingly used to promote pollinator abundance and diversity and enhance pollination services in agriculture, although results are also relevant for natural ecosystems. We used tent mesocosms with canola as a focal crop and included three flowering strip treatments: canola only, high-infection strip, and low-infection strip, crossed with presence/absence of infection. This experiment assesses the consequences of specific plant communities for bee colony infection and reproduction, which is critical for making informed decisions about pollinator habitat management as well as understanding how plant community composition structures species interactions.

Results

Pathogen Infection. *C. bombi* cells replicated in hosts; in the infected treatment final raw counts averaged 39.7 ± 30.4 cells per $0.02 \mu\text{L}$ (mean \pm SD; range 0 to 130), nearly 100 times more *C. bombi* cells than the initial inoculation (34).

We asked how flowering strips affected infection, including only bees in the infected treatment. There was no effect of flowering strips on the proportion of infected bees ($\chi^2 = 2.68$, $P = 0.262$), but flowering strips did affect mean infection intensity ($\chi^2 = 7.99$, $P = 0.018$), with higher mean infection in high-infection than low-infection tents. Mean infection intensity in canola tents was intermediate and not significantly different from either flowering strip treatment (Fig. 1A). When nectar availability per tent was included as a covariate, the effect of flowering strip was still significant ($\chi^2 = 8.97$, $P = 0.011$) and the effect of nectar was not ($\chi^2 = 2.32$, $P = 0.128$; Fig. 1B), indicating that the effect of flowering strips on infection intensity was not mediated by nectar availability.

Microcolony Performance. Surviving adult workers, number of larvae produced, and mean egg weight were significantly affected by flowering strips (Table 1), with more adults and larvae and heavier eggs with high- and low-infection strips than canola alone (Fig. 2A and C). Flowering strips did not significantly affect egg number or larval weight (Table 1). Infection had a nonsignificant tendency to reduce egg number; infection and the infection by flowering strip interaction did not affect any other performance measures (Table 1).

When nectar resources were included as a covariate (Table 1), they were positively related to number of surviving workers (Fig. 2B), larvae produced (Fig. 2D), number of eggs, and egg weight. The effect of flowering strips was still borderline




Fig. 1. Effect of flowering strip treatment on pathogen infection in bumble bees in a field tent experiment. (A) Mean infection intensity (*Crithidia* cells per $0.02 \mu\text{L}$) per tent in bumble bees (*B. impatiens*) that were originally infected and placed in microcolonies in tents that had only canola, or included flowering strips of predominantly high-infection or low-infection plant species. Values are least-square means, error bars indicate SE, and different letters above bars indicate significantly different means in post hoc tests at $P < 0.05$. Sample sizes are 15, 14, and 15 for low, canola, and high tents, respectively. (B) Mean infection intensity when available nectar resources were included as a covariate; the main effect of flowering strip treatments is still significant. Each point represents the mean value for one tent. Sample sizes are 15, 13, and 15 for low, canola, and high tents, respectively. Yellow triangles represent tents with only canola, orange squares represent high-infection, and green circles represent low-infection tents.

significant for number of larvae but not other responses, and there was a borderline significant interaction between infection and flowering strips for larval weight (Table 1). No other effects were significant (Table 1).

Pollinator Foraging Behavior. Based on "quick observations" in which observers recorded the number of bees foraging on each plant species in a tent at that moment, both infection ($\chi^2 = 13.95$, $P < 0.001$) and flowering strips ($\chi^2 = 7.81$, $P = 0.02$) reduced the number of foragers on canola, with 38% fewer foragers in tents with infected bees and 30% fewer foragers in tents with flowering strips (Fig. 3A). There was no significant interaction between infection and flowering strips ($\chi^2 = 2.42$, $P = 0.30$). However, when considering foragers to all plants, the infection by flowering strip interaction was significant ($\chi^2 = 6.16$, $P = 0.046$); infection reduced foraging on all plants in the canola and low-infection strips, but not high-infection strips (Fig. 3B).

When considering bee behavior and movement between plants, we found that infection, flowering strips, and their interaction did not affect the number of plant switches per minute, the proportion of switches to a new plant species, or the number or proportion of visits to sunflower or thyme plants. Treatments also did not affect the proportion of low- vs. high-infection plants or flowers visited ($\chi^2 < 2.05$, $P > 0.15$ for all), except that bees visited a greater proportion of low-infection plants and flowers in low-infection than high-infection flowering strips (plants, $\chi^2 = 37.96$, $P < 0.0001$; flowers, $\chi^2 = 29.23$, $P < 0.0001$). Similarly, total visits and proportion of visits to sunflowers was greater in low-infection flowering strips where sunflowers were more numerous (total visits, $\chi^2 = 6.53$, $P = 0.011$; proportion of visits, $\chi^2 = 5.29$, $P = 0.022$).

Discussion

Although previous studies have shown that floral resources can increase bumble bee parasites in some contexts (17, 18), our study demonstrates that plant species composition shapes colony-level infection intensity in free-foraging bees. All infected bees began trials with equal pathogen exposure, but after 2 wk, bees in tents with high-infection plant species had nearly twice the infection intensity as bees in tents with low-infection species,

Table 1. Effect of *C. bombi* infection (yes/no), flowering strip treatment (high-infection wildflower, low-infection wildflower, or canola only) and their interaction (all fixed effects) on multiple measures of *B. impatiens* microcolony performance using generalized linear mixed models

	Surviving workers		Number of larvae		Larval weight		Number of eggs		Egg weight	
	χ^2	<i>P</i>	χ^2	<i>P</i>	χ^2	<i>P</i>	χ^2	<i>P</i>	χ^2	<i>P</i>
Infection	0.04	0.845	0.97	0.326	0.01	0.930	3.01	0.083	1.12	0.289
Treatment	15.77	<0.001	13.70	0.001	0.84	0.657	2.79	0.248	10.72	0.005
Inf × treat	3.32	0.190	2.22	0.33	5.03	0.081	0.37	0.833	0.65	0.722
With nectar as a covariate:										
Nectar	15.39	<0.001	5.84	0.016	0.95	0.330	7.68	0.021	7.23	0.007
Infection	0.28	0.600	1.81	0.178	0.01	0.940	3.26	0.071	1.47	0.225
Treatment	3.34	0.188	5.92	0.052	4.97	0.083	0.06	0.97	3.49	0.175
Inf × treat	3.84	0.146	2.42	0.298	5.95	0.051	0.41	0.816	0.66	0.719

Surviving workers, number of larvae, and number of eggs were analyzed with Poisson distributions, and larval and egg weight with Gaussian distributions. The lower values show the same analysis with available nectar resources per tent included as a covariate. Bold indicates $P < 0.05$; italics indicate $P < 0.055$.

with intermediate infection in canola-only tents (Fig. 14). These results are particularly important since flowering strips can provide important season-long nutrients in the boom-bust resource context of agricultural fields (35). Effects of floral composition on pathogen intensity could be due to changes in pathogen transmission at flowers (20) or impacts of floral resources on bee–pathogen dynamics in the microcolonies. For example, lack of food resources can suppress immune function (36) but also decrease pathogen counts (37–39). Secondary compounds in nectar can reduce bee pathogens, as can certain types of pollen (e.g., refs. 26 and 30). Importantly, although pollen diversity can improve honey bee health and pathogen tolerance (40), the effect of floral composition on infection intensity in our study was not due to species diversity since high- and low-infection flowering strips included the same plant species, just in reversed ratios. If plant diversity drove bee–pathogen interactions, we would expect the canola-only tents to have qualitatively different results than tents with flowering strips, but this was not the case (Fig. 14). These results are important because they suggest that choice of plant species in flowering strips can influence bee disease dynamics, potentially increasing or decreasing pathogen infection intensity depending on the plant species chosen.

Even though the highest pathogen intensity occurred in high-infection tents, having enhanced floral resources from either flowering strip treatment improved bee performance compared to canola-only tents (Fig. 2). Because experimental bees were confined to tents, they were likely more food limited than wild bees and so these results should be interpreted cautiously. Nonetheless, it’s interesting that bees in high-infection and low-infection tents had similar survival and reproduction despite differences in infection intensity, suggesting that food resources play a more critical role than *C. bombi* infection for reproduction in this bee species under these experimental conditions. In support of this, we found little effect of infection treatment on most reproductive measures, with a marginal tendency for infection to reduce egg production. We note that other studies often find little impact of *C. bombi* infection on individual or colony performance in the laboratory when bees have abundant food resources, but more negative effects when bees are food stressed (41, 42). Furthermore, wild colonies of *B. terrestris* with higher *C. bombi* infection intensities were less likely to produce daughter queens (43), indicating reproductive consequences of infection under natural conditions. Experimental infection did reduce foraging in our study, consistent with a survey of wild foraging *Bombus* (44). Since bees were resource limited in our tents (demonstrated by greater microcolony reproduction with more nectar availability), it is surprising that reduced foraging due to infection treatment did not affect reproduction more

strongly. Perhaps we would have seen more negative reproductive consequences if we had conducted longer trials. Alternatively, high pathogen intensity could signal bees to invest in reproduction prior to their demise. Longer-term studies are needed to determine impacts of plant species on bee reproductive fitness via changes in resources versus pathogen infection intensity.

Although flowering strips affected both pathogen infection intensity and bee reproduction, these effects were mediated by different mechanisms. The effect of flowering strips on bee reproduction was explained by estimated nectar availability in each tent (Fig. 2 *B* and *D*), while the effect of flowering strips on pathogen infection intensity remained after accounting for nectar availability (Fig. 1*B*). It is not surprising that reproduction correlated strongly with floral resources in bees confined to tents, although this may still reflect larger-scale patterns since resources increase bee abundance and diversity in widespread agricultural settings (4). However, this contrasts with pathogen infection intensity, which was not related to nectar resources. The original trials that designated plant species as “high-” or “low-” infection allowed single uninfected workers to forage on an experimentally inoculated inflorescence for less than 20 min, and then assessed infection 1 wk later (20). It is remarkable that these categorizations of plant species based on brief foraging bouts still predicted infection intensities for microcolonies foraging in tents over a 2-wk period, when bees could forage for pollen, revisit flowers over several days, and interact with each other inside the colony.

Our previous trials identified surprisingly few floral traits associated with likelihood of acquiring disease other than number of reproductive structures (20). However, subsequent work suggests that floral architecture or surface compounds/trichomes, traits we did not measure in our original study, may play a role in disease dynamics, since the location of inoculum placement (inside flowers, outside flowers, on bracts) has variable effects on acquisition across plant species (23). We also did not measure floral volatiles, some of which reduce *C. bombi* viability (45). Nectar or pollen chemistry could also influence *C. bombi* infection intensity (26, 28, 29, 33, 45). Since the designation of high- and low-infection species accurately predicted colony infection intensity over 2 wk, this suggests that whatever traits influence short-term transmission dynamics also play important roles over longer time periods in determining infection intensities. Short-term dynamics can predict longer-term patterns in other systems; for example, greater exposure to bacterial wilt (3 h vs. 24 h) increased vector beetle infection likelihood after 5 d and persisted at 28 d, suggesting that interaction intensity affected both short- and long-term ability to acquire this pathogen (46).




Fig. 2. Effect of flowering strip treatment on bumble bee adult survival and colony reproduction in a field tent experiment. (A) Number of surviving adult workers per tent in each flowering strip treatment; sample sizes are 30, 26, and 30 for low, canola, and high tents, respectively. (B) Number of surviving adult workers when nectar resources were included as a covariate; sample sizes are 29, 22, and 27 for low, canola, and high tents, respectively. (C) Number of larvae produced per tent in each flowering strip treatment; sample sizes are 30, 28, and 30 for low, canola, and high tents, respectively. (D) Number of larvae produced when nectar resources were included as a covariate; sample sizes are 28, 22, and 27 for low, canola, and high tents, respectively. For A and C, values are back-transformed least-square means, error bars indicate SE, and different letters above bars indicate significantly different means in post hoc tests at $P < 0.05$. For B and D, each point represents the mean value for one tent. Yellow triangles represent tents with only canola, orange squares represent high-infection, and green circles represent low-infection tents.

We saw little effect of infection treatment or flowering strips on pollination service to canola, most likely because confining bees in tents resulted in excess pollen delivery to all plants. However, infection reduced the number of workers foraging at any given time, which could have more significant consequences for pollination on larger spatial scales, given that *Crithidia* can infect up to 80% of *Bombus* in western Massachusetts (47) and nearly 50% of *B. terrestris* workers in Switzerland and the United Kingdom (43, 44). Flowering strips also reduced the number of foragers on canola, but the high ratio of flowering strip to crop plants in our tents may overestimate the magnitude of this effect in the field. Designs that incorporate flowering strips or hedges on farms or in field trials would be more appropriate to assess effects on pollinator service; such studies often find benefits of flowering strips for pollination services (7–9).

In conclusion, we found that bee colony-level infection intensity nearly doubled with the addition of plant species known to increase pathogen acquisition, compared to low-infection plant species. However, bee colony reproduction was substantially improved with any flowering strips compared to canola alone. Variation in colony reproduction was explained by nectar resources, but infection intensity was not. Thus, both flowering strip treatments benefited colony reproduction by adding floral resources, but certain plant species may also increase pathogen infection intensity. This should be a consideration when selecting plant species for augmenting pollinator habitat.

Materials and Methods

Study System. *B. impatiens* (Apidae, the common eastern bumble bee) is one of the most prevalent bumble bee species in the eastern United States (e.g., refs. 47 and 48), a generalist pollinator, and available commercially. We used microcolonies of ~15 workers as our unit of replication; microcolonies have been used successfully to estimate colony performance as a function of diet (49).

C. bombi (Zoomastigophora: Trypanosomatidae) is a protozoan gut parasite that can be contracted at flowers via fecal–oral transmission (14, 50). *C. bombi* reduces learning and foraging efficiency in workers (51, 52), slows colony growth rates (53), and is associated with reduced likelihood of reproduction in wild colonies (43). Stressful conditions increase the mortality of infected workers (41) and reduce infected queen fitness (42). *C. bombi* infection is locally common; for example, *C. bombi* infected over 60% of *B. impatiens* in western Massachusetts (47).

Canola (*Brassica rapa* cultivar O. Eclipse; Spectrum Crop Development, Ritzville, WA) is obligately outcrossed and has improved yield from insect pollinators (54, 55). Canola was grown on more than 1 million acres in the United States with a production value of \$357 billion in 2011 (National Agricultural Statistics Service, US Department of Agriculture [USDA]), and so is a major foraging source for bees as well as a valuable US crop. *Bombus* exhibit natural foraging behaviors in tent enclosures, are common natural pollinators of canola (e.g., ref. 56), and have been successfully used to pollinate canola in greenhouse experiments previously (57). Thus, *Bombus* are both relevant and tractable to examine the effects of pathogens and flowering strips on pollination in canola.

We designated low- and high-infection plant species based on previous work in which we inoculated flowers of each species with a known quantity of *C. bombi* inoculum, allowed an uninfected bee to forage (typically less than 20 min), and then individually reared the bees for 1 wk and assessed pathogen cell counts (20). In most cases species were designated as low or high infection based on both the probability of a bee becoming infected and the severity of infection; mean *C. bombi* cell counts 1 wk after foraging (including zeros) were 9.0 to 13.1 cells/0.02 μ L for low- and 18.9 to 36.2 cells/0.02 μ L for high-infection plant species (20). High-infection species used in the current experiment were *Antirrhinum majus* (Plantaginaceae), *Asclepias incarnata* (Asclepiadaceae), *Lobelia siphilitica* (Campanulaceae), *Lythrum salicaria* (Lythraceae), *Penstemon digitalis* (Plantaginaceae), and *Solidago altissima* ssp. *altissima* (Asteraceae). Low-infection species were *Digitalis purpurea* (Plantaginaceae), *Helianthus annuus* (Asteraceae), *Linaria vulgaris* (Plantaginaceae), and multiple varieties of *Thymus* (Lamiaceae). See *SI Appendix* for plant source and propagation details.

Experimental Design Summary. We manipulated *C. bombi* infection (yes/no) and flowering strips in a 2×3 factorial design using tents in the field at the University of Massachusetts Center for Agriculture (South Deerfield, MA, 42° 28.6' N, 72° 34.8' W). We used tents to constrain our experimental bees to only forage on their provided treatment plants, although we acknowledge that this likely resulted in more food limitation than free-foraging bees would experience. Our flowering strip treatments were canola only,




Fig. 3. Effect of infection, flowering strip treatment (low-infection plants, high-infection plants, or only canola) and their interaction on bumble bee foraging behavior. (A) Number of foragers observed on canola inflorescences during frequent quick observations of each tent; there are 73 to 77 data points per infection/flowering strip combination. (B) Number of foragers observed on all plants during quick observations; there are 76 to 79 data points per infection/flowering strip combination. Values are least-square means and error bars indicate SE.

high-infection strip, and low-infection strip. These six treatment combinations were replicated three times per round (18 mesh-sided tents; 2.44 m × 2.44 m; Delta Canopy Inc., McKinney, TX) and we conducted five, 2-wk rounds between June 3 and August 27, 2015. This provided a total of 88 replicate tents (two replicates of uninfected canola were lost due to tent collapse). Tents were arranged in three blocks (rows) of six, with one replicate of each treatment combination per block, and treatments were randomly assigned to tents within blocks. The length of each round was approximately 2 wk, long enough for colonies to produce larvae but not newly emerged adults. Round length varied slightly across blocks, which were always set up on the same day but taken down over 2 to 3 d. Tents each contained a single microcolony of ~15 *B. impatiens* workers (see *SI Appendix* for microcolony construction methods), bins of canola plants, and flowering strips when appropriate. Microcolonies of bees were initially infected with *C. bombi* or received a sham infection. For tents with infected microcolonies, our goal was to assess how flowering strips affected final infection levels; our design did not allow us to distinguish between on-flower and within-colony transmission pathways.

At the end of each round we removed microcolonies, counted *C. bombi* cells of all surviving workers, and recorded worker survival and microcolony reproduction. We then randomly reassigned tents within blocks to new treatments, with the constraint that tents with infected bees in the first round were maintained as “infected” tents for the entire experiment to avoid contamination, even though contamination is unlikely because *C. bombi* does not survive long outside the host (23). Plants were rearranged or replaced as needed, again with the constraint that plants from infected tents were only used in other infected tents. New microcolonies were used in each round.

Flowering Strip Treatments. All tents contained bins of canola as the focal crop, with ~20 plants per bin. We included 15 bins of canola per tent in the canola-only treatment, and 12 bins per tent in the flowering strip treatments to reflect that flowering strips could result in less space for a crop (see *SI Appendix*, Fig. S1 for physical layout). Flowering strips included 12 potted plants; there were typically three low- and three high-infection species per round, but in some cases a single species was used twice (i.e., at double concentration) per round due to flowering plant availability (*SI Appendix*, Table S1). Tents with a low-infection flowering strip contained three individuals of each low-infection species and one of each high-infection species; tents with a high-infection flowering strip used three of each high-infection species and one of each low-infection species. Thus, high- and low-infection tents included the same plant species, just in reversed ratios. Manipulating the ratio of high:low-infection plant species, rather than species composition, allowed us to avoid confounding plant species composition and diversity with flowering strip treatments. High- and low-infection species varied between rounds based on phenology (*SI Appendix*, Table S1). See *SI Appendix*, Table S2 for plant sources and propagation methods.

Assessing Infection and Microcolony Performance. Microcolonies were created from commercial colonies of *B. impatiens* (Biobest, Leamington, ON, Canada) approximately 1 wk before deploying in field tents. All colonies were confirmed to be free of *C. bombi* before use by dissecting a subsample of workers. We used three colonies of origin per round. Each block in the field site contained microcolonies from the same colony of origin, so that the block factor includes genetic as well as spatial variation. Each microcolony was initiated with 11 to 17 workers, plus 2 to 9 pupal cells in the first two rounds only. We chose this range of initial workers to be large enough to facilitate microcolony survival, but small enough to have sufficient floral resources in the tents. Microcolonies were randomly assigned to infected or uninfected treatments and inoculated with *C. bombi* at microcolony initiation. In infected colonies, each bee was fed 10 μ L of inoculum with 6,000 *C. bombi* cells in 25% sucrose solution. This concentration is well within the range of natural variation (39, 58). We made inoculum fresh daily following Richardson et al. (30). Bees in uninfected microcolonies were fed a 10- μ L droplet of 25% sucrose without *C. bombi* cells to control for handling effects. Bees that did not consume their entire droplet in either treatment were discarded. Microcolonies were maintained in a growth chamber in darkness at 27 °C for approximately 1 wk before deployment at the field site. We only used microcolonies in the field that had initiated egg laying; we recorded the number of workers in each microcolony when they were moved to the field. See *SI Appendix* for microcolony inoculation, construction and field deployment details.

At the end of each round, we assessed microcolony infection and performance. We quantified *C. bombi* infection by macerating bee guts and counting cells in a 0.02- μ L sample in a hemocytometer (*SI Appendix*). Overall,

in the uninfected treatment only 2.3% of bees had detectable *C. bombi* at the end of trials compared to 66% of bees in the infected treatment, indicating that the infection treatment was effective. Uninfected bees in the infected treatment may be because our inoculation did not successfully infect, or because bees recovered from their infection. The small number of infected bees in the “uninfected” treatment may be due to a few bees we observed escaping and returning to their tents. The number of living workers per microcolony was used as a measure of survival, scaled by the number at the start of the round. In addition, we froze each microcolony at -20 °C and then counted eggs and counted and weighed all larvae and pupae. Since each round ran for only 14 d, no adults emerged.

Measuring Foraging Behavior and Pollination Service. We conducted two types of pollinator observations to ask how infection and flowering strips affected visits to plants, and whether infection shifted foraging preferences. Detailed observations were conducted two to three times per round for 30-min periods per tent by observing individual bees and recording the plant species (high- and low-infection treatments only), flowers probed per plant visit, and time per plant visit in seconds. Time per flower was calculated by dividing the flowers probed by the time per plant visit. For canola, it was not possible to distinguish individual plants and so inflorescences were the unit of observation. Whenever a tent was observed, we also recorded floral resource availability by counting or estimating the number of open flowers of each species. Beginning in round 3, we also conducted quick observations two to three times per observation day, in which a single observer would visit each tent and record the number of bees foraging on each plant species at that moment. This provided a larger dataset to ask whether infection and flowering strips affected the total number of foragers and their foraging preferences at a given time point.

We found no evidence that treatments affected pollination service, measured as pollen deposition and pollen limitation for fruit and seed set, and therefore report methods and results in the *SI Appendix*.

Statistical Analysis.

Overview. We used R version 3.5.2 for all analyses (59). Most models were generalized linear mixed models (GLMMs) using the *glmer* function in *lme4* (60), including the fixed effects of flowering strips (canola, low, or high infection), infection (yes/no), and their interaction as fixed effects. Count data (e.g., number of eggs) were analyzed with Poisson distributions, proportional data (e.g., proportion of bees infected per microcolony) with binomial distributions, and continuous data (e.g., larval weight) with Gaussian distributions. For responses with a single value per tent (e.g., larvae per microcolony), we used block nested within round as a random effect. For responses with multiple values per tent (e.g., *C. bombi* cells per bee per microcolony), we included tent nested within block within round as a random effect. When Poisson or binomial data were overdispersed, we included an observation-level random effect (61, 62). We used likelihood ratio tests to evaluate fixed factors (the interaction term, then flowering strip, then infection treatment).

Pathogen infection. Effects of flowering strips on the proportion of bees infected and mean infection level (cells per 0.02 μ L) were analyzed in the infected treatment only. To determine whether flowering strip effects on infection were due to resource availability, we reanalyzed responses including estimated nectar availability in each tent as a covariate. Nectar availability was estimated as the average number of open flowers per species in each tent (recorded during pollinator observations) multiplied by the previously determined mean nectar volume per flower for each species (20). We did not consider variation in sugar concentration since these values were unavailable for most species.

Microcolony performance. Microcolony performance was measured as number of eggs, larvae, and surviving workers (including number of workers at the start of the round as a covariate) and mean egg and larval weight. As with pathogen infection, we reanalyzed responses with estimated nectar availability in each tent as a covariate.

Pollinator foraging behavior. We analyzed quick observations of total bees on each plant species per tent and observations of individual bee behavior. We used the quick observations to ask whether infection and flowering strips affected mean foragers on canola and on all plants. We used our observations of individual foraging bouts to ask whether infection and flowering strips changed foraging behavior, including only tents with flowering strips. Responses included the total switches between plants (any species) per minute, the proportion of switches to a new plant species, the proportion of flowering strip species visited or flowers probed that were low vs. high infection, and the number and proportion of visits to sunflower and thyme plants (using the number of open sunflower or thyme flowers as covariates). We singled



out sunflower, since its pollen dramatically reduced *C. bombi* infection (26), and thyme, since thymol reduced *C. bombi* at natural nectar concentrations in vivo (30) and in vitro (45).

Data Availability. All data and R scripts are deposited in Dryad (<https://doi.org/10.5061/dryad.p2ngf1vn1>) (34).

ACKNOWLEDGMENTS. We thank Biobest for donating bumble bee colonies; S. J. Giacomini for managing laboratory components; P. Anderson, D. Delaney, J. Giacomini, A. Hachey, R. Holley, E. Mann, S. Mitchell, Y. Moon,

R. Papapietro, R. Pasquale, C. Shuja, and M. Stypulkoski for research assistance; N. Woodard and the University of Massachusetts Center for Agriculture for field site support; and three anonymous reviewers plus R. Malfi, J. Van Wyk, and other members of the L.S.A. laboratory for providing feedback on the manuscript. Research was funded by USDA/Agriculture and Food Research Initiative 2013-02536, NIH Grant 1 R01 GM122062-01, National Science Foundation/Division of Environmental Biology Grant 1258096 (all L.S.A. and R.E.I.) and USDA/Cooperative State Research, Education, and Extension Service Grant MAS000411 (L.S.A.). Any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the funding agencies.

1. S. G. Potts *et al.*, Safeguarding pollinators and their values to human well-being. *Nature* **540**, 220–229 (2016).
2. D. Goulson, E. Nicholls, C. Botías, E. L. Rotheray, Bee declines driven by combined stress from parasites, pesticides, and lack of flowers. *Science* **347**, 1255957 (2015).
3. D. Kleijn *et al.*, Scaling up effects of measures mitigating pollinator loss from local- to landscape-level population responses. *Methods Ecol. Evol.* **9**, 1727–1738 (2018).
4. N. M. Williams *et al.*, Native wildflower plantings support wild bee abundance and diversity in agricultural landscapes across the United States. *Ecol. Appl.* **25**, 2119–2131 (2015).
5. R. Winfree, “The conservation and restoration of wild bees” in *Year in Ecology and Conservation Biology 2010*, R. S. Ostfeld, W. H. Schlesinger, Eds. (Wiley-Blackwell, 2010), vol. 1195, pp. 169–197.
6. L. A. Morandin, C. Kremen, Hedgerow restoration promotes pollinator populations and exports native bees to adjacent fields. *Ecol. Appl.* **23**, 829–839 (2013).
7. B. R. Blaauw, R. Isaacs, Flower plantings increase wild bee abundance and the pollination services provided to a pollination-dependent crop. *J. Appl. Ecol.* **51**, 890–898 (2014).
8. L. G. Carvalheiro, C. L. Seymour, S. W. Nicolson, R. Veldtman, Creating patches of native flowers facilitates crop pollination in large agricultural fields: Mango as a case study. *J. Appl. Ecol.* **49**, 1373–1383 (2012).
9. L. A. Morandin, R. F. Long, C. Kremen, Pest control and pollination cost-benefit analysis of hedgerow restoration in a simplified agricultural landscape. *J. Econ. Entomol.* **109**, 1020–1027 (2016).
10. C. Carvell *et al.*, Bumblebee family lineage survival is enhanced in high-quality landscapes. *Nature* **543**, 547–549 (2017).
11. T. H. Roulston, J. H. Cane, Pollen nutritional content and digestibility for animals. *Plant Syst. Evol.* **222**, 187–209 (2000).
12. S. W. Nicolson, M. Nepi, E. Pacini, Eds., *Nectaries and Nectar* (Springer, Dordrecht, The Netherlands, 2007), pp. 395.
13. W. Kämper *et al.*, How landscape, pollen intake and pollen quality affect colony growth in *Bombus terrestris*. *Landsc. Ecol.* **31**, 2245–2258 (2016).
14. S. Durrer, P. Schmid-Hempel, Shared use of flowers leads to horizontal pathogen transmission. *Proc. Biol. Sci.* **258**, 299–302 (1994).
15. P. Graystock, D. Goulson, W. O. H. Hughes, Parasites in bloom: Flowers aid dispersal and transmission of pollinator parasites within and between bee species. *Proc. Biol. Sci.* **282**, 20151371 (2015).
16. S. H. McArt, H. Koch, R. E. Irwin, L. S. Adler, Arranging the bouquet of disease: Floral traits and the transmission of plant and animal pathogens. *Ecol. Lett.* **17**, 624–636 (2014).
17. N. Piot *et al.*, Establishment of wildflower fields in poor quality landscapes enhances micro-parasite prevalence in wild bumble bees. *Oecologia* **189**, 149–158 (2019).
18. H. Cohen, R. D. Quistberg, S. M. Philpott, Vegetation management and host density influence bee-parasite interactions in urban gardens. *Environ. Entomol.* **46**, 1313–1321 (2017).
19. S. A. Alger, P. A. Burnham, H. F. Boncristiani, A. K. Brody, RNA virus spillover from managed honeybees (*Apis mellifera*) to wild bumblebees (*Bombus* spp.). *PLoS One* **14**, e0217822 (2019).
20. L. S. Adler *et al.*, Disease where you dine: Plant species and floral traits associated with pathogen transmission in bumble bees. *Ecology* **99**, 2535–2545 (2018).
21. D. T. Peck, M. L. Smith, T. D. Seeley, *Varroa destructor* mites can nimbly climb from flowers onto foraging honey bees. *PLoS One* **11**, e0167798 (2016).
22. L. L. Truitt, S. H. McArt, A. H. Vaughn, S. P. Ellner, Trait-based modeling of multihost pathogen transmission: Plant-pollinator networks. *Am. Nat.* **193**, E149–E167 (2019).
23. L. L. Figueroa *et al.*, Bee pathogen transmission dynamics: Deposition, persistence and acquisition on flowers. *Proc. Biol. Sci.* **286**, 20190603 (2019).
24. J. Abbott, Self-medication in insects: Current evidence and future perspectives. *Ecol. Entomol.* **39**, 273–280 (2014).
25. J. Moore, An overview of parasite-induced behavioral alterations - and some lessons from bats. *J. Exp. Biol.* **216**, 11–17 (2013).
26. J. J. Giacomini *et al.*, Medicinal value of sunflower pollen against bee pathogens. *Sci. Rep.* **8**, 14394 (2018).
27. G. M. LoCascio, L. Aguirre, R. E. Irwin, L. S. Adler, Pollen from multiple sunflower cultivars and species reduces a common bumblebee gut pathogen. *R. Soc. Open Sci.* **6**, 190279 (2019).
28. D. Baracchi, M. J. F. Brown, L. Chittka, Behavioural evidence for self-medication in bumblebees? *F1000Res* **4**, 73 (2015).
29. J. S. Manson, M. C. Otterstatter, J. D. Thomson, Consumption of a nectar alkaloid reduces pathogen load in bumble bees. *Oecologia* **162**, 81–89 (2010).
30. L. L. Richardson *et al.*, Secondary metabolites in floral nectar reduce parasite infections in bumblebees. *Proc. Biol. Sci.* **282**, 20142471 (2015).
31. O. M. Biller, L. S. Adler, R. E. Irwin, C. McAllister, E. C. Palmer-Young, Possible synergistic effects of thymol and nicotine against *Crithidia bombi* parasitism in bumble bees. *PLoS One* **10**, e0144668 (2015).
32. L. P. Thorburn, L. S. Adler, R. E. Irwin, E. C. Palmer-Young, Variable effects of nicotine and anabasine on parasitized bumble bees. *F1000 Res.* **4**, 880 (2015).
33. L. L. Richardson, M. D. Bowers, R. E. Irwin, Nectar chemistry mediates the behavior of parasitized bees: Consequences for plant fitness. *Ecology* **97**, 325–337 (2016).
34. N. A. Barber, L. S. Adler, R. E. Irwin, O. M. Biller, Data from: Flowering plant composition shapes pathogen infection intensity and reproduction in bumble bee colonies. Dryad. <https://doi.org/10.5061/dryad.p2ngf1vn1>. Deposited 24 April 2020.
35. A. G. Dolezal, A. L. St Clair, G. Zhang, A. L. Toth, M. E. O’Neal, Native habitat mitigates feast-famine conditions faced by honey bees in an agricultural landscape. *Proc. Natl. Acad. Sci. U.S.A.* **116**, 25147–25155 (2019).
36. N. Roger, D. Michez, R. Wattiez, C. Sheridan, M. Vanderplanck, Diet effects on bumblebee health. *J. Insect Physiol.* **96**, 128–133 (2017).
37. T. J. Conroy, E. C. Palmer-Young, R. E. Irwin, L. S. Adler, Food limitation affects parasite load and survival of *Bombus impatiens* (Hymenoptera: Apidae) infected with *Crithidia* (Trypanosomatida: Trypanosomatidae). *Environ. Entomol.* **45**, 1212–1219 (2016).
38. C. J. Jack, S. S. Uppala, H. M. Lucas, R. R. Sagili, Effects of pollen dilution on infection of *Nosema ceranae* in honey bees. *J. Insect Physiol.* **87**, 12–19 (2016).
39. A. Logan, M. X. Ruiz-González, M. J. Brown, The impact of host starvation on parasite development and population dynamics in an intestinal trypanosome parasite of bumble bees. *Parasitology* **130**, 637–642 (2005).
40. G. Di Pasquale *et al.*, Influence of pollen nutrition on honey bee health: Do pollen quality and diversity matter? *PLoS One* **8**, e72016 (2013).
41. M. J. F. Brown, R. Loosli, P. Schmid-Hempel, Condition-dependent expression of virulence in a trypanosome infecting bumblebees. *Oikos* **91**, 421–427 (2000).
42. M. J. F. Brown, R. Schmid-Hempel, P. Schmid-Hempel, Strong context-dependent virulence in a host-parasite system: Reconciling genetic evidence with theory. *J. Anim. Ecol.* **72**, 994–1002 (2003).
43. D. Goulson, S. O’Connor, K. J. Park, The impacts of predators and parasites on wild bumblebee colonies. *Ecol. Entomol.* **43**, 168–181 (2018).
44. J. A. Shykoff, P. Schmid-Hempel, Incidence and effects of four parasites in natural populations of bumble bees in Switzerland. *Apidologie (Celle)* **22**, 117–125 (1991).
45. E. C. Palmer-Young, B. M. Sadd, P. C. Stevenson, R. E. Irwin, L. S. Adler, Bumble bee parasite strains vary in resistance to phytochemicals. *Sci. Rep.* **6**, 37087 (2016).
46. L. R. Shapiro, I. Seidl-Adams, C. M. De Moraes, A. G. Stephenson, M. C. Mescher, Dynamics of short- and long-term association between a bacterial plant pathogen and its arthropod vector. *Sci. Rep.* **4**, 4155 (2014).
47. S. Gillespie, Factors affecting parasite prevalence among wild bumblebees. *Ecol. Entomol.* **35**, 737–747 (2010).
48. N. Cordes *et al.*, Interspecific geographic distribution and variation of the pathogens *Nosema bombi* and *Crithidia* species in United States bumble bee populations. *J. Invertebr. Pathol.* **109**, 209–216 (2012).
49. J. N. Tasei, P. Aupinel, Validation of a method using queenless *Bombus terrestris* micro-colonies for testing the nutritive value of commercial pollen mixes by comparison with queenright colonies. *J. Econ. Entomol.* **101**, 1737–1742 (2008).
50. P. Schmid-Hempel, S. Durrer, Parasites, floral resources and reproduction in natural populations of bumblebees. *Oikos* **62**, 342–350 (1991).
51. R. J. Gegear, M. C. Otterstatter, J. D. Thomson, Does parasitic infection impair the ability of bumblebees to learn flower-handling techniques? *Anim. Behav.* **70**, 209–215 (2005).
52. R. J. Gegear, M. C. Otterstatter, J. D. Thomson, Bumble-bee foragers infected by a gut parasite have an impaired ability to utilize floral information. *Proc. Biol. Sci.* **273**, 1073–1078 (2006).
53. J. A. Shykoff, P. Schmid-Hempel, Parasites delay worker reproduction in bumblebees: Consequences for eusociality. *Behav. Ecol.* **2**, 242–248 (1991).
54. L. A. Morandin, M. L. Winston, Wild bee abundance and seed production in conventional, organic, and genetically modified canola. *Ecol. Appl.* **15**, 871–881 (2005).
55. L. A. Morandin, M. L. Winston, Pollinators provide economic incentive to preserve natural land in agroecosystems. *Agric. Ecosyst. Environ.* **116**, 289–292 (2006).

56. K. E. Hayter, J. E. Cresswell, The influence of pollinator abundance on the dynamics and efficiency of pollination in agricultural *Brassica napus*: Implications for landscape-scale gene dispersal. *J. Appl. Ecol.* **43**, 1196–1202 (2006).
57. I. Ahman, A. Lehrman, B. Ekbom, Impact of herbivory and pollination on performance and competitive ability of oilseed rape transformed for pollen beetle resistance. *Arthropod Plant Interact.* **3**, 105–113 (2009).
58. M. C. Otterstatter, J. D. Thomson, Within-host dynamics of an intestinal pathogen of bumble bees. *Parasitology* **133**, 749–761 (2006).
59. R Core Team, *R: A Language and Environment for Statistical Computing* (R Foundation for Statistical Computing, Vienna, Austria, 2018).
60. D. Bates, M. Machler, B. M. Bolker, S. C. Walker, Fitting linear mixed-effects models using lme4. *J. Stat. Softw.* **67**, 1–48 (2015).
61. B. M. Bolker et al., Generalized linear mixed models: A practical guide for ecology and evolution. *Trends Ecol. Evol. (Amst.)* **24**, 127–135 (2009).
62. X. A. Harrison, Using observation-level random effects to model overdispersion in count data in ecology and evolution. *PeerJ* **2**, e616 (2014).