
Theses and Dissertations

Spring 2013

Genetic analyses of microbial polychlorinated biphenyl degradation in natural and engineered systems

Yi Liang
University of Iowa

Follow this and additional works at: <https://ir.uiowa.edu/etd>



Part of the [Civil and Environmental Engineering Commons](#)

Copyright 2013 Yi Liang

This dissertation is available at Iowa Research Online: <https://ir.uiowa.edu/etd/3129>

Recommended Citation

Liang, Yi. "Genetic analyses of microbial polychlorinated biphenyl degradation in natural and engineered systems." PhD (Doctor of Philosophy) thesis, University of Iowa, 2013.

<https://doi.org/10.17077/etd.3r6xzntw>

Follow this and additional works at: <https://ir.uiowa.edu/etd>



Part of the [Civil and Environmental Engineering Commons](#)

GENETIC ANALYSES OF MICROBIAL POLYCHLORINATED BIPHENYL
DEGRADATION IN NATURAL AND ENGINEERED SYSTEMS

by
Yi Liang

A thesis submitted in partial fulfillment
of the requirements for the Doctor of
Philosophy degree in Civil and Environmental Engineering
in the Graduate College of
The University of Iowa

May 2014

Thesis Supervisor: Associate Professor Timothy E. Mattes

Copyright by

YI LIANG

2014

All Rights Reserved

Graduate College
The University of Iowa
Iowa City, Iowa

CERTIFICATE OF APPROVAL

PH.D. THESIS

This is to certify that the Ph.D. thesis of

Yi Liang

has been approved by the Examining Committee
for the thesis requirement for the Doctor of Philosophy
degree in Civil and Environmental Engineering at the May 2014 graduation.

Thesis Committee: _____
Timothy E. Mattes, Thesis Supervisor

Keri Hornbuckle

Larry Robertson

Tonya Peoples

Jerald Schnoor

ACKNOWLEDGMENTS

I extend a sincere thank you to my advisor, Dr. Timothy Mattes, who has patiently provided guidance and support throughout my Ph.D. training. I also extend my thanks to my committee members, Dr. Keri Hornbuckle, Dr. Jerald Schnoor, Dr. Larry Robertson and Dr. Tonya Peeples.

I would like to thank the former and current members in the Mattes lab, Yang Oh Jin, Adina Howe, Anne Alexander, Joshua Livermore, Carmen Owens, Meredith Dobson, Meng Chen Lee, Xikun Liu, Carly Lintner, and Caitlin Krause, for always being friendly and willing to help.

I would like to thank Andres Martinez, Dingfei Hu and Richard Meggo, for providing me the valuable PCB congener data for this research. I would also like to thank the former and current lab managers, Collin Just and Jon Durst, for keeping the lab organized. I also want to thank Judy Holland, Jennifer Rumping, Angela Schenkel for providing excellent administrative services.

I give my special thanks to my family, my parents, my grandparents, for the support and unconditional love. I would also like to thank my friends in US and in China, for staying with me during the most difficult times.

This research was funded by the National Institute of Environmental Health Sciences (NIEHS) Superfund Research Program (SRP) Grant No. P42ES13661 and by a fellowship from the Center for Biocatalysis and Bioprocessing at the University of Iowa.

ABSTRACT

Polychlorinated biphenyls (PCBs) are carcinogenic, persistent, and bioaccumulative contaminants that pose risks to human and environmental health. PCB biodegradation by indigenous microbial communities could be a cost-effective and an environmental-friendly bioremediation strategy for *in situ* PCB removal. A comprehensive understanding of the microbial PCB degradation at the contaminated site is required for the acceptance and optimization of using microbial PCB degradation as the site clean-up strategy. This thesis describes investigations of the aerobic and anaerobic microbial degradation of PCBs under both field and laboratory conditions.

The microbial PCB degradation potential in sediments from Indiana Harbor and Ship Canal (IHSC), a site that was historically contaminated by PCBs, was explored by analyzing the PCB congener distributions and microbial communities in two core sediment samples. PCB congener analysis suggested the possibility of *in situ* dechlorination in deep sediments. Molecular analysis of biomarker genes revealed the potential of both aerobic and anaerobic PCB degradation in sediments. Microbial communities were characterized by the combination use of terminal restriction fragment length polymorphism (T-RFLP), clone library, and pyrosequencing. These methods elucidated the dominant role of *Proteobacteria*, especially *Acidovorax* and *Acinetobacter* in sediments.

To improve the microbial PCB degradation, phytoremediation with switchgrass (*Panicum virgatum*) was employed under laboratory conditions. Congener analysis showed that both phytoextraction and microbial PCB degradation contributed to the enhanced PCB removal in the presence of switchgrass. Bioaugmentation with *Burkholderia xenovorans* LB400 was performed to further promote aerobic PCB degradation. The presence of LB400 was associated with improved degradation of PCB 52, but not PCB77 or PCB 153. Increased abundance of the biphenyl dioxygenase gene,

which is indicative of aerobic PCB degradation, and its transcript were observed after bioaugmentation, suggesting active aerobic PCB degradation.

To promote the anaerobic PCB degradation, redox cycling (alternating flooding and non-flooding) was performed. Redox cycling was found to improve the removal of PCB 153 in unplanted soils and to increase the dechlorinating *Chloroflexi* population. Characterization of the microbial community by T-RFLP and clone library revealed that *Proteobacteria* and *Acidobacteria* were dominant. Species that contain dechlorination potential were identified, including *Geobacter* and *Clostridium*, suggesting that their possible role in PCB dechlorination.

The research described in this thesis provides scientific knowledge and evidence for the feasibility of employing bioremediation including natural attenuation, phytoremediation, and bioaugmentation to clean up PCB contamination. Such information will be critical in selecting and optimizing remediation strategies for PCB contaminated sites.

Table of Contents

LIST OF TABLES	vii
LIST OF FIGURES	viii
CHAPTER I. RESEARCH OVERVIEW AND HYPOTHESES.....	1
CHAPTER II. BACKGROUND	4
Polychlorinated biphenyls	4
Microbial PCB biodegradation	6
Microbial PCB degradation in sediments and soils	10
Phytoremediation of PCBs	13
CHAPTER III. MICROBIAL PCB DEGRADATION POTENTIAL IN CORE 1 FROM IHSC	18
Introduction.....	18
Materials and Methods	19
Results and Discussion	23
Conclusions.....	27
CHAPTER IV. MICROBIAL PCB DEGRADATION POTENTIAL IN CORE 2 FROM IHSC	39
Introduction.....	39
Materials and Methods	41
Results and discussion	45
Conclusions.....	51
CHAPTER V. ENHANCED POLYCHLORINATED BIPHENYL REMOVAL A SWITCHGRASS PLANTED RHIZOSPHERE BY BIOAUGMENTATION WITH <i>BURKHOLDERIA XENOVORANS</i> LB400	62
Introduction.....	62
Materials and Methods	64
Results and Discussion	69
Conclusions.....	74
CHAPTER VI. MICROBIAL COMMUNITY ANALYSIS OF A SWITCHGRASS RHIZOSPHERE WITH PCB SPIKING AND REDOX CYCLING	84
Introduction.....	84
Materials and Methods	85
Results and Discussion	91
Conclusions.....	98
CHAPTER VII. ENGINEERING AND SCIENTIFIC SIGNIFICANCE.....	117

REFERENCES121

LIST OF TABLES

2.1. Positions of chlorines removed by different dechlorination patterns.	10
3.1. PCR analysis of DNA extracted from different depths of PCB-contaminated core 1 sediment sample at IHSC.....	29
3.2. Summary of 454 sequence library sizes, operational taxonomic unites (OTUs).	30
4.1. Primer sets used for qPCR analysis in Chapter IV.. .	53
4.2. Pertinent qPCR parameters (primer concentration, template mass, linear range, PCR efficiency, and y-intercept of the qPCR standard curve) with core 2 sediment DNA as template.	54
4.3. RDP classification of clone libraries for qPCR primer quality assurance with core 2 sediment DNA as template.	55
4.4. 16S rRNA gene clones recovered from sediment at the depth of 3.35-3.66 m. Clone identification was made by RDP classifier.	56
5.1. Primer sets used in qPCR analysis in Chapter V.....	75
5.2. qPCR reaction set up, linear range and efficiency for qPCR reaction in Chapter V.....	76
5.3. Molar mass balance of PCBs in unplanted soil (UP), switchgrass treated soil (SG), unplanted with bioaugmentation soil (UP B), switchgrass treated with bioaugmentation soil (SG B), and switchgrass treated with bioaugmentation with dead LB400 soil (SG BD). (n=3).....	77
6.1. Primer sets used for qPCR analysis in Chapter VI.....	99
6.2. Pertinent qPCR parameters (primer concentration, template mass, linear range, PCR efficiency, and y-intercept of the qPCR standard curve) in Chapter VI.....	100
6.3. RDP classification of clone libraries for quality assurance of qPCR primers used in Chapter VI.	101
6.4. 16S rRNA gene clones recovered from unplanted soil (UP), switchgrass treated soil (SG), unplanted soil with redox cycling (UPF), switchgrass treated soil with redox cycling (SG F). Clone identification was made by RDP classifier.	102
6.5. Significant indicator T-RFs determined by indicator species analysis.....	103

LIST OF FIGURES

2.1. Structure formula of PCBs.....	4
2.2. Biphenyl pathway of microbial PCB biodegradation under aerobic condition. BphA: biphenyl-2,3-dioxygenase; BphB: <i>cis</i> -2,3-dihydroxybiphenyl dehydrogenase; BphC: 2,3-dihydroxybiphenyl 1,2-dioxygenase; BphD: 2- hydroxy-6-phenyl-6-oxohexa-2,4-dieneoate (HOPDA) hydrolase.	8
3.1. Aerial view of Indiana Harbor depicting the location of sediment samples analyzed in this study (Blue open square: core sediment samples; Red open circle: surficial sediment samples).....	31
3.2. Sediment and pore water MDPH values against sediment depth for core 1.....	32
3.3. Fraction of light, medium and heavy PCBs versus sediment depth in core 1 (A) and pore water (B), and fraction of <i>ortho</i> -, <i>meta</i> -, <i>para</i> -chlorines versus sediment depth in core sediment (C) and pore water (D).....	33
3.4. Fraction of each PCB homolog group in sediment (A) and pore water (B) against sediment depth for core 1.	34
3.5. Phylogenetic analysis of twelve amino acids of translated sequences of BphA genes retrieved from PCB-contaminated surficial sediment sample 11, 21, 56 and core 1 sediment sample (0.16-0.32 m) at IHSC. The alignment was made with ClustalX and converted into a neighboring-joining tree, which was visualized with MEGA4 with <i>Bacillus subtilis</i> Putative ring-cleaving dioxygenase as an outgroup. The filled circles at nodes indicate bootstrap values higher than 95%, while the open circles indicate bootstrap values of 75-95%. Bootstrap values lower than 75% are not shown. The bar represents 1% sequence difference.	35
3.6. Phylogenetic analysis of 16S rRNA genes retrieved from surficial and core sediment samples at IHSC. The alignment was made with ClustalX and converted into a neighboring-joining tree, which was visualized with MEGA4 with <i>Chlorofexus aurantiacus</i> J-10-fl as an outgroup. The filled diamonds represent known PCB degraders. The filled circles at nodes indicate bootstrap values of 95%, while open circles indicate bootstrap values of 75~95%. Bootstrap values lower than 75% are not shown. The bar represents 0.5% sequence difference.....	36
3.7. NMDS ordination of Wisconsin transformed T-RFLP profiles of core 1 (C01: 0-0.16 m, C02: 0.16-0.32 m...).	37
3.8. Relative abundance of phyla and proteobacterial classes for surficial sediment samples (S11, S21 and S56) and core 1 sediment samples (depth 0.16-0.32 m and 1.44-1.60 m).....	38

4.1. Fraction of light, medium and heavy PCBs versus sediment depth in core 2 (A) and pore water (B), and fraction of <i>ortho</i> -, <i>meta</i> -, <i>para</i> -chlorines versus sediment depth in core sediment (C) and pore water(D).....	57
4.2. Fraction of each PCB homolog group in sediment (A) and pore water (B) against sediment depth for core 2.	58
4.3. Sediment and pore water MDPK values against sediment depth for core 2.....	59
4.4. Abundances of total bacteria (bac), <i>bphA</i> , putative <i>Chloroflexi</i> 16S rRNA genes (chl) and <i>Dehalococcoides</i> -like 16S rRNA (dhc) against sediment depth by gene copy No. per g sediment for core 2.	60
4.5. NMDS ordination and cluster analysis of T-RFLP profiles of core 2 sediment (C01: 0-0.30 m, C02: 0.30-0.61 m...) (A), the abundance of T-RF 196 bp (B) and 198 bp+199 bp + 252 bp (C) along sediment depth.	61
5.1. The percentage of initial molar concentrations for total PCB (A), PCB 52 (B), PCB 77 (C), and PCB 153(D) after 12 weeks and 24 weeks incubation in unplanted soil (UP), switchgrass treated soil (SG), unplanted with bioaugmentation soil (UP B), switchgrass treated with bioaugmentation soil (SG B), and switchgrass treated with bioaugmentation with dead LB400 soil (SG BD). Error bars indicate the standard deviation of three soil subsamples from the same reactor.	78
5.2. qPCR analysis of (A) bacterial 16S rRNA genes and (B) <i>bphA</i> in blank (BLK), PCB spiked and unplanted soil (UP), PCB spiked and switchgrass treated soil (SG). Error bars indicate the range of two soil subsamples from the same reactor.	79
5.3. qPCR analysis of bacterial 16S rRNA gene (A) and <i>bphA</i> (B) in switchgrass treated soil (SG), switchgrass treated soil with bioaugmentation (SG B), switchgrass treated soil with autoclaved LB400 bioaugmentation (SG BD); and bacterial 16S rRNA gene (C)and <i>bphA</i> (D) in unplanted soil (UP), unplanted soil with LB400 bioaugmentation (UP B), switchgrass treated soil with bioaugmentation (SG B). Error bars indicate the range of two soil subsamples from the same reactor.	80
5.4. qPCR analysis of <i>bphA</i> transcripts in unplanted soil with bioaugmentation (UP B), and switchgrass treated soil with bioaugmentation (SG B). Error bars indicate the range of two soil subsamples from the same reactor.	81
5.5. qPCR analysis of LB400 16S-23S ITS genes in unplanted soil with bioaugmentation (UP B), and switchgrass treated soil with bioaugmentation (SG B). Error bars indicate the range of two soil subsamples from the same reactor.	82
5.6. CFU counting for unplanted soil with bioaugmentation (UP B), switchgrass treated soil with bioaugmentation (SG B) and switchgrass treated soil with dead LB400 bioaugmentation (SG BD) before and after the application of bioaugmentation. Error bars indicate the standard deviation of three soil subsamples from the same reactor.	83

6.1. The percentage of initial concentration of PCB 52, 77 and 153 after 12 and 24 weeks incubation in unplanted soil (UP), switchgrass treated soil (SG), unplanted soil with redox cycling (UPF), switchgrass treated soil with redox cycling (SG F). Error bars indicate the standard deviation of three soil subsamples from the same reactor.....	104
6.2. Transformation products detected after 12 and 24 weeks of incubation in unplanted soil (UP), switchgrass treated soil (SG), unplanted soil with redox cycling (UPF), switchgrass treated soil with redox cycling (SG F). Error bars indicate the standard deviation of three soil subsamples from the same reactor.	105
6.3. Scheme of the possible dechlorination pathways for PCB 52, 77, and 153. Dotted arrows indicate possible pathways but no intermediate was observed.....	106
6.4. Redox potential change in unplanted soil (UP), switchgrass treated soil (SG), unplanted soil with redox cycling (UPF), switchgrass treated soil with redox cycling (SG F).....	107
6.5. Moisture content change in unplanted soil (UP), switchgrass treated soil (SG), unplanted soil with redox cycling (UPF), switchgrass treated soil with redox cycling (SG F).....	108
6.6. Percentage of total PCB mass in soil, switchgrass roots, and leaves after 24 week incubation in switchgrass treated soil (SG) and switchgrass treated soil with redox cycling (SG F). Error bars indicate the standard deviation of three soil subsamples from the same reactor.	109
6.7. Phylogenetic analysis of 16S rRNA genes retrieved from switchgrass treated soils with redox cycling after 24 weeks. The alignment was made with ClustalX and converted into a neighboring-joining tree, which was visualized with MEGA4 with <i>Oscillochloris trichoides</i> as an outgroup. The filled circles at nodes indicate bootstrap values of 95%, while open circles indicate bootstrap values of 75~95%. Bootstrap values lower than 75% are not shown. The bar represents 0.5% sequence difference.....	110
6.8. qPCR analysis of <i>Dehalococcoides</i> -like 16S rRNA gene in unplanted soil with redox cycling (UPF) and switchgrass treated soil with redox cycling (SG F). Error bars indicate the range of two soil subsamples from the same reactor....	111
6.9. qPCR analysis of bacterial 16S rRNA gene in unplanted soil (UP), switchgrass treated soil (SG), unplanted soil with redox cycling (UPF), switchgrass treated soil with redox cycling (SG F). Error bars indicate the range of two soil subsamples from the same reactor.....	112
6.10. NMDS ordination of T-RFLP profiles from blank soil without PCB spiking (BLK), unplanted soil (UP), switchgrass treated soil (SG), unplanted soil with redox cycling (UPF), switchgrass treated soil with redox cycling (SG F).	113
6.11. Clone library analysis of bacterial 16S rRNA gene in unplanted soil (UP), switchgrass treated soil (SG), unplanted soil with redox cycling (UPF), switchgrass treated soil with redox cycling (SG F). Error bars indicate the range of two soil subsamples from the same reactor.....	114

6.12. T-RF importance for classification as determined by random forest algorithm.....	115
6.13. The abundance of T-RFs that were identified by random forest algorithm or corresponded to dehalorespiring bacteria in unplanted soil (UP), switchgrass treated soil (SG), unplanted soil with redox cycling (UPF), switchgrass treated soil with redox cycling (SG F).....	116

CHAPTER I. RESEARCH OVERVIEW AND HYPOTHESES

Polychlorinated biphenyls (PCBs) are a group of 209 synthetic chemicals that were widely used in industry during the last century and are still in use in closed application today. They pose a threat to environmental and public health because they are persistent, bioaccumulative and carcinogenic. Microbial PCB degradation could be a feasible way to detoxify and remove PCBs in the environment. The main objective of this work was to investigate the microbial PCB biodegradation in the contaminated soils and sediments.

The first part of this work explored the microbial PCB degradation potential in sediments from Indiana Harbor and Ship Canal (IHSC), a site that was historically contaminated by PCBs, by analyzing the PCB congener distributions and microbial communities from two core sediment samples (Chapter III and Chapter IV). The following hypothesis was tested:

Hypothesis 1: IHSC sediments contain the potential for aerobic and anaerobic microbial PCB biodegradation.

To test this hypothesis, the specific aims are to:

1. Evaluate the PCB dechlorination potential by sediment congener profile analysis.
2. Enumerate biomarker genes (biphenyl dioxygenase genes and putative dechlorinating *Chloroflexi* 16S rRNA genes) by qPCR.
3. Identify possible PCB degrading populations by T-RFLP, clone library and pyrosequencing of bacterial 16S rRNA gene.

While the field studies provided valuable information about *in situ* microbial PCB degradation, the lack of knowledge of the complex environment conditions at the contaminated site usually makes it challenging to explain the microbial activities. To better understand microbial PCB biodegradation and to examine how PCB degradation can be improved, soil microcosms spiked with PCB 52, 77 and 153 were investigated under laboratory conditions. Phytoremediation with switchgrass (*Panicum virgatum*) was employed to improve PCB removal efficiency and PCB degrading bacteria *Burkholderia xenovorans* LB400 was introduced into the switchgrass planted rhizosphere to further enhance the aerobic PCB degradation (Chapter V). The following hypotheses were tested:

Hypothesis 2: The microbial PCB degradation in soil is improved in the presence of switchgrass (*Panicum virgatum*).

Hypothesis 3: Bioaugmentation with biphenyl-grown *Burkholderia xenovorans* LB400 improves the aerobic microbial PCB biodegradation in the switchgrass rhizosphere as compared with the rhizosphere without bioaugmentation treatment.

For these hypotheses, the specific aims are to:

1. Evaluate the PCB removal efficiency by measuring PCB concentrations in unplanted, switchgrass treated, and *Burkholderia xenovorans* LB400 bioaugmented soils.
2. Evaluate the abundance of biphenyl dioxygenase genes and transcripts in unplanted, switchgrass treated, and *Burkholderia xenovorans* LB400 bioaugmented soils.

To enhance the anaerobic microbial PCB degradation, a sequential anaerobic-aerobic environment was created by redox cycling (two weeks of flooding and two weeks of non-flooding) in PCB-spiked soil microcosms (Chapter VI). The following hypothesis was tested:

Hypothesis 4: Redox cycling improves anaerobic microbial PCB biodegradation in the switchgrass rhizosphere.

The hypothesis was addressed with the following specific aims:

1. Evaluate the transformation products of PCBs in unplanted, switchgrass treated, and redox cycled soils.
2. Evaluate the abundance of dechlorinating *Chloroflexi* 16S rRNA genes in unplanted, switchgrass treated, and redox cycled soils.
3. Evaluate the microbial community structure variation and identify key members associated with PCB degradation in unplanted, switchgrass treated, and redox cycled soils.

CHAPTER II. BACKGROUND

Polychlorinated biphenyls

Polychlorinated biphenyls (PCBs) are a group of 209 synthetic chemicals consisting of 1-10 chlorine atoms on a biphenyl backbone (Figure 2.1) (Gustafson, 1970, Wiegel & Wu, 2000). PCBs were discovered over 100 years ago and were widely used in industry as transformer dielectric fluids, heat transfer fluids, hydraulic fluids, coolants and plasticizers because of their excellent chemical stability and electrical insulating properties (Erickson and Kaley, 2011). It is estimated that about 1.3 million tons of PCBs were produced worldwide between the 1930s and the 1990s, 48% of which was in US. About 30% of the production has entered the environment (Abraham et al., 2002; Breivik et al., 2002a, 2007).




Figure 2.1. Structure formula of PCBs (Wiegel and Wu, 2000).

PCBs are among the 12 worldwide priority persistent organic pollutants (POPs) listed in the Stockholm Convention and are ranked the fifth on the EPA Priority List of Hazardous Compounds (ATSDR, 1994; UNEP, 2001). PCBs has been detected in the atmosphere, water, sediment and soil from urban and rural areas (Jeremiason et al., 1994; Motelay-Massei et al., 2004; Simcik et al., 1997; Zhang et al., 2007) and even from

remote places such as Arctic (Hung *et al.*, 2010). Because of the high hydrophobicity, PCBs, especially highly chlorinated congeners, tend to adsorb to soil and sediment particles. Lightly chlorinated congeners (1-4 chlorine substituents) are more volatile and have long-range atmospheric transport potential (MacDonald *et al.*, 2000). The current total PCB burden in global background soils was estimated to be 21,000 tons (Meijer *et al.*, 2003). The PCB concentrations in contaminated sediments can be as high as several hundred ppm, while the cleanup level is 50 ppm determined by EPA.

The production, processing, distribution and commercial sale of PCBs in US were prohibited by the EPA in 1979 under Toxic Substances Control Act. Nowadays PCBs may be released into the environment by accidental spills, uncontrolled landfills and hazardous waste sites, incineration of PCB-containing wastes, and leakage from old electrical equipment in use (Breivik *et al.*, 2002b). PCBs is resistant to degradation and can accumulate through the food web, posing a potential risk to public health and ecosystem (Campfens and MacKay, 1997). People may be exposed to PCBs mainly by food consumption, including meat, milk and fish, and by inhaling contaminated air (Huwe *et al.*, 2009). Some old building materials such as pigment and caulk contain PCBs which can migrate into air, dust and surrounding material and cause potential exposure (Herrick *et al.*, 2007; Hu and Hornbuckle, 2010).

PCBs are categorized as probably carcinogenic to humans by EPA and as Group I human carcinogen by the International Agency for Research on Cancer (IARC), mainly based on the evidence in animal studies and mechanistic considerations. PCBs are also endocrine disruptors that are capable of interfering with natural hormones in the human body (Grimm *et al.*, 2013). Other adverse health effects include alternation of the immunological system, the neurological system, skin, liver, and the cardiovascular system. PCBs also present a potential reproductive hazard to humans and prenatal exposure to PCBs may result in developmental deficiency in the offspring (Longnecker *et al.*, 1997; Ross, 2004; Van den Berg *et al.*, 2006). The tolerable daily intake (TDI) of

PCB is 1 µg/kg body weight (MinRat-Wirtschaftsministerium et al., 1995). Currently the EPA has established a maximum contaminant level goal of zero and a maximum contaminant level of 0.5 ppb for PCBs in public drinking water supplies (USEPA, 1991).

Microbial PCB biodegradation

Microorganisms are known to degrade PCBs via two general processes: aerobic oxidative mineralization and anaerobic reductive dechlorination. Aerobic PCB biodegradation is usually restricted to lightly chlorinated congeners (no more than 4 chlorine atoms per molecule), while PCBs with five or more chlorine atoms per molecule are generally resistant to aerobic degradation. Chlorine removal via anaerobic reductive dechlorination processes could make these congeners more amenable to aerobic degradation (Borja *et al.*, 2005).

Under aerobic conditions, the PCB congener biphenyl ring can be opened by an oxidative ring cleavage mechanism (Pieper and Seeger, 2008). A variety of aerobic bacteria that oxidize PCBs in this manner have been identified, including Gram-negative strains of *Pseudomonas*, *Alcaligenes*, *Achromobacter*, *Janibacter*, *Burkholderia*, *Acinetobacter*, *Comamonas*, *Sphingomonas*, *Ralstonia*, and *Enterobacter* and Gram-positive strains of *Corynebacterium*, *Rhodococcus*, *Bacillus*, *Paenibacillus*, *Arthrobacter* and *Micrococcus* (Bedard et al., 1986; Bopp, 1986; Furukawa, 2000; Macková et al., 2010; Pieper and Seeger, 2008).

PCBs can be used as carbon and energy sources by some strains (such as *Burkholderia xenovorans* LB400 and *Burkholderia cepacia* P166) and be degraded by biphenyl-degrading bacteria via co-metabolism (Field and Sierra-Alvarez, 2008; Gibson et al., 1993; Hernandez et al., 1995). Most aerobic PCB-degraders employ an enzyme system called the upper biphenyl degradation pathway. This pathway is catalyzed by a series of enzymes designated BphA, BphB, BphC and BphD. Biphenyl 2,3-dioxygenase

(BphA), a Rieske non-heme iron dioxygenase, catalyzes the incorporation of two oxygen atoms into the aromatic ring to form an arene *cis*-diol (Gibson and Parales, 2000). This initial step in the upper biphenyl pathway transforms susceptible PCB congeners to the corresponding *cis*-dihydrodiol (Furukawa, 1994). Microbes can typically metabolize one of the biphenyl rings to 2-hydroxypenta-2,4-dienoate, which is further degraded to metabolites that can enter the citric acid cycle, a central metabolic pathway. However, this process often leaves behind a chlorobenzoate, which most mediating PCB-degrading microbes cannot degrade further (Pieper and Seeger, 2008). Therefore a population of chlorobenzoate-degraders would be required to achieve mineralization of PCB congeners. Chlorobenzoate-degraders (such as *Pseudomonas. cepacia* and *Pseudomonas. aeruginosa*) have been previously isolated from PCB-contaminated soil and sediment (Pavlů *et al.*, 1999). PCB-degrading and chlorobenzoate-degrading microorganisms can be genetically modified and thus degrade PCBs without accumulation of chlorobenzoates (Rodrigues *et al.*, 2006; Saavedra *et al.*, 2010).

Biodegradation efficiencies for different PCB congeners vary based on the number and position of the chlorine substituents. PCB congeners with chlorines on one aromatic ring are more easily degraded than those with chlorine substituents on both rings (Pieper, 2005). Highly chlorinated and *ortho*-chlorinated PCB congeners appear to be particularly resistant to biodegradation (Field and Sierra-Alvarez, 2008). Higher chlorination levels result in a highly hydrophobic PCB molecule, a property which reduces their accessibility to enzymes. The bulkiness of the chlorine atoms may also prevent access to the enzyme's active site (Bedard *et al.*, 1986). Chlorine atoms on the *ortho* position may prevent oxygenases from attacking the aromatic ring. It is also reported that *ortho*-chlorinated PCB metabolites strongly inhibit the dioxygenase that catalyzes the third step of the biphenyl degradation pathway, promoting its suicide inactivation and interfering with the degradation of other compounds (Dai *et al.*, 2002). Furthermore, some metabolites of PCB degradation via the upper biphenyl pathway,

including dihydrodiols, dihydroxybiphenyls and chlorinated 2-hydroxy-6-phenyl-6-oxohexa-2,4-dieneoates (HOPDAs), can be very toxic and accumulate in bacteria, adversely affecting cell viability and growth rates even more than the parent PCB congeners (Camara *et al.*, 2004).




Figure 2.2. Biphenyl pathway of microbial PCB biodegradation under aerobic condition. BphA: biphenyl-2,3-dioxygenase; BphB: *cis*-2,3-dihydroxybiphenyl dehydrogenase; BphC: 2,3-dihydroxybiphenyl 1,2-dioxygenase; BphD: 2-hydroxy-6-phenyl-6-oxohexa-2,4-dieneoate (HOPDA) hydrolase (Pieper and Seeger, 2008).

Under anaerobic conditions, some members of the *Chloroflexi* phylum, including *Dehalococcoides* spp. , and the o-17/DF-1 group, can use certain PCB congeners as electron acceptors and transform them into less chlorinated congeners by reductive dechlorination (Adrian *et al.*, 2009; Cutter *et al.*, 2001; Fagervold *et al.*, 2007; Fagervold *et al.*, 2005; Wiegel and Wu, 2000; Yan *et al.*, 2006). Only several PCB dechlorinating bacteria have been isolated, including *Dehalococcoides maccartyi* strain CBDB1, *Dehalococcoides maccartyi* strain 195 and *Dehalobium chlorocoeria* DF-1 (Adrian *et al.*, 2009; Fennell *et al.*, 2004; May *et al.*, 2008). *Dehalobacter* phylotypes have been found

in an sediment-free culture (KFL culture) which dechlorinated 2,3,4,5-tetrachlorobiphenyl and 2,3,4-trichlorobiphenyl where no *Dehalococcoides* or o-17/DF-1 group were detected, which makes *Dehalobacter* a new candidate for anaerobic PCB-degrader (Yoshida *et al.*, 2009).

During microbial PCB dechlorination, chlorines are replaced by hydrogen ions and released as chloride ions, a process catalyzed by membrane-associated reductive dehalogenases (RDases) (Hiraishi, 2008; Holliger *et al.*, 1998; Janssen *et al.*, 2001; Wiegel and Wu, 2000). However, so far no reductive dehalogenase is identified for PCB dechlorination (Sowers and May, 2013). Currently at least eight distinct PCB dechlorination patterns have been identified in sediments and were referred to as M, Q, H', H, P, N, LP and T (Table 2.1) (Wiegel and Wu, 2000). Chlorine atoms at the *meta*- and *para*- positions are preferentially removed in many of these dechlorination patterns, which result in the generation of *ortho*-substituted PCB congeners (Tiedje *et al.*, 1993). Dechlorination of chlorine at *ortho*-position is rare but has been reported (Berkaw *et al.*, 1996; Vandort and Bedard, 1991; Wu *et al.*, 1998). Microbial dechlorination is strongly influenced by various environmental factors such as temperature, pH, carbon sources, and the presence of other electron acceptors (Wiegel and Wu, 2000).

Table 2.1. Positions of chlorines removed by different dechlorination patterns. (Wiegel and Wu, 2000)

Dechlorination process	Susceptible chlorines
M	Flanked and unflanked <i>meta</i>
Q	Flanked and unflanked <i>para</i> , <i>meta</i> of 23-group
H'	Flanked <i>para</i> , <i>meta</i> of 23-, and 234-groups
H'	Flanked <i>para</i> , doubly flanked <i>meta</i>
P	Flanked <i>para</i>
N	Flanked <i>meta</i>
LP	Flanked and unflanked <i>para</i>
T	Flanked <i>meta</i> of 2345-group, in hepta- and octachlorobiphenyls

Microbial PCB degradation in sediments and soils

Studies have been conducted to investigate the structure and function of bacterial communities in PCB-contaminated soil and sediment systems. Microcosms are a common approach to investigate both aerobic and anaerobic *in situ* PCB biodegradation potential and PCB degrading communities. Microbial diversity in the surface of an Aroclor 1260-laden sandy soil in Quebec, Canada was analyzed, and sequences of twenty clones were found to be related to low G+C gram-positives, *Cytophaga/Flexibacter/Bacteroides*, and the α -, β -, γ -*Proteobacteria* (Lloyd-Jones and Lau, 1998). In another study, metabolically active bacteria were identified by 16S rRNA gene clone library analysis in surface samples of a PCB-contaminated sandy soil in Germany. Clone sequences were found to cluster within the *Proteobacteria*, the *Holophage-Acidobacterium* phylum, the *Actinobacteria*, and the *Planctomycetales*. The β -*Proteobacteria* were abundant, mostly the genera *Burkholderia* and *Variovorax*. Sequences similar to known PCB-degraders and other organic pollutant degraders were also identified (Nogales *et al.*, 1999). In a microcosm study using aerobic agitated soil slurries with PCB congener spiking, an increase of β -*Proteobacteria* and *Actinobacteria* was observed in all microcosms exposed to PCBs (Correa *et al.*, 2010). Stable isotope

probing (SIP) was also used to explore the bacterial groups in PCB-contaminated environment. *Burkholderia* species were found dominant and active in aerobic PCB biodegradation in a biofilm community grown directly on PCB droplets, as revealed by DNA-SIP analysis of 16S rRNA (Tillmann *et al.*, 2005). In microcosms containing River Raisin sediment with ^{13}C -biphenyl addition, genera *Achromobacter*, *Pseudomonas*, *Acidovorax*, *Ramlibacter*, *Azoarcus* and *Hydrogenophaga* were detected after 14 days incubation but not at the beginning of incubation. The dominant bacterial groups were found closely related to previously known PCB oxidizers, such as *Achromobacter xylooxidans* KF701, *Acidovorax* sp strain KKS102, and *Hydrogenophaga taeniospiralis* IA3-A. Analysis of aromatic-ring-hydroxylating dioxygenase genes revealed two sequence groups similar to *bphA* of *Comamonas testosteroni* strain B-356 and *Rhodococcus* sp. RHA1 (Sul *et al.*, 2009). The impact of bioremediation of PCB-contaminated sites on the indigenous microbial community was also studied. Changes in the bacterial community structures were observed, after long-term biostimulation treatment with carvone, soya lecithin and xylose, and bioaugmentation treatments with TSZ7 mixed culture and with a *Rhodococcus* sp Z6 pure strain. *Actinobacteria*, *Bacteroidetes*, α - and γ - *Proteobacteria* were more abundant under all three bioremediation treatments, with *Actinobacteria* representing the dominant phylum (Petrić *et al.*, 2011a).

Experiments were also conducted to describe the anaerobic PCB-degrading communities in PCB-contaminated environment. Microorganisms from sediments of Hudson River, NY, and Silver Lake, MA were found capable of removing *meta*- and *para*-chlorines of four Aroclors (Aroclor 1242, 1248, 1254 and 1260) (Quensen *et al.*, 1990). Similar results were described for enrichment cultures from Housatonic River sediments (Bedard *et al.*, 1996; Bedard *et al.*, 1997). Efforts were also made to characterize the dechlorinating populations. 16S rRNA sequences similar to the theta, low-G+C, Gram-positive, and *Thermotogales* subgroups and one sequence similar to

Dehalococcoides ethenogenes were found to be associated with PCB *ortho*-dechlorination in a sediment microbial community from Baltimore Harbor, MD (Holoman *et al.*, 1998). In PCB dechlorinating enrichment cultures that selectively dechlorinated double-flanked chlorines in 2,3,4,5-tetrachlorobiphenyl, a low G+C Gram-positive eubacterium, an organism similar to green non-sulfur bacteria, an *Aminobacterium* sp. and a *Desulfovibrio* sp. were identified (Watts *et al.*, 2001). *Dehalococcoides* and o-17/DF-1 type *Chloroflexi*, *Spirochetes* and *Bacteroidetes* were identified in enriched Baltimore Harbor sediment microcosms, and a twenty five fold increase of putative dechlorinating *Chloroflexi* members (e.g. *Dehalococcoides* and o-17/DF-1 type *Chloroflexi*) was observed during the dechlorination of Aroclor 1260, which suggests the dechlorination activity of *Chloroflexi* members in Baltimore Harbor sediments (Fagervold *et al.*, 2007; Fagervold *et al.*, 2005; Watts *et al.*, 2005). In another study, active indigenous PCB dechlorinating communities were identified in sediments from three PCB-contaminated sites, Anacostia River, DC, Buffalo River, NY, and Grasse River, NY. Notably, an enriched dechlorinating population consisting exclusively of *Dehalococcoides*-like phylotypes was detected in Grasse River sediment and was found to be associated with a significantly higher dechlorination rate. PCB congener profiles, concentrations, and total organic carbon concentrations (TOC) were found to be associated with *in situ* attenuation rates (Kjellerup *et al.*, 2008). A *Dehalococcoides*-like population was found in three anaerobic cultures capable of reductively dechlorinating 2,3,4,5-tetrachlorobiphenyl but not in PCB-free microcosm. The role of the *Dehalococcoides*-like populations in the removal of 2,3,4,5-tetrachlorobiphenyl was evidenced by their disappearance when dechlorination was inhibited or when PCBs was consumed (Yan *et al.*, 2006). Sequence phylotypes related to the genus *Sulfurovum*, and the species *Desulfococcus multivorans* and two *Chloroflexi* were found to be enriched throughout incubation time in PCB-spiked microcosm consisting of marine sediments from Venice Lagoon, suggesting such bacteria were involved in PCB dechlorination in

sediments (Zanaroli *et al.*, 2010). Relatively higher abundance of dechlorinating *Chloroflexi*, *Dehalococcoides*, and o-17/DF-1 populations were observed in deep PCB contaminated sediments (Xu *et al.*, 2012). For the above studies, *Dehalococcoides*-specific 16S rRNA genes were usually targeted for the characterization of dechlorinating microbial population, since no PCB dehalogenase has been identified yet.

Bioaugmentation with both aerobic and anaerobic PCB degraders has been applied to remediate PCB contaminated soils and sediments. Various aerobic PCB degraders, including *Burkholderia xenovorans* LB400, *Ralstonia eutrophus* H850, *Arthrobacter* sp. strain B1B, have been used and achieved significantly greater PCB removal in the bioaugmented soil compared with native soil (Egorova *et al.*, 2013; Luepromchai *et al.*, 2002; Singer *et al.*, 2000). The efficiency of bioaugmentation with dehalorespiring bacteria, such as *Dehalococcoides mccartyi* strain 195 and *Dehalobium chlorocoercia* DF1, were tested in microcosms containing PCB-contaminated sediments, and it was found that bioaugmentation enhanced PCB dechlorination rates (Krumins *et al.*, 2009; Payne *et al.*, 2011). The efficiency of bioaugmentation with both *Dehalobium chlorocoercia* DF1 and *Burkholderia xenovorans* LB400 was also tested and a significantly greater PCB mass loss compared with non-bioaugmented sediments was observed. DF 1 was found to reduce the lag time for PCB degradation, but had no significant effect on the final PCB concentration (Payne *et al.*, 2013).

Phytoremediation of PCBs

Remediation of PCBs has always been a challenge because of the refractory nature of PCB congeners. Many methods including thermal, oxidative, reductive, microbial approaches have been evaluated to enhance PCB degradation (Gomes *et al.*, 2013). Among those methods, phytoremediation, the use of plants and associated bacteria

to remediate contaminated environment, is more cost effective and sustainable (Van Aken *et al.*, 2010).

Phytoremediation comprises a range of processes including phytoextraction, rhizofiltration, phytotransformation, phytovolatilization, rhizoremediation, and phytostabilization (Dietz and Schnoor, 2001; Schnoor *et al.*, 1995; Van Aken *et al.*, 2010). Plants can uptake PCBs from root systems and translocate them to other plant parts (Macková *et al.*, 2006). Studies of PCB congener fate within whole poplar plants (*Populus deltoids x nigra*) in hydroponic exposures revealed that PCBs initially sorbed to the root system were then translocated to the stem and the woody stem accumulated more PCBs than the shoots (Liu and Schnoor, 2008). The root systems of some plant species such as *Cucurbita pepo* (including pumpkin and zucchini), *Carex normalis*, *Brassica nigra* can accumulate PCBs from soils with bioaccumulation factors (BAFs = PCB concentrations in plant tissues / PCB concentrations in soils) higher than one, while the PCB concentrations in shoots were usually lower than those in the roots (Ficko *et al.*, 2010; Greenwood *et al.*, 2011; White *et al.*, 2006; Zeeb *et al.*, 2006). Shoot BAFs ranged from 0.14 – 0.45 were observed for *Cucurbita pepo* and *Carex normalis* (White *et al.*, 2006; Whitfield Åslund *et al.*, 2007; Zeeb *et al.*, 2006). However, shoot BAFs greater than one were observed for *Polygonum persicaria* and *Vicia cracca* (Ficko *et al.*, 2010), and BAFs of lower parts of plant shoot achieved as high as two were reported for *Cucurbita pepo* (Whitfield Åslund *et al.*, 2008).

PCB metabolism by plants is different from that of microorganisms, and varies among plant species (Van Aken *et al.*, 2010). The substitution pattern and the degree of chlorination of PCB congeners also affect transformation by plants (Macková *et al.*, 2006). PCB transformation in plants was found to result in the formation of hydroxylated PCB metabolites (Liu *et al.*, 2009; Macková *et al.*, 2001; Rezek *et al.*, 2007; Zhai *et al.*, 2010a; Zhai *et al.*, 2010b).

Plant-microbial interactions are known to play important roles in enhancing the degradation of xenobiotics, including PCBs, within the rhizosphere, a dynamic environment for microorganisms, which is defined as the zone of soil under direct influence of plant roots that usually extends a few millimeters from the root surface (Macková *et al.*, 2007). Microorganisms can enhance the plant uptake of certain mineral nutrients and can modify the rhizosphere by producing extracellular enzymes and plant growth factors. Plants play critical roles by supplying organic compounds in exudates that can be utilized by microorganisms as carbon and energy sources and regulating deposits of soil water. Some secondary plant metabolites could induce the genes encoding enzymes involved in the degradation of xenobiotics. Plant released compounds can also solubilize contaminants and make them more available to both plant and microbial degradative enzymes (Leigh *et al.*, 2006; Macková *et al.*, 2009; Singer *et al.*, 2003a).

Several studies have demonstrated that the presence of specific plants enhances PCB biodegradation and increases microbial PCB-degrading populations. Austrian pine (*Pinus nigra*) and goat willow (*Salix caprea*) were found to increase the abundance of microbial PCB-degraders in the root zones (Leigh *et al.*, 2006). Crushed fine roots of felt-leaf willow (*Salix alaxensis*) added to soils spiked with PCB congeners were found to effectively increase PCB biodegradation rate (Slater *et al.*, 2011). A stable isotope probing study (SIP) revealed that seventy-five different biphenyl-degrading genera and twenty seven associated functional genes were active in the root zone of an Austrian pine (*Pinus nigra L.*) grown in PCB-contaminated soil (Leigh *et al.*, 2007). Another SIP study demonstrated that the cultivation of horseradish (*Armoracia rusticana*) may affect the microorganisms that are potentially responsible for PCB degradation (Uhlík *et al.*, 2009). Increased numbers of active PCB-degraders in the rhizosphere of goat willow (*Salix caprea*) and horseradish (*Armoracia rusticana*) have also been observed, indicating these

plant species are promising candidates for rhizoremediation of PCB-contaminated soils (Ionescu *et al.*, 2009).

Some plant-released compounds have the capacity to support the growth of PCB-degraders and thereby improve PCB degradation via co-metabolism (Singer *et al.*, 2003a). For example, flavonoids fostered the growth of PCB degraders (such as *Burkholderia xenovorans* LB400 and *Rhodococcus erythropolis* U23A) and were able to induce the biphenyl catabolic pathway (Donnelly *et al.*, 1994; Leigh *et al.*, 2002; Toussaint *et al.*, 2012). Plant terpenes are also natural substrates that can support PCB oxidation (Hernandez *et al.*, 1997; Park *et al.*, 1999; Tandlich *et al.*, 2001). Transformation of Aroclor 1242 occurred completely in PCB-contaminated soils amended with orange peel, ivy leaves, pine needles, or eucalyptus leaves, but not in the non-amended soil. Also, the abundance of biphenyl-degrading microbes in the amended soils were five orders of magnitude higher than in the non-amended soils, indicating terpenes are used as a natural substrate by PCB degraders (Dudášová *et al.*, 2012; Focht, 1995; Hernandez *et al.*, 1997). In another study, S-carvone, a terpenoid compound, was reported to induce the biotransformation of Aroclor 1242 by *Arthrobacter* sp. strain B1B (Gilbert and Crowley, 1997).

Plant-microbial mutual co-operation within the rhizosphere on the level of the intermediates transformation were also studied. Chlorobenzoates are a common end product of upper biphenyl degradation pathway that usually are not degraded further by aerobic PCB degraders and thus may accumulate to inhibit microbial PCB biodegradation (Pieper and Seeger, 2008). Plant tissues of horseradish and black nightshade were reported to show significant transformation of this bacterial product of PCB degradation (Macková *et al.*, 2007). Similarly, hydroxylated PCBs, the product of PCB degradation by plants, can be transformed by PCB degrading microbes. *Ortho*-chlorinated hydroxybiphenyls were degraded by purified biphenyl dioxygenases from *Burkholderia xenovorans* LB400 and *Comamonas testosteroni* B-356 (Francova *et al.*, 2004). Despite

this, hydroxylated PCBs still appear to exhibit higher toxicity than PCBs to microorganisms (Camara et al., 2004; Purkey et al., 2004).

CHAPTER III. MICROBIAL PCB DEGRADATION POTENTIAL IN CORE 1 FROM IHSC

Introduction

Polychlorinated biphenyls (PCBs) are a family of 209 congeners that were produced and used during the last century. They remain a major concern to the public and environmental health because of their high stability and potential toxicity (Bedard, 2004).

Despite their environmental persistence, PCBs can be removed and detoxified by anaerobic and aerobic microorganisms. Microorganisms are known to degrade PCBs via two general processes: anaerobic reductive dechlorination and aerobic oxidative biphenyl ring cleavage (Borja *et al.*, 2005). Under anaerobic conditions, some members of the phylum *Chloroflexi*, including *Dehalococcoides* spp. and the o-17/DF1 group, use certain PCB congeners as electron acceptors and transform them into less chlorinated congeners (Adrian *et al.*, 2009; Wu *et al.*, 2002). Under aerobic conditions, some aerobic PCB-degraders employ the upper biphenyl degradation pathway, and biphenyl 2,3-dioxygenase (BphA) catalyzes the first step in this pathway, transforming susceptible PCB congeners into the corresponding *cis*-dihydrodiol (Furukawa and Fujihara, 2008; Gibson and Parales, 2000).

The Indiana Harbor and Ship Canal (IHSC) in the East Chicago, Indiana, is located on the southern shore of Lake Michigan. IHSC is heavily contaminated with a variety of pollutants, including heavy metals, polycyclic aromatic hydrocarbons (PAHs) and PCBs due to years of industrial operations in the area (Ingersoll *et al.*, 2002; Martinez *et al.*, 2010). It is reported to be a large source of PCBs to Lake Michigan, with the PCB concentration in surficial sediments of IHSC ranging from 53 to 35,000 ng g⁻¹ dry weight according to an intensive survey (Martinez *et al.*, 2010). To remain viable for large industrial vessels, a long-term dredging project is carried out by the U.S. Army Corps of Engineers, Chicago District to ensure an adequate navigational depth.

PCB-contaminated IHSC sediments are being dredged and permanently stored in a confined disposal facility (CDF) located north of Lake George Branch (USACE, 2013).

The objective of this study is to assess the *in situ* PCB degradation potential in IHSC sediments. The vertical pattern of a core sediment sample (core 1) was analyzed and suggests the possibility of *in situ* dechlorination. The presence of dechlorinating groups and *bphA* indicates that microbial communities in the sediments had the potential to degrade PCBs.

Materials and Methods

Sediment sample collection

In August 2006, surficial sediment samples (Figure 3.1) were collected from IHSC from aboard the U.S. Environmental Protection Agency's research vessel *Mudpuppy*. A standard ponar dredge sampler (top 10 cm layer) was used (Martinez *et al.*, 2010). In May 2009, one 4.57 m core sample (core 1) was collected using a submersible vibro-coring system with a PVC tube (length 457 cm, internal diameter 9.5 cm) (Figure 3.1). To examine microorganisms at different sediment depths, core 1 was sectioned every 0.152 m. Samples were placed in plastic bags and kept on ice during transportation. Sediment samples were stored at 4°C in the lab until analysis.

PCB congener analysis¹

Preparation, extraction and clean-up steps for measuring PCB sediment concentrations have been described previously (Martinez and Hornbuckle, 2011). Briefly, sediments were extracted using pressurized fluid extraction (Accelerated Solvent

¹ The PCB concentration measurement of sediment samples was performed by Andres Martinez, Ph.D., Department of Civil & Environmental Engineering, The University of Iowa.

Extractor, Dionex ASE-300). The extracts were concentrated and eluted through a multilayer silica gel column. Activated granulate copper was used to remove sulfur in solution. Poly-dimethylsiloxane (PDMS) coated fibers were used as passive samplers to determine the sediment pore water PCB concentration. PCB extraction and quantification procedures were also reported previously (Martinez *et al.*, 2013). PCB identification and quantification were conducted employing a modified US EPA method 1668C (USEPA, 2010). Tandem mass spectrometry GC/MS/MS (Quattro Micro GC, Micromass MS Technologies) in multiple reaction monitoring mode was utilized to quantify all 209 congeners in 161 individual or coeluting congener peaks.

The molar dechlorination product ratio (MDPR) was used to examine possible PCB dechlorination in core sediments. When determining the MDPR, it was assumed that exclusively *ortho*-chlorinated PCB congeners undergo no further dechlorination (TAMS Consultants, 1997). In this study, five exclusively *ortho*-chlorinated PCBs (PCB 1, 4, 10, 19, 54) and PCB 8 were selected as the ultimate dechlorination products. The ratio of the sum of the molar concentrations of selected congener over total PCB molar concentration was calculated and defined as MDPR. PCB 8 was considered as a dechlorination product because the proportions of PCB 8 in core sediments (averaged 1.4% of total PCB) were much higher than that in other Aroclor commercial mixtures (0.48%, 0%, 0% of Aroclor 1248, 1242, 1016, respectively), which indicates the production of this congener in IHSC sediments.

Terminal-Restriction Fragment Length Polymorphism (T-RFLP) analysis

Total chromosomal DNA from sediments was extracted using the MoBio Ultraclean Soil DNA Isolation Kit (MoBio Laboratories, Inc., Carlsbad, CA) and stored them at -20°C until samples were analyzed with T-RFLP, pyrosequencing, and functional gene sequencing. To characterize the bacterial community composition, T-RFLP was

performed on PCR-amplified 16S rRNA genes. After initial PCR with primer set 8F/1492R and another round of PCR with a fluorescence-labeled primer set (6-FAM 8F/533R), we purified the PCR products using QIAquick PCR purification kit (Qiagen Inc, Valencia, CA), and digested purified PCR products with the restriction enzyme *MspI* (New England BioLabs, Inc., Ipswich, MA). Then we precipitated DNA by centrifugation at $17,800 \times g$ for 15 min after 2h incubation at -20°C with Glycogen (Fermentas International Inc., Vilnius, Lithuania), sodium acetate (pH 5.2), and ethanol and resuspended the pellet in distilled water (Invitrogen Corp., Carlsbad, CA). We sent digested DNA to the University of Iowa DNA facility for electrophoresis using an Applied Biosystems 3730 DNA analyzer (Life Technologies Corporation, Carlsbad, CA) with the GeneScan 500 LIZ size standard.

After processing T-RFLP profiles with Peak Scanner software (Applied Biosystems, Carlsbad, CA) and T-REX software (Culman *et al.*, 2009), a TRF size matrix containing 29 samples (rows) and 490 unique 16S rRNA gene TRFs (columns) was generated and was ordered by non-metric multidimensional scaling (NMDS) with the Wisconsin transformation and the Bray-Curtis dissimilarity index. The final stress was 9.93. PCB congener profiles were ordered by principal components analysis (PCA) and the correlation between PCB congener profile and T-RFLP profile was tested by Procrustes test. The Mantel test was also applied to assess the correlation between T-RFLP profiles and PCB profiles, where a Euclidean dissimilarity index was calculated for PCB congener profiles and the Bray-Curtis dissimilarity index was calculated for T-RFLP profiles. Pearson correlation coefficient between each T-RF peak and MDP values were calculated (González *et al.*, 2000).

Bacterial tag-encoded amplicon pyrosequencing²

To get more detailed sequence information about bacterial communities, pyrosequencing for bacterial 16S rRNA gene was performed for three surficial sediment samples and two sections from core 1 (0.15-0.30 m and 1.37-1.52 m). Bacterial 16S rRNA gene fragments were amplified using barcoded primer sets BSF8/USR515 under conditions previously described (Bibby *et al.*, 2010). PCR products were purified with QIAquick PCR purification kit (Qiagen Inc, Valencia, CA) and sent for pyrosequencing on a 454 GS-FLX sequencer (Roche Diagnostics Corporation, Indianapolis, IN) using the Titanium Sequencing Kit (Roche, Branford, CT).

Raw pyrosequencing data was processed on RDP pyrosequencing pipeline (<http://pyro.cme.msu.edu/pyro/>) (Cardenas *et al.*, 2009). After quality check, trimming of barcodes and filtering of bad reads, sequences were aligned and assigned identities. Operational taxonomic units (OTUs) were generated based on 97% sequence similarity.

PCR amplification, cloning and sequencing of biphenyl dioxygenase gene (BphA) and *Dehalococcoides*-specific 16S rRNA

BphA genes were amplified using various primer sets, including bphA10, bphA20, bphA30, bphA40, and bphA50 (Hoostal *et al.*, 2002). Each reaction contained 4 μ M of each primer and 25-50 ng of DNA template. PCR reaction conditions were as follows: initial denaturing step for 4 min at 94°C, followed by 40 cycles of denaturing for 1 min at 94°C, annealing for 3 min at 40-45°C, and extending for 4 min at 72°C with a final extension step at 72°C for 10 min.

² The pyrosequencing experiment was performed by Joshua Livermore, Ph.D., Department of Civil & Environmental Engineering, The University of Iowa

Dehalococcoides-specific 16S rRNA genes were amplified with previously published DHC primer sets (Hendrickson *et al.*, 2002). Each reaction contained 20 ng of DNA template and 20 pM of each primer. Reaction conditions were as follows: 2min denature at 95°C, followed by 40 cycles of 1 min denature at 94°C, 1 min anneal at 55°C, and 1 min extend at 72°C and a final extension of 5 min at 72°C. All PCR experiments were conducted using the Qiagen Master Mix PCR Kit (Qiagen Inc, Valencia, CA).

PCR products were purified by Qiaquick Gel Extraction Kit (Qiagen Inc, Valencia, CA), cloned into the pDRIVE vector using the Qiagen PCR Cloning Kit (Qiagen Inc, Valencia, CA), and transformed into NEB 5-alpha F'I^d competent *E.coli* cells (New England BioLabs, Inc., Ipswich, MA). Recombinant *E.coli* were plated on Luria Broth agar with kanamycin (30 mg/L), X-gal (0.4 mg/plate) and IPTG (10⁻⁴ mmol/plate) and incubated overnight at 37°C. Plasmid DNA was extracted from white colonies using the Qiagen Miniprep Plasmid Extraction Kit (Qiagen Inc, Valencia, CA) and the inserts were sequenced at the University of Iowa DNA facility. We then analyzed sequence data using Ribosomal Database Project (RDP) pipeline (Cole *et al.*, 2007) and Basic Local Alignment Search Tool (BLAST) (Altschul *et al.*, 1997) and performed multiple sequence alignments and phylogenetic tree construction using ClustalX2 programme (Thompson *et al.*, 1997). MEGA 4 was used to graphic present the results (Tamura *et al.*, 2007).

Results and Discussion

PCB congener analysis

The vertical pattern of PCB congener distribution of core 1 was examined to search for the evidence of *in situ* PCB dechlorination. Changes of PCB concentration and congener distribution along sediment depth could be the result of the discharge of different commercial mixtures, sediment resuspension and transport, and PCB

biodegradation (Li *et al.*, 2009). Anaerobic PCB biodegradation usually leads to congener profile shift from higher chlorinated congeners to lower chlorinated ones. Furthermore, anaerobic PCB degraders preferentially remove chlorine atoms at *meta*- and *para*-position, resulting in the increase of *ortho* chlorine fraction. Sediment suspension and transportation may lead to accumulation of highly chlorinated congeners because lightly chlorinated PCBs are relatively less hydrophobic and more volatile, and tend to migrate from sediment to water and atmosphere.

For core 1, the fraction of lesser chlorinated PCBs (1-3 chlorine substituents) increased along sediment depth till 2.4 m, and then decreased. Sediment depth 0-2.4 m is where the total PCB concentrations were high (averaged 52,000 ng/g⁻¹ d.w.). Sediment deeper than 2.4 m had much lower PCB concentrations (averaged 2,500 ng/g⁻¹ d.w.), which section corresponded to the time period before PCBs were widely used (before 1930s) (Martinez and Hornbuckle, 2011). Thus the accumulation of lesser chlorinated PCBs in sediments 0-2.4 m may suggest the microbial dechlorination. At the same sediment depth, the fraction of *ortho*-Cl kept relative stable, but MDP values increased, which suggests that PCB dechlorination has occurred (Figure 3.2, Figure 3.3). Furthermore, although the sediment PCB profile was reported to resemble the Aroclor 1248, the average MDP value for sediments of 0-2.4 m (0.031) was much greater than that of Aroclor 1248 (0.009), which also indicates the *in situ* PCB dechlorination (Frame *et al.*, 1996; Martinez and Hornbuckle, 2011).

Detection of *bphA* and *Dehalococcoides*-specific

16S rRNA genes in IHSC sediments

The signal of *bphA* gene was detected by PCR in core sediment samples water-sediment surface to 2.4 m deep (Table 3.1). In deeper sediments, *bphA* genes were not amplified, possibly because of the intense anaerobic conditions. Eight different partial *bphA* sequences were identified by clone library, six of which were 93-98% identical to

BphA from *Acidovorax* sp. KKS102 (GenBank Acc. No. BAJ72245). These sequences were also 91-97% similar to BphA from *Comamonas testosteroni* TK102 (GenBank Acc. No. BAC01052), also a known PCB degrader. The other two sequences were 98-99% similar to BphA in *Pseudomonas putida* B2-6 (GenBank Acc. No. ACN62349).

We also identified four partial *bphA* sequences from S11, five partial *bphA* sequences from S21 and one sequence from S56. Two sequences of S21 were 95~96% identical to BphA from uncultured bacterium (GenBank Acc. No. ACL99824). The two sequences were 95~96% similar to BphA from *Pseudomonas alcaligenes* B357 (GenBank Acc. No. ABR08355), respectively (Uhlík et al., 2009). Two other sequences were 98~99% identical to ring hydroxylating dioxygenase alpha subunit in *Polaromonas naphthalenivorans* CJ2 (GenBank Acc. No. ABM39439), which is a naphthalene-degrading bacterium (Jeon et al., 2004). Another five sequences were 98~99% identical to BphA in *Pseudomonas putida* B6-2 (GenBank Acc. No. ACN62349). One sequence was 48~54% similar to BphA from *Acidovorax* sp. KKS102 (GenBank Acc. No. BAJ72245), a soil PCB degrading bacterium (Kimbara et al., 1988). A phylogenetic tree of translated *bphA* sequences shows the relationship among various aerobic PCB-degrading organisms and the *bphA* sequences retrieved from the sediment samples, revealing the diversity of those obtained *bphA* sequences (Figure 3.5). The successful identification of *bphA* genes from S11, S21, S56 and core 1 (0.16-0.32 m) indicates possible occurrence of aerobic PCB degradation.

The signal for *Dehalococcoides*-specific 16S rRNA gene was negative till 0.91 m, and was positive between 0.91 - 3.05 m, then disappeared till the end of the core. Five unique *Dehalococcoides*-specific 16S rRNA sequences were obtained from core sediments at the depth 1.44-1.60 m. Three sequences were 99-100% identical to 16S rRNA gene from *Dehalococcoides* sp. enrichment culture clone CT3-2 (GenBank Acc. No. JF698644), which was isolated from a 2,4,6-trichlorophenol dechlorinating microcosm. The other two sequences were 96% similar to uncultured *Chloroflexi*

bacterium 16S rRNA gene (GenBank Acc. No.CU923702). These *Dehalococcoides* sequences were closely related to known PCB degraders and were within the dechlorinating *Chloroflexi* group (Figure 3.6), indicating the presence of microbial dechlorinating bacteria in contaminated sediments.

Bacterial community structure assessed by T-RFLP and pyrosequencing

Bacterial communities in upper sediments were different from those in deeper sediments according to the NMDS ordination of T-RFLP profiles, primarily on NMDS axis 1, suggesting depth-associated shifts in bacterial communities (Figure 3.7). A strong correlation between T-RFLP profile and PCB congener profile was detected by Mantel test ($p=0.028$). The correlation was also confirmed by Procrustes test ($p=0.002$). The correlation suggests that sediment sections with similar PCB congener profiles tend to have similar bacterial community structures.

Several T-RFs that co-varied with the PCB congener patterns along depths were identified (Pearson correlation, $p < 0.001$). T-RFs 235 bp, 250 bp, 258 bp, 273 bp, 278 bp, 302 bp, and 398 bp were found to have strong correlation with sediment MDP values, while T-RFs 252 bp and 515 bp were correlated with pore water MDP values. Those T-RF peaks were not abundant, usually 0.01~0.1% of total T-RFs, which is not surprising, because microbes associated with PCB degradation do not necessarily dominate the bacterial community (Petrić et al., 2011a).

The bacterial communities were also assessed by pyrosequencing. In total, pyrosequencing generated 144,163 16S rRNA sequences from five sediment samples, among which 9,119 unique OTUs were identified (Table 3.2). At least twelve bacterial phyla were detected in all the five sediment samples, among which, *Proteobacteria* was found to be the most abundant taxa in all the samples, accounting for 95~99% of each sample's membership. The *Betaproteobacteria* were the dominant class among the

Proteobacteria in all the samples except for S56, where *Alphaproteobacteria* were most abundant. Overall, each sample had different community composition (Figure 3.8).

Many known aerobic PCB degraders belong to genera within the gamma- and beta- subclasses of *Proteobacteria*, such as *Burkholderia*, *Pseudomonas*, and *Sphingomonas* (Furukawa and Fujihara, 2008). In this study, the *Betaproteobacteria* were the dominant class in all the samples except for S56. Even in S56, *Betaproteobacteria* occupied 36% of total sequences. *Gammaproteobacteria* also took considerable percentage, ranging from 1.38~19.1% of total sequences. This finding was consistent with Correa and colleagues' microcosm results, where the addition of PCB congeners into soil samples resulted in statistically significant increase of *Betaproteobacteria* (Correa et al., 2010).

Genera that include possible aerobic PCB degraders (such as *Acidovorax*, *Acinetobacter*, *Burkholderia*, *Comamonas*, *Polaromonas*, *Pseudomonas*) were detected in all five sediment samples, but they were not abundant. *Chloroflexi* phylum was detected, 94% of which were found in at depth of 1.44-1.60 m of core 1 (C10). However, BLAST results show that the sequences belong to Class *Anaerolineae* rather than *Dehalococcoides*. The *Dehalococcoides* was not detected probably because of the low abundance of *Dehalococcoides* species in the sediments.

Conclusions

The microbial PCB degradation potential was evaluated for a 4.57 m core sediment sample collected from IHSC by exploring the congener profiles and microbial communities. The vertical pattern of the congener profile suggested some extent of *in situ* dechlorination. The successful identification of *bphA* and *Dehalococcoides*-specific 16S rRNA genes in sediments indicated both aerobic and anaerobic PCB degradation

potential. The presence of microorganisms that are known to degrade PCBs may also take part in the microbial PCB metabolism.

Table 3.1. PCR analysis of DNA extracted from different depths of PCB-contaminated core 1 sediment sample at IHSC.

Depth, m	PCR test results	
	bphA 10/30	dhc 946/1212
0-0.16	+	-
0.16-0.32	+	-
0.32-0.48	+	+
0.48-0.64	+	-
0.64-0.80	+	-
0.80-0.96	+	-
0.96-1.12	+	+
1.12-1.28	+	+
1.28-1.44	+	+
1.44-1.60	+	+
1.60-1.76	+	+
1.76-1.92	+	-
1.92-2.08	+	+
2.08-2.24	+	+
2.24-2.40	+	+
2.40-2.56	+	+
2.56-2.72	+	+
2.72-2.88	+	+
2.88-3.04	-	+
3.04-3.20	+	+
3.20-3.36	-	+
3.36-3.52	-	-
3.52-3.68	-	+
3.68-3.84	-	-
3.84-4.00	-	-
4.00-4.16	-	-
4.16-4.32	-	-
4.32-4.48	+	-
4.48-4.57	+	-

Table 3.2. Summary of 454 sequence library sizes, operational taxonomic unites (OTUs).

Sample	454 sequence libraries	
	Sequence library size	Number of OTUs
S11	4192	3413
S21	11584	10019
S56	4109	3756
Core 1 (0.16-0.32m)	3806	3282
Core 1 (1.44-1.60m)	120472	110402



Figure 3.1. Aerial view of Indiana Harbor depicting the location of sediment samples analyzed in this study (Blue open square: core sediment samples; Red open circle: surficial sediment samples).




Figure 3.2. Sediment and pore water MDPR values against sediment depth for core 1.




Figure 3.3. Fraction of light, medium and heavy PCBs versus sediment depth in core sediment (A) and pore water (B), and fraction of *ortho*-, *meta*-, *para*-chlorines versus sediment depth in core sediment (C) and pore water(D).




Figure 3.4. Fraction of each PCB homolog group in sediment (A) and pore water (B) against sediment depth for core 1.




Figure 3.5. Phylogenetic analysis of twelve amino acids of translated sequences of BphA genes retrieved from PCB-contaminated surficial sediment sample 11, 21, 56 and core 1 sediment sample (0.16-0.32 m) at IHSC. The alignment was made with ClustalX and converted into a neighboring-joining tree, which was visualized with MEGA4 with *Bacillus subtilis* Putative ring-cleaving dioxygenase as an outgroup. The filled circles at nodes indicate bootstrap values higher than 95%, while the open circles indicate bootstrap values of 75-95%. Bootstrap values lower than 75% are not shown. The bar represents 1% sequence difference.




Figure 3.6. Phylogenetic analysis of 16S rRNA genes retrieved from surficial and core sediment samples at IHSC. The alignment was made with ClustalX and converted into a neighboring-joining tree, which was visualized with MEGA4 with *Chloroflexus aurantiacus* J-10-fl as an outgroup. The filled diamonds represent known PCB degraders. The filled circles at nodes indicate bootstrap values of 95%, while open circles indicate bootstrap values of 75~95%. Bootstrap values lower than 75% are not shown. The bar represents 0.5% sequence difference.




Figure 3.7. NMDS ordination of Wisconsin transformed T-RFLP profiles of core 1 (C01: 0-0.16 m, C02: 0.16-0.32 m...).




Figure 3.8. Relative abundance of phyla and proteobacterial classes for surficial sediment samples (S11, S21 and S56) and core 1 sediment samples (depth 0.16-0.32 m and 1.44-1.60 m).

CHAPTER IV. MICROBIAL PCB DEGRADATION POTENTIAL IN CORE 2 FROM IHSC

Introduction

Polychlorinated biphenyls (PCBs) are a group of 209 synthetic chemicals that were widely used in industry during the last century. The production and commercial use of PCBs were banned by the EPA in 1979 because they were found to be toxic, carcinogenic, and bioaccumulative in the food web (Ross, 2004). Nevertheless, due to their high stability, PCBs still exist in the environment, and pose a potential risk to public health more than 30 years later (Bedard, 2004). Despite their environmental persistence, PCBs can be transformed to less toxic forms or even mineralized by microorganisms. Field and laboratory studies indicate two predominant processes: anaerobic dechlorination and aerobic biphenyl ring cleavage (Borja *et al.*, 2005).

Under anaerobic conditions, PCBs are used as electron acceptors, resulting in the removal of chlorine atoms. The less chlorinated congeners are more amenable to aerobic degradation (Alder *et al.*, 1993). Anaerobic PCB-degraders include *Dehalococcoides* spp., the o-17/DF-1 group, as well as some other members of the *Chloroflexi* (Adrian *et al.*, 2009; Wiegel and Wu, 2000; Wu *et al.*, 2002). Reductive dehalogenase (RDase) is considered to be the key enzyme catalyzing the PCB dechlorination process (Hiraishi, 2008; Pieper and Seeger, 2008). Nonidentical RDase genes were found in PCB degrading strains such as *Dehalococcoides mccartyi* strain 195 and CBDB1, but no RDase gene was identified to dechlorinate PCBs (Fung *et al.*, 2007; Hölscher *et al.*, 2004; Wagner *et al.*, 2009).

Under aerobic conditions, PCBs can be used as electron donors, or fortuitously oxidized by oxygenase enzymes. A variety of PCB-oxidizing bacteria have been identified, including Gram-negative strains of *Pseudomonas*, *Burkholderia*, *Acinetobacter*, and *Comamonas* and Gram-positive strains of *Corynebacterium*,

Rhodococcus and *Bacillus* (Furukawa and Fujihara, 2008; Pieper and Seeger, 2008). These microbes harbor biphenyl dioxygenase (Bph), the key enzyme catalyzing the first step of aerobic biphenyl ring cleavage, generating a *cis*-dihydrodiol intermediate, which is further degraded to a chlorobenzoate (Gibson and Parales, 2000). PCB-oxidizing microbes usually cannot metabolize the chlorobenzoate, thus a consortium of chlorobenzoate-degraders are required for complete PCB mineralization (Pavlů *et al.*, 1999).

The site of this study is the Indiana Harbor and Ship Canal (IHSC), a heavily industrial area of southern Lake Michigan contaminated with a variety of pollutants, including heavy metals, polycyclic aromatic hydrocarbons (PAHs) and PCBs. It is reportedly a major source of PCBs to Lake Michigan, with PCB concentrations in IHSC surficial sediments ranging from 53 to 35,000 ng g⁻¹ dry weight (Martinez *et al.*, 2010). PCB-contaminated IHSC sediments are currently being dredged and permanently stored in a confined disposal facility (CDF) (USACE, 2013).

The purpose of this study is to evaluate both anaerobic and aerobic PCB biodegradation potential in IHSC sediments. We characterized PCB congener profiles and bacterial communities in a 4.57-m long core sediment sample, which revealed evidence of *in situ* aerobic and anaerobic PCB degradation in upper 1.83 m sediments. We also explored the correlation between the bacterial communities and PCB congener profiles and found that sediment sections with similar PCB congener profiles tend to have similar bacterial community structure. We conclude that microbial communities in IHSC sediments have the potential for aerobic and anaerobic PCB biodegradation. This suggests that natural attenuation of PCBs could continue in IHSC sediments after they are transferred to the CDF.

Materials and Methods

Site description and sampling

In May 2009, a 4.57-m core sediment sample from IHSC was collected using a submersible vibro-coring system with a PVC tube (length 457 cm, internal diameter 9.5 cm) from aboard the U.S. EPA's research vessel, *Mudpuppy* (Figure 3.1). The core was sectioned every 0.305 m, and each section was homogenized, placed in plastic bags and kept on ice during transportation. Sediment samples were stored at 4°C in the lab until analysis.

PCB congener analysis³

Preparation, extraction and clean-up steps for measuring PCB sediment concentrations were performed as previously described (Martinez and Hornbuckle, 2011). Briefly, sediments were extracted using pressurized fluid extraction (Accelerated Solvent Extractor, Dionex ASE-300). The extracts were concentrated and eluted through a multilayer silica gel column. Sulfur in solution was removed by activated granulate copper. Poly-dimethylsiloxane (PDMS) coated fibers were used as passive sampler to measure PCB concentration of the sediment pore water. PCB extraction and quantification procedures were also reported previously (Martinez *et al.*, 2013). PCB identification and quantification were conducted employing a modified US EPA method 1668C (USEPA, 2010). Tandem mass spectrometry GC/MS/MS (Quattro Micro GC, Micromass MS Technologies) in multiple reaction monitoring mode was utilized to quantify all 209 congeners in 161 individual or coeluting congener peaks.

The molar dechlorination product ratio (MDPR) was used to examine possible PCB dechlorination in core sediments. When determining the MDPR, it is assumed that

³ The PCB concentration measurement of sediment samples was performed by Andres Martinez, Ph.D., Department of Civil & Environmental Engineering, The University of Iowa.

exclusively *ortho*-chlorinated PCB congeners undergo no further dechlorination (TAMS Consultants, 1997). In this study, five exclusively *ortho*-chlorinated PCBs (PCB 1, 4, 10, 19, 54) and PCB 8 were selected as the ultimate dechlorination products. The ratio of the sum of the molar concentrations of selected congener over total PCB molar concentration was calculated and defined as MDP. PCB 8 was considered as a dechlorination product because the proportions of PCB 8 in core sediments (averaged 2.7% of total PCB) were much higher than that in other Aroclor commercial mixtures (0.48%, 0%, 0% of Aroclor 1248, 1242, 1016, respectively), which indicates the production of this congener in IHSC sediments.

Quantitative PCR

Total DNA from sediments was isolated with the MoBio Ultraclean Soil DNA Isolation Kit (MoBio Laboratories, Inc., Carlsbad, CA) and stored at -20°C until analysis. The abundance of total bacteria, aerobic PCB-degrading bacteria, and putative PCB-dechlorinating bacteria were estimated using qPCR targeting bacterial 16S rRNA gene (primer set 16SU f/r) (Nadkarni *et al.*, 2002), *bphA* (primer set *bphA* 463f/674r) (Petrić *et al.*, 2011b), putative dechlorinating *Chloroflexi* 16S rRNA genes (primer set chl348f/dehal884r) (Fagervold *et al.*, 2005), and *Dehalococcoides*-like 16S rRNA genes (primer set dhc793f/946r) (Yoshida *et al.*, 2005) (Table 4.1). PCR conditions were as follows: 10 min at 95°C , 40 cycles of 15s at 95°C and 1 min at 60°C followed by a dissociation step. Each 25 μl reaction contained 12.5 μl Power SYBR Green PCR Master Mix (Invitrogen, Carlsbad, CA), 2.5 pmol primers and variable amounts of primers and template (Table 4.2). Bovine serum albumin (0.5 μg) was added to relieve possible PCR inhibition (Kreader, 1996).

For total bacterial 16S rRNA gene qPCR, the standard DNA template was PCR products amplified from *Burkholderia xenovorans* LB400 with primer set 8F/1492R (Grabowski *et al.*, 2005). For *bphA*, the standard DNA template was the LB400 *bphA*

(amplified with the 463f/674r primer set) cloned into the pCR 2.1-TOPO vector. For putative dechlorinating *Chloroflexi* 16S rRNA genes and *Dehalococcoides*-like 16S rRNA gene, standard curves were prepared from pCR 2.1-TOPO vector containing the target PCR products with primer chl3487f/dehal884r and dhc793f/946r, respectively. All qPCRs were performed with an ABI 7000 Sequence Detection System (Applied Biosystem, Grand Island, NY) and fluorescence data was analyzed by ABI 7000 System SDS Software (Applied Biosystems, Grand Island, NY). With each primer set, the target gene was not detected in no template (DI water) controls (Ct value > 35). Additional qPCR information, such as primer concentrations, template concentrations, qPCR linear range, qPCR efficiency range of the standard curves, and Y-intercepts are provided in Table 4.2, in accordance with MIQE guidelines (Bustin *et al.*, 2009).

qPCR quality assurance

To verify the specificity of qPCR primer sets chl348f/dehal884 and dhc793f/946r, two clone libraries were constructed from the amplification products in DNA extracted from sediments at 0-0.30m and 0.91- 1.22 m. Clone libraries were prepared as described in section 2.6. From the chl348f/dehal884r PCR product clone library, 21 unique sequences were obtained from 22 clones and from the dhc793f/946r PCR product clone library, 10 unique sequences were obtained from 10 clones. All of the sequences were determined to be from *Chloroflexi*, using RDP classifier (Cole *et al.*, 2007), with 91% classified as *Dehalogenimonas* (Table 4.3). The specificity of the SYBR green based qPCR is also validated by dissociation curve analysis, which shows similar melting temperature for each primer set.

Terminal-Restriction Fragment Length

Polymorphism (T-RFLP) analysis

PCR with fluorescently-labeled 6-FAM 8Fm and 533R was performed with 1 ng

DNA (Schütte *et al.*, 2008). PCR conditions were: 4 min at 94°C, 30 cycles of 1 min at 94°C, 2 min at 55°C and 1 min at 72°C followed by a final extension step of 10 min at 72°C. PCR product (12µl) was digested with the restriction enzyme *Hae*III (New England BioLabs, Inc., Ipswich, MA), and then precipitated with glycogen (Fermentas International Inc., Vilnius, Lithuania), sodium acetate (pH 5.2), and ethanol at -20°C for 2h. DNA was recovered by centrifugation at $17,800 \times g$ for 15 min and resuspended in distilled water (Invitrogen Corp., Carlsbad, CA). The digested DNA was sent to the University of Iowa DNA facility for electrophoresis for sizing using an Applied Biosystems 3730 DNA analyzer (Life Technologies Corporation, Carlsbad, CA) with the GeneScan 500 LIZ size standard.

Terminal restriction fragment (T-RF) sizes were estimated with Peak Scanner software (Applied Biosystems, Carlsbad, CA). The TRF size matrix containing 15 samples (rows) and 354 unique 16S rRNA gene TRFs (columns) was generated by T-REX software after filtering background peak noise and rounding fragment sizes to the nearest whole number (Culman *et al.*, 2009). TRF profiles were analyzed by non-metric multidimensional scaling (NMDS) in R (da C Jesus *et al.*, 2009). The Bray-Curtis dissimilarity index was calculated with a random starting configuration, and a two-dimensional solution was reached after seven runs. The final stress was 7.42.

Correlations between PCB congener profiles and T-RFLP profiles were assessed with the Procrustes test, where PCB congener profiles were analyzed by principal components analysis (PCA) and T-RFLP profiles were ordinated by NMDS. The Mantel test was applied to assess the correlation between T-RFLP profiles and PCB profiles, where a Euclidean dissimilarity index was calculated for PCB congener profiles and the Bray-Curtis dissimilarity index was calculated for T-RFLP profiles. Pearson correlation coefficients between each TRF and molar dechlorination product ratio (MDPR) values were also calculated (González *et al.*, 2000).

PCR amplification, cloning and sequencing of bacterial 16S rRNA

A clone library was constructed with bacterial 16S rRNA genes amplified from sediment at the depth 3.35 – 3.66m. Relatively high MDP values (0.08 for sediment and 0.37 for pore water) were observed at this depth. Bacterial 16S rRNA genes were amplified using primer set 8F/1492R (Grabowski *et al.*, 2005). PCR conditions were as described in the T-RFLP analysis section. PCR products were purified with the Qiaquick PCR purification Kit (Qiagen Inc., Valencia, CA), cloned into the pCR 2.1-TOPO vector using the TOPO TA cloning Kit (Invitrogen Corp., Carlsbad, CA), and transformed into One Shot TOP10 chemically competent *E. coli* cells (Invitrogen Corp., Carlsbad, CA). The transformation efficiency was checked by plating recombinant *E. coli* on Luria Broth agar with kanamycin (50 mg/L), and X-gal (0.4 mg/plate) and incubated overnight at 37°C. Clones were Sanger sequenced at HTSeq.org (Seattle, WA) with M13F (5'-TGTAACGACGGCCAGT-3') and M13R (5'-CAGGAAACAGCTATGAC-3') primers. Sequence identification and classification were performed using RDP Classifier (Cole *et al.*, 2007) and Basic Local Alignment Search Tool (BLAST) (Altschul *et al.*, 1997). Predicted TRFs were generated with TRiFLE (Junier *et al.*, 2008).

Results and discussion

Evidence for PCB dechlorination in IHSC sediment congener profiles

Changes in PCB congener distribution patterns with sediment depth can be explained by changes in production, use and discharge of different commercial mixtures, sediment resuspension and transport, and *in situ* microbial PCB degradation (Li *et al.*, 2009). Sediment resuspension and PCB desorption would be expected to result in accumulation of highly chlorinated PCBs in deeper sediments. However, anaerobic PCB

dechlorination in sediments would lead to decreased abundance of higher chlorinated congeners and increased abundance of less chlorinated ones. With this in mind, we examined PCB congener profiles in sediment samples to search for evidence of PCB dechlorination. Both sediment and pore water PCB profiles showed the accumulation of lesser chlorinated PCBs (1-3 chlorine substituents) in the deeper sediment of the core, along with a decrease of medium chlorinated congener abundance (4-6 chlorine substituents) (Figure 4.1A, 4.1B). There was also an increasing trend of *tri*-chlorinated PCBs and a corresponding decreasing fraction of *tetra*- and *penta*-chlorinated congeners with increasing depth (Figure 4.2A, 4.2B).

Anaerobic PCB dechlorination in sediments preferentially removes chlorines at *para*- and *meta*- positions, resulting in the enrichment of *ortho*-chlorinated congeners (Alder et al., 1993; Bedard et al., 1997; Fava et al., 2003). Therefore, if PCB dechlorination had occurred, we would expect to see an increase in the abundance of PCB congeners with *ortho*-chlorine substituents. Indeed, analysis of the relative abundance of *ortho*-, *meta*-, and *para*-chlorines against core depth revealed an accumulation of *ortho*-chlorines in deep sediments, in accordance with a decrease in *meta*- and *para*-chlorinated PCBs (Figure 4.1C). In pore water, the enrichment of *ortho*-chlorinated congeners was also observed with depth (Figure 4.1D).

MDPR values were estimated to examine potential PCB dechlorination in the core sediment. Accumulation of the six selected *ortho*-chlorinated congeners in the core sediments was observed, which further supports that PCB dechlorination has occurred (Figure 4.3). Although sediment PCB profiles in the core reportedly mainly resemble the Aroclor 1248 mixture, the average sediment MDPR value (0.052) was much greater than that of Aroclor 1248 (0.009) (Frame et al., 1996; Martinez and Hornbuckle, 2011), which is also a sign of *in situ* dechlorination. MDPR values of pore water were higher than those of corresponding sediment sections (Figure 4.3). This can be explained by the tendency for *ortho*-chlorine substituents to increase the aqueous solubility of the PCB

molecule (Huang and Hong, 2002), and that the lower chlorinated congeners as the six selected ones are more soluble (Van Noort *et al.*, 2010). Another possible reason is that PCBs dissolved in pore water are more bioavailable than those sorbed to sediment particles and thus are more vulnerable to *in situ* dechlorination (Reid *et al.*, 2000).

Aerobic and anaerobic PCB biodegradation potential in sediment

With the PCB congener-based evidence for reductive dechlorination in IHSC sediments, we aimed next to characterize the PCB biodegradation potential by estimating the abundance of total 16S rRNA genes, putative *Chloroflexi* 16S rRNA genes, and *bphA*, a functional gene associated with aerobic PCB oxidation. The total bacterial 16S gene abundance was relative stable with sediment depth ($2.37 \times 10^9 \pm 2.41 \times 10^9$ 16S rRNA genes per g sediment) (Figure 4.4). The lowest 16S rRNA gene abundance was from 0-0.30 m sediment, possibly because this section was at the water-sediment surface and had relatively high water content compared with the other sediment samples.

The abundance of *bphA* averaged $7.39 \times 10^6 \pm 3.96 \times 10^6$ genes per g sediment (0-1.83m depth), and $3.20 \times 10^5 \pm 4.58 \times 10^5$ gene copy per g sediment (1.83 - 4.57m depth). Especially, *bphA* gene abundances were significantly correlated with total sediment PCB concentrations in the upper 1.83 m, which could indicate a microbial response to PCB contamination in the sediments (Pearson's R, $p < 0.05$).

Although the upper 1.83m was anaerobic as suggested by negative ORP data (-13- -200mv) (Martinez *et al.*, 2013), micro-aerobic conditions could possibly exist in some portions of the sediment. The sharp decrease in *bphA* abundance below 1.83m could be explained by the presence of a stricter anaerobic environment. A strong positive correlation was observed between the *bphA* abundance and the ORP along depth (Pearson's R, $p < 0.01$).

Two primer sets (chl384f/884r and dhc793f/946r) were used to quantify the

dechlorinating *Chloroflexi* group in sediments. According to the clone library results, both of the two primer sets primarily amplified *Dehalogenimonas* sequences. (Table S3). Some isolates of *Dehalogenimonas* strains can reductively dehalogenate polychlorinated alkanes (Maness et al., 2012; Yan et al., 2009). Also, recent study found that *Dehalogenimonas* can couple their growth with PCB dechlorination in a sediment free culture, indicating their role in PCB dechlorination (Wang and He, 2013).

Putative dechlorinating *Chloroflexi* 16S rRNA gene abundances were $2.53 \times 10^5 \pm 2.45 \times 10^5$ genes per g sediment (0-1.83m depth) and $1.53 \times 10^4 \pm 2.80 \times 10^4$ genes per g sediment (1.83-4.57 m depth) as assessed by chl384f/884r, while the gene abundance averaged $2.15 \times 10^5 \pm 2.07 \times 10^5$ gene copy per g sediment (0-1.83m depth) and $7.64 \times 10^3 \pm 1.30 \times 10^4$ gene copy per g sediment (1.83-4.57 m depth) as assessed by dhc793f/946r. These gene abundances are lower than observed in sediments from other PCB-contaminated sites, which were on the order of 10^6 - 10^8 genes per g sediment, possibly because these other sites contained higher PCB concentrations (Kjellerup et al., 2008; Xu et al., 2012).

The co-occurrence of putative dechlorinating *Chloroflexi* 16S rRNA genes and *bphA* indicate the potential for simultaneous aerobic and anaerobic PCB degradation in the upper 1.83 m of the sediment. However, the pattern of putative dechlorinating *Chloroflexi* population does not appear to be consistent with the congener profile, which displayed greater dechlorination potential in sediments deeper than 1.83 m. A possible explanation is that some other *Chloroflexi* members were degrading PCBs in deep sediments and were not detected by the primers used in this study. It is also possible that the congener profiles in the deep sediments were the record of dechlorination from the past. In addition, the 16S rRNA gene based phylogenetic analysis of *Chloroflexi* doesn't directly identify the dechlorinating capability of the microbial community. Reductive dehalogenase gene would be a better indicator of microbial dechlorination, but due to the lack of knowledge about the PCB dehalogenase, no primer targeting PCB dehalogenase

is currently available.

Relationships between *bphA*, putative dechlorinating *Chloroflexi* 16S rRNA genes, and total bacterial 16S rRNA gene abundance with environmental variables were explored by calculating Pearson correlations. No positive correlations were observed between any of these gene abundances and the total PCB concentration, or with total organic carbon.

Relationships between microbial community structure and PCB congener profiles

To explore the possibility that other microbial community members could be relevant to PCB biodegradation potential in IHSC sediments, we evaluated microbial community structure with depth using T-RFLP analysis. A significant correlation between the overall sediment T-RFLP profile and PCB congener profile was noted when using the Mantel test ($p=0.001$ for sediment, $p=0.001$ for pore water). This correlation was confirmed with the Procrustes test ($p=0.002$ for sediment, $p=0.001$ for pore water). This suggests that sediment sections with similar PCB congener profiles tend to have similar bacterial community structures.

NMDS ordination of the T-RFLP profiles revealed that bacterial community structures in the shallow sediment sections (0-1.83 m depth) were distinct from the structure in deeper sediments (1.83-4.57 m depth) (Figure 4.5A). The disparity between shallow and deep sediments was also found by qPCR, which show much less abundant *bphA*, putative *Chloroflexi* 16S rRNA genes, and *Dehalococcoides*-like 16S rRNA genes in sediments of 1.83-4.57 m deep.

The microbial community composition in sediments at 3.35-3.66 m depth was assessed by sequencing 91 clones from a 16S rRNA gene clone library. This indicated that the community was dominated by *Proteobacteria* (11.0% *Alphaproteobacteria*, 34.1% *Betaproteobacteria* and 54.9% *Gammaproteobacteria*). At the genus level, *Acinetobacter*

spp. and *Acidovorax* spp. were dominant, comprising 45.1% and 20.9% of the clones, respectively.

A 196bp T-RF, which was abundant in sediments deeper than 1.22 m (26.9% \pm 13.0%) (Figure 4.5B), was significantly correlated with sediment and pore water MDPH values (Pearson's R, $p < 0.001$). This T-RF was identified to be from an *Acidovorax* sp. when compared with virtually digested T-RFs from a 16S rRNA gene clone library (91 clones) from sediments of 3.35-3.66 m. The corresponding T-RFs of *Acinetobacter* were 198bp, 199bp and 252bp (Table 4.4). A 199 bp T-RF comprised 20.0% of all T-RFs in 3.35-3.66 m section, but was only 0-2.5% in other sections. Clones identified as *Methylotenera* and unclassified *Comamonadaceae* also generated a 199 bp T-RF, but the majority of the 199 bp T-RFs in the clone library were classified as *Acinetobacter*. Two clones from the clone library generated a 198 bp T-RF. This T-RF was not found in T-RFLP profiles of 3.35-3.66 m sediments, but was found in 0-1.52m sediments (1.97-9.51% relative abundance) and had a relative abundance of 18.74% in sediments at 4.27-4.57 m depth. A 252 bp T-RF was also identified as coming from an *Acinetobacter* sp., and had a relative abundance range of 0.62-8.10% along the core. Overall, this analysis revealed that although *Acinetobacter* was an abundant (and thus potentially important) community member in the core sediments. However, there was no correlation between TRFs identified as *Acinetobacter* and sediment or pore water MDPH values.

Proteobacteria are often found abundant in soils and sediments (de Cárcer et al., 2007; Petrić et al., 2011a; Spain et al., 2009). However, an increase in *Betaproteobacteria* and *Alphaproteobacteria* abundance was observed in PCB-exposed samples in a microcosm study (Correa et al., 2010). Many known PCB-degraders are *Proteobacteria* (e.g. *Pseudomonas*, *Acidovorax*, *Acinetobacter*, *Comamonas* and *Burkholderia* (Bedard et al., 1986; Furukawa and Fujihara, 2008)). Both *Acidovorax* and *Acinetobacter* are known to degrade PCBs and were commonly found in PCB-contaminated sites (de Cárcer et al., 2007; Ionescu et al., 2009; Shuai et al., 2010; Slater

et al., 2011). Aerobic PCB biodegradation was likely not active in sediments deeper than 1.83 m, as indicated by negative ORP data and the low abundance of *bphA*. So it is unlikely that *Acinetobacter*, *Acidovorax* or other possible PCB degraders were carrying out aerobic PCB degradation in deep sediments.

Some *Acidovorax* species are anaerobic and are involved in nitrate reduction, but are not known to be associated with dechlorination (Byrne-Bailey et al., 2010; Hohmann et al., 2010). *Acinetobacter* has been detected in sediments deeper than 100 m (Breuker et al., 2011), and studies have reported that some *Acinetobacter* species are capable of dechlorinating and degrading chlorobenzoate, chlorophenols and chloroanilines (Adriaens and Focht, 1991; Copley and Crooks, 1992; Hongsawat and Vangnai, 2011; Kim and Hao, 1999). *Acinetobacter* has been found in PCB-dechlorinating JN cultures (Bedard et al., 2007), although no evidence yet suggests that *Acinetobacter* can dechlorinate PCBs.

Chloroflexi was not detected in the 16S rRNA gene clone library, probably because of its relatively low abundance in the sediment sample with respect to the total 16S rRNA gene abundance. This was verified by our qPCR results. Yet PCB congener profile analysis suggests extensive PCB dechlorination in deep sediments, indicating that there might be unknown dechlorinating species other than *Chloroflexi* were degrading PCBs.

Conclusions

In general, both aerobic and anaerobic PCB degradation potential at IHSC were assessed by the combination of the PCB congener analysis and the microbial analysis. PCB congener profiles of the core sediment revealed that *in situ* dechlorination has occurred. The presence of microbial dechlorinating groups and *bphA* in the upper 2-m sediments suggested that the indigenous microbial communities contained the potential

for both aerobic and anaerobic PCB degradation. In deeper sediments, the microbial communities were different from those in upper sediments as suggested by T-RFLP profiles. *Acinetobacter* was highly abundant in deep sediments, suggesting their possible role in PCB dechlorination. Overall our results provide a comprehensive understanding about the *in situ* PCB degradation potential at the IHSC sediment, and suggest that natural attenuation of PCBs could continue in IHSC sediments stored at the CDF after being dredged.

Table 4.1. Primer sets used for qPCR analysis in Chapter IV.

Target gene	Primer name	Sequences	Expected product size	Source
total bacterial 16S rRNA	16SU f	5'-TCCTACGGGAGGCAGCAGT-3'	466 bp	(Nadkarni <i>et al.</i> , 2002)
	16SU r	5'-GGACTACCAGGGTATCTAATCCTGT T-3'		
<i>bphA</i>	<i>bphA</i> 463f	5'-CGCGTSGMVACCTACAARG-3'	211 bp	(Petrić <i>et al.</i> , 2011b)
	<i>bphA</i> 674r	5'-GGTACATGTCRCTGCAGAAYTGC-3'		
putative dechlorinating <i>Chloroflexi</i> 16S rRNA	chl348f	5'-GAGGCAGCAGCAAGGAA-3'	514 bp	(Fagervold <i>et al.</i> , 2005)
	dehal884r	5'-GGCGGGACACTTAAAGCG-3'		
<i>Dehalococcoides</i> -like 16S rRNA	dhc 793f	5'-GGGAGTATCGACCCTCTCTG-3'	193 bp	(Yoshida <i>et al.</i> , 2005)
	dhc 946r	5'-CGTTYCCCTTTCRGTTCACT-3'		

Table 4.2. Pertinent qPCR parameters (primer concentration, template mass, linear range, PCR efficiency, and y-intercept of the qPCR standard curve) with core 2 sediment DNA as template.

Target gene	primer concentration, μM	template	qPCR linear range, gene copies/reaction	qPCR efficiency	Y-intercept
total bacterial 16S rRNA	0.1	1 ng DNA	30 - 30×10^7	90 - 100%	37.07
<i>bphA</i>	1	10 ng DNA	40 - 40×10^7	95 - 100%	37.55
putative dechlorinating <i>Chloroflexi</i> 16S rRNA	0.4	10 ng DNA	40 - 40×10^5	90 - 100%	38.02
<i>Dehalococcoides</i> -like 16S rRNA	0.8	10 ng DNA	30 - 30×10^7	95 - 100%	36.55

Table 4.3. RDP classification of clone libraries for qPCR primer quality assurance with core 2 sediment DNA as template.

Primer set	Closest classified relative (% certainty)	No. of clones detected
chl348f/dehal884r	<i>Dehalogenimonas</i> (80-100)	21/22
	Unclassified <i>Chloroflexi</i> (80)	1/22
dhc793f/946r	<i>Dehalogenimonas</i> (83-100)	8/10
	Unclassified <i>Chloroflexi</i> (83-85)	2/10

Table 4.4. 16S rRNA gene clones recovered from sediment at the depth of 3.35-3.66 m.
Clone identification was made by RDP classifier.

Observed T-RF (bp)	Predicted T-RF (bp)	Closest classified relative (% certainty)	No. of clones	NMDS axis 1	NMDS axis 2
196	198	<i>Acidovorax</i> (93-100)	19	0.717	-0.052
196	198	Unclassified <i>Comamonadaceae</i> (99)	1	0.717	-0.052
197	198	<i>Methylotenera</i> (100)	1	-0.241	1.070
198	199	<i>Acinetobacter</i> (100)	2	-0.430	0.276
199	200	<i>Acinetobacter</i> (100)	27	0.316	-0.393
199	200	Unclassified <i>Comamonadaceae</i> (90)	1	0.316	-0.393
199	200	<i>Methylotenera</i> (100)	2	0.316	-0.393
217	219	<i>Janthinobacterium</i> (97)	1	-0.138	0.358
223	225	<i>Rhizobium</i> (95-97)	2	-0.176	-0.444
226	227	<i>Novosphingobium</i> (100)	1	-0.644	-0.056
252	253	<i>Acinetobacter</i> (99-100)	10	0.276	-0.151
316	317	<i>Comamonas</i> (99-100)	5	0.625	-0.301




Figure 4.1. Fraction of light, medium and heavy PCBs versus sediment depth in core 2 (A) and pore water (B), and fraction of *ortho*-, *meta*-, *para*-chlorines versus sediment depth in core sediment (C) and pore water(D).




Figure 4.2. Fraction of each PCB homolog group in sediment (A) and pore water (B) against sediment depth for core 2.




Figure 4.3. Sediment and pore water MDPR values against sediment depth for core 2.




Figure 4.4. Abundances of total bacteria (*bac*), *bphA*, putative *Chloroflexi* 16S rRNA genes (*chl*) and *Dehalococcoides*-like 16S rRNA (*dhc*) against sediment depth by gene copy No. per g sediment for core 2.




Figure 4.5. NMDS ordination and cluster analysis of T-RFLP profiles of core 2 sediment (C01: 0-0.30 m, C02: 0.30-0.61 m...) (A), the abundance of T-RF 196 bp (B) and 198 bp+199 bp + 252 bp (C) along sediment depth.

CHAPTER V. ENHANCED POLYCHLORINATED BIPHENYL
REMOVAL IN A SWITCHGRASS PLANTED RHIZOSPHERE BY
BIOAUGMENTATION WITH *BURKHOLDERIA XENOVORANS*

LB400

Introduction

Conventional remediation strategies such as soil excavation and the following incineration or landfilling are usually expensive and unsustainable (Amend and Lederman, 1992). Phytoremediation, the use of plants and associated microbes to remove and detoxify PCBs, represents a potentially cost effective and less disruptive approach to clean up PCB contaminated soils and sediments (Van Aken *et al.*, 2010).

The presence of specific plant species is known to enhance the PCB removal efficiency in soils and sediments (Campanella *et al.*, 2002; Chekol *et al.*, 2004; Ionescu *et al.*, 2009; Slater *et al.*, 2011; Smith *et al.*, 2007). Plants can uptake PCBs from soil and convert lower chlorinated congeners into hydroxylated products (Ficko *et al.*, 2010; Greenwood *et al.*, 2011; Kučerová *et al.*, 2000; Lee and Fletcher, 1992; Rezek *et al.*, 2007; Zhai *et al.*, 2010b). Also, plants contribute to PCB biodegradation by providing a favorable environment for PCB-degrading microorganisms and can thus increase PCB-degrading populations (de Cárcer *et al.*, 2007; Leigh *et al.*, 2006; Slater *et al.*, 2011; Uhlík *et al.*, 2009). Plants not only increase soil permeability and oxygen diffusion in the rhizosphere, but also release organic exudates as inducers, surfactants and microbial growth factors thereby stimulating and facilitating microbial PCB biodegradation in the root zone (Gilbert and Crowley, 1997; Hernandez *et al.*, 1997; Leigh *et al.*, 2002; Van Aken *et al.*, 2010). For example, augmentation of soil with plant-derived compounds such as carvone and salicylic acid assisted aerobic PCB degraders and increased PCB removal (Singer *et al.*, 2000; Singer *et al.*, 2003b).

Microorganisms are known to degrade PCBs via two general processes: aerobic oxidative biphenyl ring cleavage and anaerobic reductive dechlorination (Adrian et al., 2009; Borja et al., 2005; Gibson and Parales, 2000; Pieper, 2005; Wiegel and Wu, 2000). Most aerobic PCB-degraders employ the upper biphenyl degradation pathway, and biphenyl 2,3-dioxygenase (BphA) catalyzes the first step in this pathway, transforming susceptible PCB congeners into the corresponding *cis*-dihydrodiol (Furukawa, 1994). Under anaerobic conditions, some members of the phylum *Chloroflexi* use certain PCB congeners as electron acceptors and transform them into less chlorinated congeners (Adrian et al., 2009; Fagervold et al., 2007; Fagervold et al., 2005; Wiegel and Wu, 2000). Some efficient microbial PCB degraders, such as *Burkholderia xenovorans* strain LB400, *Arthrobacter* sp. strain B1B, *Ralstonia eutropha* H850, *Dehalococcoides chlorocoercia* DF1 have been isolated and introduced to contaminated soils and sediments to improve PCB removal (Payne et al., 2013; Payne et al., 2011; Petrić et al., 2011b; Singer et al., 2000).

In this study, we monitor the removal of three selected PCB congeners (PCB 52, PCB 77 and PCB 153) in switchgrass-planted soil microcosms and attempt to improve PCB degradation by introducing biphenyl-grown *Burkholderia xenovorans* strain LB400 into the rhizosphere of selected microcosms. The PCB congeners in this study were chosen because they represent bottlenecks in environmental PCB degradation (Bedard *et al.*, 1986). PCB 52 and 153 are both *ortho*-chlorinated, while PCB 77 has a dioxin-like structure, and is considered one of the most toxic congeners (Van den Berg *et al.*, 2006). All three congeners are quite ubiquitous in the environment (Meggo et al., 2013). Enhanced degradation of these congeners was observed in switchgrass treated soil, and following bioaugmentation with LB400 PCB 52 was significantly improved. The combined use of switchgrass and LB400 bioaugmentation was found to be most efficient for specific PCB congener removal in this experiment.

Materials and Methods

Soil reactor set up⁴

The soil microcosm set-up procedure has been described previously (Meggo and Schnoor, 2013). Briefly, PCB-free soil from the village of Middle Amana in Iowa, USA, was passed through a 60 mesh sieve before experiments. The soil was spiked with a mixture of PCB 52, PCB 77 and PCB 153 (99% pure) (Accustandard Inc., New Haven, CT) at the concentration of 500 ng g⁻¹ each. PCB congeners were dissolved in hexane before adding to the soil. A quartering technique was used to homogenize soil. The soil was divided into four quadrants on a quartering canvas, and 20 diagonal trajectories were used to mix the soil components. The contaminated soil was aged for two months at 25°C in sealed tubs to allow PCBs sequestration into the soil matrix and thereby better represent field conditions. To establish the initial PCB concentration, 20 subsets of soil collected from different locations were homogenized and measurements were performed in triplicates (Meggo and Schnoor, 2013).

Plastic containers (33.8cm × 21.6cm × 21.9cm) and lids with aluminum foil covers were used for soil reactor set-up. Each reactor was filled with 2500g sieved and homogenized soil. The spiked soil was planted with switchgrass (*Panicum virgatum*) seeds (Adams-Briscoe Seed Co., Jackson, GA). Switchgrass in all six reactors was grown at 25°C in a plant growth chamber under a 16 hour light/8 hour dark photo period (light intensity of 200 mmol m⁻²s⁻¹ and 60% humidity). Soil samples were taken for PCB congener analysis and microbial analysis. Soil samples were taken at 12 week and 24 week for PCB congener analysis. For microbial analysis, soil samples were taken every four weeks starting at 12 week (Meggo and Schnoor, 2013).

⁴ Soil reactor set-up was performed by Richard Meggo, Ph.D., Department of Civil & Environmental Engineering, The University of Iowa.

Burkholderia xenovorans strain

LB400 bioaugmentation

Bioaugmentation with *Burkholderia xenovorans* strain LB400 was conducted with both switchgrass-treated and unplanted microcosms every each month. Prior to bioaugmentation LB400 was grown on solid biphenyl (1 g) as sole carbon and energy source in K1 medium until exponential phase (OD_{600} 0.6~0.8) (Zaitsev and Karasevich, 1985). Cells were harvested by centrifugation ($5000 \times g$, 15 min), washed once with sterile K1 medium, resuspended in K1 medium and inoculated into reactors (approximately 10^9 CFU per g soil). Autoclaved LB400 was added to a switchgrass planted reactor as control.

Abundance of cultivable aerobic biphenyl-degrading bacteria was estimated by CFU enumeration one day after each bioaugmentation. Soil samples (1 g) were mixed with 9 ml of 0.9% sodium chloride solution and shaken for 1 h at 225 rpm on a platform shaker (New Brunswick Scientific, Pittsburgh, PA) at 25°C. Serially diluted supernatant was spread on K1 agar plates with biphenyl crystals as sole carbon and energy source. CFUs were enumerated after 72h of incubation at 30°C.

PCB extraction and quantification⁵

The PCB extraction procedure has been previously described (Meggo *et al.*, 2013). Briefly, PCBs in soil and plant material (roots and shoots) was extracted by adding a 1:1 hexane/acetone mixture (3 ml/g) to homogenized soil or plant material (5 g) and sonicated for 1 h. Before sonication, surrogate standards including PCB 14 (3,5-dichlorobiphenyl), deuterated PCB 65 (2,3,5,6-tetrachlorobiphenyl) and PCB 166 (2,3,4,4',5,6-hexachlorobiphenyl) (Cambridge Isotope Laboratories, Inc.) were added into

⁵ PCB extraction was performed by Richard Meggo, Ph.D., Department of Civil & Environmental Engineering, The University of Iowa. PCB quantification was performed by Dingfei Hu, Ph.D., Department of Civil & Environmental Engineering, The University of Iowa.

the samples (50 ng of each surrogate) to account for any loss during the extraction. Surrogate recoveries were $103 \pm 17.2\%$ (PCB 14), $97.9 \pm 15.2\%$ (PCB 65) and $97.7 \pm 17.7\%$ (PCB 166).

The sonicated material was centrifuged at $1500 \times g$ for 5 min, after which the supernatant was transferred to a fresh vial. The precipitates were subjected to a second extraction. The combined supernatants of first and second extraction were evaporated to dryness using rotary evaporation and the solvent was changed to hexane. Double extraction with concentrated sulfuric acid and hexane was performed to remove of lipids and other polar substances. This hexane extract was concentrated to approximately 0.5 ml under a gentle stream of nitrogen. The concentrate was eluted with 10 ml of hexane through a filter consisting of 0.1 g of silica (70-230 mesh, Fisher Scientific, Inc.), 0.1 g of anhydrous sodium sulfate and 0.9 g silica gel acidified with H_2SO_4 (silica: H_2SO_4 = 2:1) (Meggo *et al.*, 2013).

The concentrated extracts were spiked with the internal standard containing 100 ng of PCB 204 (2,2',3,4,4',5,6,6'-octachlorobiphenyl). The samples were analyzed for PCB congeners using a gas chromatograph with mass selective detection (GC-MS/MS) modified from the EPA method 1668A (U.S.EPA 1999). The quantification of PCBs was performed by an Agilent 6890N gas chromatograph with an Agilent 7683 series auto sampler coupled to a Waters Micromass Quattro micro GC mass spectrometer (Milford, MA, USA) operating under electron impact (EI) positive mode at 70 eV and multiple reaction monitoring (MRM), and the trap current was 200 μA . The retention windows were defined by PCB parent/daughter ion pairs from mono- to deca- homologs which were 188/152, 222/152.10, 255.96/186, 291.92/222, 325.88/255.90, 359.84/289.90, 393.80/323.90, 427.76/357.80, 461.72/391.83, 497.68/427.70, respectively (Hu *et al.*, 2010).

DNA and RNA extraction,
reverse transcription, and quantitative PCR

Total RNA and DNA were extracted immediately after sampling from 2 g soil using the MoBio RNA PowerSoil Total RNA Isolation Kit (Mobio, Carlsbad, CA) and RNA PowerSoil DNA Elution Accessory Kit (Mobio, Carlsbad, CA) and stored at -80°C prior to further analysis. Before quantitative PCR (qPCR), RNA samples were subjected to contaminating DNA removal, RNA clean-up and reverse transcription. Contaminating DNA was removed by DNase I treatment according to the manufacturer's instructions (Biolab, Ipswich, MA). Each 100 µl DNase I treatment reaction contained 30 µl RNA and 20U DNase I. RNA was then purified with the RNeasy Mini Kit (Qiagen, Germantown, MD) and reversed-transcribed to cDNA by SuperScript II Reverse Transcriptase (Invitrogen, Carlsbad, CA) according to manufacturer's instructions. Each 80 µl reverse transcription reaction contained 40 µl purified RNA, 10mM dNTP mix, and 3 µg random primers.

To evaluate the RNA recovery efficiency during DNaseI treatment, RNA purification and reverse transcription, 1 ng of luciferase control mRNA (Promega, Madison, WI) was added to RNA sample before DNaseI treatment. After reverse transcription, luciferase cDNA was quantified by qPCR with the ref primer set (Table 5.1) as described previously (Johnson *et al.*, 2005). The RNA recovery efficiency averaged $14.2\% \pm 11.5\%$ among all RNA samples processed in this study.

The abundances of total bacteria, aerobic PCB-degrading bacteria, and *Burkholderia xenovorans* strain LB400 were estimated by qPCR using the bacterial 16S rRNA gene primer set 16SU f/r (Nadkarni *et al.*, 2002), *bphA* primer set *bphA* 463f/674r (Petrić *et al.*, 2011b), and LB400 16S-23S rRNA internal transcribed spacers(ITS) primer set LB400 84f/278r (Norini *et al.*, 2013), respectively (Table 5.1). Each 25 µl qPCR reaction contained 12.5 µl Power SYBR Green PCR Master Mix (Invitrogen, Carlsbad, CA), and variable amounts of primers and templates (Table C.2). Bovine serum

albumin(0.5 µg) was added to relieve possible PCR inhibition (Kreader, 1996). PCR conditions were as follows: 10 min at 95°C, followed by 40 cycles at 95°C (15 s) and 60°C (1 min) followed by a dissociation step. All qPCR was performed with an ABI 7000 Sequence Detection System (Applied Biosystem, Grand Island, NY) and fluorescence data was analyzed by ABI 7000 System SDS Software (Applied Biosystems, Grand Island, NY). With each primer set, the target gene was not detected in no template (DI water) controls (Ct value > 35). Additional qPCR information, such as primer concentrations, template concentrations, qPCR linear range, qPCR efficiency range of the standard curves, and Y-intercepts are provided in Table 5.2, in accordance with MIQE guidelines (Bustin *et al.*, 2009).

For luciferase mRNA qPCR, the standard DNA template was PCR products amplified from luciferase DNA with primer set refST (Table 5.1). For the bacterial 16S rRNA gene, the standard DNA template was prepared from PCR products amplified from *Burkholderia xenovorans* strain LB400 with primer set 8F/1492R (Klappenbach *et al.*, 2000). For *bphA*, the standard DNA template was the LB400 *bphA* (amplified with the 463f/674r primer set) cloned into the pCR 2.1-TOPO vector. For LB400 16S-23S rRNA ITS gene, the standard DNA template was the LB400 16S-23S rRNA ITS gene (amplified with the LB400 84f/278r primer set) cloned into the pCR 2.1-TOPO vector.

Statistical analyses

PCB congener data was analyzed with an independent sample t test using R. Total bacterial 16S rRNA gene abundance, *bphA* gene and transcript abundance and LB400 16S-23S rRNA ITS abundance were analyzed with a two factor analysis of variance after log transformation using R (Rieu and Powers, 2009).

Results and Discussion

Enhanced degradation of specific PCB congeners in soil with switchgrass treatment

Significantly higher removal of PCB parent congeners (PCB 52, 77, 153) was achieved in the switchgrass-treated soil than in the unplanted soil after 24 weeks of incubation ($p < 0.05$) (Figure 5.1). The presence of switchgrass enhanced PCB 52 removal more than the other two congeners (16.7%), while PCB 77 and PCB 153 removal was improved by 7.9% and 9.6%, respectively. Overall, $39.9 \pm 0.4\%$ of the total PCB molar mass loss was observed in switchgrass treated soil after 24 weeks of incubation, compared with $29.5 \pm 3.4\%$ in unplanted soil (Figure 5.1). The increase of PCB removal efficiency by switchgrass in this study is comparable to some previous studies, although different plants were used (Ding et al., 2009; Ionescu et al., 2009; Secher et al., 2013; Smith et al., 2007).

Switchgrass has been observed previously to enhance PCB removal in soil and promote microbial activity in the rhizosphere (Chekol et al., 2004; Dzantor et al., 2000; Meggo et al., 2013). The enhanced PCB removal could be the result of rhizosphere biostimulation, rhizofiltration, phytoextraction, phytotransformation and phytovolatilization (Van Aken *et al.*, 2010). We investigated this possibility of phytoextraction by quantifying total PCBs in switchgrass roots and shoots. At the end of the 24 week experiment, we found 4.3-7.3% of total PCBs in switchgrass roots, and 0.11-0.24% in shoots, suggesting that plant extraction were active as PCB removal mechanisms (Table 5.3). All three parent congeners and some transformation products were detected in switchgrass roots and shoots. Especially, the detection of PCB 153 in switchgrass shoots demonstrated its capability to take up and translocate highly chlorinated PCB congeners. The uptake of PCBs by switchgrass and other plants has been reported (Ficko et al., 2010; Greenwood et al., 2011; Liu et al., 2009), although in some studies the effect of phytoextraction was considered to be negligible in removing

PCBs from soil because of the low percentage of total PCB mass observed in plant roots and shoots (<1% of total PCBs) (Di Gregorio et al., 2013; Huesemann et al., 2009; Li et al., 2011). In this study, the fraction of total PCB mass in switchgrass was higher than previous studies, probably because of the high plant biomass for each switchgrass treated soil microcosm (560 ± 51 g roots and 360 ± 41 g shoots per soil microcosm).

Phytovolatilization and phytotransformation were not considered to affect the PCB removal much, given the low PCB concentrations in switchgrass shoots. Previous study also showed that hydroxylated metabolite was not detected in switchgrass hydroponically exposed to PCBs, suggesting that PCB transformation in switchgrass was not active (Liu et al., 2009).

Biostimulation of microbial PCB degradation was considered to be another mechanism contributing to enhanced PCB removal in the rhizosphere soil. The occurrence of aerobic microbial PCB degradation in both unplanted and switchgrass treated soils was verified by the decrease in total PCB molar mass (Figure 5.1A). The enhanced microbial activity was estimated to be responsible for about 3.05% total PCB removal from soil (Table 5.3). Different PCB congeners were affected differently by the enhanced microbial PCB degradation. It is estimated that 7.1% of PCB 52 removal was due to the enhanced microbial activity, while this number was 1.8% and 1.3% for PCB 77 and PCB 153, respectively. Microbial degradation of PCB 52 was promoted more than the other two congeners probably because PCB 52 is more susceptible for aerobic degradation by indigenous microorganisms. PCB 77 has a dioxin-like structure and was relatively more toxic than other tetrachlorinated biphenyls. The high chlorination level of PCB 153 results in high hydrophobicity and reduces its accessibility to enzymes. The bulkiness of the chlorine atoms may also prevent access to the enzyme's active site (Bedard et al., 1986).

Molecular biology analyses of unplanted and switchgrass planted soils

We estimated the abundance of total bacteria, biphenyl dioxygenase gene (*bphA*) and *bphA* transcripts as an independent means of tracking aerobic microbial PCB biodegradation during the experiment. Total bacterial abundance (as estimated by the abundance of bacterial 16S rRNA genes) was significantly higher in switchgrass reactors than in unplanted reactors ($p < 0.05$, Figure 5.2A). We suspect that the switchgrass serves to improve oxygen diffusion and soil permeability. Switchgrass root exudates could also increase the microbial population in the rhizosphere by serving as a carbon and energy source.

Detecting *bphA* in both unplanted and switchgrass-treated soils suggests that the indigenous soil microbial community harbored the potential for aerobic PCB degradation. In the presence of switchgrass, *bphA* abundance was about 20 times higher ($p < 0.05$, Figure 5.2B). However, *bphA* transcripts in both switchgrass planted and unplanted reactors were below our quantification limit of RT-qPCR (40 copies/qPCR reaction), suggesting that the *bphA* expression was not promoted much by the switchgrass. It is possible that the presence of other carbon sources in the rhizosphere limited the *bphA* expression (Parnell *et al.*, 2010). The availability of a variety of naturally-occurring organic carbon and plant exudates in the rhizosphere may make PCBs less favorable for indigenous microorganisms (Luo *et al.*, 2008). Soil extract may also inhibit *bphA* expression (Master and Mohn, 2001). The low level of *bphA* expression could also explain why microbial PCB degradation was not improved much by switchgrass as suggested by congener data.

Enhanced microbial PCB degradation in soil with LB400 bioaugmentation

To further enhance aerobic degradation of PCB congeners in the rhizosphere,

biphenyl-grown *Burkholderia xenovorans* strain LB400 was introduced to the soil (at a loading of approximately 10^9 CFU per g soil). LB400 is an effective aerobic PCB-degrader, capable of degrading a broad spectrum of PCB congeners including those used in this study (PCB 52, 77, 153) (Bedard et al., 1986; Bopp, 1986). With LB400 bioaugmentation, total PCB molar mass loss increased to $39.4 \pm 1.4\%$ and $47.3 \pm 1.2\%$ in the unplanted and switchgrass-treated soil, respectively, after 24 weeks of incubation, suggesting enhanced aerobic PCB degradation (Figure 5.1). Removal of PCB 52, 77 and 153 was not significantly improved in the control switchgrass reactor, which was amended with autoclaved LB400 (Figure 5.1).

Specifically, PCB 52 removal appeared to be most improved by LB400 bioaugmentation, achieving 43.7% and 24.9% more removal in unplanted and switchgrass-treated soils, respectively ($p < 0.05$). PCB 153 removal in switchgrass-treated soil with bioaugmentation increased by 3.3% as compared with soil with only switchgrass treatment ($p < 0.05$). In unplanted soil with bioaugmentation, PCB 77 and 153 removal was not improved (Figure 5.1). This observation is consistent with previous LB400 PCB biodegradation studies, where LB400 displayed a stronger capability to degrade PCB 52 than PCB 77 and 153 (Bedard et al., 1986; Bopp, 1986; Rein et al., 2007). Compared with PCB 52, PCB 77 and 153 have higher K_{ow} values and lower water solubilities. Although LB400 could degrade PCB 77 and 153 in the resting cell assays, in the soil microcosms, the two congeners may be sorbed strongly to soil particles and thus are less bioavailable.

Molecular biology analyses of unplanted, switchgrass planted, and LB400 bioaugmented soils

The enhanced PCB removal in bioaugmented soil was associated with elevated abundance of *bphA* and its transcripts, suggesting that aerobic PCB degradation was actively occurring (Figure 5.3, Figure 5.4). In bioaugmented soil, *bphA* abundance

increased throughout incubation time, and was two orders of magnitude higher than in reactors without bioaugmentation ($p < 0.05$, Figure 5.3B, Figure 5.3D). The transcripts of *bphA* were detected (3.17×10^5 - 8.90×10^5 transcripts per g soil in unplanted soil with bioaugmentation, 4.73×10^4 - 4.14×10^6 transcripts per g soil in switchgrass treated soil with bioaugmentation), suggesting the successful *bphA* expression (Figure 5.4).

Bioaugmentation did not significantly increase the total microbial abundance ($p > 0.05$, Figure 5.3A, Figure 5.3C). Yet the reactor with autoclaved LB400 injection displayed greater total 16S rRNA gene abundance than the reactor with live LB400 bioaugmentation ($p < 0.05$, Figure 5.3A), probably because debris from the autoclaved LB400 cells was used as carbon and energy source by indigenous microorganisms.

The survival of bioaugmented bacteria is crucial to the successful bioaugmentation and contaminant removal (Bouchez *et al.*, 2000). To improve LB400 survival and keep a relatively stable LB400 population, we performed bioaugmentation once a month. The survival of LB400 after bioaugmentation was estimated by quantifying biphenyl-degrading bacteria with CFU enumeration. After bioaugmentation, the biphenyl-degrading bacterial abundance increased (4.7×10^8 - 1.9×10^9 CFU per g soil for unplanted reactor, 2.1×10^8 - 2.3×10^9 CFU per g soil for switchgrass planted reactor). Yet this number decreased to the level of approximately 10^7 CFU per g soil after four weeks, prior to the next bioaugmentation event (Figure 5.6). The decrease may result from the stressed environment caused by PCB (Kim *et al.*, 2001; Parnell *et al.*, 2006), and the competition with indigenous microbial community. LB400 16-23S ITS gene was quantified approximately two weeks after bioaugmentation, and ranged from 1.16×10^7 to 4.15×10^8 gene copy per g soil (unplanted soil), and 9.61×10^7 to 7.04×10^8 gene copy per g soil (switchgrass treated soil) (Figure 5.5). An accumulation of ITS genes was observed throughout the 24 week experiment, suggesting that after each bioaugmentation, a percentage of LB400 adapted to the soil environment and balance with indigenous microorganisms (Figure 5.5).

LB400 16-23S ITS gene quantification shows higher LB400 abundance with the presence of switchgrass (Figure 5.5). Also, the most PCB removal, the highest *bphA* and transcript abundances were observed in switchgrass treated soil with bioaugmentation after 24 weeks of treatment (Figure 5.1, Figure 5.3 and Figure 5.4), indicating switchgrass not only benefited LB400 survival, but also helped LB400 degrade PCBs. The support of plants to bioaugmentation was reported (Juhanson et al., 2009; Secher et al., 2013; Tam and Wong, 2008). Switchgrass helped LB400 survival may be by improving aeration and providing root exudates as extra carbon and energy source, the same way as it benefited indigenous microorganisms. The root system can help spread LB400 through the PCB contaminated soil. Furthermore, switchgrass may increase the bioavailability of PCB molecules by releasing organic compounds as complexing agents.

Conclusions

In this study, switchgrass was found to be able to improve the removal of PCB 52, 77 and 153 by phytoextraction and biostimulation. PCB 52 was more susceptible to microbial degradation, and its removal was more improved by switchgrass and bioaugmentation with LB400 than PCB 77 and 153. Switchgrass was beneficial for LB400 survival and its ability to perform aerobic degradation. The combination of phytoremediation and bioaugmentation offers an efficient and environmental-friendly strategy to eliminate recalcitrant PCB congeners and remediate PCB contaminated soil.

Table 5.1. Primer sets used in qPCR analysis in Chapter V.

Target gene	Primer name	Sequences	Expected product size	Source
luciferase mRNA	ref f	5'-TACAACACCCCAACATCTTCGA-3'	150 bp	(Johnson <i>et al.</i> , 2005)
	ref r	5'-GGAAGTTCACCGGCGTCAT-3'		
total bacterial 16S rRNA	16SU f	5'-TCCTACGGGAGGCAGCAGT-3'	466 bp	(Nadkarni <i>et al.</i> , 2002)
	16SU r	5'-GGACTACCAGGGTATCTAATCCTGTT-3'		
<i>bphA</i>	<i>bphA</i> 463f	5'-CGCGTSGMVACCTACAARG-3'	211 bp	(Petrić <i>et al.</i> , 2011b)
	<i>bphA</i> 674r	5'-GGTACATGTCRCTGCAGAAYTGC-3'		
LB400 16S-23S ITS	LB400 84f	5'-GAAATTGAAGACAGAAACGCA-3'	191 bp	(Norini <i>et al.</i> , 2013)
	LB400 278r	5'-AGTCATGCACACACCAGAT-3'		

Table 5.2. qPCR reaction set up, linear range and efficiency for qPCR reaction in Chapter V.

Target gene	primer conc, μM	Template	qPCR linear range, gene copies/reaction	qPCR efficiency	Y-intercept
luciferase mRNA	0.1	2 μl cDNA	30 - 30×10^7	95 - 100%	37.65 - 38.61
total bacterial 16S rRNA	0.1	1 ng DNA	30 - 30×10^7	95 - 100%	35.10 - 35.84
<i>bphA</i>	1	10 ng DNA or 2 μl cDNA	40 - 40×10^7	95 - 100%	36.43 - 37.64
LB400 16S-23S ITS	0.2	10 ng DNA	40 - 40×10^7	90 - 100%	36.54 - 37.44

Table 5.3. Molar mass balance of PCBs in unplanted soil (UP), switchgrass treated soil (SG), unplanted with bioaugmentation soil (UP B), switchgrass treated with bioaugmentation soil (SG B), and switchgrass treated with bioaugmentation with dead LB400 soil (SG BD). (n=3).

		PCB fraction in soils	PCB fraction in roots	PCB fraction in shoots	Total recovery
Overall	UP	70.5 ± 3.4%	-	-	70.5 ± 3.4%
	SG	60.1 ± 0.4%	7.3 ± 0.2%	0.11 ± 0.01%	67.5 ± 0.6%
	UP B	60.6 ± 1.4%	-	-	60.6 ± 1.4%
	SG B	52.7 ± 1.2%	4.3 ± 0.1%	0.24 ± 0.03%	57.3 ± 1.4%
	SG DB	57.1 ± 10.4%	-	-	57.1 ± 10.4%
PCB 52	UP	98.6 ± 3.9%	-	-	98.6 ± 3.9%
	SG	81.9 ± 0.6%	9.4 ± 0.2%	0.17 ± 0.01%	91.4 ± 0.8%
	UP B	54.8 ± 0.8%	-	-	54.8 ± 0.8%
	SG B	57.0 ± 1.7%	4.7 ± 0.1%	0.40 ± 0.04%	62.2 ± 1.9%
	SG DB	76.7 ± 10.7%	-	-	76.7 ± 10.7%
PCB 77	UP	61.0 ± 4.0%	-	-	61.0 ± 4.0%
	SG	53.2 ± 1.2%	6.0 ± 0.2%	0.06 ± 0.01%	59.2 ± 1.4%
	UP B	62.1 ± 1.8%	-	-	62.1 ± 1.8%
	SG B	53.1 ± 1.4%	3.8 ± 0.2%	0.16 ± 0.01%	57.1 ± 1.6%
	SG DB	51.1 ± 9.4%	-	-	51.1 ± 9.4%
PCB 153	UP	66.1 ± 2.4%	-	-	66.1 ± 2.4%
	SG	56.5 ± 0.0%	8.2 ± 0.2%	0.11 ± 0.02%	64.8 ± 0.25%
	UP B	69.7 ± 1.5%	-	-	69.7 ± 1.5%
	SG B	53.2 ± 0.6%	5.0 ± 0.1%	0.17 ± 0.02%	58.4 ± 0.7%
	SG DB	53.7 ± 9.0 %	-	-	53.7 ± 9.0 %




Figure 5.1. The percentage of initial molar concentrations for total PCB (A), PCB 52 (B), PCB 77 (C), and PCB 153(D) after 12 weeks and 24 weeks incubation in unplanted soil (UP), switchgrass treated soil (SG), unplanted with bioaugmentation soil (UP B), switchgrass treated with bioaugmentation soil (SG B), and switchgrass treated with bioaugmentation with dead LB400 soil (SG BD). Error bars indicate the standard deviation of three soil subsamples from the same reactor.




Figure 5.2. qPCR analysis of (A) bacterial 16S rRNA genes and (B) *bphA* in blank (BLK), PCB spiked and unplanted soil (UP), PCB spiked and switchgrass treated soil (SG). Error bars indicate the range of two soil subsamples from the same reactor.




Figure 5.3. qPCR analysis of bacterial 16S rRNA gene (A) and *bphA* (B) in switchgrass treated soil (SG), switchgrass treated soil with bioaugmentation (SG B), switchgrass treated soil with autoclaved LB400 bioaugmentation (SG BD); and bacterial 16S rRNA gene (C) and *bphA* (D) in unplanted soil (UP), unplanted soil with LB400 bioaugmentation (UP B), switchgrass treated soil with bioaugmentation (SG B). Error bars indicate the range of two soil subsamples from the same reactor.




Figure 5.4. qPCR analysis of *bphA* transcripts in unplanted soil with bioaugmentation (UP B), and switchgrass treated soil with bioaugmentation (SG B). Error bars indicate the range of two soil subsamples from the same reactor.




Figure 5.5. qPCR analysis of LB400 16S-23S ITS genes in unplanted soil with bioaugmentation (UP B), and switchgrass treated soil with bioaugmentation (SG B). Error bars indicate the range of two soil subsamples from the same reactor.




Figure 5.6. CFU counting for unplanted soil with bioaugmentation (UP B), switchgrass treated soil with bioaugmentation (SG B) and switchgrass treated soil with dead LB400 bioaugmentation (SG BD) before and after the application of bioaugmentation. Error bars indicate the standard deviation of three soil subsamples from the same reactor.

CHAPTER VI. MICROBIAL COMMUNITY ANALYSIS OF A SWITCHGRASS RHIZOSPHERE WITH PCB SPIKING AND REDOX CYCLING

Introduction

Bioremediation of PCB contaminated soils and sediments has always been challenging because of the hydrophobicity and recalcitrance of PCB congeners (Agarwal et al., 2007; Amend and Lederman, 1992). Phytoremediation of PCBs, the use of plants and associated microorganisms to remediate PCB contaminated soils and sediments, is more cost-effective and less disruptive to the environment (Van Aken *et al.*, 2010).

Microbial PCB degradation in the rhizosphere plays an important part in the phytoremediation of PCBs (Borja et al., 2005; Macková et al., 2010; Pieper and Seeger, 2008). Microorganisms are capable of degrading and detoxifying PCBs via anaerobic PCB dechlorination and anaerobic PCB mineralization (Borja et al., 2005; Gibson and Parales, 2000; Pieper and Seeger, 2008; Wiegel and Wu, 2000). Anaerobic PCB degraders can use PCB congeners as electron acceptors and remove chlorine atoms from the biphenyl ring or dechlorinate PCBs via co-metabolism (Adrian et al., 2009; Wiegel and Wu, 2000; Wu et al., 1998). So far all the known PCB degraders are within the *Chloroflexi* phylum, including the *Dehalococcoides* spp. and the o-17/DF-1 group (Fagervold et al., 2007; Fennell et al., 2004; Yan et al., 2006). Under aerobic conditions, some microorganism can mineralize PCBs via upper biphenyl pathway, leaving behind a chlorobenzoate as product (Gibson and Parales, 2000; Macková et al., 2010).

Many studies have confirmed the effect of plants on enhancing PCB degradation and increasing microbial PCB degrading populations (de Cárcer et al., 2007; Leigh et al., 2006; Macková et al., 2006; Slater et al., 2011). However, most of the current studies focus on the influence of plants on aerobic PCB degradation. The anaerobic dechlorination in the rhizosphere is rarely investigated. Here in this study, we set up soil

reactors with switchgrass treatment and created an anaerobic environment by flooding the soil to facilitate dechlorination. PCB congener analysis confirmed the occurrence of dechlorination in switchgrass treated soils, with or without redox cycling. Putative dechlorinating *Chloroflexi* was more abundant in redox cycled soils, as shown by qPCR. Other bacteria capable of dechlorination such as *Geobacter* and *Clostridium* were detected by clone library analysis, indicating their possible role in PCB dechlorination.

Materials and Methods

Soil reactor set up⁶

The soil microcosm set-up procedure was similar as previously described (Meggo and Schnoor, 2013). Briefly, PCB-free soil from the village of Middle Amana in Iowa, USA, was passed through a 60 mesh sieve before experiments. The soil was spiked with a mixture of PCB 52, PCB 77 and PCB 153 (99% pure) (Accustandard Inc., New Haven, CT) at the concentration of 500 ng g⁻¹ each. PCB congeners were dissolved in hexane before adding to the soil. A quartering technique was used to homogenize soil. The soil was divided into four quadrants on a quartering canvas, and 20 diagonal trajectories were used to mix the soil components. The contaminated soil was aged for two months at 25°C in sealed tubs to allow PCBs sequestration into the soil matrix and thereby better represent field conditions. To establish the initial PCB concentration, 20 subsets of soil collected from different locations were homogenized and measurements were performed in triplicates (Meggo and Schnoor, 2013).

Plastic containers (33.8cm × 21.6cm × 21.9cm) and lids with aluminum foil covers were used for soil reactor set-up. Each reactor was filled with 2500g sieved and homogenized soil. The spiked soil was planted with switchgrass (*Panicum virgatum*)

⁶ Soil reactor set-up was performed by Richard Meggo, Ph.D., Department of Civil & Environmental Engineering, The University of Iowa.

seeds (Adams-Briscoe Seed Co., Jackson, GA). Switchgrass in all four reactors was grown at 25°C in a plant growth chamber under a 16 hour light/8 hour dark photo period (light intensity of 200 $\text{mmol m}^{-2}\text{s}^{-1}$ and 60% humidity). Switchgrass was allowed to establish for 8 weeks. After 8 weeks, alternate cycles (two weeks of flooding and two weeks of non-flooding) were performed to one unplanted soil reactor and one switchgrass treated soil reactor. During the flooding period, soils were submerged by deionized water. After two weeks of flooding, the extra water was withdrawn. Four flooding was performed in total. Another unplanted soil reactor and switchgrass treated soil reactor with normal moisture content and without flooding treatment were used as control. For each soil reactor, redox measurement was performed with a Redox/pH combination meter (Hanna Instruments, Smithfield, RI). Volumetric soil moisture content was monitored with Vernier soil moisture sensors interfaced with the Vernier LabQuest data collection system (Vernier Software & Technology, Beaverton, OR). Soil samples were taken at 12 week and 24 week for PCB congener analysis. For microbial analysis, soil samples were taken every two weeks starting at 8 week (Meggo and Schnoor, 2013).

PCB congener analysis⁷

The PCB extraction procedure has been described previously (Meggo *et al.*, 2013). PCBs in soil and plant material (roots and shoots) was extracted by adding a 1:1 hexane/acetone mixture (3 ml/g) to homogenized soil or plant material (5 g) and sonicated for 1 h. Before sonication, surrogate standards including PCB 14 (3,5-dichlorobiphenyl), deuterated PCB 65 (2,3,5,6-tetrachlorobiphenyl) and PCB 166 (2,3,4,4',5,6-hexachlorobiphenyl) (CambridgeIsotope Laboratories, Inc.) were added into the samples (50 ng of each surrogate) to account for any loss during the extraction.

⁷ PCB extraction was performed by Richard Meggo, Ph.D., Department of Civil & Environmental Engineering, The University of Iowa. PCB quantification was performed by Dingfei Hu, Ph.D., Department of Civil & Environmental Engineering, The University of Iowa.

Surrogate recoveries were $103 \pm 17.2\%$ (PCB 14), $97.9 \pm 15.2\%$ (PCB 65) and $97.7 \pm 17.7\%$ (PCB 166).

The sonicated material was centrifuged at $1500 \times g$ for 5 min, after which the supernatant was transferred to a fresh vial. The precipitates were subjected to a second extraction. The combined supernatants of first and second extraction were evaporated to dryness using rotary evaporation and the solvent was changed to hexane. Double extraction with concentrated sulfuric acid and hexane was performed to remove of lipids and other polar substances. This hexane extract was concentrated to approximately 0.5 ml under a gentle stream of nitrogen. The concentrate was eluted with 10 ml of hexane through a filter consisting of 0.1 g of silica (70-230 mesh, Fisher Scientific, Inc.), 0.1 g of anhydrous sodium sulfate and 0.9 g silica gel acidified with H_2SO_4 (silica: H_2SO_4 = 2:1) (Meggo and Schnoor, 2013).

The concentrated extracts were spiked with the internal standard containing 100 ng of PCB 204 (2,2',3,4,4',5,6,6'-octachlorobiphenyl). The samples were analyzed PCB congeners using a gas chromatograph with mass selective detection (GC-MS/MS) modified from the EPA method 1668A (U.S.EPA 1999). The quantification of PCBs was performed by an Agilent 6890N gas chromatograph with an Agilent 7683 series autosampler coupled to a Waters Micromass Quattro micro GC mass spectrometer (Milford, MA, USA) operating under electron impact (EI) positive mode at 70 eV and multiple reaction monitoring (MRM), and the trap current was 200 μA . The retention windows were defined by PCB parent/daughter ion pairs from mono- to deca- homologs which were 188/152, 222/152.10, 255.96/186, 291.92/222, 325.88/255.90, 359.84/289.90, 393.80/323.90, 427.76/357.80, 461.72/391.83, 497.68/427.70, respectively (Hu *et al.*, 2010).

Quantitative PCR

DNA was extracted from 2 g soil immediately after sampling using the RNA

PowerSoil DNA Elution Accessory Kit (Mobio, Carlsbad, CA) and stored at -80 C prior to further analysis. The abundances of total bacterial and Putative dechlorinating *Chloroflexi* were estimated using qPCR targeting bacterial 16S rRNA gene (primer set 16SU f/r) (Nadkarni *et al.*, 2002) and putative dechlorinating *Chloroflexi* 16S rRNA genes (primer set dhc793f/946r) (Yoshida *et al.*, 2005), respectively. PCR conditions were as follows: 10 min at 95°C, 40 cycles of 15s at 95°C and 1 min at 60°C followed by a dissociation step. Each 25 µl qPCR reaction contained 12.5 µl Power SYBR Green PCR Master Mix (Invitrogen, Carlsbad, CA), 2.5 pmol primers and various amount of primers and templates (Table 6.1). Bovine serum albumin (0.5 µg) was added to relieve possible PCR inhibition (Kreader, 1996).

For total bacterial 16S rRNA gene qPCR, the standard DNA template was a PCR product amplified from *Burkholderia xenovorans* strain LB400 with primer set 8F/1492R (Grabowski *et al.*, 2005). For putative dechlorinating *Chloroflexi* 16S rRNA gene, standard curves were prepared from pCR 2.1-TOPO vector containing the target PCR products with primer dhc793f/946r. All qPCRs were performed with an ABI 7000 Sequence Detection System (Applied Biosystem, Grand Island, NY) and fluorescence data was analyzed by ABI 7000 System SDS Software (Applied Biosystems, Grand Island, NY). With each primer set, the target gene was not detected in no template (DI water) controls (Ct value > 35). Additional qPCR information, such as primer concentrations, template concentrations, qPCR linear range, qPCR efficiency range of the standard curves, and Y-intercepts are provided in Table 6.2, in accordance with MIQE guidelines (Bustin *et al.*, 2009).

qPCR quality assurance

Several published qPCR primer sets targeting putative dechlorinating *Chloroflexi* 16S rRNA gene and *Dehalococoides*-like 16S rRNA gene were initially tested in this study. Primer sets chl348f/884r (Fagervold *et al.*, 2005), dhc1f/264r (Grostern and

Edwards, 2006) , dhc793f/946r (Yoshida *et al.*, 2005) and dhc1154f/1286r (Krzmarzick *et al.*, 2012) produced single bands of the expected size with DNA template derived from soil. To verify the specificity of these qPCR primer sets, clone libraries were constructed from the amplification products in DNA extracted from soil with redox cycling treatment. From the dhc793f/946r PCR product clone library, 6 unique sequences were obtained from 12 clones and all of these sequences were identified to be *Dehalogenimonas* by RDP classifier (Table 6.3) (Cole *et al.*, 2007). Clone libraries show that primer sets chl348f/884r, dhc1f/264r and dhc1154f/1286r amplified sequences from the class *Anaerolineae* of *Chloroflexi* and also from *Actinobacteria*, both of which are not associated with dechlorination (Table 6.3). Only the primer set dhc793f/946r was used to quantify putative dechlorinating *Chloroflexi* 16S rRNA gene in soil samples. The specificity of the SYBR green based qPCR was also validated by dissociation curve analysis, which shows similar melting temperature for each sample.

Terminal-Restriction Fragment Length

Polymorphism (T-RFLP) analysis

PCR with a fluorescent-labeled primer set 6-FAM 8Fm/533R was performed with 1 ng DNA as template (Schütte *et al.*, 2008). PCR conditions were as follows: 4 min at 94°C, 30 cycles of 1 min at 94°C, 2 min at 55°C and 1.5 min at 72°C followed by a final extension step of 10 min at 72°C. PCR product (12 µl) was digested with the restriction enzyme HaeIII (New England BioLabs, Inc., Ipswich, MA), and then precipitated with glycogen (Fermentas International Inc., Vilnius, Lithuania), sodium acetate (pH 5.2), and ethanol at -20°C for 2h. DNA was recovered by centrifugation at 17,800 × g for 15 min and resuspended in distilled water (Invitrogen Corp., Carlsbad, CA). The digested DNA was sent to the University of Iowa DNA facility for electrophoresis for sizing using an Applied Biosystems 3730 DNA analyzer (Life Technologies Corporation, Carlsbad, CA) with the GeneScan 500 LIZ size standard.

Terminal restriction fragment (TRF) sizes were estimated with Peak Scanner software (Applied Biosystems, Carlsbad, CA). A terminal restriction fragment (T-RF) size matrix containing 84 samples (rows) and 439 unique 16S rRNA gene TRFs (columns) was generated by T-REX software after filtering background peak noise and rounding fragment sizes to the nearest whole number (Culman *et al.*, 2009). TRF profiles were analyzed by nonmetric multidimensional scaling (NMDS) in R (da C Jesus *et al.*, 2009). The Bray-Curtis dissimilarity index was calculated with a random starting configuration, and a two-dimensional solution was reached after two runs. The final stress was 12.77. The T-RF composition differences among groups were assessed by multi-response permutation procedure (MRPP), which evaluates grouping with real and randomized data based on Bray-Curtis distances and 5000 permutations, and permutational multivariate analyses of variance (PERMANOVA) based on Bray-Curtis distances with 5000 permutations. Random forests classification was used to estimate the importance of the individual T-RFs in the classification (Cutler *et al.*, 2007). An indicator species analysis was performed to assess the treatment preference of T-RFs using *multipatt* function with R (Cáceres and Legendre, 2009).

PCR amplification, cloning and sequencing of bacterial 16S rRNA

Clone libraries were constructed with bacterial 16S rRNA genes amplified from unplanted soil, switchgrass treated soil, unplanted and cycled soil, switchgrass treated and cycled soil at 24 wk. Bacterial 16S rRNA genes were amplified using primer set 8F/1492R (Grabowski *et al.*, 2005). PCR conditions were as described in T-RFLP analysis section. PCR products were purified with the Qiaquick PCR purification Kit (Qiagen Inc., Valencia, CA), cloned into the pCR 2.1-TOPO vector using the TOPO TA cloning Kit (Invitrogen Corp., Carlsbad, CA), and transformed into One Shot TOP10 chemically competent *E.coli* cells (Invitrogen Corp., Carlsbad, CA). The transformation

efficiency was checked by plating recombinant *E.coli* on Luria Broth agar with kanamycin (50 mg/L), and X-gal (0.4 mg/plate) and incubated overnight at 37°C. Clones were Sanger sequenced at HTSeq.org (Seattle, WA) with M13F (5'-TGTAACGACGGCCAGT-3') and M13R (5'-CAGGAAACAGCTATGAC-3') primers. Sequence identification and classification were performed using RDP Classifier (Cole *et al.*, 2007) and Basic Local Alignment Search Tool (BLAST) . Predicted TRFs were generated with TRiFLE (Junier *et al.*, 2008).

Statistical analyses

PCB congener data was analyzed with an independent sample t test using R. The total bacterial 16S rRNA gene copy numbers and dechlorinating *Chloroflexi* 16S rRNA gene copy numbers were analyzed with a two factor analysis of variance after log transformation using R (Rieu and Powers, 2009).

Results and Discussion

PCB congener analysis

Significantly higher total PCB removal (39.9 ± 0.4 %) was achieved in switchgrass treated soil as compared with unplanted soil (29.5 ± 3.4 %) ($p < 0.05$, Figure 6.1). PCB 52 removal was enhanced most (16.7%) with the presence of switchgrass, while PCB 77 and PCB 153 removal was improved by 7.9% and 9.6%, respectively ($p < 0.05$, Figure 6.1). The enhanced PCB removal in switchgrass rhizosphere could be the result of rhizosphere biostimulation, rhizofiltration, phytoextraction, phytotransformation and phytovolatilization (Van Aken *et al.*, 2010). At the end of the 24 week experiment, we found 4.5-10.8% of total PCBs in switchgrass roots, and 0.11-0.17% in shoots, suggesting the uptake and translocation of PCBs within switchgrass. PCB 153 was less

favorable by switchgrass, which has significantly lower concentration in the roots than PCB 52 and 77 ($p < 0.05$), probably because of the higher hydrophobicity.

The experiment design involving redox cycling treatment was intended to create alternating anaerobic-aerobic conditions in the soil reactors by flooding and draining soils. During the non-flooding period, redox potential measurement suggests that anoxic conditions prevailed in both unplanted and switchgrass treated soils ($E_h < 350\text{mV}$). During the two weeks of flooding period, the redox potential could drop as low as -400mV (Figure 6.4). In redox cycled and unplanted soil, PCB 153 removal was increased by 5.3% after 24 weeks of incubation ($p < 0.05$), which could be the result of the enhanced anaerobic dechlorination. The removal of PCB 52 and 77 was not significantly improved. In redox cycled soils with switchgrass treatment, PCB removal was not improved as compared with soil with only switchgrass treatment (Figure 6.1). It is possible that switchgrass roots enhanced the oxygen diffusion in the flooded soils, making it difficult to enhance dechlorination.

The occurrence of dechlorination was verified by the detection transformation products including PCB1, PCB3, PCB 35, PCB 48, and PCB 118 in unplanted, switchgrass treated, and redox cycled soils (Figure 6.2). As suggested by redox measurements, the soils without cycling of flooded and unflooded conditions were under anoxic conditions, which made it possible for microorganisms to use alternative electron acceptors (Figure 6.4). The observed dechlorination is encouraging, because PCB dechlorination was typically observed in anaerobic environments (Bedard et al., 1996; Krumins et al., 2009; Quensen et al., 1990; Rodenburg et al., 2010). However, the transformation products were not very abundant and didn't accumulate significantly, possibly because of simultaneous aerobic oxidation. PCB 3 was relatively more abundant at 12 weeks, and was at lower level at 24 weeks, indicating the possible PCB 3 removal by aerobic microbial degradation (Figure 6.2).

PCB 101 and PCB 118 could be the dechlorination products of PCB 153 (Figure

6.3). The detection of PCB 118 was notable, which requires *ortho* chlorine removal from PCB 153. PCB 101 could be transformed into PCB 52 by *para* chlorine removal, and PCB 118 can be dechlorinated to PCB 77 by *ortho* chlorine removal (Figure 6.3). Thus PCB 153 can be dechlorinated to PCB 52 and 77. The transformation from PCB 153 to PCB 52 is more likely, which can be achieved by removing two *para*-chlorines, while PCB 77 production requires *ortho* dechlorination and is more difficult (Tiedje *et al.*, 1993). PCB 35 could be the dechlorination product of PCB 77 by *para* chlorine removal, and PCB 26 could be generated from PCB 52 by *ortho* chlorine removal (Figure 6.3).

Some PCB congeners, such as PCB 95 and 110, were also detected. They are not the direct dechlorination product from the three parent PCB congener, and were probably produced by chlorine rearrangement, although the mechanism is unclear. Chlorine rearrangement of PCB congeners have been observed before in poplars, switchgrass and maize (Liu *et al.*, 2009; Wang *et al.*, 2011).

Quantification of putative dechlorinating *Chloroflexi* 16S rRNA genes

While the detection of transformation products verified the occurrence of dechlorination in the soils, we then aimed to quantify the presence of putative dechlorinating *Chloroflexi*, which phylum includes the only known anaerobic PCB degrader, in the soil samples. The clone library constructed from DNA extracted from switchgrass treated soil with redox cycling treatment shows that the sequences amplified in the qPCR targeting putative dechlorinating *Chloroflexi* were all *Dehalogenimonas* species (Table 6.3). Strains of *Dehalogenimonas* species are able to reductively dechlorinate polychlorinated aliphatic alkanes (Maness *et al.*, 2012; Yan *et al.*, 2009), and a recent study found that *Dehalogenimonas* can couple their growth with PCB dechlorination in a sediment free culture CG3 (Wang and He, 2013). Our sequences share 98-100% similarity with the *Dehalogenimonas* sequence retrieved from the culture CG3

(Figure 6.7). The presence of *Dehalogenimonas* in our soil samples strongly suggests that they may be responsible for the dechlorination in the rhizosphere.

Before the redox cycling treatment, the abundances of putative dechlorinating *Chloroflexi* 16S rRNA genes were low in both unplanted and switchgrass treated soils. After two weeks of flooding, putative dechlorinating *Chloroflexi* 16S rRNA gene abundances increased (Figure 6.8). However, despite of the repeated flooding and draining afterwards, the abundances of putative dechlorinating *Chloroflexi* 16S rRNA genes were stable till the end of the experiment ($2.91 \times 10^5 \pm 8.64 \times 10^4$ gene copy per g soil in unplanted soil with redox cycling, $5.63 \times 10^5 \pm 4.98 \times 10^5$ gene copy per g soil in switchgrass treated soil with redox cycling), and were not correlated to the redox potential or moisture content change (Pearson's r , $p > 0.05$). It appears that the first two weeks of flooding successfully created an environment suitable for the growth of possible PCB-dechlorinating *Chloroflexi* on available electron donors in the soil. In the subsequent non-flooding period, the putative dechlorinating *Chloroflexi* were able to survive in the unplanted soil and the switchgrass rhizosphere. Actually the abundance of putative dechlorinating *Chloroflexi* 16S rRNA genes in switchgrass treated soil was significantly higher than that in unplanted and redox cycled soils ($p < 0.05$). It is likely that anaerobic microenvironments existed in the rhizosphere, which could be where dechlorination took place. In the soils without redox recycling treatment, the putative dechlorinating *Chloroflexi* 16S rRNA genes were at or below the detection limit of qPCR (30 gene copies per reaction).

Despite the increase after the first flooding session, putative dechlorinating *Chloroflexi* 16S rRNA genes were still at low abundance with respect to the total bacterial 16S rRNA genes (0.0028 ± 0.0008 % and 0.0029 ± 0.0025 % of total 16S rRNA genes in unplanted soil with redox cycling and switchgrass treated soil with redox cycling, respectively). The abundance of dechlorinating bacteria in PCB-dechlorinating sediment free culture was higher. For example, *Dehalococcoides* was 3.74% of total bacterial

community in highly enriched sediment-free JN culture (Bedard *et al.*, 2006).

Dehalogenimonas was 2.16% in a sediment-free culture that dechlorinated Aroclor 1260 (Wang and He, 2013). The total bacterial abundance was not affected by redox cycling ($p>0.05$), while the switchgrass treatment significantly increased the total bacterial abundance ($p<0.05$) (Figure 6.9).

Microbial community analysis

To explore the effect of switchgrass and redox cycling treatment on microbial communities, the T-RFLP analysis was performed to rapidly assess microbial community structure during the incubation time. NMDS ordination of the T-RFLP profiles revealed that the blank soil (i.e. soil without PCB spiking), unplanted soil, switchgrass treated soil, unplanted soil with redox cycling, and switchgrass treated soil with redox cycling displayed different bacterial community structure. At the beginning of the experiment, unplanted soil and blank soil had similar microbial communities (Figure 6.10). The differences of T-RF compositions in soils with different treatments were confirmed by MRPP and PERMANOVA ($p=0.001$). No significant correlation was found between bacterial communities and the incubation time (Pearson's R, $p>0.05$). Indicator species analysis shows that some T-RFs were statistically associated with different treatments (Table 6.5). Those T-RFs were rare members of the microbial community in the soil samples ($< 1\%$ of total T-RFs), and were not identified by clone library analysis.

The microbial community structure in soils with different treatments at 24 weeks were assessed by sequencing in total 187 clones from four 16S rRNA gene clone libraries. All soil communities were dominated by *Proteobacteria* and *Acidobacteria* (comprising $46.9 \pm 13.5\%$ and $13.9 \pm 5.8\%$ of total clones in the four clone libraries) (Figure 6.11). The rest of phylotypes were affiliated to *Bacteroidetes*, *Actinobacteria*, *Chloroflexi*, *Chloroplast*, *Firmicutes*, *Gemmatimonadetes*, *Nitrospira*, *OP11*, *Planctomycetes* and *Verrucomicrobia*. Clone libraries in unplanted and switchgrass treated soils were similar,

while increased abundance of *Proteobacteria* and decreased *Acidobacteria* were observed in redox cycled soils.

Proteobacteria are commonly found in soils and sediments (de Cárcer et al., 2007; Petrić et al., 2011a; Spain et al., 2009), and many known aerobic PCB-degraders are *Proteobacteria* (e.g. *Pseudomonas*, *Sphingomonas*, *Acinetobacter*, *Comamonas* and *Burkholderia* (Nováková et al., 2002)), which may be involved in PCB degradation in our soil samples. Possible aerobic PCB degraders other than *Proteobacteria*, such as *Bacillus* and *Rhodococcus*, were also identified (Masai et al., 1995; Shimura et al., 1999). The high abundance of *Acidobacteria* was previously found in other PCB contaminated soils (Nogales et al., 1999; Petrić et al., 2011a), indicating their possible role in PCB biodegradation.

Within *Proteobacteria*, *Delta*- and *Gamma*-*proteobacteria* include some species that can degrade chlorinated compounds via dehalorespiration or co-metabolic reductive dehalogenation, such as *Desulfuromonas* spp (Krumholz, 1997), *Geobacter lovleyi* (Sung et al., 2006) and *Shewanella putrefaciens* (Picardal et al., 1995). *Geobacter*, known as sulfate/sulfur and iron reducers, were only detected in the clone library from the unplanted soil with redox cycling treatment. *Geobacter* is able to dechlorinate trichloroacetic acid and tetrachlorethene (de Wever et al., 2000; Strycharz et al., 2008; Sung et al., 2006), and have been observed in PCB dechlorinating enrichment cultures in previous studies (Bedard et al., 2006; Bedard et al., 2007; Holoman et al., 1998). The T-RFs corresponding to *Geobacter* spp. were 203bp and 215bp (Table 6.4). However the 203 bp T-RF is not specific to *Geobacter* sp. – it can also be generated from other *Deltaproteobacteri*, and the 215 bp T-RF could be produced from 16S sequences from other *Betaproteobacteria*, which may explain the relatively high abundances of the two T-RFs in soils with switchgrass treatment (Figure 6.13). Furthermore, T-RF 203 bp was identified as an the important T-RF in the classification of T-RFLP profiles by random

forest algorithm, as well as T-RF 235 bp, another T-RF corresponding to *Deltaproteobacteria* (Figure 6.12).

Clostridium is another genera identified in the clone libraries that has the potential to dechlorinate PCBs. For instance, some certain *Clostridium* strains can dechlorinate tetrachloroethylene (Chang et al., 2000; Okeke et al., 2001). *Clostridium* spp. were detected in a PCB-dechlorinating enrichment culture derived from St. Lawrence River sediment (Oh et al., 2008). In another study, *Clostridium* species were abundant in sediments that were experiencing *para*- and *meta*- PCB dechlorination (Hou and Dutta, 2000). *Clostridium* spp. can also produce hydrogen, an important and common electron donor for dechlorination (Bowman et al., 2009; Nandi and Sengupta, 1998). *Clostridium* clones generated 288 bp and 298 bp T-RFs, which comprised 0.25% and 0.54% of the total T-RF peak areas (Figure 6.13). *Geobacter* and *Clostridium* were not abundant in the four clone library, yet the presence of such dehalorespiring species in the soils experiencing PCB dechlorination suggests their possible role in PCB degradation.

Chloroflexi detected by clone library in switchgrass treated soils, unplanted soil with redox cycling, and switchgrass treated soil with redox cycling were identified to *Anaerolineae* and *Caldilineae*, which were not associated to dechlorination (Gupta et al., 2013). Putative dechlorinating *Chloroflexi* was not detected by clone library, probably because of their low abundance in the soil samples with respect to total bacteria, which was verified by the qPCR results.

Redox cycling had an effect on the microbial community structure in all, according to the T-RFLP analysis. In clone libraries, higher abundance of *Proteobacteria*, especially *Gammaproteobacteria*, was found in soils with redox cycling. Also, an increase of putative dechlorinating *Chloroflexi* 16S rRNA gene abundances in soils was observed after redox cycling treatment as shown by qPCR. The redox cycling was repeated for in total four times, but the T-RFLP profile didn't show much difference before and after the flooding session, as well as the abundances of putative dechlorinating

Chloroflexi 16S rRNA genes, suggesting that the redox cycling resulted in bacterial community change, but the bacterial community was resilient and stable despite of the sequential flooding and non-flooding cycles.

Conclusions

In this study, we investigated the microbial communities in the PCB spiked and switchgrass treated soils that experienced PCB dechlorination. The occurrence of dechlorination was verified by the detection of transformation products of parent PCB congeners in all soil microcosms, some of which were under anoxic conditions. Redox cycling was found to significantly enhance the removal of PCB 153 in unplanted soil. The first two weeks of flooding of the redox cycling treatment successfully promoted the putative dechlorinating *Chloroflexi* populations, which sequences were confirmed to be *Dehalogenimonas*, a potential PCB-dechlorinating species. Although no putative dechlorinating *Chloroflexi* was detected in soils without redox cycling, the identification of other dehalorespiring bacteria such as *Geobacter* and *Clostridium* suggests the dechlorination potential of the indigenous microbial community and their possible role in PCB dechlorination.

Table 6.1. Primer sets used for qPCR analysis in Chapter VI.

Target gene	Primer name	Sequences	Expected product size	Source
total bacterial 16S rRNA	16SU f	5'-TCCTACGGGAGGCAGCAGT-3'	466 bp	(Nadkarni <i>et al.</i> , 2002)
	16SU r	5'- GGACTACCAGGGTATCTAATCCTGTT- 3'		
<i>Dehalococcoides</i> - like 16S rRNA	dhc 793f	5'-GGGAGTATCGACCCTCTCTG-3'	193 bp	(Yoshida <i>et al.</i> , 2005)
	dhc 946r	5'-CGTTYCCCTTTCRGTTCACT-3'		

Table 6.2. Pertinent qPCR parameters (primer concentration, template mass, linear range, PCR efficiency, and y-intercept of the qPCR standard curve) in Chapter VI.

Target gene	primer concentration, μM	template	qPCR linear range, gene copies/reaction	qPCR efficiency	Y-intercept
total bacterial 16S rRNA	0.1	1 ng DNA	30 - 30×10^7	95 - 100%	36.11-38.38
<i>Dehalococcoides</i> -like 16S rRNA	0.8	10 ng DNA	30 - 30×10^7	95 - 100%	33.89-35.48

Table 6.3. RDP classification of clone libraries for quality assurance of qPCR primers used in Chapter VI.

Primer set	Target group	RDP classification (% certainty)	No. of clones detected	Sources
dhc1f/264r	<i>Dehalococcoides</i> spp.	<i>Rubrobacteridae</i> (96-100)	3/9	(Grostern and Edwards, 2006)
		TM7 (93)	1/9	
		unclassified Bacteria (98-100)	5/9	
chl384f/dehal884r	<i>Chloroflexi</i>	<i>Dehalogenimonas</i> (84-100)	5/13	(Fagervold <i>et al.</i> , 2005)
		<i>Rubrobacter</i> (100)	5/13	
		unclassified Bacteria (89-100)	3/13	
dhc793f/946r	<i>Dehalococcoides</i> spp.	<i>Dehalogenimonas</i> (99-100)	12/12	(Yoshida <i>et al.</i> , 2005)
		<i>Anaerolineaceae</i> (84-89)	3/12	
dhc1154f/1286r	<i>Dehalococcoides</i> -like <i>Chloroflexi</i>	unclassified <i>Chloroflexi</i> (87-92)	5/12	(Krzmarzick <i>et al.</i> , 2012)
		unclassified Bacteria (86-100)	3/12	
		unclassified	1/12	

Table 6.4. 16S rRNA gene clones recovered from unplanted soil (UP), switchgrass treated soil (SG), unplanted soil with redox cycling (UPF), switchgrass treated soil with redox cycling (SG F). Clone identification was made by RDP classifier.

Observed T-RF (bp)	Predicted T-RF (bp)	Closest classified relative (% certainty)	NMDS axis 1	NMDS axis 2
71	75	<i>Sorangium</i> (95%)	-0.010	-0.165
187	194	unclassified <i>Betaproteobacteria</i> (100%)	-0.156	0.065
195	196	<i>Luteolibacter</i> (100%)	-0.045	-0.331
197	200	<i>Variovorax</i> (100%)	0.267	-0.229
196	200	unclassified <i>Betaproteobacteria</i> (96%)	-0.004	0.307
203	204	<i>Geobacter</i> (100%)	0.295	-0.429
203	204	unclassified <i>Desulfuromonadales</i> (100%)	0.295	-0.429
204	206	unclassified <i>Xanthomonadaceae</i> (97%)	-0.285	0.202
205	206	<i>Rhodanobacter</i> (100%)	-0.198	-0.014
210	212	unclassified <i>Betaproteobacteria</i> (87%)	0.567	-0.109
210	214	<i>Dokdonella</i> (94%)	0.567	-0.109
213	214	<i>Caldilinea</i> (100%)	0.561	-0.168
213	216	<i>Gp3</i> (100%)	0.561	-0.168
215	217	<i>Geobacter</i> (100%)	1.389	0.242
216	217	unclassified <i>Comamonadaceae</i> (100%)	0.154	-0.056
216	219	unclassified <i>Comamonadaceae</i> (100%)	0.154	-0.056
225	227	<i>Novosphingobium</i> (95%)	-0.176	0.230
228	230	<i>Gracilibacter</i> (100%)	-0.109	0.178
229	231	<i>Bacillus</i> (100%)	-0.045	-0.114
232	234	<i>Gp6</i> (100%)	-0.286	0.046
235	237	unclassified <i>Deltaproteobacteria</i> (97%)	-0.298	-0.129
248	250	unclassified <i>Ruminococcaceae</i> (100%)	0.307	-0.187
260	262	<i>Flavisolibacter</i> (99%)	0.085	0.047
279	281	<i>Planctomyces</i> (98%)	-0.218	-0.048
288	290	<i>Clostridium XI</i> (100%)	0.433	-0.141
298	300	<i>Clostridium sensu stricto</i> (100%)	-0.142	-0.028
314	316	unclassified <i>Anaerolineaceae</i> (100%)	0.247	-0.051
328	331	<i>Nitrospira</i> (100%)	-0.348	-0.298

Table 6.5. Significant indicator T-RFs determined by indicator species analysis.

	Indicator T-RF	Statistic	p value
UP	383 bp	0.722	0.0002
	286 bp	0.663	0.0008
	385 bp	0.632	0.0006
	384 bp	0.594	0.0004
UPF	128 bp	0.84	0.0002
	245 bp	0.836	0.0002
	166 bp	0.73	0.0002
	396 bp	0.667	0.0002
	339 bp	0.663	0.0004
	349 bp	0.594	0.0006
SGF	108 bp	0.851	0.0002
	406 bp	0.804	0.0002
	215 bp	0.793	0.001
	60 bp	0.706	0.0002
	514 bp	0.606	0.0008
	464 bp	0.557	0.001
UPF+SGF	263 bp	0.817	0.0002
	231 bp	0.791	0.0002
	502 bp	0.778	0.0002
SG+SGF	258 bp	0.98	0.0002
	265 bp	0.966	0.0002
	198 bp	0.918	0.0002
	209 bp	0.909	0.0002




Figure 6.1. The percentage of initial concentration of PCB 52, 77 and 153 after 12 and 24 weeks of incubation in unplanted soil (UP), switchgrass treated soil (SG), unplanted soil with redox cycling (UPF), switchgrass treated soil with redox cycling (SG F). Error bars indicate the standard deviation of three soil subsamples from the same reactor.




Figure 6.2. Transformation products detected after 12 and 24 weeks of incubation in unplanted soil (UP), switchgrass treated soil (SG), unplanted soil with redox cycling (UPF), switchgrass treated soil with redox cycling (SG F). Error bars indicate the standard deviation of three soil subsamples from the same reactor.




Figure 6.3. Scheme of the possible dechlorination pathways for PCB 52, 77, and 153. Dotted arrows indicate possible pathways but no intermediate was observed.




Figure 6.4. Redox potential change in unplanted soil (UP), switchgrass treated soil (SG), unplanted soil with redox cycling (UPF), switchgrass treated soil with redox cycling (SG F).




Figure 6.5. Moisture content change in unplanted soil (UP), switchgrass treated soil (SG), unplanted soil with redox cycling (UPF), switchgrass treated soil with redox cycling (SG F).




Figure 6.6. Percentage of total PCB mass in soil, switchgrass roots, and leaves after 24 week incubation in switchgrass treated soil (SG) and switchgrass treated soil with redox cycling (SG F). Error bars indicate the standard deviation of three soil subsamples from the same reactor.




Figure 6.7. Phylogenetic analysis of 16S rRNA genes retrieved from switchgrass treated soils with redox cycling after 24 weeks. The alignment was made with ClustalX and converted into a neighboring-joining tree, which was visualized with MEGA4 with *Oscillochloris trichoides* as an outgroup. The filled circles at nodes indicate bootstrap values of 95%, while open circles indicate bootstrap values of 75~95%. Bootstrap values lower than 75% are not shown. The bar represents 0.5% sequence difference.




Figure 6.8. qPCR analysis of *Dehalococcoides*-like 16S rRNA gene in unplanted soil with redox cycling (UPF) and switchgrass treated soil with redox cycling (SG F). Error bars indicate the range of two soil subsamples from the same reactor.




Figure 6.9. qPCR analysis of bacterial 16S rRNA gene in unplanted soil (UP), switchgrass treated soil (SG), unplanted soil with redox cycling (UPF), switchgrass treated soil with redox cycling (SG F). Error bars indicate the range of two soil subsamples from the same reactor.




Figure 6.10. NMDS ordination of T-RFLP profiles from blank soil without PCB spiking (BLK), unplanted soil (UP), switchgrass treated soil (SG), unplanted soil with redox cycling (UPF), switchgrass treated soil with redox cycling (SG F).




Figure 6.11. Clone library analysis of bacterial 16S rRNA gene in unplanted soil (UP), switchgrass treated soil (SG), unplanted soil with redox cycling (UPF), switchgrass treated soil with redox cycling (SG F). Error bars indicate the range of two soil subsamples from the same reactor.




Figure 6.12. T-RF importance for classification as determined by random forest algorithm.




Figure 6.13. The abundance of T-RFs that were identified by random forest algorithm or corresponded to dehalorespiring bacteria in unplanted soil (UP), switchgrass treated soil (SG), unplanted soil with redox cycling (UPF), switchgrass treated soil with redox cycling (SG F).

CHAPTER VII. ENGINEERING AND SCIENTIFIC SIGNIFICANCE

This work contributes to the understanding of microbial PCB degradation as a remediation strategy. The research objectives included evaluating the *in situ* microbial PCB degradation potential at a PCB contaminated site, improving PCB removal by manipulating switchgrass rhizosphere, and investigating the microbial community variation during the remediation processes.

The selection and optimization of bioremediation strategies for PCB contaminated site clean-up requires a comprehensive understanding about the *in situ* PCB degradation potential. In Chapter III and IV, the aerobic and anaerobic microbial PCB degradation potential was evaluated for two PCB contaminated sediment cores from IHSC (core 1 and core 2) by the combination use of PCB congener analysis and molecular analysis of microbial communities along sediment depth. The correlation between PCB congener profiles and microbial communities was explored, in an attempt to identify the key members that might be involved in the PCB degradation in the deep sediments.

Core 2 exhibited a higher PCB dechlorination potential than core 1, as suggested by congener analysis. The increasing MDPR along core 2 sediment depth suggested a higher degree of dechlorination in deep sediments. The microbial PCB degradation potential was estimated by quantifying biomarker genes along sediment depth, including *bphA*, the gene indicative of aerobic PCB degradation, and the putative dechlorinating *Chloroflexi* 16S rRNA genes. Both *bphA* and putative dechlorinating *Chloroflexi* 16S rRNA genes were more abundant in upper 1.83 m sediments, suggesting the occurrence of both aerobic and anaerobic PCB degradation in the upper sediments. The low abundance of *bphA* in deep sediments may be the result of the stricter anaerobic environment. The discrepancy between PCB congener data and putative dechlorinating *Chloroflexi* 16S rRNA gene data revealed that targeting solely the *Chloroflexi* may be not enough for the evaluation of anaerobic PCB degradation potential. Other dehalorespiring

microorganisms may also be involved in the PCB dechlorination process at IHSC. To identify possible dechlorinating bacteria, microbial communities were characterized by T-RFLP and clone library. *Proteobacteria*, which include many known aerobic PCB degraders, were found dominant. Especially, *Acidovorax* and *Acinetobacter* were abundant. And the abundance of *Acidovorax* was correlated with sediment and pore water MDP values.

The results from the studies described in Chapter III and IV confirmed the potential of PCB degradation at the contaminated sediments at IHSC, suggesting that the indigenous microbial communities possessed the capability to degrade PCBs in sediments, and that monitored natural attenuation could be a feasible remediation strategy. Another question is how to enhance microbial PCB degradation during the bioremediation process. In Chapters V and VI, phytoremediation with switchgrass was performed on soil microcosms with PCB spiking (PCB 52, 77 and 153). To further enhance the aerobic PCB degradation, bioaugmentation with *Burkholderia xenovorans* LB400 was performed, and to improve the anaerobic PCB degradation, soil microcosms were subjected to redox cycling (two weeks of flooding and two weeks of non-flooding).

Chapter V described the results of phytoremediation and bioaugmentation. The presence of switchgrass significantly enhanced the PCB removal in the soil after 24 weeks. The improved PCB removal was found to be the result of phytoextraction and enhanced microbial activity. The microbial PCB degradation was not improved as much as expected, which was also indicated by RT-qPCR of *bphA*. The reason may be the availability of other carbon sources in the rhizosphere. It is also noted that microbial activity had different effect on different PCB congeners. PCB 52 is more susceptible to aerobic degradation by indigenous microorganisms, despite of its *ortho* chlorine substituents. After bioaugmentation, PCB 52 removal was improved the most, while the removal of PCB 77 and 153 was not significantly improved. The active aerobic microbial PCB degradation was also confirmed by the successful detection of *bphA* transcripts. The

presence of switchgrass was found to benefit LB400 survival and its ability to degrade PCBs. Thus the combination use of phytoremediation and bioaugmentation was recommended.

While Chapter V shows aerobic PCB degradation can be enhanced by bioaugmentation with LB400, in Chapter VI, an alternating anaerobic-aerobic condition was created by redox cycling, in an attempt to promote the anaerobic PCB degradation. The removal of PCB 153 was significantly improved in redox cycled and unplanted soils, as compared with unplanted soil, suggesting the enhanced anaerobic PCB degradation. The first two weeks of flooding of the redox cycling treatment successfully promoted the putative dechlorinating *Chloroflexi* population, which may be associated with the improved PCB 153 removal. However, the transformation products were detected in all soils. Especially, *ortho* dechlorination product from PCB 153 was observed. Then the microbial community analysis was performed to search for the presence of taxa that contain a dechlorination potential in the rhizosphere. *Proteobacteria* was again found to be the most dominant, followed by *Acidobacteria*. *Geobacter*, known as iron reducers, and *Clostridium*, which was able to dechlorinate other chlorinated compounds, were detected, indicating their possible role in the PCB dechlorination.

One innovation of this work is assessing the *in situ* microbial PCB degradation by congener analysis in conjunction with culture-independent microbial analysis for core sediment samples. The depth pattern of congener data can provide evidence that anaerobic dechlorination has occurred, while the microbial analysis targeting biomarker genes shows the potential for both aerobic and anaerobic PCB degradation by indigenous microbial communities. The obtained information is valuable to decide whether microbial PCB degradation (natural attenuation) is feasible for the specific contaminated site.

Another innovation of this work is the detection of dehalorespiring bacteria in the sediments and soils undergoing PCB dechlorination. Currently most microbial PCB dechlorination research focuses on *Chloroflexi*, which include the only confirmed PCB

dechlorinators (*Dehalococcoides* spp. and o-17/DF1). Although the capability of dechlorinating PCBs is unknown for many other dehalorespiring bacteria, the lack of analysis of these bacteria can lead to incomplete understanding of the PCB degradation potential in contaminated sediments and soils, and a biased selection of remediation strategies. In this research, the presence of *Acinetobacter*, *Geobacter*, *Clostridium* and *Dehalogenimonas* in the contaminated sediments and soils may suggest their potential in PCB dechlorination. *Dehalogenimonas* was found present in PCB contaminated sediments from IHSC and in the soil microcosms under laboratory condition, making them a promising candidate for PCB dechlorination. Especially, flooding treatment was found to be able to increase *Dehalogenimonas* population, suggesting that flooding was effective in promoting the dechlorinating group. Effort was also made to identify the taxa that might be important to PCB biodegradation by linking microbial data and congener data. For example, the abundance of *Acidovorax* was found to be correlated with an indicator of PCB dechlorination. These observations provide the starting point for future studies about microbial PCB dechlorination. It is hoped that the use of high-throughput sequencing technologies could provide more insight into the microbial communities involved in the dechlorination process.

REFERENCES

- Abraham, W.-R., Nogales, B., Golyshin, P.N., Pieper, D.H., Timmis, K.N., 2002. Polychlorinated biphenyl-degrading microbial communities in soils and sediments. *Curr Opin Microbiol* 5, 246-253.
- Adriaens, P., Focht, D.D., 1991. Cometabolism of 3,4-dichlorobenzoate by *Acinetobacter* sp. strain 4-CB1. *Appl Environ Microbiol* 57, 173-179.
- Adrian, L., Dudková, V., Demnerová, K., Bedard, D.L., 2009. "*Dehalococcoides*" sp. strain CBDB1 extensively dechlorinates the commercial polychlorinated biphenyl mixture Aroclor 1260. *Appl Environ Microbiol* 75, 4516-4524.
- Agarwal, S., Al-Abed, S.R., Dionysiou, D.D., 2007. *In situ* technologies for reclamation of PCB-Contaminated sediments: Current challenges and research thrust areas. *J Environ Eng* 133, 1075-1078.
- Alder, A.C., Hägglom, M.M., Oppenheimer, S.R., Young, L.Y., 1993. Reductive dechlorination of polychlorinated biphenyls in anaerobic sediments. *Environ Sci Technol* 27, 530-538.
- Altschul, S.F., Madden, T.L., Schäffer, A.A., Zhang, J., Zhang, Z., Miller, W., Lipman, D.J., 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* 25, 3389-3402.
- Amend, L.J., Lederman, P.B., 1992. Critical evaluation of PCB remediation technologies. *Environ Prog* 11, 173-177.
- ATSDR, 1994. CERCLA Priority list of hazardous substances that will be the subject of toxicological profiles (support document). In: services, U.d.o.h.a.h. (Ed.), Atlanta, GA.
- Bedard, D.L., 2004. Polychlorinated biphenyls in aquatic sediments: environmental fate and outlook for biological treatment, Dehalogenation. Springer US, pp. 443-465.
- Bedard, D.L., Bailey, J.J., Reiss, B.L., Jerzak, G.V., 2006. Development and characterization of stable sediment-free anaerobic bacterial enrichment cultures that dechlorinate Aroclor 1260. *Appl Environ Microbiol* 72, 2460-2470.
- Bedard, D.L., Bunnell, S.C., Smullen, L.A., 1996. Stimulation of microbial *para*-dechlorination of polychlorinated biphenyls that have persisted in Housatonic River sediment for decades. *Environ Sci Technol* 30, 687-694.
- Bedard, D.L., Ritalahti, K.A., Löffler, F.E., 2007. The *Dehalococcoides* population in sediment-free mixed cultures metabolically dechlorinates the commercial polychlorinated biphenyl mixture Aroclor 1260. *Appl Environ Microbiol* 73, 2513-2521.
- Bedard, D.L., Unterman, R., Bopp, L.H., Brennan, M.J., Haberl, M.L., Johnson, C., 1986. Rapid assay for screening and characterizing microorganisms for the ability to degrade polychlorinated biphenyls. *Appl Environ Microbiol* 51, 761-768.
- Bedard, D.L., Van Dort, H.M., May, R.J., Smullen, L.A., 1997. Enrichment of microorganisms that sequentially *meta*, *para*-dechlorinate the residue of Aroclor 1260 in Housatonic River sediment. *Environ Sci Technol* 31, 3308-3313.

Berkaw, M., Sowers, K.R., May, H.D., 1996. Anaerobic *ortho* dechlorination of polychlorinated biphenyls by estuarine sediments from Baltimore Harbor. *Appl Environ Microbiol* 62, 2534-2539.

Bibby, K., Viau, E., Peccia, J., 2010. Pyrosequencing of the 16S rRNA gene to reveal bacterial pathogen diversity in biosolids. *Water Res* 44, 4252-4260.

Bopp, L.H., 1986. Degradation of highly chlorinated PCBs by *Pseudomonas* strain LB400. *J Ind Microbiol Biot* 1, 23-29.

Borja, J., Taleon, D.M., Auresenia, J., Gallardo, S., 2005. Polychlorinated biphenyls and their biodegradation. *Process Biochem* 40, 1999-2013.

Bouchez, T., Patureau, D., Dabert, P., Juretschko, S., Dore, J., Delgenes, P., Moletta, R., Wagner, M., 2000. Ecological study of a bioaugmentation failure. *Environ Microbiol* 2, 179-190.

Bowman, K.S., Rainey, F.A., Moe, W.M., 2009. Production of hydrogen by *Clostridium* species in the presence of chlorinated solvents. *FEMS Microbiol Lett* 290, 188-194.

Breivik, K., Sweetman, A., Pacyna, J.M., Jones, K.C., 2002a. Towards a global historical emission inventory for selected PCB congeners - a mass balance approach 1. global production and consumption. *Sci Total Environ* 290, 181-198.

Breivik, K., Sweetman, A., Pacyna, J.M., Jones, K.C., 2002b. Towards a global historical emission inventory for selected PCB congeners - a mass balance approach 2. emissions. *Sci Total Environ* 290, 199-224.

Breivik, K., Sweetman, A., Pacyna, J.M., Jones, K.C., 2007. Towards a global historical emission inventory for selected PCB congeners - a mass balance approach-3. an update. *Sci Total Environ* 377, 296-307.

Breuker, A., Köweker, G., Blazejak, A., Schippers, A., 2011. The deep biosphere in terrestrial sediments in the Chesapeake Bay area, Virginia, USA. *Front Microbiol* 2, 156.

Bustin, S.A., Benes, V., Garson, J.A., Hellems, J., Huggett, J., Kubista, M., Mueller, R., Nolan, T., Pfaffl, M.W., Shipley, G.L., Vandesompele, J., Wittwer, C.T., 2009. The MIQE guidelines: minimum information for publication of quantitative real-time PCR experiments. *Clin Chem* 55, 611-622.

Byrne-Bailey, K.G., Weber, K.A., Chair, A.H., Bose, S., Knox, T., Spanbauer, T.L., Chertkov, O., Coates, J.D., 2010. Completed genome sequence of the anaerobic iron-oxidizing bacterium *Acidovorax ebreus* strain TPSY. *J Bacteriol* 192, 1475-1476.

Cáceres, M.D., Legendre, P., 2009. Associations between species and groups of sites: indices and statistical inference. *Ecology* 90, 3566-3574.

Camara, B., Herrera, C., Gonzalez, M., Couve, E., Hofer, B., Seeger, M., 2004. From PCBs to highly toxic metabolites by the biphenyl pathway. *Environ Microbiol* 6, 842-850.

Campanella, B.F., Bock, C., Schroder, P., 2002. Phytoremediation to increase the degradation of PCBs and PCDD/Fs - Potential and limitations. *Environ Sci Pollut R* 9, 73-85.

- Campfens, J., MacKay, D., 1997. Fugacity-based model of PCB bioaccumulation in complex aquatic food webs. *Environ Sci Technol* 31, 577-583.
- Cardenas, E., Cole, J.R., Tiedje, J.M., Park, J., 2009. Microbial community analysis using RDP 2 (Ribosomal Database Project 2): methods, tools and new advances. *Environ Eng Res* 14, 3-9.
- Chang, Y.C., Hatsu, M., Jung, K., Yoo, Y.S., Takamizawa, K., 2000. Isolation and characterization of a tetrachloroethylene dechlorinating bacterium, *Clostridium bifermentans* DPH-1. *J Biosci Bioeng* 89, 489-491.
- Chekol, T., Vough, L.R., Chaney, R.L., 2004. Phytoremediation of polychlorinated biphenyl-contaminated soils: the rhizosphere effect. *Environ Int* 30, 799-804.
- Cole, J.R., Chai, B., Farris, R.J., Wang, Q., Kulam-Syed-Mohideen, A.S., McGarrell, D.M., Bandela, A.M., Cardenas, E., Garrity, G.M., Tiedje, J.M., 2007. The ribosomal database project (RDP-II): introducing myRDP space and quality controlled public data. *Nucleic Acids Res* 35, 169-172.
- Copley, S.D., Crooks, G.P., 1992. Enzymic dehalogenation of 4-chlorobenzoyl coenzyme A in *Acinetobacter* sp. strain 4-CB1. *Appl Environ Microbiol* 58, 1385-1387.
- Correa, P.A., Lin, L., Just, C.L., Hu, D., Hornbuckle, K.C., Schnoor, J.L., Van Aken, B., 2010. The effects of individual PCB congeners on the soil bacterial community structure and the abundance of biphenyl dioxygenase genes. *Environ Int* 36, 901-906.
- Culman, S.W., Bukowski, R., Gauch, H.G., Cadillo-Quiroz, H., Buckley, D.H., 2009. T-REX: software for the processing and analysis of T-RFLP data. *BMC Bioinformatics* 10, 171.
- Cutler, D.R., Edwards, T.C., Beard, K.H., Cutler, A., Hess, K.T., 2007. Random forests for classification in ecology. *Ecology* 88, 2783-2792.
- Cutter, L.A., Watts, J.E.M., Sowers, K.R., May, H.D., 2001. Identification of a microorganism that links its growth to the reductive dechlorination of 2,3,5,6-chlorobiphenyl. *Environ Microbiol* 3, 699-709.
- da C Jesus, E., Marsh, T.L., Tiedje, J.M., de S Moreira, F.M., 2009. Changes in land use alter the structure of bacterial communities in Western Amazon soils. *ISME J* 3, 1004-1011.
- Dai, S., Vaillancourt, F.H., Maaroufi, H., Drouin, N.M., Neau, D.B., Snieckus, V., Bolin, J.T., Eltis, L.D., 2002. Identification and analysis of a bottleneck in PCB biodegradation. *Nat Struct Biol* 9, 934-939.
- de Cárcer, D.A., Martín, M., Karlson, U., Rivilla, R., 2007. Changes in bacterial populations and in biphenyl dioxygenase gene diversity in a polychlorinated biphenyl-polluted soil after introduction of willow trees for rhizoremediation. *Appl Environ Microbiol* 73, 6224-6232.
- de Wever, H., Cole, J.R., Fettig, M.R., Hogan, D.A., Tiedje, J.M., 2000. Reductive dehalogenation of trichloroacetic acid by *Trichlorobacter thiogenes* gen. nov., sp. nov. *Appl Environ Microbiol* 66, 2297-2301.

- Di Gregorio, S., Azaizeh, H., Lorenzi, R., 2013. Biostimulation of the autochthonous microbial community for the depletion of polychlorinated biphenyls (PCBs) in contaminated sediments. *Environ Sci Pollut R* 20, 3989-3999.
- Dietz, A.C., Schnoor, J.L., 2001. Advances in phytoremediation. *Environ Health Persp* 109, 163-168.
- Ding, N., Guo, H.C., Hayat, T., Wu, Y.P., Xu, J.M., 2009. Microbial community structure changes during Aroclor 1242 degradation in the rhizosphere of ryegrass (*Lolium multiflorum* L.). *FEMS Microbiol Ecol* 70, 305-314.
- Donnelly, P.K., Hegde, R.S., Fletcher, J.S., 1994. Growth of PCB-degrading bacteria on compounds from photosynthetic plants. *Chemosphere* 28, 981-988.
- Dudášová, H., Lukáčová, L., Murínová, S., Dercová, K., 2012. Effects of plant terpenes on biodegradation of polychlorinated biphenyls (PCBs). *Int Biodeter Biodegr* 69, 23-27.
- Dzantor, E.K., Chekol, T., Vough, L.R., 2000. Feasibility of using forage grasses and legumes for phytoremediation of organic pollutants. *J Environ Sci Heal A* 35, 1645-1661.
- Egorova, D.O., Demakov, V.A., Plotnikova, E.G., 2013. Bioaugmentation of a polychlorobiphenyl contaminated soil with two aerobic bacterial strains. *J Hazard Mater* 261, 378-386.
- Erickson, M.D., Kaley, R.G., 2011. Applications of polychlorinated biphenyls. *Environ Sci Pollut R* 18, 135-151.
- Fagervold, S.K., May, H.D., Sowers, K.R., 2007. Microbial reductive dechlorination of Aroclor 1260 in Baltimore harbor sediment microcosms is catalyzed by three phylotypes within the phylum *Chloroflexi*. *Appl Environ Microbiol* 73, 3009-3018.
- Fagervold, S.K., Watts, J.E., May, H.D., Sowers, K.R., 2005. Sequential reductive dechlorination of *meta*-chlorinated polychlorinated biphenyl congeners in sediment microcosms by two different *Chloroflexi* phylotypes. *Appl Environ Microbiol* 71, 8085-8090.
- Fava, F., Gentilucci, S., Zanaroli, G., 2003. Anaerobic biodegradation of weathered polychlorinated biphenyls (PCBs) in contaminated sediments of Porto Marghera (Venice Lagoon, Italy). *Chemosphere* 53, 101-109.
- Fennell, D.E., Nijenhuis, I., Wilson, S.F., Zinder, S.H., Häggblom, M.M., 2004. *Dehalococcoides ethenogenes* strain 195 reductively dechlorinates diverse chlorinated aromatic pollutants. *Environ Sci Technol* 38, 2075-2081.
- Ficko, S.A., Rutter, A., Zeeb, B.A., 2010. Potential for phytoextraction of PCBs from contaminated soils using weeds. *Sci Total Environ* 408, 3469-3476.
- Field, J.A., Sierra-Alvarez, R., 2008. Microbial transformation and degradation of polychlorinated biphenyls. *Environ Pollut* 155, 1-12.
- Focht, D.D., 1995. Strategies for the improvement of aerobic metabolism of polychlorinated-biphenyls. *Curr Opin Biotechnol* 6, 341-346.

- Frame, G.M., Cochran, J.W., Bøwadt, S.S., 1996. Complete PCB congener distributions for 17 Aroclor mixtures determined by 3 HRGC systems optimized for comprehensive, quantitative, congener-specific analysis. *J High Res Chrom* 19, 657-668.
- Francova, K., Macková, M., Macek, T., Sylvestre, M., 2004. Ability of bacterial biphenyl dioxygenases from *Burkholderia* sp. LB400 and *Comamonas testosteroni* B-356 to catalyse oxygenation of *ortho*-hydroxychlorobiphenyls formed from PCBs by plants. *Environ Pollut* 127, 41-48.
- Fung, J.M., Morris, R.M., Adrian, L., Zinder, S.H., 2007. Expression of reductive dehalogenase genes in *Dehalococcoides ethenogenes* strain 195 growing on tetrachloroethene, trichloroethene, or 2,3-dichlorophenol. *Appl Environ Microbiol* 73, 4439-4445.
- Furukawa, K., 1994. Molecular genetics and evolutionary relationship of PCB-degrading bacteria. *Biodegradation* 5, 289-300.
- Furukawa, K., 2000. Biochemical and genetic bases of microbial degradation of polychlorinated biphenyls (PCBs). *J Gen Appl Microbiol* 46, 283-296.
- Furukawa, K., Fujihara, H., 2008. Microbial degradation of polychlorinated biphenyls: biochemical and molecular features. *J Biosci Bioeng* 105, 433-449.
- Gibson, D.T., Cruden, D.L., Haddock, J.D., Zylstra, G.J., Brand, J.M., 1993. Oxidation of polychlorinated-biphenyls by *Pseudomonas* sp strain LB400 and *Pseudomonas Pseudoalcaligenes* KF707. *J Bacteriol* 175, 6735-6735.
- Gibson, D.T., Parales, R.E., 2000. Aromatic hydrocarbon dioxygenases in environmental biotechnology. *Curr Opin Biotechnol* 11, 236-243.
- Gilbert, E.S., Crowley, D.E., 1997. Plant compounds that induce polychlorinated biphenyl biodegradation by *Arthrobacter* sp. strain B1B. *Appl Environ Microbiol* 63, 1933-1938.
- Gomes, H.I., Dias-Ferreira, C., Ribeiro, A.B., 2013. Overview of *in situ* and *ex situ* remediation technologies for PCB-contaminated soils and sediments and obstacles for full-scale application. *Sci Total Environ* 445-446, 237-260.
- González, J.M., Simó, R., Massana, R., Covert, J.S., Casamayor, E.O., Pedrós-Alió, C., Moran, M.A., 2000. Bacterial community structure associated with a dimethylsulfoniopropionate-producing North Atlantic algal bloom. *Appl Environ Microbiol* 66, 4237-4246.
- Grabowski, A., Nercessian, O., Fayolle, F., Blanchet, D., Jeanthon, C., 2005. Microbial diversity in production waters of a low-temperature biodegraded oil reservoir. *FEMS Microbiol Ecol* 54, 427-443.
- Greenwood, S.J., Rutter, A., Zeeb, B.A., 2011. The absorption and translocation of polychlorinated biphenyl congeners by *Cucurbita pepo* ssp *pepo*. *Environ Sci Technol* 45, 6511-6516.
- Grimm, F.A., Lehmler, H.J., He, X., Robertson, L.W., Duffel, M.W., 2013. Sulfated metabolites of polychlorinated biphenyls are high-affinity ligands for the thyroid hormone transport protein transthyretin. *Environ Health Perspect* 121, 657-662.

- Grosterm, A., Edwards, E.A., 2006. Growth of *Dehalobacter* and *Dehalococcoides* spp. during degradation of chlorinated ethanes. *Appl Environ Microbiol* 72, 428-436.
- Gupta, R.S., Chander, P., George, S., 2013. Phylogenetic framework and molecular signatures for the class *Chloroflexi* and its different clades; proposal for division of the class *Chloroflexi* class. nov into the suborder *Chloroflexineae* subord. nov., consisting of the emended family *Oscillochloridaceae* and the family *Chloroflexaceae* fam. nov., and the suborder *Roseiflexineae* subord. nov., containing the family *Roseiflexaceae* fam. nov. *Anton Leeuw Int J G* 103, 99-119.
- Hölscher, T., Krajmalnik-Brown, R., Ritalahti, K.M., von Wintzingerode, F., Görisch, H., Löffler, F.E., Adrian, L., 2004. Multiple nonidentical reductive-dehalogenase-homologous genes are common in *Dehalococcoides*. *Appl Environ Microbiol* 70, 5290-5297.
- Hendrickson, E.R., Payne, J.A., Young, R.M., Starr, M.G., Perry, M.P., Fahnestock, S., Ellis, D.E., Ebersole, R.C., 2002. Molecular analysis of *Dehalococcoides* 16S ribosomal DNA from chloroethene-contaminated sites throughout North America and Europe. *Appl Environ Microbiol* 68, 485-495.
- Hernandez, B., Arensdorf, J., Focht, D., 1995. Catabolic characteristics of biphenyl-utilizing isolates which cometabolize PCBs. *Biodegradation* 6, 75-82.
- Hernandez, B.S., Koh, S.C., Chial, M., Focht, D.D., 1997. Terpene-utilizing isolates and their relevance to enhanced biotransformation of polychlorinated biphenyls in soil. *Biodegradation* 8, 153-158.
- Herrick, R.F., Lefkowitz, D.J., Weymouth, G.A., 2007. Soil contamination from PCB-containing buildings. *Environ Health Persp* 115, 173-175.
- Hiraishi, A., 2008. Biodiversity of dehalorespiring bacteria with special emphasis on polychlorinated biphenyl/dioxin dechlorinators. *Microbes Environ* 23, 1-12.
- Hohmann, C., Winkler, E., Morin, G., Kappler, A., 2010. Anaerobic Fe(II)-oxidizing bacteria show As resistance and immobilize As during Fe(III) mineral precipitation. *Environ Sci Technol* 44, 94-101.
- Holliger, C., Wohlfarth, G., Diekert, G., 1998. Reductive dechlorination in the energy metabolism of anaerobic bacteria. *Fems Microbiol Rev* 22, 383-398.
- Holoman, T.R., Elberson, M.A., Cutter, L.A., May, H.D., Sowers, K.R., 1998. Characterization of a defined 2,3,5, 6-tetrachlorobiphenyl-ortho-dechlorinating microbial community by comparative sequence analysis of genes coding for 16S rRNA. *Appl Environ Microbiol* 64, 3359-3367.
- Hongsawat, P., Vangnai, A.S., 2011. Biodegradation pathways of chloroanilines by *Acinetobacter baylyi* strain GFJ2. *J Hazard Mater* 186, 1300-1307.
- Hoostal, M.J., Bullerjahn, G.S., McKay, R.M.L., 2002. Molecular assessment of the potential for *in situ* bioremediation of PCBs from aquatic sediments. *Hydrobiologia* 469, 59-65.

- Hou, L.H., Dutta, S.K., 2000. Phylogenetic characterization of several *para*- and *meta*-PCB dechlorinating *Clostridium* species: 16s rDNA sequence analyses. *Lett Appl Microbiol* 30, 238-243.
- Hu, D.F., Hornbuckle, K.C., 2010. Inadvertent polychlorinated biphenyls in commercial paint pigments. *Environ Sci Technol* 44, 2822-2827.
- Hu, D.F., Lehmler, H.J., Martinez, A., Wang, K., Hornbuckle, K.C., 2010. Atmospheric PCB congeners across Chicago. *Atmos Environ* 44, 1550-1557.
- Huang, Q.D., Hong, C.S., 2002. Aqueous solubilities of non-*ortho* and *mono-ortho* PCBs at four temperatures. *Water Res* 36, 3543-3552.
- Huesemann, M.H., Hausmann, T.S., Fortman, T.J., Thom, R.M., Cullinan, V., 2009. *In situ* phytoremediation of PAH- and PCB-contaminated marine sediments with eelgrass (*Zostera marina*). *Ecol Eng* 35, 1395-1404.
- Hung, H., Kallenborn, R., Breivik, K., Su, Y.S., Brorstrom-Lunden, E., Olafsdottir, K., Thorlacius, J.M., Leppanen, S., Bossi, R., Skov, H., Mano, S., Patton, G.W., Stern, G., Sverko, E., Fellin, P., 2010. Atmospheric monitoring of organic pollutants in the Arctic under the Arctic Monitoring and Assessment Programme (AMAP): 1993-2006. *Sci Total Environ* 408, 2854-2873.
- Huwe, J., Pagan-Rodriguez, D., Abdelmajid, N., Clinch, N., Gordon, D., Holterman, J., Zaki, E., Lorentzen, M., Dearfield, K., 2009. Survey of polychlorinated dibenzo-p-dioxins, polychlorinated dibenzofurans, and non-*ortho*-polychlorinated biphenyls in US meat and poultry, 2007-2008: effect of new toxic equivalency factors on toxic equivalency levels, patterns, and temporal trends. *J Agr Food Chem* 57, 11194-11200.
- Ingersoll, C.G., MacDonald, D.D., Brumbaugh, W.G., Johnson, B.T., Kemble, N.E., Kunz, J.L., May, T.W., Wang, N., Smith, J.R., Sparks, D.W., Ireland, D.S., 2002. Toxicity assessment of sediments from the Grand Calumet River and Indiana Harbor Canal in northwestern Indiana, USA. *Arch Environ Con Tox* 43, 156-167.
- Ionescu, M., Beranova, K., Dudkova, V., Kochankova, L., Demnerova, K., Macek, T., Mackova, M., 2009. Isolation and characterization of different plant associated bacteria and their potential to degrade polychlorinated biphenyls. *Int Biodeter Biodegr* 63, 667-672.
- Janssen, D.B., Oppentocht, J.E., Poelarends, G.J., 2001. Microbial dehalogenation. *Curr Opin Biotechnol* 12, 254-258.
- Jeon, C.O., Park, W., Ghiorse, W.C., Madsen, E.L., 2004. *Polaromonas naphthalenivorans* sp. nov., a naphthalene-degrading bacterium from naphthalene-contaminated sediment. *Int J Syst Evol Microbiol* 54, 93-97.
- Jeremiason, J.D., Hornbuckle, K.C., Eisenreich, S.J., 1994. PCBs in Lake-Superior, 1978-1992 - decreases in water concentrations reflect loss by volatilization. *Environ Sci Technol* 28, 903-914.
- Johnson, D.R., Lee, P.K.H., Holmes, V.F., Alvarez-Cohen, L., 2005. An internal reference technique for accurately quantifying specific mRNAs by real-time PCR with a application to the *tceA* reductive dehalogenase gene. *Appl Environ Microbiol* 71, 3866-3871.

Juhanson, J., Truu, J., Heinaru, E., Heinaru, A., 2009. Survival and catabolic performance of introduced *Pseudomonas* strains during phytoremediation and bioaugmentation field experiment. *FEMS Microbiol Ecol* 70, 446-455.

Junier, P., Junier, T., Witzel, K.P., 2008. TRiFLe, a program for *in silico* terminal restriction fragment length polymorphism analysis with user-defined sequence sets. *Appl Environ Microbiol* 74, 6452-6456.

Kim, I.S., Lee, H., Trevors, J.T., 2001. Effects of 2,2',5,5'-tetrachlorobiphenyl and biphenyl on cell membranes of *Ralstonia eutropha* H850. *FEMS Microbiol Lett* 200, 17-24.

Kim, M.H., Hao, O.J., 1999. Cometabolic degradation of chlorophenols by *Acinetobacter* species. *Water Res* 33, 562-574.

Kimbara, K., Hashimoto, T., Fukuda, M., Koana, T., Takagi, M., Oishi, M., Yano, K., 1988. Isolation and characterization of a mixed culture that degrades polychlorinated biphenyls. *Agr Biol Chem Tokyo* 52, 2885-2891.

Kjellerup, B.V., Sun, X.L., Ghosh, U., May, H.D., Sowers, K.R., 2008. Site-specific microbial communities in three PCB-impacted sediments are associated with different *in situ* dechlorinating activities. *Environ Microbiol* 10, 1296-1309.

Klappenbach, J.A., Dunbar, J.M., Schmidt, T.M., 2000. rRNA operon copy number reflects ecological strategies of bacteria. *Appl Environ Microb* 66, 1328-1333.

Kreader, C.A., 1996. Relief of amplification inhibition in PCR with bovine serum albumin or T4 gene 32 protein. *Appl Environ Microbiol* 62, 1102-1106.

Krumholz, L.R., 1997. *Desulfuromonas chloroethenica* sp. nov. uses tetrachloroethylene and trichloroethylene as electron acceptors. *Int J Syst Bacteriol* 47, 1262-1263.

Krumins, V., Park, J.W., Son, E.K., Rodenburg, L.A., Kerkhof, L.J., Häggblom, M.M., Fennell, D.E., 2009. PCB dechlorination enhancement in Anacostia River sediment microcosms. *Water Res* 43, 4549-4558.

Krzmarzick, M.J., Crary, B.B., Harding, J.J., Oyerinde, O.O., Leri, A.C., Myneni, S.C.B., Novak, P.J., 2012. Natural niche for organohalide-respiring *Chloroflexi*. *Appl Environ Microbiol* 78, 393-401.

Kučerová, P., Macková, M., Chromá, L., Burkhard, J., Tríska, J., Demnerová, K., Macek, T., 2000. Metabolism of polychlorinated biphenyls by *Solanum nigrum* hairy root clone SNC-90 and analysis of transformation products. *Plant Soil* 225, 109-115.

Lee, I., Fletcher, J.S., 1992. Involvement of mixed-function oxidase systems in polychlorinated biphenyl metabolism by plant-cells. *Plant Cell Rep* 11, 97-100.

Leigh, M.B., Fletcher, J.S., Fu, X.O., Schmitz, F.J., 2002. Root turnover: an important source of microbial substrates in rhizosphere remediation of recalcitrant contaminants. *Environ Sci Technol* 36, 1579-1583.

Leigh, M.B., Pellizari, V.H., Uhlík, O., Sutka, R., Rodrigues, J., Ostrom, N.E., Zhou, J., Tiedje, J.M., 2007. Biphenyl-utilizing bacteria and their functional genes in a pine root zone contaminated with polychlorinated biphenyls (PCBs). *ISME J* 1, 134-148.

- Leigh, M.B., Prouzová, P., Macková, M., Macek, T., Nagle, D.P., Fletcher, J.S., 2006. Polychlorinated biphenyl (PCB)-degrading bacteria associated with trees in a PCB-contaminated site. *Appl Environ Microbiol* 72, 2331-2342.
- Li, A., Rockne, K.J., Sturchio, N., Song, W., Ford, J.C., Wei, H., 2009. PCBs in sediments of the Great Lakes--distribution and trends, homolog and chlorine patterns, and *in situ* degradation. *Environ Pollut* 157, 141-147.
- Li, H.S., Liu, L.Y., Lin, C.X., Wang, S.R., 2011. Plant uptake and in-soil degradation of PCB-5 under varying cropping conditions. *Chemosphere* 84, 943-949.
- Liu, J., Schnoor, J.L., 2008. Uptake and translocation of lesser-chlorinated polychlorinated biphenyls (PCBs) in whole hybrid poplar plants after hydroponic exposure. *Chemosphere* 73, 1608-1616.
- Liu, J.Y., Hu, D.F., Jiang, G.B., Schnoor, J.L., 2009. *In vivo* biotransformation of 3,3',4,4'-tetrachlorobiphenyl by whole plants - poplars and switchgrass. *Environ Sci Technol* 43, 7503-7509.
- Lloyd-Jones, G., Lau, P.C., 1998. A molecular view of microbial diversity in a dynamic landfill in Quebec. *FEMS Microbiol Lett* 162, 219-226.
- Longnecker, M.P., Rogan, W.J., Lucier, G., 1997. The human health effects of DDT (dichlorodiphenyltrichloroethane) and PCBs (polychlorinated biphenyls) and an overview of organochlorines in public health 1. *Annu Rev Publ Health* 18, 211-244.
- Luepromchai, E., Singer, A.C., Yang, C.H., Crowley, D.E., 2002. Interactions of earthworms with indigenous and bioaugmented PCB-degrading bacteria. *FEMS Microbiol Ecol* 41, 191-197.
- Luo, W., D'Angelo, E.M., Coyne, M.S., 2008. Organic carbon effects on aerobic polychlorinated biphenyl removal and bacterial community composition in soils and sediments. *Chemosphere* 70, 364-373.
- MacDonald, R.W., Barrie, L.A., Bidleman, T.F., Diamond, M.L., Gregor, D.J., Semkin, R.G., Strachan, W.M.J., Li, Y.F., Wania, F., Alaee, M., Alexeeva, L.B., Backus, S.M., Bailey, R., Bewers, J.M., Gobeil, C., Halsall, C.J., Harner, T., Hoff, J.T., Jantunen, L.M.M., Lockhart, W.L., Mackay, D., Muir, D.C.G., Pudykiewicz, J., Reimer, K.J., Smith, J.N., Stern, G.A., Schroeder, W.H., Wagemann, R., Yunker, M.B., 2000. Contaminants in the Canadian Arctic: 5 years of progress in understanding sources, occurrence and pathways. *Sci Total Environ* 254, 93-234.
- Macková, M., Barriault, D., Francová, K., Sylvestre, M., Möder, M., Vrchotová, B., Lovecká, P., Najmanová, J., Demnerová, K., Nováková, M., 2006. Phytoremediation of polychlorinated biphenyls, Phytoremediation rhizoremediation. Springer, pp. 143-167.
- Macková, M., Chroma, L., Kučerová, P., Burkhard, J., Demnerová, K., Macek, T., 2001. Some aspects of PCB metabolism by Horseradish cells. *Int J Phytoremediat* 3, 401-414.
- Macková, M., Prouzova, P., Stursa, P., Ryslava, E., Uhlík, O., Beranova, K., Rezek, J., Kurzawova, V., Demnerová, K., Macek, T., 2009. Phyto/rhizoremediation studies using long-term PCB-contaminated soil. *Environ Sci Pollut R* 16, 817-829.

- Macková, M., Uhlík, O., Lovecká, P., Viktorova, J., Nováková, M., Demnerová, K., Sylvestre, M., Macek, T., 2010. Bacterial degradation of polychlorinated biphenyls, In: Barton, L.L., Mandl, M., Loy, A. (Eds.), *Geomicrobiology: Molecular and Environmental Perspective*. Springer Netherlands, pp. 347-366.
- Macková, M., Vrchotová, B., Francová, K., Sylvestre, M., Tomaniová, M., Lovecká, P., Demnerová, K., Macek, T., 2007. Biotransformation of PCBs by plants and bacteria – consequences of plant-microbe interactions. *Eur J Soil Biol* 43, 233-241.
- Maness, A.D., Bowman, K.S., Yan, J., Rainey, F.A., Moe, W.M., 2012. *Dehalogenimonas* spp. can reductively dehalogenate high concentrations of 1,2-dichloroethane, 1,2-dichloropropane, and 1,1,2-trichloroethane. *AMB Express* 2, 1-7.
- Martinez, A., Hornbuckle, K.C., 2011. Record of PCB congeners, sorbents and potential toxicity in core samples in Indiana Harbor and Ship Canal. *Chemosphere* 85, 542-547.
- Martinez, A., Norstrom, K., Wang, K., Hornbuckle, K.C., 2010. Polychlorinated biphenyls in the surficial sediment of Indiana Harbor and Ship Canal, Lake Michigan. *Environ Int* 36, 849-854.
- Martinez, A., O'Sullivan, C., Reible, D., Hornbuckle, K.C., 2013. Sediment pore water distribution coefficients of PCB congeners in enriched black carbon sediment. *Environ Pollut* 182, 357-363.
- Masai, E., Yamada, A., Healy, J.M., Hatta, T., Kimbara, K., Fukuda, M., Yano, K., 1995. Characterization of biphenyl catabolic genes of Gram-positive polychlorinated biphenyl degrader *Rhodococcus* sp strain RHA1. *Appl Environ Microbiol* 61, 2079-2085.
- Master, E.R., Mohn, W.W., 2001. Induction of *bphA*, encoding biphenyl dioxygenase, in two polychlorinated biphenyl-degrading bacteria, psychrotolerant *Pseudomonas* strain Cam-1 and mesophilic *Burkholderia* strain LB400. *Appl Environ Microbiol* 67, 2669-2676.
- May, H.D., Miller, G.S., Kjellerup, B.V., Sowers, K.R., 2008. Dehalorespiration with polychlorinated biphenyls by an anaerobic ultramicrobacterium. *Appl Environ Microb* 74, 2089-2094.
- Meggo, R.E., Schnoor, J.L., 2013. Rhizosphere redox cycling and implications for rhizosphere biotransformation of selected polychlorinated biphenyl (PCB) congeners. *Ecol Eng* 57, 285-292.
- Meggo, R.E., Schnoor, J.L., Hu, D., 2013. Dechlorination of PCBs in the rhizosphere of switchgrass and poplar. *Environ Pollut* 178, 312-321.
- Meijer, S.N., Ockenden, W.A., Sweetman, A., Breivik, K., Grimalt, J.O., Jones, K.C., 2003. Global distribution and budget of PCBs and HCB in background surface soils: implications or sources and environmental processes. *Environ Sci Technol* 37, 667-672.
- MinRat-Wirtschaftsministerium, H., Baden-Württemberg, I.B., Bork, H., Köln, G., Goßler-Landesgewerbeanstalt, H.D., Bremen, F.H., Horneffer, H., 1995. Richtlinie für die bewertung und sanierung PCB-belasteter baustoffe und bauteile in Gebäuden (PCB-richtlinie).

- Motelay-Massei, A., Ollivon, D., Garban, B., Teil, M.J., Blanchard, M., Chevreuil, M., 2004. Distribution and spatial trends of PAHs and PCBs in soils in the Seine River basin, France. *Chemosphere* 55, 555-565.
- Nadkarni, M.A., Martin, F.E., Jacques, N.A., Hunter, N., 2002. Determination of bacterial load by real-time PCR using a broad-range (universal) probe and primers set. *Microbiol* 148, 257-266.
- Nandi, R., Sengupta, S., 1998. Microbial production of hydrogen: an overview. *Crit Rev Microbiol* 24, 61-84.
- Nogales, B., Moore, E.R., Abraham, W.R., Timmis, K.N., 1999. Identification of the metabolically active members of a bacterial community in a polychlorinated biphenyl-polluted moorland soil. *Environ Microbiol* 1, 199-212.
- Norini, M.P., Secher, C., Lollier, M., Jezequel, K., Cornu, J.Y., Lebeau, T., 2013. Quantification of the 16S-23S rRNA internal transcribed spacers of *Burkholderia xenovorans* strain LB400 using real-time PCR in soil samples. *Lett Appl Microbiol* 56, 366-372.
- Nováková, H., Vošahlíková, M., Pazlarová, J., Macková, M., Burkhard, J., Demnerová, K., 2002. PCB metabolism by *Pseudomonas* sp P2. *Int Biodeter Biodegr* 50, 47-54.
- Oh, K.H., Ostrofsky, E.B., Cho, Y.C., 2008. Molecular characterization of polychlorinated biphenyl-dechlorinating populations in contaminated sediments. *J Microbiol* 46, 165-173.
- Okeke, B.C., Chang, Y.C., Hatsu, M., Suzuki, T., Takamizawa, K., 2001. Purification, cloning, and sequencing of an enzyme mediating the reductive dechlorination of tetrachloroethylene (PCE) from *Clostridium bifermentans* DPH-1. *Can J Microbiol* 47, 448-456.
- Park, Y.I., So, J.S., Koh, S.C., 1999. Induction by carvone of the polychlorinated biphenyl (PCB)-degradative pathway in *Alcaligenes eutrophus* H850 and its molecular monitoring. *J Microbiol Biotechnol* 9, 804-810.
- Parnell, J.J., Denef, V.J., Park, J., Tsoi, T., Tiedje, J.M., 2010. Environmentally relevant parameters affecting PCB degradation: carbon source- and growth phase-mitigated effects of the expression of the biphenyl pathway and associated genes in *Burkholderia xenovorans* LB400. *Biodegradation* 21, 147-156.
- Parnell, J.J., Park, J., Denef, V., Tsoi, T., Hashsham, S., Quensen, J., 3rd, Tiedje, J.M., 2006. Coping with polychlorinated biphenyl (PCB) toxicity: Physiological and genome-wide responses of *Burkholderia xenovorans* LB400 to PCB-mediated stress. *Appl Environ Microbiol* 72, 6607-6614.
- Pavlu, L., Vosáhllová, J., Klierová, H., Prouza, M., Demnerová, K., Brenner, V., 1999. Characterization of chlorobenzoate degraders isolated from polychlorinated biphenyl-contaminated soil and sediment in the Czech Republic. *J Appl Microbiol* 87, 381-386.
- Payne, R.B., Fagervold, S.K., May, H.D., Sowers, K.R., 2013. Remediation of polychlorinated biphenyl impacted sediment by concurrent bioaugmentation with anaerobic halorespiring and aerobic degrading bacteria. *Environ Sci Technol* 47, 3807-3815.

- Payne, R.B., May, H.D., Sowers, K.R., 2011. Enhanced reductive dechlorination of polychlorinated biphenyl impacted sediment by bioaugmentation with a dehalorespiring bacterium. *Environ Sci Technol* 45, 8772-8779.
- Petrić, I., Bru, D., Udiković-Kolić, N., Hršak, D., Philippot, L., Martin-Laurent, F., 2011a. Evidence for shifts in the structure and abundance of the microbial community in a long-term PCB-contaminated soil under bioremediation. *J Hazard Mater* 195, 254-260.
- Petrić, I., Hršak, D., Fingler, S., Udiković-Kolić, N., Bru, D., Martin-Laurent, F., 2011b. Insight in the PCB-degrading functional community in long-term contaminated soil under bioremediation. *J Soil Sediment* 11, 290-300.
- Picardal, F., Arnold, R.G., Huey, B.B., 1995. Effects of electron-donor and acceptor conditions on reductive dehalogenation of tetrachloromethane by *Shewanella Putrefaciens* 200. *Appl Environ Microbiol* 61, 8-12.
- Pieper, D.H., 2005. Aerobic degradation of polychlorinated biphenyls. *Appl Microbiol Biot* 67, 170-191.
- Pieper, D.H., Seeger, M., 2008. Bacterial metabolism of polychlorinated biphenyls. *J Mol Microbiol Biotechnol* 15, 121-138.
- Purkey, H.E., Palaninathan, S.K., Kent, K.C., Smith, C., Safe, S.H., Sacchettini, J.C., Kelly, J.W., 2004. Hydroxylated polychlorinated biphenyls selectively bind transthyretin in blood and inhibit amyloidogenesis: rationalizing rodent PCB toxicity. *Chem Biol* 11, 1719-1728.
- Quensen, J.F., Boyd, S.A., Tiedje, J.M., 1990. Dechlorination of four commercial polychlorinated biphenyl mixtures (Aroclors) by anaerobic microorganisms from sediments. *Appl Environ Microbiol* 56, 2360-2369.
- Reid, B.J., Jones, K.C., Semple, K.T., 2000. Bioavailability of persistent organic pollutants in soils and sediments - a perspective on mechanisms, consequences and assessment. *Environ Pollut* 108, 103-112.
- Rein, A., Fernqvist, M.M., Mayer, P., Trapp, S., Bittens, M., Karlson, U.G., 2007. Degradation of PCB congeners by bacterial strains. *Appl Microbiol Biotechnol* 77, 469-481.
- Rezek, J., Macek, T., Mackova, M., Triska, J., 2007. Plant metabolites of polychlorinated biphenyls in hairy root culture of black nightshade *Solanum nigrum* SNC-90. *Chemosphere* 69, 1221-1227.
- Rieu, I., Powers, S.J., 2009. Real-time quantitative RT-PCR: design, calculations, and statistics. *Plant Cell* 21, 1031-1033.
- Rodenburg, L.A., Du, S.Y., Fennell, D.E., Cavallo, G.J., 2010. Evidence for widespread dechlorination of polychlorinated biphenyls in groundwater, landfills, and wastewater collection systems. *Environ Sci Technol* 44, 7534-7540.
- Rodrigues, J.L., Kachel, C.A., Aiello, M.R., Quensen, J.F., Maltseva, O.V., Tsoi, T.V., Tiedje, J.M., 2006. Degradation of aroclor 1242 dechlorination products in sediments by *Burkholderia xenovorans* LB400(ohb) and *Rhodococcus* sp. strain RHA1(fcb). *Appl Environ Microbiol* 72, 2476-2482.

- Ross, G., 2004. The public health implications of polychlorinated biphenyls (PCBs) in the environment. *Ecotoxicol Environ Saf* 59, 275-291.
- Saavedra, J.M., Acevedo, F., Gonzalez, M., Seeger, M., 2010. Mineralization of PCBs by the genetically modified strain *Cupriavidus necator* JMS34 and its application for bioremediation of PCBs in soil. *Appl Microbiol Biotechnol* 87, 1543-1554.
- Schütte, U.M.E., Abdo, Z., Bent, S.J., Shyu, C., Williams, C.J., Pierson, J.D., Forney, L.J., 2008. Advances in the use of terminal restriction fragment length polymorphism (T-RFLP) analysis of 16S rRNA genes to characterize microbial communities. *Appl Microbiol Biot* 80, 365-380.
- Schnoor, J.L., Licht, L.A., Mccutcheon, S.C., Wolfe, N.L., Carreira, L.H., 1995. Phytoremediation of organic and nutrient contaminants. *Environ Sci Technol* 29, A318-A323.
- Secher, C., Lollier, M., Jezequel, K., Cornu, J.Y., Amalric, L., Lebeau, T., 2013. Decontamination of a polychlorinated biphenyls-contaminated soil by phytoremediation-assisted bioaugmentation. *Biodegradation* 24, 549-562.
- Shimura, M., Mukerjee-Dhar, G., Kimbara, K., Nagato, H., Kiyohara, H., Hatta, T., 1999. Isolation and characterization of a thermophilic *Bacillus* sp JF8 capable of degrading polychlorinated biphenyls and naphthalene. *FEMS Microbiol Lett* 178, 87-93.
- Shuai, J.J., Tian, Y.S., Yao, Q.H., Peng, R.H., Xiong, F., Xiong, A.S., 2010. Identification and analysis of polychlorinated biphenyls (PCBs)-biodegrading bacterial strains in Shanghai. *Curr Microbiol* 61, 477-483.
- Simcik, M.F., Zhang, H.X., Eisenreich, S.J., Franz, T.P., 1997. Urban contamination of the Chicago coastal Lake Michigan atmosphere by PCBs and PAHs during AEOLOS. *Environ Sci Technol* 31, 2141-2147.
- Singer, A.C., Crowley, D.E., Thompson, I.P., 2003a. Secondary plant metabolites in phytoremediation and biotransformation. *Trends Biotechnol* 21, 123-130.
- Singer, A.C., Gilbert, E.S., Luepromchai, E., Crowley, D.E., 2000. Bioremediation of polychlorinated biphenyl-contaminated soil using carvone and surfactant-grown bacteria. *Appl Microbiol Biotechnol* 54, 838-843.
- Singer, A.C., Smith, D., Jury, W.A., Hathuc, K., Crowley, D.E., 2003b. Impact of the plant rhizosphere and augmentation on remediation of polychlorinated biphenyl contaminated soil. *Environ Toxicol Chem* 22, 1998-2004.
- Slater, H., Gouin, T., Leigh, M.B., 2011. Assessing the potential for rhizoremediation of PCB contaminated soils in northern regions using native tree species. *Chemosphere* 84, 199-206.
- Smith, K.E., Schwab, A.R., Banks, M.K., 2007. Phytoremediation of polychlorinated biphenyl (PCB)-contaminated sediment: a greenhouse feasibility study. *J Environ Qual* 36, 239-244.
- Sowers, K.R., May, H.D., 2013. *In situ* treatment of PCBs by anaerobic microbial dechlorination in aquatic sediment: are we there yet? *Curr Opin Biotechnol* 24, 482-488.

- Spain, A.M., Krumholz, L.R., Elshahed, M.S., 2009. Abundance, composition, diversity and novelty of soil *Proteobacteria*. *ISME J* 3, 992-1000.
- Strycharz, S.M., Woodard, T.L., Johnson, J.P., Nevin, K.P., Sanford, R.A., Löffler, F.E., Lovley, D.R., 2008. Graphite electrode as a sole electron donor for reductive dechlorination of tetrachlorethene by *Geobacter lovleyi*. *Appl Environ Microbiol* 74, 5943-5947.
- Sul, W.J., Park, J., Quensen, J.F., Rodrigues, J.L.M., Seliger, L., Tsoi, T.V., Zylstra, G.J., Tiedje, J.M., 2009. DNA-stable isotope probing integrated with metagenomics for retrieval of biphenyl dioxygenase genes from polychlorinated biphenyl-contaminated river sediment. *Appl Environ Microbiol* 75, 5501-5506.
- Sung, Y., Fletcher, K.F., Ritalaliti, K.M., Apkarian, R.P., Ramos-Hernandez, N., Sanford, R.A., Mesbah, N.M., Löffler, F.E., 2006. *Geobacter lovleyi* sp nov strain SZ, a novel metal-reducing and tetrachloroethene-dechlorinating bacterium. *Appl Environ Microbiol* 72, 2775-2782.
- Tam, N.F.Y., Wong, Y.S., 2008. Effectiveness of bacterial inoculum and mangrove plants on remediation of sediment contaminated with polycyclic aromatic hydrocarbons. *Mar Pollut Bull* 57, 716-726.
- TAMS Consultants, I., The CADMUS Group, Inc., and Gradient Corporation, 1997. Phase 2 report (review copy): Further site characterization and analysis volume 2C - Data evaluation and interpretation report Hudson River PCBs reassessment RI/FS. USEPA, Washington, DC.
- Tamura, K., Dudley, J., Nei, M., Kumar, S., 2007. MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Mol Biol Evol* 24, 1596-1599.
- Tandlich, R., Brezna, B., Dercova, K., 2001. The effect of terpenes on the biodegradation of polychlorinated biphenyls by *Pseudomonas stutzeri*. *Chemosphere* 44, 1547-1555.
- Thompson, J.D., Gibson, T.J., Plewniak, F., Jeanmougin, F., Higgins, D.G., 1997. The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res* 25, 4876-4882.
- Tiedje, J.M., Quensen, J.F., 3rd, Chee-Sanford, J., Schimel, J.P., Boyd, S.A., 1993. Microbial reductive dechlorination of PCBs. *Biodegradation* 4, 231-240.
- Tillmann, S., Strompl, C., Timmis, K.N., Abraham, W.R., 2005. Stable isotope probing reveals the dominant role of Burkholderia species in aerobic degradation of PCBs. *FEMS Microbiol Ecol* 52, 207-217.
- Toussaint, J.P., Thi, T.M.P., Barriault, D., Sylvestre, M., 2012. Plant exudates promote PCB degradation by a rhodococcal rhizobacteria. *Appl Microbiol Biotechnol* 95, 1589-1603.
- Uhlík, O., Jecna, K., Macková, M., Vlcek, C., Hroudova, M., Demnerová, K., Paces, V., Macek, T., 2009. Biphenyl-metabolizing bacteria in the rhizosphere of Horseradish and bulk soil contaminated by polychlorinated biphenyls as revealed by stable isotope probing. *Appl Environ Microbiol* 75, 6471-6477.

- UNEP, 2001. Final act of the conference of plenipotentiaries on the Stockholm Convention on Persistent Organic Pollutants, Stockholm, Sweden.
- USACE, 2013. Indiana Harbor and Canal dredging and disposal project. Available from <http://www.lrc.usace.army.mil/Missions/CivilWorksProjects/IndianaHarbor.aspx>.
- USEPA, 1991. National Primary Drinking Water Regulations, Final Rule 40 CFR Parts 141-143. Fed Regist. 56, (20). Washington, DC.
- USEPA, 2010. Method 1668C: Chlorinated biphenyl congeners in water, soil, sediment, biosolids, and tissue by HRGC/HRMS, Washington, DC.
- Van Aken, B., Correa, P.A., Schnoor, J.L., 2010. Phytoremediation of polychlorinated biphenyls: new trends and promises. *Environ Sci Technol* 44, 2767-2776.
- Van den Berg, M., Birnbaum, L.S., Denison, M., De Vito, M., Farland, W., Feeley, M., Fiedler, H., Hakansson, H., Hanberg, A., Haws, L., Rose, M., Safe, S., Schrenk, D., Tohyama, C., Tritscher, A., Tuomisto, J., Tysklind, M., Walker, N., Peterson, R.E., 2006. The 2005 World Health Organization reevaluation of human and mammalian toxic equivalency factors for dioxins and dioxin-like compounds. *Toxicol Sci* 93, 223-241.
- Van Noort, P.C.M., Haftka, J.J.H., Parsons, J.R., 2010. Updated Abraham solvation parameters for polychlorinated biphenyls. *Environ Sci Technol* 44, 7037-7042.
- Vandort, H.M., Bedard, D.L., 1991. Reductive *ortho*-dechlorination and *meta*-dechlorination of a polychlorinated biphenyl congener by anaerobic microorganisms. *Appl Environ Microbiol* 57, 1576-1578.
- Wagner, A., Adrian, L., Kleinstaub, S., Andreesen, J.R., Lechner, U., 2009. Transcription analysis of genes encoding homologues of reductive dehalogenases in "*Dehalococcoides*" sp strain CBDB1 by using terminal restriction fragment length polymorphism and quantitative PCR. *Appl Environ Microbiol* 75, 1876-1884.
- Wang, S., Zhang, S.Z., Huang, H.L., Zhao, M.M., Lv, J.T., 2011. Uptake, translocation and metabolism of polybrominated diphenyl ethers (PBDEs) and polychlorinated biphenyls (PCBs) in maize (*Zea mays L.*). *Chemosphere* 85, 379-385.
- Wang, S.Q., He, J.Z., 2013. Phylogenetically distinct bacteria involve extensive dechlorination of Aroclor 1260 in sediment-free cultures. *Plos One* 8, e59178.
- Watts, J.E.M., Fagervold, S.K., May, H.D., Sowers, K.R., 2005. A PCR-based specific assay reveals a population of bacteria within the *Chloroflexi* associated with the reductive dehalogenation of polychlorinated biphenyls. *Microbiology* 151, 2039-2046.
- Watts, J.E.M., Wu, Q.Z., Schreier, S.B., May, H.D., Sowers, K.R., 2001. Comparative analysis of polychlorinated biphenyl-dechlorinating communities in enrichment cultures using three different molecular screening techniques. *Environ Microbiol* 3, 710-719.
- White, J.C., Parrish, Z.D., Isleyen, M., Gent, M.P.N., Iannucci-Berger, W., Eitzer, B.D., Kelsey, J.W., Mattina, M.I., 2006. Influence of citric acid amendments on the availability of weathered PCBs to plant and earthworm species. *Int J Phytoremediat* 8, 63-79.

- Whitfield Åslund, M.L., Rutter, A., Reimer, K.J., Zeeb, B.A., 2008. The effects of repeated planting, planting density, and specific transfer pathways on PCB uptake by *Cucurbita pepo* grown in field conditions. *Sci Total Environ* 405, 14-25.
- Whitfield Åslund, M.L., Zeeb, B.A., Rutter, A., Reimer, K.J., 2007. *In situ* phytoextraction of polychlorinated biphenyl - (PCB) contaminated soil. *Sci Total Environ* 374, 1-12.
- Wiegel, J., Wu, Q., 2000. Microbial reductive dehalogenation of polychlorinated biphenyls. *FEMS Microbiol Ecol* 32, 1-15.
- Wu, Q., Sowers, K.R., May, H.D., 1998. Microbial reductive dechlorination of aroclor 1260 in anaerobic slurries of estuarine sediments. *Appl Environ Microbiol* 64, 1052-1058.
- Wu, Q.Z., Watts, J.E.M., Sowers, K.R., May, H.D., 2002. Identification of a bacterium that specifically catalyzes the reductive dechlorination of polychlorinated biphenyls with doubly flanked chlorines. *Appl Environ Microbiol* 68, 807-812.
- Xu, Y., Yu, Y., Gregory, K.B., VanBriesen, J.M., 2012. Comprehensive assessment of bacterial communities and analysis of PCB congeners in PCB-contaminated sediment with depth. *J Environ Eng* 138, 1167-1178.
- Yan, J., Rash, B.A., Rainey, F.A., Moe, W.M., 2009. Isolation of novel bacteria within the *Chloroflexi* capable of reductive dechlorination of 1,2,3-trichloropropane. *Environ Microbiol* 11, 833-843.
- Yan, T., LaPara, T.M., Novak, P.J., 2006. The reductive dechlorination of 2,3,4,5-tetrachlorobiphenyl in three different sediment cultures: evidence for the involvement of phylogenetically similar *Dehalococcoides*-like bacterial populations. *FEMS Microbiol Ecol* 55, 248-261.
- Yoshida, N., Takahashi, N., Hiraishi, A., 2005. Phylogenetic characterization of a polychlorinated-dioxin-dechlorinating microbial community by use of microcosm studies. *Appl Environ Microbiol* 71, 4325-4334.
- Yoshida, N., Ye, L., Baba, D., Katayama, A., 2009. Reductive dechlorination of polychlorinated biphenyls and dibenzo-p-dioxins in an enrichment culture containing *Dehalobacter* species. *Microbes Environ* 24, 343-346.
- Zaitsev, G., Karasevich, Y., 1985. Primary steps in metabolism of 4-chlorobenzoate in *Arthrobacter globiformis*. *Mikrobiologiya* 50, 423-428.
- Zanaroli, G., Balloi, A., Negroni, A., Daffonchio, D., Young, L.Y., Fava, F., 2010. Characterization of the microbial community from the marine sediment of the Venice lagoon capable of reductive dechlorination of coplanar polychlorinated biphenyls (PCBs). *J Hazard Mater* 178, 417-426.
- Zeeb, B.A., Amphlett, J.S., Rutter, A., Reimer, K.J., 2006. Potential for phytoremediation of polychlorinated biphenyl-(PCB-)contaminated soil. *Int J Phytoremediat* 8, 199-221.
- Zhai, G., Lehmler, H.-J., Schnoor, J.L., 2010a. Hydroxylated metabolites of 4-monochlorobiphenyl and its metabolic pathway in whole poplar plants. *Environ Sci Technol* 44, 3901-3907.

Zhai, G.S., Lehmler, F.J., Schnoor, J.L., 2010b. Identification of hydroxylated metabolites of 3,3',4,4'-tetrachlorobiphenyl and metabolic pathway in whole poplar plants. *Chemosphere* 81, 523-528.

Zhang, S.Y., Zhang, Q.A., Darisaw, S., Ehie, O., Wang, G.D., 2007. Simultaneous quantification of polycyclic aromatic hydrocarbons (PAHs), polychlorinated biphenyls (PCBs), and pharmaceuticals and personal care products (PPCPs) in Mississippi river water, in New Orleans, Louisiana, USA. *Chemosphere* 66, 1057-1069.