

2016

Guidelines are not evidence


James P. Meza MD, PhD

Wayne State University School of Medicine, jmeza@med.wayne.edu

Nicholus H. Yee MD

Oakwood Annapolis Hospital Family Medicine Residency, nicholus.h@gmail.com

Follow this and additional works at: <https://digitalcommons.wayne.edu/crp>

 Part of the [Medical Education Commons](#), and the [Translational Medical Research Commons](#)

Recommended Citation

Meza JP & Yee NH. Guidelines are not evidence. *Clin. Res. Pract.* June 10, 2016;2(1);eP1219. doi: [10.22237/crp/1465591836](https://doi.org/10.22237/crp/1465591836)

This Editorial is brought to you for free and open access by the Open Access Journals at DigitalCommons@WayneState. It has been accepted for inclusion in Clinical Research in Practice: The Journal of Team Hippocrates by an authorized editor of DigitalCommons@WayneState.

EDITORIAL:

Guidelines are not evidence

JAMES P. MEZA, MD, PhD, Wayne State University School of Medicine, jmeza@med.wayne.edu
NICHOLUS H. YEE, MD, Oakwood Annapolis Hospital Family Medicine Residency, nicholus.h@gmail.com

Guidelines are inherently flawed when caring for an individual patient.

When doctors use guidelines in clinical care, it is far too easy to make the cognitive error that they are helping that patient. Guidelines are misleading because they imply that there is one right management that applies to all patients; and conversely, that all other approaches are wrong. We don't think doctors should use guidelines as a direct mandate for individual patient care. Guidelines, ostensibly, are based on evidence that applies to populations. Physicians are charged with caring for individuals. There is a risk of benefit and a risk of harm in following a guideline. A more accurate portrayal of the role of guidelines is to initiate shared decision-making.

The Patient Centered Medical Home is an oxymoron because it is a hybrid of population management, patient communication protocols, and patient care processes: population management is not patient-centered. Population management follows guidelines that may or may not be in the best interest of our individual patients. For example, at our clinical practice site, our patients receive a requisition for a mammogram and are told to schedule an appointment before ever being seen by a doctor—there is no shared decision making. We are not using guidelines appropriately.

Being able to explain the evidence supporting a guideline is part of the doctor's job. Evidence from clinical research must be applied to an individual patient's circumstance. It is perfectly legitimate to use the same evidence and come to diametrically opposed decisions for two separate patients. Patient values are important when applying evidence. We don't think abrogating that responsibility is in anyone's best interest—neither the patient nor the healthcare system.

Without understanding clinical evidence, an algorithm, flow diagram or computer can replace the doctor. Such a lack of critical thinking is dangerous. An example of blindly following guidelines occurred recently in the precepting room. A resident said, "...and I'm going to check an LDL." When asked why, the resident replied, "I check an LDL on all my patients every year." The resident had internalized an out-of-date guideline and continued to use it for patient care.¹ Additionally, the resident couldn't conceptualize the reason for the recommendation; the resident was unaware of the evidence on which the guideline was based. The preceptor gave a provocative response, saying, "Oh, you must still believe in the cholesterol hypothesis," and gave the resident a flyer describing a medication that failed to improve cardiovascular deaths despite significantly lowering cholesterol levels.²

Then there is the issue of guidelines being out-of-date³, based on poor-quality or biased interpretation of clinical research⁴, or simply a marketing ploy of Big Pharma.⁵⁻¹¹ Doctors need to be familiar with major practice guidelines of their chosen specialty and decide which are trustworthy in practice. To do this, doctors need to be able to read and interpret clinical research. Only then can doctors understand when it's in the best interest of the patient to deviate from the recommendations of the guideline. Guidelines have to be viewed as any other intervention, having a number needed to treat and a number needed to harm.

We are proud that our contributors have demonstrated mastery of these skills.^{12,13}

Clinical Research in Practice Faculty Editor JAMES P. MEZA, MD, PhD teaches Translational Medicine and Evidence Based Practice at Wayne State University School of Medicine. He is also the Director of Research at Oakwood Annapolis Hospital Family Medicine Residency. NICHOLUS H. YEE, MD is a faculty member from Beaumont Health.

References

1. Stine NW, Chokshi DA. Elimination of lipid levels from quality measures: implications and alternatives. *JAMA*. 2014;312(19):1971-1972. doi:[10.1001/jama.2014.15600](https://doi.org/10.1001/jama.2014.15600)
2. Eli Lilly and Company. A study of Evacetrapib (LY2484595) in participants with high cholesterol (ACCENTUATE). In: ClinicalTrials.gov [Internet]. Bethesda (MD): National Library of Medicine (US). 2000 [cited 2016 June 8]. Available from: <https://clinicaltrials.gov/ct2/show/NCT02227784> NLM Identifier: NCT02227784.
3. Mandell LA, Wunderink RG, Anzueto A, et al. Infectious Diseases Society of America/American Thoracic Society consensus guidelines on the management of community-acquired pneumonia in adults. *Clin Infect Dis*. 2007;44 Suppl 2:S27-72. doi:[10.1086/511159](https://doi.org/10.1086/511159)
4. American Diabetes Association. Glycemic targets. Sec. 5. In *Standards of Medical Care in Diabetes—2016*. Diabetes Care 2016;39(Suppl. 1):S39–S46. doi:[10.2337/dc16-S008](https://doi.org/10.2337/dc16-S008)
5. Hutchison G. Guidelines can harm patients too. *BMJ*. 2012 Apr 18;344:e2685. doi:[10.1136/bmj.e2685](https://doi.org/10.1136/bmj.e2685)
6. Kung J, Miller RR, Mackowiak PA. Failure of clinical practice guidelines to meet Institute of Medicine standards: Two more decades of little, if any, progress. *Arch Intern Med*. 2012;172(21):1628-1633. doi:[10.1001/2013.jamainternmed.56](https://doi.org/10.1001/2013.jamainternmed.56)
7. Lenzer J. Why we can't trust clinical guidelines. *BMJ*. 2013;346:f3830. doi:[10.1136/bmj.f3830](https://doi.org/10.1136/bmj.f3830)
8. Lenzer J, Hoffman JR, Furberg CD, Ioannidis JP, Guideline Panel Review Working G. Ensuring the integrity of clinical practice guidelines: a tool for protecting patients. *BMJ*. 2013;347:f5535. doi:[10.1136/bmj.f5535](https://doi.org/10.1136/bmj.f5535)
9. McAlister FA, van Diepen S, Padwal RS, Johnson JA, Majumdar SR. How evidence-based are the recommendations in evidence-based guidelines? *PLoS Med*. 2007 Aug 7;4(8):e250. doi:[10.1371/journal.pmed.0040250](https://doi.org/10.1371/journal.pmed.0040250)
10. Mendelson TB, Meltzer M, Campbell EG, Caplan AL, Kirkpatrick JN. Conflicts of interest in cardiovascular clinical practice guidelines. *Arch Intern Med*. 2011 Mar 28;171(6):577-84. doi:[10.1001/archinternmed.2011.96](https://doi.org/10.1001/archinternmed.2011.96)
11. Sinuff T, Patel RV, Adhikari NK, Meade MO, Schunemann HJ, Cook DJ. Quality of professional society guidelines and consensus conference statements in critical care. *Critical care medicine*. 2008 Apr;36(4):1049-58. doi:[10.1097/CCM.0b013e31816a01ec](https://doi.org/10.1097/CCM.0b013e31816a01ec)
12. Mychajlowycz M. Abnormal uterine bleeding guidelines and clinical practice are based on poor quality evidence. *Clinical Research in Practice: The Journal of Team Hippocrates*. Feb 5, 2016;2(1):eP1077. <http://digitalcommons.wayne.edu/crp/vol2/iss1/4/>
13. Simmer S. Intravenous estrogen is not proven effective for abnormal uterine bleeding: guidelines are based on flawed evidence. *Clinical Research in Practice: The Journal of Team Hippocrates*. 2015;1(1):eP1003. <http://digitalcommons.wayne.edu/crp/vol1/iss1/3/>