

## IONIC MECHANISMS UNDERLYING SYNCHRONIZED OSCILLATIONS AND TRAVELING WAVES IN A MODEL OF FERRET THALAMIC SLICES

A. Destexhe\*, T. Balj, D.A. McCormick† and T.J. Sejnowski

The Salk Institute, Computational Neurobiology Laboratory, PO Box 85800, San Diego, CA 92186, USA; † Section of Neurobiology, Yale University School of Medicine, 333 Cedar Street, New Haven CT 06510, USA.

A network model of thalamocortical (TC) and thalamic reticular (RE) neurons was developed based on electrophysiological measurements in ferret thalamic slices. Single-compartment TC and RE cells included voltage- and calcium-sensitive currents described by Hodgkin-Huxley type of kinetics. Synaptic currents were modeled by kinetic models of AMPA, GABA<sub>A</sub> and GABA<sub>B</sub> receptors. The model reproduced successfully the characteristics of spindle and slow bicuculline-induced (SBI) oscillations. The most critical parameters were: (a) the mutual recruitment between TC and RE cells, due to their reciprocal connectivity. The time course of GABAergic IPSPs from RE to TC determined the frequency of oscillations (8-12 Hz spindle or 2-4 Hz SBI). (b) Cooperativity in the activation of GABA<sub>B</sub> responses, which are absent during spindles. If the intra-RE GABA<sub>A</sub> IPSPs were suppressed, the stronger discharges of RE cells enhanced GABA<sub>B</sub> IPSPs in TC cells, recruiting TC and RE neurons in the slower highly synchronized SBI oscillations. (c) A TC origin for the "waning" of these oscillations, through the upregulation of  $I_h$  by intracellular calcium, leading to a refractory period in these cells. (d) Local axonal arborization of the TC-RE and RE-TC projections, which allowed traveling phenomena to occur. The model predicts that intrinsic properties of TC and RE neurons and their reciprocal connectivity establish a highly excitable structure; stimulation at any location can induce an oscillatory travelling wave through the reciprocal recruitment of TC and RE cells. The presence of a refractory period in TC cells accounts for the waxing and waning properties of the traveling waves, collisions between waves and the slower propagation of highly synchronized SBI oscillations.