

In vivo imaging identifies temporal signature of D1 and D2 medium spiny neurons in cocaine reward

Erin S. Calipari^a, Rosemary C. Bagot^a, Immanuel Purushothaman^a, Thomas J. Davidson^b, Jordan T. Yorgason^c, Catherine J. Peña^a, Deena M. Walker^a, Stephen T. Pirpinias^a, Kevin G. Guise^a, Charu Ramakrishnan^b, Karl Deisseroth^{b,d,e}, and Eric J. Nestler^{a,1}

^aFishberg Department of Neuroscience and Friedman Brain Institute, Icahn School of Medicine at Mount Sinai, New York, NY 10029; ^bDepartment of Bioengineering, Stanford University, Stanford, CA 94305; ^cVollum Institute, Oregon Health and Science University, Portland, OR 97239; ^dHoward Hughes Medical Institute, Stanford University, Stanford, CA 94305; and ^eDepartment of Psychiatry & Behavioral Sciences, Stanford University, Stanford, CA 94305

Edited by Susan G. Amara, National Institutes of Health, Bethesda, MD, and approved January 8, 2016 (received for review October 27, 2015)

The reinforcing and rewarding properties of cocaine are attributed to its ability to increase dopaminergic transmission in nucleus accumbens (NAc). This action reinforces drug taking and seeking and leads to potent and long-lasting associations between the rewarding effects of the drug and the cues associated with its availability. The inability to extinguish these associations is a key factor contributing to relapse. Dopamine produces these effects by controlling the activity of two subpopulations of NAc medium spiny neurons (MSNs) that are defined by their predominant expression of either dopamine D1 or D2 receptors. Previous work has demonstrated that optogenetically stimulating D1 MSNs promotes reward, whereas stimulating D2 MSNs produces aversion. However, we still lack a clear understanding of how the endogenous activity of these cell types is affected by cocaine and encodes information that drives drug-associated behaviors. Using fiber photometry calcium imaging we define D1 MSNs as the specific population of cells in NAc that encodes information about drug associations and elucidate the temporal profile with which D1 activity is increased to drive drug seeking in response to contextual cues. Chronic cocaine exposure dysregulates these D1 signals to both prevent extinction and facilitate reinstatement of drug seeking to drive relapse. Directly manipulating these D1 signals using designer receptors exclusively activated by designer drugs prevents contextual associations. Together, these data elucidate the responses of D1- and D2-type MSNs in NAc to acute cocaine and during the formation of context–reward associations and define how prior cocaine exposure selectively dysregulates D1 signaling to drive relapse.

cocaine | reward | medium spiny neuron | calcium imaging | associative learning

Drug seeking associated with addiction is mediated in part by strong associations between the rewarding effects of drugs and the environment within which they are administered (1). The nucleus accumbens (NAc) is a key neural substrate of context–reward associations because it mediates multiple aspects of this process, including learning, selecting, and executing goal-oriented behaviors (2–5). Over time, contextual cues in the absence of drug can elicit neural responses that resemble that of the drug itself and trigger drug craving and seeking, resulting in relapse in drug-addicted individuals (1). Furthermore, an inability to extinguish previously formed associations is thought to contribute to pathological drug seeking that persists despite periods of abstinence (6–9). Thus, identifying the specific neuronal populations that drive these behaviors, and determining therapeutic strategies to manipulate the cell populations to promote extinction of these associations, would be advantageous to improving treatment outcomes in drug-addicted individuals.

Increased activity of the mesolimbic dopamine system is a central mechanism underlying the reinforcing and rewarding actions of drugs of abuse, including cocaine, as well as the compulsive drug seeking that develops over time and characterizes an addicted state (10–12). Dopamine action in NAc is mediated predominantly via activation of D1 or D2 dopamine receptors that are expressed by

largely nonoverlapping populations of medium spiny neurons (MSNs) (13). These two subtypes of MSNs exert opposite effects on behavior, with optogenetic activation of D1-type neurons promoting positive reinforcement and increasing the formation of cocaine reward–context associations and activation of D2-type neurons being aversive and decreasing cocaine reward (14, 15); related differences in behavioral responses are seen in response to D1 vs. D2 receptor agonists or antagonists (16). Changes in NAc MSN activity in vivo have been documented in response to acute and chronic cocaine (17–19); however, it has not been possible to obtain this information in a cell-type-specific manner.

The aim of the current study is to overcome this major gap in knowledge and define changes in D1- and D2-type MSN activity in vivo driven by acute and chronic exposures to cocaine and to determine how these cell types respond during context–reward associations. Thus, whereas the optogenetic studies cited above define the consequences of D1 or D2 MSN stimulation on reward-related behaviors, how the cell types respond under physiological conditions remains unknown. Furthermore, because optogenetic stimulations are experimenter-defined they eliminate the temporally specific nature of the endogenous signaling that encodes information. Advances in Ca²⁺ imaging have made it possible to record from genetically distinct subpopulations of neurons in awake behaving animals (20, 21). Here, we combine Ca²⁺ imaging and fiber photometry with conditioned place preference (CPP) to establish the patterns of activity of D1- and D2-type MSNs in NAc during formation of reward–context associations and to determine how these patterns are dysregulated by prior chronic exposure to cocaine. We then causally establish

Significance

Strong associations between cocaine and the environmental contexts where cocaine is administered are thought to drive relapse. The nucleus accumbens (NAc) encodes these cue–reward associations, and here we determined how cocaine alters the ability of cells in NAc to respond to drug-associated environmental stimuli to drive drug seeking. Using fiber photometry calcium imaging we define the specific population of cells, dopamine D1 receptor-expressing neurons, that encodes information about drug associations and show that these cells can be manipulated to attenuate the strength of drug associations and prevent relapse. Together, these data define a basic circuit mechanism underlying drug–context associations and suggest that pharmacotherapeutic agents aimed at D1-type neurons may help to promote sustained abstinence in cocaine abusers.

Author contributions: E.S.C. and E.J.N. designed research; E.S.C., R.C.B., I.P., C.J.P., D.M.W., and S.T.P. performed research; I.P., T.J.D., J.T.Y., K.G.G., C.R., and K.D. contributed new reagents/analytic tools; E.S.C. analyzed data; and E.S.C. and E.J.N. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

¹To whom correspondence should be addressed. Email: eric.nestler@mssm.edu.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1521238113/-DCSupplemental.

the role of altered firing of a given cell type to key aspects of context–reward associations by use of chemogenetic approaches.

Results

Biphasic Ca^{2+} Responses in NAc to Cocaine-Associated Contextual Cues.

To examine real-time activity of NAc neurons in vivo during associative learning for drug rewards, we injected wild-type C57BL/6J mice with AAV-GCaMP6f—which infects all types of neurons but not nonneuronal cells—into NAc core and recorded Ca^{2+} transients through an optic fiber (20) while mice were trained for CPP (Fig. 1 *A* and *B*). During pairing, cocaine reduced the frequency of Ca^{2+} transients in NAc (Fig. 1 *C* and *D*). In a choice test 24 h after the final pairing, animals spent more time on the side that was previously paired with drug, indicating that cocaine had induced a place preference via an association between the context and the rewarding effects of cocaine (Fig. 1*E*). Robust spikes in Ca^{2+} activity were recorded immediately preceding entry into the drug-paired, but not saline-paired, chamber (Fig. 1 *F* and *G*).




Fig. 1. Biphasic responses of NAc neurons to cocaine-associated cues. (A) Timeline of the experimental design. (B) Viral expression of AAV-GCaMP6f and placement of the fiber-optic probe in NAc core. (C) Representative Ca^{2+} traces from a single animal over pairing sessions. Data are represented as the percent change in fluorescence over the mean fluorescence ($\Delta F/F$). (D) Peak analysis of Ca^{2+} imaging traces. Cocaine reduces the number of events [Student's *t* test; $t(5) = 3.48$, $P < 0.05$, $n = 6$]. (E) (Left) Animals formed a preference for the chamber associated with cocaine [Student's *t* test; $t(5) = 4.04$, $P < 0.01$, $n = 6$]. (Right) Heat maps showing time spent in each area of the chamber. (F) Representative traces demonstrating the temporal profile of NAc activity around paired and unpaired chamber entry. (G) Quantification of the peak amplitude of the Ca^{2+} signal in the five seconds preceding paired and unpaired chamber entry [Student's *t* test; $t(5) = 4.38$, $P < 0.01$, $n = 6$]. (H) Representative trace showing the change in event frequency when the animal is in the paired or unpaired chamber. (I) Quantification of event frequency [Student's *t* test; $t(5) = 2.68$, $P < 0.05$, $n = 6$]. * $P < 0.05$, ** $P < 0.01$.

Further, firing was suppressed while the mouse remained in the drug-paired chamber (Fig. 1 *H* and *I*).

Specific Contribution of D1 and D2 MSNs to Learning Cocaine-Associated Cues.

These changes in NAc neuronal activity presumably reflect altered firing of MSNs, which comprise >95% of all NAc neurons. To determine the MSN subtypes that mediate the two distinct phases of NAc activity during context–reward associations, we injected AAV-DIO-GCaMP6f into NAc of mice expressing Cre-recombinase in either D1 or D2 MSNs to induce GCaMP6f expression specifically in each cell type (Fig. S1). At baseline, D2 MSNs displayed greater than fivefold higher Ca^{2+} transient frequency than D1 MSNs (Fig. 2 *C–E*). Expression levels of GCaMP6f were not different between the neuron subtypes, indicating that the effect is physiological and not an artifact of the mouse line or viral expression (Fig. S1 *A* and *B*). Because MSNs make up the vast majority of NAc neurons, the most parsimonious explanation of these findings is that D2 MSNs are more active at baseline, although one important caveat is that D2 receptors are also expressed by certain NAc interneurons. Single cocaine injections, during training, increased D1 MSN Ca^{2+} transient frequency, while reducing D2 frequency (Fig. 2*A*). During the subsequent choice test (Fig. 2*B*), entry into the drug-paired context similarly increased D1 and decreased D2 firing frequency (Fig. 2 *C* and *D*), eliciting the same physiological response as cocaine had previously, suggesting that these contextual cues acquire rewarding properties (22). D1 and D2 MSNs displayed distinct temporal profiles of firing: increased D1 MSN activity immediately preceded entry into the drug-paired context (Fig. 2*C* and *Movie S1*), whereas D2 activity was suppressed only after entry (Fig. 2*D* and *Movie S2*). It is possible that D1 activity drives the motivation to enter a paired compartment, with the reduced activity of D2 cells after entry driving the motivation to remain in that compartment. Further, these data suggest that the biphasic neural signature of global NAc neuronal activity (Fig. 1 *H* and *I*) is mediated by the two distinct populations of MSNs. Additionally, during choice testing, D1 MSN peak amplitude in the 5 s preceding entry into the drug-paired context correlated with time spent in the drug-paired side (Fig. 2*E*), suggesting that D1, but not D2 (Fig. 2*F*), signaling specifically drives expression of place preference.

Cocaine Pretreatment Alters D1 MSN Signaling in Association with Reduced Extinction.

Animals were injected for 7 d with 10 mg/kg i.p. cocaine in their home cage followed by 7 d of withdrawal (Fig. S2) and then baseline MSN activity was recorded followed by CPP (Fig. 3*A*). Prior cocaine administration selectively increased baseline D1 MSN activity without affecting D2 activity (Fig. S3). Further, although cocaine administration did not alter initial CPP strength (Fig. 3 *B* and *C*) it impaired later extinction of the place preference, indicating a latent potentiation of the cocaine–context association (Fig. 3 *C* and *D*). This effect associated with heightened D1 activity (Fig. 3*E*) preceding drug-paired chamber entry that uniquely persisted throughout extinction in cocaine-pre-exposed animals (Fig. 3*F*). Cocaine pretreatment did not alter the magnitude of D2 MSN responses to the drug-associated contexts during choice testing, extinction, or reinstatement (Fig. S4). These findings suggest that cocaine experience alters associative learning processes by selectively enhancing D1 MSN activity, to potentiate the association between drugs and cues and promote relapse during periods of abstinence.

Cocaine Pretreatment Facilitates Reinstatement of Conditioned Place Preferences.

Previous work has shown that prior cocaine exposure induces time-dependent increases in cocaine seeking (6, 23). We found that a subthreshold challenge dose of cocaine [5 mg/kg i.p.; i.e., a dose insufficient to induce preference in cocaine-naïve animals (24)] induced reinstatement of CPP only in animals with a history of prior chronic cocaine exposure and that the reinstated preference exceeded the initial preference for the drug-paired chamber (Fig. 3*D*). This increased preference was accompanied by an augmented spike in D1, but no change in D2




Fig. 2. Specific contribution of D1 and D2 NAc MSNs to learning cocaine-associated cues. (A) Representative Ca^{2+} traces and peak analysis from a D1-Cre (Left, green) and D2-Cre (Right, blue) mouse showing higher activity of D2-MSNs under baseline conditions. Cocaine increases D1 events [Student's t test; $t(3) = 5.84$, $P < 0.05$, $n = 4$] and decreases D2 events [Student's t test; $t(4) = 2.92$, $P < 0.05$, $n = 4$] in NAc. (B) (Left) Animals formed a conditioned preference [Student's t test; $t(21) = 3.96$, $P < 0.001$, $n = 22$]. (Right) Heat maps showing time spent in each area of the CPP chamber. (C) (Top) Representative heat map of D1-MSN-mediated Ca^{2+} signaling during successive entries into the drug-paired chamber. (Bottom) Averaged D1 traces (also see Movie S1). Quantification of peak amplitude of Ca^{2+} events 5 s around entry [Student's t test; $t(10) = 2.61$, $P < 0.05$, $n = 11$]. (D) (Top) Heat map of D2-MSN-mediated Ca^{2+} signaling during successive entries into the drug-paired chamber. (Bottom) Averaged D2 traces (also see Movie S2). Quantification of peak amplitude of Ca^{2+} events [Student's t test; $t(9) = 2.70$, $P < 0.05$, $n = 10$]. (E) Correlation analysis showing the relationship between CPP and the amplitude of D1 events preceding drug-paired chamber entry ($r = 0.63$, $P < 0.05$, $n = 11$). (F) Correlation analysis showing no relationship with D2 events ($r = -0.15$, $P =$ not significant, $n = 10$). * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

(Fig. S4), MSN activity preceding drug-paired chamber entry and this persisted over many entries (Fig. 3 G and H). Further, this potentiated reinstatement was associated with selective enhancement of the acute effects of cocaine—opposite in the two MSN subtypes—on both D1 MSNs (Fig. 3 I and J) and D2 MSNs (Fig. S5), only when the animal was in the drug-paired context. Unpaired chamber entry associated with consistent suppression of D1 activity, an effect not seen in animals without prior cocaine experience, and this was accompanied by reduced time spent in the unpaired chamber (Fig. 3 I and J). Thus, chronic cocaine produces a state in which cocaine-associated contexts activate the same neuronal mechanisms as acute cocaine administration, which may be a primary mechanism by which repeated cocaine administration strengthens cue–reward associations that drive relapse, even following periods of extended abstinence (6).

D1 MSN Activity Is Required for the Acquisition and Expression of Cocaine Conditioned Place Preferences. To confirm this causal role of D1 MSNs in associative learning, we expressed the inhibitory designer receptors exclusively activated by designer drugs (DREADD), hM4Di, selectively in D1 MSNs and administered clozapine N-oxide (CNO; 5 mg/kg i.p.) before cocaine pairing to inhibit D1 activity (Fig. 4). CNO administration at this dose inhibited D1-specific Ca^{2+} activity (Fig. 4B). When CNO was administered 1 h before each cocaine pairing it blocked the cocaine-induced increases in D1 activity (Fig. 4 C and D). Further, it blocked both the preference for the drug-paired chamber as well as the spike in D1 MSN activity that preceded entry into the drug-paired chamber (Fig. 4 E–G), showing that this signal is indeed required for associative learning for drug rewards.

To assess whether the D1 signal is also causal for the decision to enter the drug-paired chamber, we administered CNO on

choice test day (Fig. 4H). CNO administration completely abolished the temporally specific increase in D1 MSN activity that is associated with entry into the drug-paired chamber as well as the expression of a place preference (Fig. 4 I–K). Further, when animals were examined in a choice test 2 wk later, long after CNO had cleared, preference scores remained down (Fig. 4K), suggesting that inhibiting D1 firing in the presence of the associated context is sufficient to extinguish the association indefinitely. This could reflect enhanced extinction learning or blockade of reconsolidation. Regardless, the effect was specific for D1 MSNs: Inhibiting D2 MSN activity during the choice test had no effect on cocaine place preference (Fig. S6).

Discussion

Results of the present study demonstrate distinct patterns of D1 and D2 MSN signaling in NAc during cocaine reward learning, extinction, and reinstatement and provide fundamentally new insight into the circuit basis of drug–cue associations and drug seeking. We show that D2 MSNs exhibit manifold higher activity than D1 MSNs in NAc at baseline, that acute cocaine administration enhances D1 and suppresses D2 MSN activity, and that cocaine-induced facilitation of D1 MSN activity is required for formation of cocaine–context associations. Further, temporally precise, cell-type-specific signaling encodes contextual information about cocaine experiences such that increased D1 activity precedes entry into a drug-paired context, with decreased D2 activity occurring only after entry. Further, prior chronic exposure to cocaine impairs extinction of contextual associations by preventing the concomitant extinction of D1 MSN signaling that precedes drug-paired context entry. Inhibiting this D1 signal by DREADD-induced D1 MSN inhibition blocked the expression of conditioned preference, confirming that D1 signaling is the critical mediator of




Fig. 3. Chronic cocaine administration alters D1 MSN signaling in association with reduced extinction and facilitated reinstatement of CPP. (A) Experimental timeline. (B) Animals form a preference for the drug-paired chamber [two-way ANOVA; $F(1, 88) = 14.17, P < 0.001, n = 14$]; cocaine pretreatment does not change CPP. (C) Heat maps showing time spent in each chamber. (D) Chronic cocaine administration impairs extinction and facilitates cocaine-primed reinstatement [two-way ANOVA; $F(1, 147) = 20.08, P < 0.0001, n = 20$]. Data plotted as percent change from the original choice test. During extinction test 1, cocaine pretreated animals increased their preference (one-sample t test; $t_{22} = 2.56, P < 0.05$). During reinstatement, cocaine-pretreated animals reinstated above their original choice test values (one-sample t test; $t_{22} = 2.09, P < 0.05$). (E) Representative traces averaged over entries showing increased amplitude of D1 MSN Ca^{2+} activity at paired chamber entry during extinction. (F) Peak amplitude analysis. Saline-treated animals extinguish D1 MSN responses; cocaine pretreated animals do not [two-way ANOVA; $F(1, 17) = 4.492, P < 0.05, n = 10$]. (G) Representative traces averaged over entries showing increased amplitude of D1 MSN Ca^{2+} activity at paired chamber entry during cocaine-primed reinstatement. (H) Peak amplitude analysis over trials. Cocaine-treated animals reinstate D1 MSN responses, whereas saline-pretreated animals do not [two-way ANOVA; $F(1, 12) = 5.262, P < 0.05, n = 6, 9$]. (I) Representative D1 MSN traces from a cocaine-treated (Left) and saline-treated (Right) animal during reinstatement. Acute effects of the cocaine challenge are augmented in cocaine pretreated animals, but only when the animal is in the paired chamber. (J) Peak analysis: cocaine effects are enhanced in cocaine-pretreated animals in the previously drug-paired context [two-way ANOVA; $F(1, 4) = 22.16, P < 0.01, n = 3$]. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, #### $P < 0.0001$.

drug-context learning. More importantly, reducing the activity of D1 MSNs is sufficient to block (i.e., extinguish) the expression of preference, an effect that persists for weeks. Together, these data elucidate cell-type-specific engrams that contribute to the encoding of cocaine reward, and a D1 MSN-mediated mechanism by which cocaine exposure retards extinction of drug-cue associations to promote relapse.

The precise temporal profile of the encoding of cue-reward associations in MSNs provided here now enables elucidation of the circuit-wide signaling that controls these responses to drive reward- and reinforcement-related behaviors. The majority of work outlining the neural networks underlying cue-reward associations for both natural and drug rewards has shown that the ventral tegmental area (VTA) and NAc exhibit temporally specific signaling in response to unexpected rewards (25). However, once animals have learned the association between cues and reward availability the neural response transfers to the reward-predictive cue, thus guiding behavior toward environments that will result in obtaining that reward (26). Studies

using *in vivo* pharmacological approaches have demonstrated differential roles of NAc D1 and D2 receptors in drug conditioning by use of selective receptor agonists or antagonists, further supporting a role for both dopamine and D1 and D2 MSN subtypes in associative learning (27). Whereas this work has focused on the VTA-to-NAc dopamine circuit, tracking postsynaptic responses in NAc MSNs is particularly important because they integrate information not only from VTA dopamine neurons but also from numerous glutamatergic projections (28, 29). From a network perspective, D1 and D2 MSNs receive inputs from several regions known to encode and store information about context or context-drug associations such as the prefrontal cortex, basolateral amygdala, and hippocampus (30). Although dopamine plays an integral role in controlling motivated behaviors, its effects are largely modulatory, and it is the glutamatergic inputs that provide the excitatory drive for these circuits. Thus, interactions between dopamine and glutamate are critical in the case of drugs of abuse where dopaminergic involvement changes over the course of drug administration. Initially,




Fig. 4. D1 MSN activity is required for the expression of cocaine CPP. (A) Confocal image showing hM4Di and GCaMP6f coexpression in NAc. (B) CNO reduces D1-mediated Ca^{2+} activity. (C) Representative D1 Ca^{2+} traces from mCherry (red) and hM4Di (black) animals following cocaine (10 mg/kg i.p.). (D) Peak analysis of D1 MSNs. CNO reduces the frequency of events [Student's *t* test; $t(5) = 4.87$, $P < 0.01$, $n = 6$]. (E) Animals were injected with CNO (5 mg/kg i.p.) before pairing sessions. (F) Representative traces averaged over entries. (G) Time spent in each area of the CPP chamber. (H) Animals form a preference for the drug-paired chamber, with CNO reducing the time spent in the drug-paired chamber [two-way ANOVA; $F(1, 28) = 4.55$, $P < 0.05$, $n = 8$]. (I) Animals were injected with CNO before choice test. (J) Representative traces averaged over entries showing increased amplitude of D1 MSN Ca^{2+} activity at paired chamber entry. (K) Time spent in each area of the CPP chamber. (L) Animals form a preference for the drug-paired chamber, with CNO reducing the time spent in the drug-paired chamber. The reduced preference in hM4Di animals remained 2 wk after the initial choice test [two-way ANOVA; $F(1, 37) = 5.32$, $P < 0.05$, $n = 5, 10$]. * $P < 0.05$, ** $P < 0.01$.

dopaminergic responses to cues predicting drug availability drive drug seeking; however, over time, dopaminergic responses to cocaine-associated cues are blunted, whereas sensitization occurs to the motivational aspects of drug seeking and taking (28). For example, despite blunted dopaminergic responses, multiunit recordings have shown that enhanced cue-induced responses in NAc remain intact (17, 19). Further, recent findings have shown that repeated cocaine injections strengthen glutamatergic inputs onto D1 MSNs specifically (31), which could underlie some of the changes in cue responsivity that are seen in the current study. Together, these data suggest that, although dopamine is likely highly involved, it is the activity of the NAc, regardless of originating influences, that mediates these behaviors.

Although suppressing D1 MSN activity during CPP testing reduced the choice to enter the drug-paired chamber, we were surprised to find that this manipulation also eliminated the association indefinitely. Subsequent testing showed that blocking D1 MSNs while the animal was exposed to the cocaine-paired context facilitated the extinction of these associations. Thus, it is likely that the D1 signal not only guides animals to seek the reward, driving them to enter the drug-paired context, but the lack of signal also functions to update new information as associations change. In other words, eliminating the D1 signal in the drug-paired context indicates that the association no longer exists. The ability to attenuate these associations is important because we show here that prior chronic cocaine exposure inhibits the ability of this signal to be extinguished. These data suggest that pharmacological intervention could help to eliminate or attenuate these associations in drug-addicted individuals in a way that would greatly improve treatment outcomes.

Prior chronic cocaine exposure not only inhibits extinction of place conditioning but also facilitates reinstatement of place conditioning. Similar dysregulated associative learning concerning environmental stimuli and rewards occurs in human drug addicts, and an inability to extinguish previously formed associations is thought to contribute to pathological drug seeking as well as relapse (6, 7). Our findings provide a neural mechanism for the deficits in extinction and enhanced reinstatement, whereby cocaine treatment (*i*) dysregulates the ability of D1 MSNs to respond to context information and update information when the association no longer occurs and (*ii*) enhances the effects of challenge doses of cocaine on D1 MSN signaling only when animals are in the drug-paired context. These data show that, in addition to the contextual cues eliciting a temporally specific response, the context in which drug is administered can greatly enhance the pharmacological effects of the drug, as has been demonstrated in many other settings (32, 33). Enhancement of the pharmacological effects of drugs is likely one mechanism by which the presentation of drug-paired cues can increase the motivation to administer cocaine.

Our data highlight the important role played by D1 MSNs in NAc core in establishing context-reward associations and in controlling the strength of these associations after cocaine exposure. This prominent influence of D1 MSNs in cocaine action is consistent with previous studies that have shown that chronic cocaine exposure alters D1 MSNs selectively via both changes in dendritic morphology and function as well as changes in gene transcription and epigenetic mechanisms that last long into the abstinence period (34–36). Further work is needed to determine whether similar patterns of D1 MSN activity in NAc core as shown here are also

observed in NAc shell and to establish the function of cocaine-induced changes in D2 MSNs that have also been demonstrated.

Here we show that regulation of associative learning, and its dysregulation by cocaine, is driven primarily through alterations in D1 MSNs in NAc core, which both impair the extinction of previously learned associations and enhance reinstatement following abstinence. The magnitude of D1 firing to reward-associated cues may play a critical role in determining the perceived value of a reward and bias decision making toward outcomes most likely to result in obtaining that reward. Further, we show that inhibiting D1 MSNs in NAc is sufficient to prevent preference for contexts previously paired with a drug reward, an effect that lasts indefinitely. These findings suggest that therapies targeted selectively to D1 MSNs may help to normalize D1 activity following a history of cocaine use and prevent relapse. Additionally, the causal role of temporally specific cue-elicited D1 signaling in the expression of learned associations demonstrates the importance of this cell type in the development of fundamental associative learning processes. Disruption of associative learning is a hallmark of diverse psychopathologies that span developmental stages. Demonstrating the causal role of NAc D1 MSN signaling in mediating these processes provides fundamental insight to guide the development of novel targeted therapeutic interventions not only for addiction but also for a wide range of psychiatric disorders.

Methods

See *SI Methods* for more detailed explanations.

Experimental Subjects. D1-Cre and D2-Cre BAC transgenic mice on a C57BL/6J background were obtained from N. Heintz, P. Greengard, C. Gerfen, and National Institute of Neurological Disorders and Stroke/Gene Expression Nervous System Atlas (NINDS/GENSAT) (www.gensat.org/index.html) from Rockefeller University/National Institute of Mental Health. C57BL/6J wild-type mice were obtained from The Jackson Laboratory (SN: 000664). All

experiments were conducted in accordance with the guidelines of the Institutional Animal Care and Use Committee at Icahn School of Medicine at Mount Sinai.

Stereotaxic Virus Injection and Cannula Implantation. AAVdj-CaMKII α -GCaMP6f-WPRE (for pan-neuronal recordings), AAVdj-EF1 α -DIO-GCaMP6f-WPRE (for cell-type-specific recording), or AAV5/hSyn-DIO-hm4Di-mcherry [University of North Carolina at Chapel Hill (37), for DREADD inhibition studies] was infused into NAc core. Chronically implantable optic fibers (Doric Lenses) were positioned above the injection site.

Fiber Photometry Ca²⁺ Imaging. Fiber photometry uses the same fiber to both excite and record from GCaMP in real time. The system used two light-emitting diodes at 490 and 405 (Thor Labs), reflected off dichroic mirrors (Semrock, FF495), and coupled into a 400- μ m 0.48 N.A. optical fiber (BFH48-600; Thorlabs). Analysis of the resulting signal was performed with custom-written MATLAB software.

Behavioral Testing. CPP was performed in a rectangular apparatus consisting of two side chambers with different contextual cues. Mice were paired with cocaine in one of the chambers and saline in the other chamber. After pairing, during a choice test, mice were allowed to freely explore the entire apparatus and preference for each chamber was determined. During extinction animals were placed into the chamber with free access. For reinstatement, animals were injected with 5 mg/kg cocaine i.p. immediately before being placed into the chamber with free access. For DREADD experiments, CNO (5 mg/kg) was administered 1 h before the session.

Statistics. For groups of three or more, one-way ANOVA was run. For groups of two a two-tailed Student's *t* test was run. Significance was set at $P < 0.05$ for all tests. All data are expressed as mean \pm SEM.

ACKNOWLEDGMENTS. This work was supported by grants from the National Institute on Drug Abuse (R01DA014133 and P01DA008227 to E.J.N.) and Brain & Behavior Research Foundation Grant NARSAD 22713 (to R.C.B.).

- Phillips PE, Stuber GD, Heien ML, Wightman RM, Carelli RM (2003) Subsecond dopamine release promotes cocaine seeking. *Nature* 422(6932):614–618.
- Schultz W (2006) Behavioral theories and the neurophysiology of reward. *Annu Rev Psychol* 57:87–115.
- Steinberg EE, et al. (2013) A causal link between prediction errors, dopamine neurons and learning. *Nat Neurosci* 16(7):966–973.
- Schultz W, Dayan P, Montague PR (1997) A neural substrate of prediction and reward. *Science* 275(5306):1593–1599.
- du Hoffmann J, Nicola SM (2014) Dopamine invigorates reward seeking by promoting cue-evoked excitation in the nucleus accumbens. *J Neurosci* 34(43):14349–14364.
- Grimm JW, Hope BT, Wise RA, Shaham Y (2001) Neuroadaptation. Incubation of cocaine craving after withdrawal. *Nature* 412(6843):141–142.
- O'Brien CP, Childress AR, McLellan AT, Ehrman R (1992) Classical conditioning in drug-dependent humans. *Ann N Y Acad Sci* 654:400–415.
- Reid AG, Lingford-Hughes AR, Canela LM, Kalivas PW (2012) Substance abuse disorders. *Handb Clin Neurol* 106:419–431.
- Xue YX, et al. (2012) A memory retrieval-extinction procedure to prevent drug craving and relapse. *Science* 336(6078):241–245.
- Di Chiara G, Imperato A (1988) Drugs abused by humans preferentially increase synaptic dopamine concentrations in the mesolimbic system of freely moving rats. *Proc Natl Acad Sci USA* 85(14):5274–5278.
- Thomas MJ, Kalivas PW, Shaham Y (2008) Neuroplasticity in the mesolimbic dopamine system and cocaine addiction. *Br J Pharmacol* 154(2):327–342.
- Dong Y, et al. (2004) Cocaine-induced potentiation of synaptic strength in dopamine neurons: Behavioral correlates in GluRA(-/-) mice. *Proc Natl Acad Sci USA* 101(39):14282–14287.
- Gerfen CR, Surmeier DJ (2011) Modulation of striatal projection systems by dopamine. *Annu Rev Neurosci* 34:441–466.
- Lobo MK, et al. (2010) Cell type-specific loss of BDNF signaling mimics optogenetic control of cocaine reward. *Science* 330(6002):385–390.
- Kravitz AV, Tye LD, Kreitzer AC (2012) Distinct roles for direct and indirect pathway striatal neurons in reinforcement. *Nat Neurosci* 15(6):816–818.
- Self DW, Barnhart WJ, Lehman DA, Nestler EJ (1996) Opposite modulation of cocaine-seeking behavior by D1- and D2-like dopamine receptor agonists. *Science* 271(5255):1586–1589.
- Carelli RM, King VC, Hampson RE, Deadwyler SA (1993) Firing patterns of nucleus accumbens neurons during cocaine self-administration in rats. *Brain Res* 626(1-2):14–22.
- Wheeler RA, Carelli RM (2009) Dissecting motivational circuitry to understand substance abuse. *Neuropharmacology* 56(Suppl 1):149–159.
- Hollander JA, Carelli RM (2007) Cocaine-associated stimuli increase cocaine seeking and activate accumbens core neurons after abstinence. *J Neurosci* 27(13):3535–3539.
- Gunaydin LA, et al. (2014) Natural neural projection dynamics underlying social behavior. *Cell* 157(7):1535–1551.
- Cui G, et al. (2013) Concurrent activation of striatal direct and indirect pathways during action initiation. *Nature* 494(7436):238–242.
- Crombag HS, Bossert JM, Koya E, Shaham Y (2008) Review. Context-induced relapse to drug seeking: A review. *Philos Trans R Soc Lond B Biol Sci* 363(1507):3233–3243.
- Ahmed SH, Koob GF (1998) Transition from moderate to excessive drug intake: Change in hedonic set point. *Science* 282(5387):298–300.
- Vialou V, et al. (2012) Serum response factor and cAMP response element binding protein are both required for cocaine induction of Δ FosB. *J Neurosci* 32(22):7577–7584.
- Humphries MD, Prescott TJ (2010) The ventral basal ganglia, a selection mechanism at the crossroads of space, strategy, and reward. *Prog Neurobiol* 90(4):385–417.
- Fiorillo CD, Tobler PN, Schultz W (2003) Discrete coding of reward probability and uncertainty by dopamine neurons. *Science* 299(5614):1898–1902.
- Graham DL, Hoppenot R, Hendryx A, Self DW (2007) Differential ability of D1 and D2 dopamine receptor agonists to induce and modulate expression and reinstatement of cocaine place preference in rats. *Psychopharmacology (Berl)* 191(3):719–730.
- Lammel S, Lim BK, Malenka RC (2014) Reward and aversion in a heterogeneous midbrain dopamine system. *Neuropharmacology* 76(Pt B):351–359.
- Russo SJ, et al. (2010) The addicted synapse: Mechanisms of synaptic and structural plasticity in nucleus accumbens. *Trends Neurosci* 33(6):267–276.
- Robbins TW, Everitt BJ (1996) Neurobehavioural mechanisms of reward and motivation. *Curr Opin Neurobiol* 6(2):228–236.
- MacAskill AF, Cassel JM, Carter AG (2014) Cocaine exposure reorganizes cell type- and input-specific connectivity in the nucleus accumbens. *Nat Neurosci* 17(9):1198–1207.
- Bell K, Duffy P, Kalivas PW (2000) Context-specific enhancement of glutamate transmission by cocaine. *Neuropsychopharmacology* 23(3):335–344.
- Anagnostaras SG, Schallert T, Robinson TE (2002) Memory processes governing amphetamine-induced psychomotor sensitization. *Neuropsychopharmacology* 26(6):703–715.
- Ishikawa M, et al. (2013) Dopamine triggers heterosynaptic plasticity. *J Neurosci* 33(16):6759–6765.
- Robison AJ, Nestler EJ (2011) Transcriptional and epigenetic mechanisms of addiction. *Nat Rev Neurosci* 12(11):623–637.
- Grueter BA, Robison AJ, Neve RL, Nestler EJ, Malenka RC (2013) Δ FosB differentially modulates nucleus accumbens direct and indirect pathway function. *Proc Natl Acad Sci USA* 110(5):1923–1928.
- Urban DJ, Roth BL (2015) DREADDs (designer receptors exclusively activated by designer drugs): Chemogenetic tools with therapeutic utility. *Annu Rev Pharmacol Toxicol* 55:399–417.