



Indian annual ambient air quality standard is achievable by completely mitigating emissions from household sources

Sourangsu Chowdhury^{a,b}, Sagnik Dey^{a,c,1}, Sarath Guttikunda^d, Ajay Pillariseti^b, Kirk R. Smith^{b,e,1}, and Larry Di Girolamo^f

^aCentre for Atmospheric Sciences, Indian Institute of Technology Delhi, New Delhi 110016, India; ^bSchool of Public Health, University of California, Berkeley, CA 94720-7360; ^cSchool of Public Policy, Indian Institute of Technology Delhi, New Delhi 110016, India; ^dUrban Emissions, New Delhi 110019, India; ^eCollaborative Clean Air Policy Centre, Delhi 110003, India; and ^fDepartment of Atmospheric Sciences, University of Illinois at Urbana-Champaign, Urbana, IL 61801

Contributed by Kirk R. Smith, March 11, 2019 (sent for review January 17, 2019; reviewed by Achuthan Jayaraman and Christine Wiedinmyer)

Exposures to ambient and household fine-particulate matter (PM_{2.5}) together are among the largest single causes of premature mortality in India according to the Global Burden of Disease Studies (GBD). Several recent investigations have estimated that household emissions are the largest contributor to ambient PM_{2.5} exposure in the country. Using satellite-derived district-level PM_{2.5} exposure and an Eulerian photochemical dispersion model CAMx (Comprehensive Air Quality Model with Extensions), we estimate the benefit in terms of population exposure of mitigating household sources—biomass for cooking, space- and water-heating, and kerosene for lighting. Complete mitigation of emissions from only these household sources would reduce India-wide, population-weighted average annual ambient PM_{2.5} exposure by 17.5, 11.9, and 1.3%, respectively. Using GBD methods, this translates into reductions in Indian premature mortality of 6.6, 5.5, and 0.6%. If PM_{2.5} emissions from all household sources are completely mitigated, 103 (of 597) additional districts (187 million people) would meet the Indian annual air-quality standard (40 μg m⁻³) compared with baseline (2015) when 246 districts (398 million people) met the standard. At 38 μg m⁻³, after complete mitigation of household sources, compared with 55.1 μg m⁻³ at baseline, the mean annual national population-based concentration would meet the standard, although highly polluted areas, such as Delhi, would remain out of attainment. Our results support expansion of programs designed to promote clean household fuels and rural electrification to achieve improved air quality at regional scales, which also has substantial additional health benefits from directly reducing household air pollution exposures.

air pollution | PM_{2.5} | cooking | lighting | heating

Chronic exposure to PM_{2.5} (i.e., particulate matter smaller than 2.5 μm) has detrimental effects on human health, including premature mortality (1–7). Multiple studies have established evidence of causal associations between exposure to PM_{2.5} and cardiorespiratory health endpoints, including chronic obstructive pulmonary disease, ischemic heart disease, stroke, lung cancer, and child acute lower respiratory infection. Annual PM_{2.5} exposure in India has been increasing in recent years (8, 9), primarily due to the rapid rise in pollution during the dry season (October–January). In 2017, this rise in pollution led the Indian Medical Association to declare a public health emergency in the National Capital Region of Delhi. Given the current situation, strict mitigation measures need to be implemented in India to meet at least the National Ambient Air Quality Standard (NAAQS) of 40 μg m⁻³. Formulation of a successful clean-air management plan requires understanding the relative contributions of individual sources to average ambient PM_{2.5} exposure at regional scales.

Several recent studies have attributed the largest share of ambient PM_{2.5} exposure in India to household emissions, as opposed to vehicles, power plants, industries, and other sectors (10–14). Four major activities contribute mainly to household emissions: biomass burned for residential cooking, space- and water heating,

and kerosene used for lighting. Relatively small amounts of coal are used in households nationally and use of kerosene for cooking has declined substantially in recent years. Different policy frameworks are likely required to mitigate each activity and/or source.

In this work, we develop seven scenarios based on current and plausible future mitigation policies and the estimated adaptive capacity of the exposed population. We quantify the contributions of individual household activities to ambient PM_{2.5} exposure in India and examine the expected health benefits of scenarios to reduce emissions from these sources. We also identify the districts (equivalent to counties in other countries) that would meet the Indian NAAQS and World Health Organization (WHO) ambient air quality guidelines after complete mitigation of these household sources.

Scenario Development. The seven scenarios are defined in Table 1. We assess the potential benefit of each scenario by estimating (i) the decrease in population-weighted ambient PM_{2.5} exposure and (ii) the averted premature mortality burden with respect to the 2015 annual population-weighted ambient PM_{2.5}, which is referred to as “baseline.”

The first scenario (P_C) assumes that the government successfully implements a policy to provide clean fuel for cooking to all

Significance

The Indian government has initiated programs to reduce household use of biomass cookfuels, but their motivation does not include ambient pollution reductions. We estimate ambient exposure reductions that result and subsequent health benefits of seven different scenarios that mirror plausible mitigation policies to address household energy needs currently met by biomass combustion for cooking and water- and space heating, and by kerosene for lighting. If all households transitioned to clean fuels, ~13% of premature mortality in India could be averted from the reduction in ambient pollution, and the average ambient PM_{2.5} concentration in the country would meet national PM_{2.5} standards. These results will assist in assessing current policies and making decisions on future mitigation policies to tackle air pollution.

Author contributions: S.C., S.D., S.G., K.R.S., and L.D.G. designed research; S.C. and S.D. performed research; S.C., S.D., S.G., A.P., and K.R.S. analyzed data; and S.C., S.D., S.G., A.P., and K.R.S. wrote the paper.

Reviewers: A.J., National Atmospheric Research Laboratory; and C.W., University of Colorado Boulder.

The authors declare no conflict of interest.

Published under the [PNAS license](#).

See Commentary on page 10614.

¹To whom correspondence may be addressed. Email: krksmith@berkeley.edu or sagnik@cas.iitd.ac.in.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1900888116/-DCSupplemental.

Published online April 15, 2019.

Table 1. Definition of the scenarios formulated for the study

Scenarios	Cooking (%)	Space	Water	Lighting (%)
		heating (%)	heating (%)	
P_C	100	0	0	0
P_{CL}	100	0	0	100
P_{CLS}	100	100	0	100
Ideal	100	100	100	100
Progress A	75	50	50	75
Progress M	50	25	25	50
Progress S	25	0	0	25

The numeric values depict the percent mitigation of the particular household source. C = cooking, L = lighting, S = space heating, A = aspirational progress, M = moderate progress, S = slow progress.

solid-fuel-using families in the country, but does not implement any policies to counter emissions from space- and water heating and kerosene lighting. The second scenario (P_{CL}) assumes that, on top of " P_C ," the government successfully implements policies that provide cheap and continuous electricity across India, which eradicates lighting via kerosene lamps in households.

We note that the Government of India (GoI) in 2016 launched Pradhan Mantri Ujjwala Yojana (15), which intends to provide 80 million "below poverty level" households with liquefied petroleum gas (LPG) by 2019. Under the program, about 75% of the planned LPG connections had been dispersed by January 2019 (16). GoI also initiated the Deen Dayal Upadhyay Grameen Jyoti Yojana (17) in 2014, which promises to electrify all villages in India, thus eradicating emissions from kerosene lighting. In view of these GoI mitigation schemes, it is expected that in the next few years solid-fuel cooking and kerosene lighting could be entirely replaced by clean fuels if the population adapts and adheres to the above-mentioned policies perfectly [which is often not the case, as discussed in previous studies (18, 19)].

" P_{CLS} " adds policies to mitigate space heating and assumes they are also efficiently undertaken such that there are no emissions of $PM_{2.5}$ from space-heating-related activities. Finally, the "ideal" scenario assumes full adoption of household air pollution mitigation policies, resulting in elimination of all major sources that result in household $PM_{2.5}$ pollution.

We acknowledge that "perfect" policies—ones that are fully distributed and implemented—in a socioeconomically and behaviorally diverse country like India are hypothetical and aspirational. To account for this, we develop three scenarios which consider three interim targets that set more realistic goals for reach and population uptake. "Progress A" assumes that policymakers and the population are fairly successful in achieving their mitigative and adaptative goals of eradicating usage of solid fuels in household activities. It assumes that 75% of all emission from residential cooking and lighting and 50% of all emission from space- and water heating in a district are eliminated. We note that although the transitions from traditional fuels to LPG and electric lighting are aspirational for many populations, it can be difficult to transition from usage of solid fuels for space- and water heating to clean fuels, as both activities consume large amounts of energy. Using clean fuels for these activities may not be perceived as cost-effective for the rural poor, justifying our lower expectation weights on space and water heating.

"Progress S" assumes "slow" progress of policies in terms of reaching communities and less complete adoption by targeted populations. It assumes that 25% of all emissions from residential cooking and lighting are mitigated and the government fails to devise any policy for mitigating space heating and water heating. "Progress M" assumes a moderate scenario, where 50% of the residential cooking and lighting are phased out and 25% of space- and water heating are eliminated.

Results

Population-weighted mean annual ambient $PM_{2.5}$ exposure in India in the baseline year, 2015, is estimated to have been $55 \mu g m^{-3}$. Exposures vary widely across India, with the largest ($>100 \mu g m^{-3}$) exposure observed over the Delhi National Capital Region (Delhi NCR). Fig. 1 displays exceedances of annual ambient $PM_{2.5}$ exposure relative to the NAAQS and WHO-IT (Interim Target) 1 ($35 \mu g m^{-3}$), WHO-IT2 ($25 \mu g m^{-3}$), WHO-IT3 ($15 \mu g m^{-3}$), and the WHO-AQG ($10 \mu g m^{-3}$) across India in 2015. Overall in India, 58, 67, 83, 97, and 99% of districts have annual $PM_{2.5}$ exposure exceeding these guidelines, respectively. In Northwest India and across the Gangetic Plain, all districts have annual exposures higher than the NAAQS. In Central and West India, 82 and 50% of districts exceed the NAAQS, while in North, Northeast, and South India, only 22, 19, and 2% districts violate the NAAQS, respectively.

Fig. 2 depicts the percentage contribution of household sources toward ambient $PM_{2.5}$. There is a strong north-south gradient in the percentage contribution of household $PM_{2.5}$ toward ambient $PM_{2.5}$. It exceeds 40% in most of the districts in the Gangetic basin, while in the southern and central states the contribution is below 30%. Residential cooking contributes 20–50% (Fig. 3) of all household $PM_{2.5}$ emission across all districts, with higher contributions in districts with greater proportions of the population living in rural areas, which is again pronounced in the Gangetic basin. Space heating contributes more than 40% of all household emission in a few regions of North, Northeast, South, and West India and in the Himalayan foothills. Water heating contributes up to 30% of ambient $PM_{2.5}$ exposure with a north to south spatial gradient. Kerosene lighting contributes up to 10% of total emissions, without much spatial gradient. Overall, cooking is the largest contributor among all household sources, followed by space heating, water heating, and lighting.

Mitigation of Emission from Various Household Sources. In 2015, the baseline year, 58% of districts in India had $PM_{2.5}$ exposures above the NAAQS. The corresponding numbers are 68, 83, 98, and 99% for the WHO-IT1, WHO-IT2, WHO-IT3, and WHO guidelines, respectively (Fig. 4A). If the hypothetical ideal scenario (as seen in Fig. 4A) can be achieved, 103 more districts, housing 187 million people, would meet the NAAQS and ambient $PM_{2.5}$ exposure would fall below WHO-IT1, IT2, IT3, and the WHO-AQG in 104, 75, 26, and 2 additional districts, respectively. Successfully achieving the less ambitious Progress A scenario would add 72 additional districts below NAAQS. The corresponding numbers for Progress M, Progress S, P_C , P_{CL} , and P_{CLS} are 39, 11, 55, 61, and 86, respectively. In North Indian districts that do not meet Indian NAAQS standard even after achievement of the ideal scenario ($n = 248$), the annual ambient $PM_{2.5}$ exposure comes within $10 \mu g m^{-3}$ of the NAAQS in 115 districts, especially in the central and eastern parts of the Gangetic Plain (Fig. 4B). Only the Delhi NCR and some districts of Northwest India remain out of attainment with the Indian NAAQS standard by $>30 \mu g m^{-3}$ —even under the ideal scenario. This may be due to the large share of dust (20, 21) and relatively larger (compared with the rest of the country) contributions of other outdoor sources (e.g., vehicular emission, crop waste, industry, and trash burning, etc.) to ambient $PM_{2.5}$ concentrations in this region. Fig. 4B summarizes the remaining reductions needed to meet the Indian NAAQS after implementation of an ideal scenario.

The population-weighted, all-India mean annual ambient $PM_{2.5}$ exposure decreases from 55 to $45 \mu g m^{-3}$ (a 17.6% decrease) if P_C is achieved and to $45 \mu g m^{-3}$ if the P_{CL} scenario is realized (Fig. 5). If space-heating emissions are additionally mitigated as in P_{CLS} , the all-India mean annual ambient $PM_{2.5}$ exposure drops to $41 \mu g m^{-3}$; and, further, to $38 \mu g m^{-3}$ (below the NAAQS) if the ideal scenario is achieved. Achieving the least ambitious scenario,




Fig. 1. Annual ambient PM_{2.5} exposure exceedances in India with reference to Indian NAAQS and WHO air-quality guidelines in the baseline year, 2015. Indian NAAQS: 40 $\mu\text{g m}^{-3}$; WHO-IT1: 35 $\mu\text{g m}^{-3}$; WHO-IT2: 25 $\mu\text{g m}^{-3}$; WHO-IT3: 15 $\mu\text{g m}^{-3}$; WHO-AQG: 10 $\mu\text{g m}^{-3}$. Statistics (as inset figures) are shown as percentage of districts in each region exceeding the standard/guidelines.

Progress S would only reduce the average population-weighted PM_{2.5} by 2 $\mu\text{g m}^{-3}$ from the baseline year. Pursuing the more ambitious pathways—Progress M and Progress A—would reduce the population-weighted PM_{2.5} exposure by 21 and 13%, respectively.

Expected Health Benefit. We estimate expected health benefits (Fig. 5) in terms of the percentage of premature deaths, relative to baseline premature mortality, that could be averted if our scenarios are implemented. Estimates are made with both the integrated exposure-response functions (22) and a linear risk function (*SI Appendix, Attribution Method*). Scenarios are described in detail in Table 1 and the results summarized in Fig. 5. Briefly, those labeled with a P represent complete displacement of unclean fuels for cooking (C), space heating (S), and lighting (L) in various combinations.

Premature mortality due to ambient PM_{2.5} exposure is estimated to be 0.88 (95% UI 0.3–1.5) million at baseline (the method used to estimate premature mortality is described in *SI Appendix*). This material contains details of household emission estimates and modeling. Following the optimal scenarios P_C,

P_{CL}, and P_{CLS} would avert 7, 7, and 11%, respectively, of premature deaths across India annually from reduction of ambient exposures. Achieving the hypothetical ideal scenario could result in avoiding 13% of the all-India premature mortality burden. The more realistic scenarios (Progress A and Progress M) would avert an estimated 8 and 5% of the total premature mortality burden, respectively. The least ambitious Progress S scenario would avoid only 2% of the premature mortality at baseline. Using the Attribution method, we estimate that 270,000 (95% UI, 69,000–487,000) premature mortality can be averted per year under the ideal scenario.

Discussion

As of early 2016, around 43% of the Indian population (23) was still dependent on solid fuels for cooking, heating, and other household energy services. In this paper, we demonstrate the exposure and health benefits of mitigating household emissions sources on ambient air pollution exposures using several scenarios. Given that household PM_{2.5} exposures in India are about three times higher than ambient exposures in many parts of the country




Fig. 2. Percentage of ambient $PM_{2.5}$ exposure that can be attributed to household $PM_{2.5}$ sources at baseline, 2015.

(24), however, there would be additional and substantial benefits for households as well, where direct exposures to solid-fuel combustion by-products are high. These are not estimated in this assessment, however.

Household interventions may be more socially acceptable and easier to implement than other forms of interventions (25) such as traffic restrictions (25–27) and industry closures that are often perceived as a hindrance to economic growth and disruptive of daily routines. However, despite aggressive new programs to provide clean energy to households, significant challenges remain in ensuring sustained use of clean fuels in households that receive an LPG stove (28) and in ensuring reliable and consistent electricity supplies to displace kerosene lamp usage.

Several studies (10–14, 29) have confirmed that ambient $PM_{2.5}$ exposure can be reduced substantially in India by controlling household sources. Our results show that the all-India average annual ambient $PM_{2.5}$ exposure can be brought down below the Indian NAAQS standard by mitigating all household sources (our ideal scenario). While this scenario is aspirational and may seem hypothetical, it could be achieved through aggressive mitigation measures to promote clean-fuel usage across all household sources, not just cooking. Achieving comparatively pragmatic scenarios like Progress A and Progress M would also result in a large fraction of districts reaching the NAAQS standard and could help avoid considerable premature mortality. In most of the districts in the central and eastern Gangetic Plain, this would reduce the ambient $PM_{2.5}$ exposure closer to the Indian NAAQS. Districts that are out of attainment could be assisted by enforced policies to limit agricultural crop waste and open trash burning, to control industrial emissions, to restrain emissions from construction activities, to increase the frequency of mechanized cleaning of roads to limit on-road resuspension of dust, and/or to curb emissions from brick kilns.

In Delhi NCR, more stringent measures and large-scale shifts in practice have already been undertaken. These include switching the public transport system to Compressed Natural Gas (30) and

the recent implementation of Bharat Stage - VI emissions standards to curb vehicular emission in this region (31, 32). A multi-pronged approach addressing both cleaner fuels for household purposes and other emissions sources is also necessary. Similar efforts are underway to implement a series of mitigation measures recommended by a steering committee formed by the Ministry of Health and Family Welfare, Government of India (33).

In districts where ambient $PM_{2.5}$ exposure is already lower than the NAAQS, curbing household emissions could help approach WHO guidelines. Reductions in both household and ambient $PM_{2.5}$ exposures are expected to improve standards of living in many ways, for example less time spent collecting fuel and fewer episodes of impaired visibility. Therefore, ongoing social programs should continue and their effectiveness should be ensured through rigorous monitoring and evaluation, iteration, and continued public financing. The public should be made aware of the health benefits of clean air through awareness campaigns to promote the use of cleaner fuels for cooking and the adoption of cleaner lifestyles to contribute to improved air quality and sustainable development.

China, another middle-income country with substantial air pollution, also exhibits significant contribution of household sources to ambient pollution, although coal for space heating plays a bigger role than in India (34). Studies have estimated that the biggest impact on ambient pollution during 2005–2015 was actually from “natural” improvements in household fuels due economic growth and urbanization, rather than direct policies to control ambient pollution (35). China now has initiated promulgation of clean household fuels as part of its modern control strategy for addressing ambient air pollution (36). Ironically, neither in India nor China, however, are the full benefits of providing clean household fuels yet




Fig. 3. Relative contributions (%) of biomass use for (A) solid-fuel cooking, (B) space heating, and (C) water heating and kerosene use for (D) lighting to annual ambient $PM_{2.5}$ exposure at the district level at baseline, 2015.




Fig. 4. (A) Percentage of districts where ambient $PM_{2.5}$ exposure exceeds various guidelines before and after mitigation of household emissions. Definition of the scenarios in the y axis are provided in Table 1 and (B) annual $PM_{2.5}$ exposure that needs to be further mitigated in each district after complete mitigation of household $PM_{2.5}$ to achieve Indian standard in that district.

considered in national policies, which would include benefits to the households themselves from lower pollution exposures as well as lower ambient pollution levels.

Methods

Ambient $PM_{2.5}$ Exposure. We consider ambient $PM_{2.5}$ exposure for 2015 as our baseline. Due to the paucity of ground-based measurements across the country in 2015, we utilize aerosol optical depth (AOD) retrieved by MISR (Multiangle Imaging SpectroRadiometer) at 17.6-km resolution to derive annual $PM_{2.5}$ over India. This approach has been described in detail in our previous study (37). In brief, we use a spatially and temporally varying daily conversion factor (η) simulated by the GEOS-Chem chemical transport model (8, 38) to convert AOD to surface $PM_{2.5}$. Inferred $PM_{2.5}$, after bias correction against coincident measurements of $PM_{2.5}$, has $\sim 8\%$ uncertainty (8, 37). Summary statistics are derived at the district level from gridded satellite data using geographic information system.

Contribution of Individual Household Sources Toward Ambient $PM_{2.5}$ Exposure.

The basic methodology for estimating emissions, in turn used for estimating the contribution of household air pollution toward ambient $PM_{2.5}$, is as follows:

$$E_{s,d,g,f,p} = \sum_d \sum_g EC_{f,s} * EF_p,$$

where E is total emissions at the state and district level, by fuel type and pollutant type; EC is energy consumption per capita per year, varying by state; EF is emission factor by pollutant for a state s , district d , fuel f , and pollutant p .

The emissions are broken down into four categories: biomass for (i) cooking, (ii) space heating, and (iii) water heating and kerosene for (iv) lighting. The emission factors for the fuel categories listed in the census reports are obtained from ref. 39 and GAINS (www.iiasa.ac.at/web/home/research/researchPrograms/air/GAINS.html). The fuel use patterns are linked to the fields from household-level components of Census of India, 2011, specifically (i) HH10 for cooking and heating and (ii) HH7 for lighting. The data are further segregated at the district level into urban and rural cooking and inside and outside cooking. Average energy consumed by a household for cooking is calculated based on National Sample Survey Office survey dataset, which lists the amount of food varieties cooked at the state level (39, 40). Heating emissions were further adjusted to account for spatial and temporal profiles in temperature, extracted at the grid level from Weather Research and Forecasting model simulations using National Centers for Environmental Prediction Reanalysis data. Pandey et al. (39, 40) conducted an uncertainty analysis in the form of Monte Carlo simulations. Overall uncertainty in the estimated emissions is $>30\%$, mostly stemming from emission factors. Based on recent measurements, we assume minimal contribution of household biomass burning to precursors of secondary aerosols (41). No laboratory tests for the emission factors or surveys to ascertain the household energy consumption patterns were conducted under this study. More details about the methodology and data used in this study are provided in *SI Appendix*.

The Comprehensive Air Quality Model with Extensions (CAMx) (42–45), an Eulerian photochemical dispersion model, was utilized for dispersion modeling and for estimating the contribution of household emissions toward ambient $PM_{2.5}$ exposure. The emission inventory preparation and modeling framework is discussed in detail in *SI Appendix*, Figs. S1–S4. The model has been validated to perform satisfactorily in earlier studies (42, 46).




Fig. 5. Changes in population-weighted mean ($\pm 1\sigma$ shown by error bars) annual ambient $PM_{2.5}$ exposure (Top) and percentage averted premature mortality based (Bottom). The range of baseline $PM_{2.5}$ estimate is shaded in the top; the mean value is indicated by the dotted line. The dashed horizontal line in the top represents the Indian NAAQS. See text for details.

To estimate the contribution of each household activity (residential cooking, space heating, water heating, and lighting) toward ambient PM_{2.5} we use the following equation:

$$f_{i,j} = \sum_{j=1}^N fh_j \times fe_{i,j},$$

where $f_{i,j}$ is the fraction of ambient PM_{2.5} in a district i contributed by a household activity j . fh_j is the fraction of ambient PM_{2.5} contributed by household burning of solid fuels during the above-mentioned household activities simulated using the CAMx model. The percentage of ambient PM_{2.5} that can be attributed to household PM_{2.5} is depicted in Fig. 1. $fe_{i,j}$ represents the fraction of the total household emission in a district that can be attributed to a particular source j . The percentage contribution of each of the four household activities toward ambient PM_{2.5} exposure is represented in Fig. 3.

Averted Premature Mortality. We estimate the percentage changes in premature mortality (%Av) for diseases which are established to have causal relationships with exposure to ambient PM_{2.5} (2, 47–49) (chronic obstructive

pulmonary diseases, ischemic heart diseases, stroke, and lung cancer) for each of the above-mentioned scenarios, m , as follows:

$$\%Av_m = \frac{(M - M_m)}{M} \times 100,$$

where M is the estimated premature mortality due to PM_{2.5} exposure for the baseline year, 2015; M_m is the estimated premature mortality for PM_{2.5} exposure for each of the developed scenarios, m . Details about estimation of premature mortality are described in [SI Appendix](#).

ACKNOWLEDGMENTS. We appreciate comments by Chandra Venkataraman and acknowledge Aaron van Donkelaar for sharing the conversion factors used for generating PM_{2.5} estimates. MISR aerosol data are archived in NASA Langley Research Atmospheric Science Data Center. S.D. acknowledges Department of Science and Technology- Funds for Improvement of Science and Technology infrastructure in universities & higher educational institutions (DST-FIST) Grant (SR/FST/ESII-016/2014) for computing support. S.C. and S.D. acknowledge the Fulbright-Nehru Doctoral Fellowship and Fulbright-Nehru Academic and Professional Excellence Fellowship, respectively.

- Lim SS, et al. (2012) A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990-2010: A systematic analysis for the global burden of disease study 2010. *Lancet* 380:2224-2260.
- Pope CA, 3rd, et al. (2011) Lung cancer and cardiovascular disease mortality associated with ambient air pollution and cigarette smoke: Shape of the exposure-response relationships. *Environ Health Perspect* 119:1616-1621.
- Gehring U, et al. (2013) Air pollution exposure and lung function in children: The ESCAPE project. *Environ Health Perspect* 121:1357-1364.
- Gao Y, Chan EY, Li L, Lau PW, Wong TW (2014) Chronic effects of ambient air pollution on respiratory morbidities among Chinese children: A cross-sectional study in Hong Kong. *BMC Public Health* 14:105.
- Mitter SS, et al. (2016) Household fuel use and cardiovascular disease mortality: Golestan cohort study. *Circulation* 133:2360-2369.
- Alam DS, et al. (2012) Adult cardiopulmonary mortality and indoor air pollution: A 10-year retrospective cohort study in a low-income rural setting. *Glob Health* 7:215-221.
- Lipsett MJ, et al. (2011) Long-term exposure to air pollution and cardiorespiratory disease in the California teachers study cohort. *Am J Respir Crit Care Med* 184:828-835.
- Dey S, et al. (2012) Variability of outdoor fine particulate (PM_{2.5}) concentration in the Indian subcontinent: A remote sensing approach. *Remote Sens Environ* 127:153-161.
- Dey S, Di Girolamo L (2011) A decade of change in aerosol properties over the Indian subcontinent. *Geophys Res Lett* 38:1-5.
- Conibear L, Butt EW, Knotte C, Arnold SR, Spracklen DV (2018) Residential energy use emissions dominate health impacts from exposure to ambient particulate matter in India. *Nat Commun* 9:617.
- Butt EW, et al. (2016) The impact of residential combustion emissions on atmospheric aerosol, human health, and climate. *Atmos Chem Phys* 16:873-905.
- Silva RA, et al. (2016) The effect of future ambient air pollution on human premature mortality to 2100 using output from the ACCMIP model ensemble. *Atmos Chem Phys* 16:9847-9862.
- Lelieveld J, Evans JS, Fnais M, Giannadaki D, Pozzer A (2015) The contribution of outdoor air pollution sources to premature mortality on a global scale. *Nature* 525:367-371.
- Venkataraman C, et al. (2018) Source influence on emission pathways and ambient PM_{2.5} pollution over India (2015-2050). *Atmos Chem Phys* 18:8017-8039.
- Government of India, Ministry of Petroleum and Natural Gas (2016) Pradhan Mantri Ujjwala Yojana - Scheme Guidelines. Available at petroleum.nic.in/marketing/policies-and-guidelines/lpg-policies-and-guidelines. Accessed April 2, 2019.
- Dabadge A, Sreenivas A, Josey A (2018) What has the Pradhan Mantri Ujjwala Yojana achieved so far? *Econ Polit Wkly* LIII:69-75.
- Government of India, Ministry of Power (2014) Deen Dayal Upadhyaya Gram Jyoti Yojana. Available at www.ddugjy.gov.in/portal/memo/DDUGJY_Guidelines.pdf. Accessed April 2, 2019.
- Pillarsetti A, et al. (2014) Patterns of stove usage after introduction of an advanced cook-stove: The long-term application of household sensors. *Environ Sci Technol* 48:14525-14533.
- Smith KR (2018) Pradhan Mantri Ujjwala Yojana: Transformation of subsidy to social investment in India. *Making of New India: Transformation Under Modi Government*, eds Debroy B, Gangul A, Desai K (Dr Syama Prasad Mookerjee Research Foundation and Wisdom Tree, New Delhi), pp 401-410.
- Giles DM, et al. (2011) Aerosol properties over the Indo-Gangetic Plain: A mesoscale perspective from the TIGER2 experiment. *J Geophys Res Atmos* 116:1-19.
- Srivastava AK, et al. (2011) Pre-monsoon aerosol characteristics over the Indo-Gangetic Basin: Implications to climatic impact. *Ann Geophys* 29:789-804.
- Burnett RT, et al. (2014) An integrated risk function for estimating the global burden of disease attributable to ambient fine particulate matter exposure. *Environ Health Perspect* 122:397-403.
- Health Effects Institute (2018) State of Global Air 2018. Special Report, Boston, MA: Health Effects Institute. Available at https://www.stateofglobalair.org/report. Accessed January 1, 2019.
- Dandona L, et al.; India State-Level Disease Burden Initiative Collaborators (2017) Nations within a nation: Variations in epidemiological transition across the states of India, 1990-2016 in the global burden of disease study. *Lancet* 390:2437-2460.
- Chowdhury S, et al. (2017) "Traffic intervention" policy fails to mitigate air pollution in megacity Delhi. *Environ Sci Policy* 74:8-13.
- Wu Q, Wang Z, Gbaguidi A, Tang X, Zhou W (2010) Numerical study of the effect of traffic restriction on air quality in Beijing. *SOLA* 6:17-20.
- Wang L, Xu J, Qin P (2014) Will a driving restriction policy reduce car trips? The case study of Beijing, China. *Transp Res Part A Policy Pract* 67:279-290.
- Smith KR, Pillarisetti A (2017) Household air pollution from solid cookfuels and health. *Injury Prevention & Environmental Health, Disease Control Priorities*, eds Mock C, Kobusingye O, Nugent R, Smith KR (World Bank, Washington, DC), 3rd Ed.
- Chafe ZA, et al. (2014) Household cooking with solid fuels contributes to ambient PM_{2.5} air pollution and the burden of disease. *Environ Health Perspect* 122:1314-1320.
- Narain U, Krupnick A (2007) The Impact of Delhi's CNG Program on Air Quality. Available at dx.doi.org/10.2139/ssrn.969727. Accessed January 1, 2019.
- Government of India, Niti Aayog (2018) Action Plan for Clean Fuel, Report of the Task Force on Clean Fuel, CII-NITI Aayog's 'Cleaner Air Better Life Initiative'. Available at www.niti.gov.in/documents/reports. Accessed January 1, 2019.
- Government of India, Niti Aayog (2018) Action Plan for Clean Transportation, Report of the Task Force on Clean Transportation, CII-NITI Aayog's 'Cleaner Air Better Life Initiative'. Available at www.niti.gov.in/documents/reports. Accessed January 1, 2019.
- Sagar A, Balakrishnan K, Guttikunda S, Roychowdhury A, Smith KR (2016) India leads the way: A health-centered strategy for air pollution. *Environ Health Perspect* 124:A116-A117.
- Liu J, et al. (2016) Air pollutant emissions from Chinese households: A major and underappreciated ambient pollution source. *Proc Natl Acad Sci USA* 113:7756-7761.
- Zhao B, et al. (2018) Change in household fuels dominates the decrease in PM_{2.5} exposure and premature mortality in China in 2005-2015. *Proc Natl Acad Sci USA* 115:12401-12406.
- Ministry of Environmental Protection of China (2017) *Work Plan for Air Pollution Control in Beijing-Tianjin-Hebei and its Surrounding Areas in 2017* (Ministry of Environmental Protection of China, Beijing).
- Chowdhury S, Dey S (2016) Cause-specific premature death from ambient PM_{2.5} exposure in India: Estimate adjusted for baseline mortality. *Environ Int* 91:283-290.
- van Donkelaar A, et al. (2010) Global estimates of ambient fine particulate matter concentrations from satellite-based aerosol optical depth: Development and application. *Environ Health Perspect* 118:847-855.
- Pandey A, Sadavarte P, Rao AB, Venkataraman C (2014) Trends in multi-pollutant emissions from a technology-linked inventory for India: II. Residential, agricultural and informal industry sectors. *Atmos Environ* 99:341-352.
- Pandey A, Sadavarte P, Rao AB, Venkataraman C (2014) Trends in multi-pollutant emissions from a technology-linked inventory for India: I. Industry and transport sectors. *Atmos Environ* 99:353-364.
- Rooney B, et al. (December 6, 2018) Impacts of household sources on air pollution at village and regional scales in India. *Atmos Chem Phys*, 10.5194/acp-2018-1198.
- Sarkar M, Venkataraman C, Guttikunda S, Sadavarte P (2016) Indian emissions of technology-linked NMVOCs with chemical speciation: An evaluation of the SAPRC99 mechanism with WRF-CAMx simulations. *Atmos Environ* 134:70-83.
- Vijayaraghavan K, et al. (2012) Effects of light duty gasoline vehicle emission standards in the United States on ozone and particulate matter. *Atmos Environ* 60:109-120.
- Koo Y-S, Choi D-R, Cho J-S (2014) Evaluations of WRF-CMAQ/CAMx performance in East Asia focusing on the Seoul metropolitan area. *Air Pollution Modeling and its Application XXII*, eds Steyn DG, Builtjes PJH, Timmermans RMA (Springer, Dordrecht, The Netherlands), pp 441-448.
- Koo B, Wilson GM, Morris RE, Dunker AM, Yarwood G (2009) Comparison of source apportionment and sensitivity analysis in a particulate matter air quality model. *Environ Sci Technol* 43:6669-6675.
- Guttikunda SK, Jawahar P (2014) Atmospheric emissions and pollution from the coal-fired thermal power plants in India. *Atmos Environ* 92:449-460.
- Stieb DM, Szyszkowicz M, Rowe BH, Leech JA (2009) Air pollution and emergency department visits for cardiac and respiratory conditions: A multi-city time-series analysis. *Environ Health* 8:25.
- Pope CA, 3rd, et al. (2002) Lung cancer, cardiopulmonary mortality, and long-term exposure to fine particulate air pollution. *JAMA* 287:1132-1141.
- Wellenius GA, et al. (2012) Ambient air pollution and the risk of acute ischemic stroke. *Arch Intern Med* 172:358-366.