



as the amount and type of coral, algae, and other marine life. "It's kind of old-school, just getting in there and getting wet and counting stuff," says Friedlander. "After you've seen a couple of thousand reefs, you start to get a picture of what's good, what's bad, and what's affecting these systems."

The researchers combined their biological data with survey data from NOAA covering lots of physical factors, including sea surface temperature and wave power, as well as with data on the human drivers affecting reef health, such as sediment runoffs and development. "Putting this all together is probably unprecedented in the scale and the amount of data that we have," says Friedlander.

The researchers constructed detailed models of five different types of "reef regimes," based on the type and amount of marine life present, as well as physical settings and human impacts. For example, one regime tends to occur near human population centers and has very low numbers of fish and coral and high levels of macroalgae. "It's what we normally think of as a degraded reef," says Selkoe. Another regime has low coral cover and macroalgae, but lots of grazing fish, whereas three others have more fish and coral but differ in the level and types of marine organisms. The researchers are analyzing the relationships between the regimes to identify the factors leading to tipping points, in which a site shifts from one regime to another as the intensity of various biological, environmental, or human drivers increases or decreases.

"It's really powerful to see these very distinct, dramatic shifts in the ecology of the reef when the driver levels change," says Selkoe, adding that human communities see the impacts on tourism, fishing, and other cultural practices.

### Managing Trade-Offs

Selkoe and her colleagues aim to have their research inform the actions of natural and aquatic resource managers and policy makers (5). A recent study found that management strategies that considered tipping points were more effective in achieving conservation and management goals than strategies that did not (6).

Identifying the early warning signs of tipping points can help managers prioritize monitoring and interventions. "You don't need to necessarily have intensive regulation and spending, until you're close to a tipping point," says Selkoe.

Plus, interactions between different stressors, as well as the presence of positive and negative feedback processes in these ecosystems, means that the effects are not linear—stressors don't incrementally make things worse. Instead, there may be little effect until the system suddenly tips, with dramatic and undesirable consequences. So when regulating, for example, fishing and pollution, managers must be particularly mindful of the maximum thresholds beyond which the system tips over.



**Sediment runoff often has adverse impacts on coral reefs. Scientists are trying to help control land-based pollution at this West Maui watershed, which includes dense urban development and a resort area. Image courtesy of Kirsten Oleson (University of Hawaii at Manoa, Honolulu).**

For example, managers might have demands from one group that wants ample fishing and another concerned with conserving fish stocks. A seemingly reasonable compromise would be to maintain fish counts halfway between the two demands. "But the system might not be stable at that number, because there's inherent feedbacks in a system that keep it in one state or the other," Selkoe explains. "You can't just keep it at a level of stress that's putting it right on the edge."

To identify management options that minimize costs and conflicts, Kirsten Oleson, an assistant professor of ecological economics at the University of Hawaii, developed a way to perform trade-off analyses. As part of a trial in West Maui, Oleson estimated how fixing agricultural roads in the watersheds would reduce the amount of reef-damaging sediment flowing into the ocean. Most of these dirt roads are on mountain slopes, and the researchers calculated the cost of repairing them with gravel or adding small channels called "waterbars" to redirect water and sediment and reduce erosion and runoff. Some solutions provide significant sediment reduction for relatively little management cost, she says. "The managers were intrigued and really excited about it."

Oleson is working on additional management actions and considering other factors, such as nutrient inflows, that affect the reefs. The researchers are planning to release more analyses over the next year to help managers prioritize protection of sites, create effective monitoring strategies, and set meaningful management targets. "In particular, we will pinpoint the threshold levels of key drivers that cause tipping points," says Selkoe, "so that these thresholds can be avoided."

- 1 Alley RB (2000) Ice-core evidence of abrupt climate changes. *Proc Natl Acad Sci USA* 97(4):1331–1334.
- 2 Lenton TM, et al. (2008) Tipping elements in the Earth's climate system. *Proc Natl Acad Sci USA* 105(6):1786–1793.
- 3 Lenton TM (2011) Early warning of climate tipping points. *Nat Clim Chang* 1(4):201–209.
- 4 Boulton CA, Lenton TM (2015) Slowing down of North Pacific climate variability and its implications for abrupt ecosystem change. *Proc Natl Acad Sci USA* 112(37):11496–11501.
- 5 Selkoe KA, et al. (2015) Principles for managing marine ecosystems prone to tipping points. *Ecosyst Health Sustainability* 1(5):1–18.
- 6 Kelly RP, et al. (2015) Embracing thresholds for better environmental management. *Philos Trans R Soc Lond B Biol Sci* 370(1659):20130276.