

Calendula***Apium graveolens* L.*****officinalis* L.**² /

1

¹/

-

-

² /

-

-

1

.A

B

21

Minimal Inhibitory

. I.R U.V

9

B A

(MIC) Concentration

26.55

A

36.33

B

Isolation and identification two active componenets from *Calendula officinalis* L and *Apium graveolens* L. and tested their activity against some of human pathogenic fungi

¹Abdul -Ridha Akbar Alwan Al-Meyah , ²Abdullah Hmoud Al-Saadoon , ³Wathiq star abd AL-Hassen AL-Shibany & ⁴Najaw Mohammed Jameel Ali Abu- Mejdad .

^{1,2,4}Biology Department-Collage of Science – University of Basrah
³Chemistry Department-Collage of Science – University of Thi-Qar

Abstract

Two active components , one from hot alcoholic extract of *Calendula officinalis* flower called (A) component , the others component isolated from hot alcoholic extract seeds of *Apium graveolense* (named B component) isolated and identified. The component A were separated by acetic acid whil B separated by column and thin layer chromatography. (21) chemical test , UV and IR spectral analysis techniques were used for components identification . Minimal inhibitory concentration of A and B components a gainst 9 fungal isolates , were determind. The A component shows better inhibition activity which showed average of MIC 26.55 comparing with B component 36.33. The laboratory mice were used to test the acute short toxicity of the two components showed no toxicity .

Superficial mycosis

Superficial candidiasis

Midgley) Onychomycosis

Malassezia ↓

.(1997

)

(1987

(1984

)

Materials and Methods -2

. (1995) Guarro Hoogde

-1

BSRA

-2

20

(2004) Souri

Thumbles

24 %95 400 Soxhelet extractor

Whatman-No-1-(England)

° 4

A**-3**

(2001 David 1989)

(A)

(1) *Calendula officinalis*

Whatman-No-15 England

(Butanol :Acetic acid:Water)

(7.5:2.5:1.5) (B:A:W)

72

A
° 4

B -4

(T.L.C) -
(B)

Apium graveolens

.(1984 Harborn) .(4:1:7) (Butanol:Acetic acid:Water)

-

(2×40)

mesh) Silica gel G60 3 mesh) 3
35 (28) (270-120
(1) (4:1:7) (B:A:W) Eluent
(2)

/ 0.2

(1)

.(2001 David) (Rf)

-5

, , , , , , , ,
, , , , , , , ,
- , , , , , , , ,
(366 nm) u.v

Criddle ;1984 ;,Harborn ;1973 Evans Trease)
. (2001 Adedayo ;1998,Richard ;1994 Ellis

B A -6

Melting point -

Gallenkamp B A
B A thermal point apparatus
Jenway 3320 pH Meter , Jenway comp.Germany
) . Solubility test Ignition
(1983)

Infrared spectra -

FTIR-84005 Fourier Transform Infra
KBr) Red Spectrophotometer,Shimadzu,Japan
1⁻ (500-4000) (discs

Electronic spectra -

Heliosαv4.60U.V-visibleSpectrophotometer,England.
(600-200) - -
1
B A
(1997 Schwedt) . 25-5 / 0.005

- 7

/ 500
-: B A

(Oxoid) -1
 10-7 ° 27

/ ⁶10× 6 3 0.2 -2
 2 1

Spreader

(Cork borer) 6 5 -3
 100 -4

5-2 10-7 ° 27 -5

(2002 Hammer 1996 Collee)

Minimum

-: B A (MIC) Inhibition Concentration
 95 / 100 / 200 / 300 / 400 / 500
 / 70 / 75 / 80 / 85 / 90 /
 40 / 45 / 50 / 55 / 60 / 65
 / 15 / 20 / 25 / 30 / 35 /
 / 5 / 6 / 7 / 8 / 9 / 10
 / 1 / 2 / 3 / 4

Acute (short-term) toxicity () -8

()albino mice (15)
 (3) (5±25)

acclimatization

(LD50) Median Leathal Dose

Stomach tube %30

(C) Control Oral administration
/ 16 B A
48
. (1980 Doull Klaassen)

Statistical Analysis -9
) T-test
. (2000)

A (%80) B A
A A
(%50) B
B A -3
B A
(2)

. B A

(2)

B	A		
			1
		I2	2
			3
0.98	0.982	Rf	4
			5
-	-		
-	-		
-	-		
-	-		
-	-		
-	-		
			6
-	-		
-	-		
			7
-	-	%1	
-	-		
			8
-	-		
-	-		
			9
+	+		

+	+		
+	+		
			10
+	+	%1	
+	+	%1	
-	-	%1	11
			12
+	+		
+	+	%1	
+	+	%4	13
		- -	
-	-	%0.5	14
			15
+	+		
+	+) (
+	+		16
			17
+	+		
+	+		
+	+		
			18
196-194 dec.	109-107		
6.7	2.85		

❖ (+) :

❖ (-) :

(2) B A) (O-H)

.(1972 Gayon

. (1994 Ellis Criddle)

B A

-4

(4 3)


B A

(4,3)

Silverstein) $\pi-\pi^*$

(6,5)

(1991)



A


(3)

(

) A

(3)

Bond Causing Absorption	Spectral Region Wave number, $\bar{\nu}$(cm⁻¹)	Band Shap
O-H Str. Vib.	3500 – 3056	Broad Weak
Aliphatic C-H Str. Vib.	2910 & 2854	Weak
C=O & C=C Str. Vib.	1729	Medium
Aliphatic C-H bend.	1450 – 1379	Weak
C=S Str. Vib.	1293	Weak
C-O Str. Vib.	1160 & 1118	Weak
C=C-H bend.& O-H bend.	1069	Weak



B (4)

()B (4)


Bond Causing Absorptoin	Spectral Region Wave number, $\bar{\nu}$ (cm^{-1})	Band Shap
O-H Str. Vib.	3577- 3049	Broad Strong
C-H Str. Vib. Of C=C-H	3021	Very Weak
Aliphatic C-H Str. Vib.	2958 , 2924 & 283	Medium
C=C Str. Vib.	1407	Strong
C=S & C-N Str. Vib.	1264	Weak
C=C-H bend.& O-H bend.	1114	Strong
C-O Str. Vib.	1057	Strong


B A
(6,5) $\pi-\pi^*$

$n-\pi^*$

(1998 Webster Silverstein)

$\pi-\pi^*$





A (5)

B (6)

B A -6

9 / 500 B A

(P<0.05)

(5) B A

/ 500 B A 9 (5)

()				
	B	A		
3.5	2	5	<i>Aspergillus flavus</i>	1
5	3	7	<i>A. terreus</i>	2
28	20	36	<i>Candida albicans</i>	3
14.5	8	21	<i>Chrysosporium tropicum</i>	4
16	10	22	<i>Geotrichum candidum</i>	5
17.5	14	21	<i>G. capitatum</i>	6
14.5	4	25	<i>Microsporum gypseum</i>	7
24	15	33	<i>Rhodotorula rubra</i>	8

12	5	19	<i>Trichophyton rubrum</i>	9
	9	21		

❖

P<0.05 ❖

B A (2)

B A

.(2001)

(O-H) B A

Tylor (2)

(1999) Cowan (1996)

(1993)

(H₂S)

9 B A

-7

9 B A (6)

/			
B	A		
85	75	<i>Aspergillus flavus</i>	1
80	70	<i>A. terreus</i>	2
8	1	<i>Candida albicans</i>	3
35	25	<i>Chrysosporium tropicum</i>	4
10	7	<i>Geotrichum candidum</i>	5
10	8	<i>G. capitatum</i>	6
50	30	<i>Microsporum gypseum</i>	7
9	3	<i>Rhodotorula rubra</i>	8
40	20	<i>Trichophyton rubrum</i>	9
36.33	26.55		

B A (6)

(P<0.05)

(1976 Ian Ian)

() -8
B A
B A

(1987) ■
.7-4 .

(1983) ■

(1984) ■

(2000) ■

(1993) ■

520 .

(1989) ■

395

(2001)

65 .

- **Adedayo , O. ;Anderson , W.A. ; Moo-Young , M. ; Snickus , V. ; Patil , P.A. & Kolawole , D.O. (2001)** . Phytochemistry and anti bacterial activity of *Senna alata* flower. *Pharmaceutical Biology* . 39 : 1-5 .
- **Collee , J. ;Fraser , A. ; Marmion , B. & Simon , A. (1996)** . Makie and McCartney practical medical microbiology . 14th ed. Churchill Liverstone . New York .978p .
- **Cowan , M. (1999)** . Plants products as Anti microbial agents. *Clin. Microbiol . Reviews* . 12 (4) : 564-582 .
- **Criddle , W.J. & Ellis , G.P. (1994)** .Spectral and chemical characterization of organic compounds : *Laboratory Hand book* . 3th ed. John Wiley and Sons , Chichester . New York . Brisbane . Toronto. Singapore. Pp 1-62.
- **David , J. P. ;Rey ,D. ;Meyran , J. & Marigo , G. (2001)**.Involvement of lignin like compounds in toxicity of dietary alder leaf litter against Mosquito larvae. *J. of. Chemic. Ecology* .27 (1) :161-173 .
- **Gayon , G.A. (1972)** . Plant phenolic. Oliver and Boyed Edinburg . 254 p.
- **Hammer, K.A. ; Carson , C.F. & Riley , T.V. (2002)** . In vitro activity of *Melaleuca alternifolia* (tea tree oil) against dermatophytes and other filamentous fungi . *J. Of. Anti microbial . Chemotherapy* 50 :195-199.
- **Harborn , J.B. (1984)** . Phytochemical methods aguide to modern techniques of plants analysis. 2nd ed. Chapman and Hall , London , New York . 288p.
- **Hoog de , G.S. & Guarro, J. (1995)** . Atlas of clinical fungi . center albureau Voor Shimmel-cultures and universitat Rovirai Virgili . Spainb. 720p .
- **Ian , W.D. & Ian , W. S. (1976)**. Microbiol physiology . Black well scientific publications. London . pp 12-18.

-
- **Klaassen , C. D. & Doull , J. (1980)** . Evaluation of safty : Toxicology : The basic science of poisonous. Doull , J.Klaassen , C. D. & Amder , M.O. (eds) .2nd ed. Macmillan publishing co. , Inc. New York , USA . pp 11-27 .
 - **Midgley , G.;Clayton , Y.M. & Hay, R.J.(1997)** . Diagnosis in color medical mycology. Mosby- wolfe , an imprint of mosby international , Spain 155 p.
 - **Richard , J.P.C. (1998)** . Natural Products Isolation . Humana press , Totowa , New Jersey . 473p.
 - **Schwedt , G. (1997)** . The essential guide to analytical chemistry. Translated by Brooks Haderlie. John Wiley and Sons . Chichester. New York . Weinhiin .Drisbane. Singapore. Toronto. 248p .
 - **Silverstein , R.M. & Webster , F.X. (1998)** . Spectro metric identification of organic compounds. 6th ed. John Wiley and Sons, Inc. USA.
 - **Silverstein , R.M. ; Bassler , G.C. & Morrill , T. C. (1991)** . Spectrometric identification of organic compounds. 5th ed. John Wiley and Sons, Inc. USA. 419 p.
 - **Souri , E. ;Amin , G. ; Sharifabadi , A.D. ; Nazifi , A. & Firsam , H. (2004)** . Anti oxidative Activity of Sixty Plants From Iran Iranian . J. of pharmaceutical Research. 3: 55-59 .
 - **Trease , E.J. & Evans , C.W. (1973)** . Pharmacognosy . 10th ed. Bailiere tinball , London . 14 : 139-145.
 - **Tylor , R.S. ; Manadhar , N.P. ; Hudson , J. B. & Towers , G.H.N. (1996)** . Anti microbial activity of Nepalese medicinal plants . J.Ethnopharmacol . 52 : 157-163