



I κ B ζ is a key transcriptional regulator of IL-36–driven psoriasis-related gene expression in keratinocytes

Anne Müller^a, André Hennig^b, Sebastian Lorscheid^a, Paula Grondona^a, Klaus Schulze-Osthoff^{a,c}, Stephan Hailfinger^a, and Daniela Kramer^{a,1}

^aInterfaculty Institute for Biochemistry, University of Tübingen, 72076 Tübingen, Germany; ^bCenter for Bioinformatics, University of Tübingen, 72076 Tübingen, Germany; and ^cGerman Cancer Consortium, German Cancer Research Center, 69120 Heidelberg, Germany

Edited by Sankar Ghosh, College of Physicians and Surgeons, Columbia University, New York, NY, and accepted by Editorial Board Member Tadatsugu Taniguchi August 17, 2018 (received for review January 24, 2018)

Proinflammatory cytokine signaling in keratinocytes plays a crucial role in the pathogenesis of psoriasis, a skin disease characterized by hyperproliferation and abnormal differentiation of keratinocytes and infiltration of inflammatory cells. Although IL-17A and TNF α are effective therapeutic targets in psoriasis, IL-36 has recently emerged as a proinflammatory cytokine. However, little is known about IL-36 signaling and its downstream transcriptional responses. Here, we found that exposure of keratinocytes to IL-36 induced the expression of I κ B ζ , an atypical I κ B member and a specific transcriptional regulator of selective NF- κ B target genes. Induction of I κ B ζ by IL-36 was mediated by NF- κ B and STAT3. In agreement, IL-36–mediated induction of I κ B ζ was found to be required for the expression of various psoriasis-related genes involved in inflammatory signaling, neutrophil chemotaxis, and leukocyte activation. Importantly, I κ B ζ -knockout mice were protected against IL-36–mediated dermatitis, accompanied by reduced proinflammatory gene expression, decreased immune cell infiltration, and a lack of keratinocyte hyperproliferation. Moreover, expression of I κ B ζ mRNA was highly up-regulated in biopsies of psoriasis patients where it coincided with *IL36G* levels. Thus our results uncover an important role for I κ B ζ in IL-36 signaling and validate I κ B ζ as an attractive target for psoriasis therapy.

NFKBIZ | I κ B ζ | IL-36 | keratinocytes | psoriasis

Transcription factor NF- κ B has been implicated in several inflammatory diseases, including psoriasis, by activating various proinflammatory target genes (1). The classical activation of NF- κ B is controlled by cytoplasmic inhibitory proteins, such as I κ B α , which sequester NF- κ B in the cytoplasm (2). Inflammatory stimulation of cells results in the rapid activation of I κ B kinase (IKK), which triggers the phosphorylation-induced degradation of I κ B α , leading to NF- κ B's nuclear translocation and transcriptional activation. Recent evidence, however, suggests that the activation of NF- κ B target genes is more complex and is dependent on the particular gene context or stimulus, which is thought to facilitate selective gene regulation in distinct physiological settings (3). Whereas the rapid activation of primary response genes is directly induced by the classical NF- κ B pathway, expression of so-called “secondary-response genes” requires prior protein synthesis of additional NF- κ B regulators (4). In this context, we and others have identified I κ B ζ , an atypical nuclear I κ B protein, which functions not only as a repressor but, more importantly, also as an activator of a selective subset of NF- κ B target genes (5–8). The mechanisms of this differential gene regulation by I κ B ζ remain largely unknown, but increasing evidence suggests that the transcriptional activity of I κ B ζ is mainly mediated at the level of chromatin remodeling (6, 9, 10).

In keratinocytes (KCs), IL-17A and, more potently, its combination with TNF α induce I κ B ζ expression (11). Subsequently, I κ B ζ mediates the induction of important psoriasis-related gene products, including chemokines (e.g., *CXCL8* and *CCL20*), cytokines (e.g., *IL22* and *IL17C*), and antimicrobial proteins, such as S100 calcium-binding proteins (e.g., *S100A9*), β -defensin-2

(*DEFB4A*), or lipocalin-2 (*LCN2*). Antagonists of TNF α and IL-17A have therefore been approved for the treatment of psoriasis (12). Moreover, *NFKBIZ*, the gene encoding I κ B ζ , has been identified as a psoriasis-susceptibility locus (13). Global *Nfkbiz*-KO mice are resistant to imiquimod (IMQ)- or IL-23–induced psoriasis-like skin inflammation (11). In contrast, *Tnfa*- or *Il17a*-KO mice, which are only partially protected against IMQ-induced psoriasis, still show elevated I κ B ζ mRNA levels in inflamed skin areas (11). These observations imply an additional IL-17A/TNF α –independent pathway which drives I κ B ζ expression and thereby contributes to inflammatory gene expression in psoriasis.

Recently, IL-36 cytokines have received attention as therapeutic targets for psoriasis (14). This subfamily of IL-1–related cytokines consists of three proinflammatory members, IL-36 α (encoded by *IL1F6/IL36A*), IL-36 β (encoded by *IL1F8/IL36B*), and IL-36 γ (encoded by *IL1F9/IL36G*) (15–17). All family members bind to a common heterodimeric receptor, composed of IL-36R (also termed “IL-1RL2”) and IL-1RAcP, leading to the recruitment of the adapter MyD88 and subsequent activation of NF- κ B and MAPK (18). A fourth IL-36 member, IL-36RN, acts as a natural antagonist of IL-36 signaling, as it binds to IL-36R but does not recruit the coreceptor IL-1RAcP (19, 20).

Significance

Psoriasis is an autoinflammatory disease characterized by cytokine-driven keratinocyte proliferation and infiltration of immune cells. While IL-17A and TNF α are established targets in psoriasis therapy, IL-36 is emerging as an important cytokine in this disease. The mechanisms of IL-36–driven proinflammatory responses are largely unknown. Here we identified I κ B ζ , a transcriptional regulator of selective NF- κ B target genes, as a crucial mediator of IL-36 action. In keratinocytes, I κ B ζ was required for the expression of several psoriasis-related cytokines and chemokines. Moreover, genetic deletion of I κ B ζ prevented IL-36–mediated dermatitis induction in mice. Since I κ B ζ is essential not only for IL-36 but also for IL-17 signaling, our results suggest that inhibition of I κ B ζ function could be a future strategy in psoriasis therapy.

Author contributions: K.S.-O., S.H., and D.K. designed research; A.M., A.H., S.L., P.G., and D.K. performed research; A.M., A.H., S.L., and D.K. analyzed data; and K.S.-O., S.H., and D.K. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission. S.G. is a guest editor invited by the Editorial Board.

This open access article is distributed under [Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 \(CC BY-NC-ND\)](https://creativecommons.org/licenses/by-nc-nd/4.0/).

Data deposition: RNA-sequencing data have been deposited in the National Center for Biotechnology Information BioProject database (ID [PRJNA465504](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA465504); Sequence Read Archive accession no. [SRP144926](https://www.ncbi.nlm.nih.gov/sra/SRP144926)).

¹To whom correspondence should be addressed. Email: daniela.kramer@uni-tuebingen.de.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1801377115/-DCSupplemental.

Published online September 17, 2018.

Importantly, while full-length IL-36 proteins seem to be biologically inactive, activation of IL-36 signaling requires their N-terminal proteolytic processing (19, 21).

IL-36 contributes to skin inflammation by acting on KCs and immune cells. Interestingly, IL-36 can induce a subset of proinflammatory target genes similar to those induced by IL-17A in KCs, including *CXCL8*, *IL23A*, *DEFB4*, or *LCN2* (22–24). Vice versa, IL-17, which is typically expressed by immune cells, induces IL-36 γ expression in KCs (25, 26). Therefore, IL-36 appears to have a central position in the interplay between immune cells and KCs. In patients with psoriasis vulgaris, IL-36 α and IL-36 γ are overexpressed, whereas inactivating mutations of *IL36RN* are enriched in a psoriasis subtype, called “generalized pustular psoriasis” (22, 23, 27, 28). In agreement, mice overexpressing IL-36 α in basal KCs exert skin inflammation at 3 wk of age, which is augmented in an *IL36RN*-deficient background (20, 29). In contrast, mice deficient for the IL-36R are fully protected against IMQ-induced psoriasis (30).

Despite its involvement in psoriasis, little is known about IL-36 signaling and its transcriptional responses. In the present study, we found that IL-36 α and IL-36 γ are potent inducers of I κ B ζ expression. Moreover, we identified MyD88, NF- κ B and STAT3 as crucial components for IL-36-induced I κ B ζ expression. Silencing of I κ B ζ in primary human KCs prevented IL-36-mediated up-regulation of multiple psoriasis-associated genes, while a global knockout of I κ B ζ protected against IL-36-mediated psoriasis-like dermatitis in mice. These results and our finding of a strong correlation of *NFKBIZ* and *IL36G* expression in psoriatic lesions uncover an important role for I κ B ζ in IL-36 signaling and thus validate I κ B ζ as an attractive target for psoriasis therapy.

Results

IL-36 Induces I κ B ζ Expression in KCs. To investigate the relationship between IL-36 and I κ B ζ , we treated the keratinocyte cell line HaCaT and primary human KCs with recombinant IL-36 α for 1–24 h. Whereas untreated KCs lacked I κ B ζ expression, 1 h of stimulation with IL-36 α was sufficient to induce sustained I κ B ζ expression on the mRNA and protein level (Fig. 1A). As revealed by the addition of actinomycin D to IL-36 α -treated cells, the increased *NFKBIZ* mRNA levels resulted from transcriptional up-regulation of *NFKBIZ* rather than from mRNA stabilization (*SI Appendix*, Fig. S1A). Importantly, full-length IL-36 α , which supposedly lacks biological activity, failed to induce I κ B ζ expression, whereas IL-17A, either alone or combined with TNF α , induced I κ B ζ expression with kinetics similar to those of truncated IL-36 α (Fig. 1A and *SI Appendix*, Fig. S1B and C). As some reports implied distinct target gene regulation by the different IL-36 members (14, 24, 25), we also stimulated HaCaT cells and primary KCs with IL-36 γ . IL-36 γ induced *NFKBIZ* mRNA and protein expression with kinetics and potency similar to that of IL-36 α (Fig. 1B).

We next investigated whether other psoriasis-associated cytokines, such as IL-1 β , IL-17A, TNF α , or IFN γ , could potentiate the effect of IL-36 α on I κ B ζ protein expression (Fig. 1C). Although certain differences were noted between HaCaT cells and primary KCs, most of the tested cytokines enhanced IL-36 α -mediated I κ B ζ expression. Importantly, the combination of IL-17A and IL-36 α was clearly more effective in triggering I κ B ζ expression than were the single cytokines alone.

Induction of I κ B ζ by IL-36 Is Mediated by MyD88, NF- κ B, and STAT3. As I κ B ζ is also induced by IL-17A, we further dissected the mechanism of I κ B ζ expression induced by IL-36 compared to IL-17A. IL-17A binds and activates the IL-17RA/IL-17RC receptor, followed by the recruitment of the adapter protein Act1 and the activation of MAPK and NF- κ B (31). In contrast, IL-36 utilizes a divergent proximal signaling cascade by binding to the IL-36




Fig. 1. IL-36 induces I κ B ζ expression in KCs. (A and B) HaCaT cells (Left) or human primary KCs (Right) were treated with 100 ng/mL IL-36 α (amino acids 6–158) (A) or 100 ng/mL IL-36 γ (amino acids 18–169) (B) for the indicated times. I κ B ζ protein was analyzed by Western blotting. Relative mRNA levels of *NFKBIZ* were measured in parallel and normalized to the reference *RPL37A*. (C) HaCaT cells (Upper) and primary KCs (Lower) were treated for 2 h with 100 ng/mL IL-36 α alone or in combination with 100 ng/mL IL-17A, 10 ng/mL TNF α , 100 ng/mL IFN γ , or 100 ng/mL IL-1 β . I κ B ζ was detected by Western blotting. HSC70 or β -actin served as loading controls.

receptor complex, composed of IL1RL2 and its coreceptor IL1RAP, leading to the recruitment of MyD88 and activation of MAPK and NF- κ B (17). Indeed, knockdown of MyD88 revealed that it was indispensable for I κ B ζ expression upon IL-36 α stimulation, while it had no effect in IL-17A-treated cells (Fig. 2A and *SI Appendix*, Fig. S2A).

As I κ B ζ is transcriptionally induced by IL-36, we explored the *NFKBIZ* promoter region to identify relevant transcription factors. Two major I κ B ζ isoforms have been described, including a long isoform (I κ B ζ _L) of 718 aa and a N-terminally truncated isoform (I κ B ζ _S) of 618 aa that is thought to be generated by alternative splicing (8, 32). By analyzing published DNase I and Pol II ChIP-seencing (ChIP-seq) data (33), we identified that the two isoforms arise not only from alternative splicing but also from two different promoter regions with distinct transcriptional start sites (Fig. 2B). Moreover, our own RNA-seencing (RNA-seq) data revealed that KCs use only the proximal promoter 2 that is translated into the I κ B ζ _L isoform. Previous promoter




Fig. 2. Molecular dissection of $\text{I}\kappa\text{B}\zeta$ induction by IL-36. HaCaT cells were stimulated for the indicated times with 100 ng/mL IL-36 α . (A) Cells stably expressing a control (sh ctrl) or two different shRNAs targeting *MyD88* were treated for 2 h with IL-36 α and were analyzed by Western blotting. (B) Analysis of *NFKBIZ* promoter accessibility and structure. The genomic region around *NFKBIZ* was analyzed from a published DNase I dataset and a polymerase II ChIP-seq track (33). Exon reads of *NFKBIZ* were derived from our own RNA-seq data of HaCaT cells stimulated for 1.5 h with IL-36 α . (C) Analysis of the *NFKBIZ* promoter 2 region in IL-36 α -stimulated HaCaT cells using luciferase reporter constructs harboring deletions of transcription factor-binding sites. (D) P65, STAT3, and RNA polymerase II (Pol II) bind to *NFKBIZ* promoter region 2 in IL-36 α -treated cells. ChIP was performed from HaCaT cells treated for 30 min with IL-36 α . The promoter region of the muscle-specific gene *MYO1* represents a negative control. (E) Immunoblot analysis of IL-36 α -induced signaling pathways. Active NF- κ B and STAT3 were detected by the phosphorylated forms of $\text{I}\kappa\text{B}\alpha$ (p- $\text{I}\kappa\text{B}\alpha$ at Ser32), p65 (p-p65 at Ser536), and STAT3 (p-STAT3 at Tyr705). MAPK activation was detected by phosphorylated JNK (p-JNK at Thr183/Tyr185) and ERK (p-p44/42 at Thr202/Thr204). (F) Cells were treated for 1 h with IL-36 α in the presence or absence of the vehicle DMSO or 10 μM of the IKK inhibitors BMS-345541 or IMD0354. *NFKBIZ* mRNA and $\text{I}\kappa\text{B}\zeta$ protein levels were measured after 2 h of IL-36 α stimulation. Detection of p- $\text{I}\kappa\text{B}\alpha$ served as a control for NF- κ B inhibition. (G and H) Gene expression and Western blot analysis of $\text{I}\kappa\text{B}\zeta$ in control and *RELA* (p65)-knockdown (G) or *STAT3*-knockdown (H) cells after 1 h of IL-36 α treatment. Knockdown was controlled by detection of p65 or STAT3. *** $P < 0.001$.

analyses, however, had examined only promoter 1, which is located ~20 kb upstream of promoter 2 (32, 34, 35). This distal promoter is used in several cell types for transcription of *NFKBIZ* variant 2, which lacks exon 3 and thus is translated to the $\text{I}\kappa\text{B}\zeta_{\text{S}}$ variant.

Bioinformatic analysis of the *NFKBIZ* promoter 2 revealed putative binding sites for STAT3, NF- κ B, AP1, KLF4, and STAT1. To uncover the contribution of these sites to *NFKBIZ* induction, we cloned the promoter region (~1.5 kb upstream of the transcription start site of $\text{I}\kappa\text{B}\zeta_{\text{L}}$) into a luciferase construct

and generated deletions lacking one of the predicted binding sites. Expression of the constructs was analyzed after transfection of HaCaT cells followed by stimulation with IL-36 α . Indeed, expression of the *NFKBIZ* promoter 2 was significantly increased by IL-36 α , whereas deletion of the STAT3- or the NF- κ B-binding site inhibited *NFKBIZ* promoter expression (Fig. 2C). In accordance, ChIP identified a direct physical binding of NF- κ B p65 and STAT3 to *NFKBIZ* promoter 2, along with the binding of phosphorylated RNA polymerase II as a marker for active transcription (Fig. 2D). IL-36 α also triggered the early activation of STAT3, NF- κ B, and MAPK in HaCaT cells or primary KCs (Fig. 2E and *SI Appendix, Fig. S2B*). Interestingly, a similar activation of STAT3 and NF- κ B was detected in IL-17A-treated cells (*SI Appendix, Fig. S2C*). Whereas inhibition of MAPK did not affect $\text{I}\kappa\text{B}\zeta$ expression in IL-36 α -treated HaCaT cells (*SI Appendix, Fig. S2D*), the blocking of NF- κ B activation by IKK inhibition or knockdown of p65 efficiently prevented $\text{I}\kappa\text{B}\zeta$ expression upon IL-36 α stimulation (Fig. 2F and G). Moreover, depletion of STAT3 by two different shRNAs strongly inhibited $\text{I}\kappa\text{B}\zeta$ mRNA and protein expression (Fig. 2H). Similarly, depletion of p65 or STAT3 impaired $\text{I}\kappa\text{B}\zeta$ induction after stimulation with IL-17A (*SI Appendix, Fig. S2E and F*). Thus, IL-36 α and IL-17A both employ NF- κ B and STAT3 for $\text{I}\kappa\text{B}\zeta$ induction.

$\text{I}\kappa\text{B}\zeta$ Is a Key Mediator of IL-36-Induced Gene Expression in KCs. Next, we investigated the function of $\text{I}\kappa\text{B}\zeta$ in IL-36 signaling and therefore first explored the time course of $\text{I}\kappa\text{B}\zeta$ -modulated gene expression. We stimulated control and *NFKBIZ*-knockdown HaCaT cells for 0–24 h with IL-36 α and analyzed selected IL-36 target genes. IL-36 α stimulation led to the induction of *IL36G*, *IL17C*, *CXCL5*, or *S100A9* with different kinetics (*SI Appendix, Fig. S3A*). Surprisingly, *NFKBIZ* silencing not only prevented the induction of late-responsive genes such as *S100A9* but also affected early gene induction, e.g., of *IL36G* or *IL17C*.

To reveal a global picture of IL-36-driven gene expression by $\text{I}\kappa\text{B}\zeta$, we generated control and *NFKBIZ*-depleted primary KCs and performed transcriptome analyses after 1.5 and 24 h of IL-36 α stimulation (Fig. 3A and B). Silencing of $\text{I}\kappa\text{B}\zeta$ resulted in the deregulation of several hundred target genes in IL-36 α -stimulated primary human KCs (*SI Appendix, Tables S1 and S2*). Interestingly, early after IL-36 α stimulation most genes were down-regulated by $\text{I}\kappa\text{B}\zeta$, including genes for antiinflammatory phosphatases (*DUSP2* and *DUSP9*). In contrast, after 24 h most $\text{I}\kappa\text{B}\zeta$ -modulated genes were positively regulated and hence were down-regulated by the *NFKBIZ* knockdown. Many of these $\text{I}\kappa\text{B}\zeta$ -inducible genes are typically overexpressed in psoriasis, including genes for antimicrobial proteins (*DEFB4* and *LCN2*), S100 proteins (*S100A7*, *S100A8*, and *S100A9*), and chemo- and cytokines (*CSF2*, *CSF3*, *CXCL8*, *IL23A*, and *IL36A*).

Principal component analysis (PCA) revealed that the gene-expression profile not only differed between untreated and IL-36 α -stimulated cells but was also divergent after 1.5 and 24 h of IL-36 α stimulation (*SI Appendix, Fig. S3B*). Moreover, as shown in the Venn diagrams in Fig. 3C and D, only a subset of the IL-36 α -regulated genes was $\text{I}\kappa\text{B}\zeta$ -dependent (83 of 607 genes after 1.5 h and 86 of 800 genes after 24 h of IL-36 α stimulation). Gene ontology (GO) term analysis of the affected genes uncovered that $\text{I}\kappa\text{B}\zeta$ mostly regulated inflammatory responses, neutrophil chemotaxis, and leukocyte function downstream of IL-36 (Fig. 3D). We also compared our RNA-seq analyses with a previously defined IL-36 core signature comprising 182 genes that were regulated by IL-36 after 24 h in human KCs (14). The comparison not only revealed a high overlap with our RNA-seq analyses but also identified 39 of the 182 IL-36 core target genes as $\text{I}\kappa\text{B}\zeta$ -dependent (*SI Appendix, Fig. S3C and D*).

The $\text{I}\kappa\text{B}\zeta$ -dependent gene regulation by IL-36 α in primary KCs at early and late time points was confirmed by qPCR of




Fig. 3. IκBζ regulates a subset of psoriasis-related IL-36 target genes. Primary KCs or HaCaT cells were transfected with a control or *NFKBIZ*-specific shRNA. Triplicates of each time point and shRNA were analyzed by RNA-seq or qPCR and were normalized to the reference gene *RPL37A*. (A) Control of *NFKBIZ*-knockdown efficiency. (Upper) IκBζ protein was detected in primary KCs treated for 1 h with 100 ng/mL IL-36α. (Lower) *NFKBIZ* mRNA levels were measured after 1.5 h and 24 h of IL-36α stimulation. (B) After library preparation from total RNA, primary KC samples were sequenced, and reads were aligned to the human genome hg19. Depicted are two separate heatmaps with normalized z-scores of IκBζ target genes after 1.5 h and 24 h of IL-36α treatment. As a cutoff, genes with a minimum fold change of 1 and a *P* value < 0.05 were considered. (C) Venn diagrams showing the fraction of IκBζ target genes among IL-36α-regulated genes 1.5 and 24 h after stimulation of primary KCs. (D) GO term analysis of significantly enriched IκBζ-dependent gene sets after 1.5 and 24 h of IL-36α treatment. (E) Validation of selected IκBζ target genes by qPCR in primary KCs after 1.5 and 24 h of incubation with 100 ng/mL IL-36α. (F) Gene-expression analysis of IκBζ target genes in primary KCs stimulated with 100 ng/mL IL-17A and 10 ng/mL TNFα for 1.5 and 24 h. **P* < 0.05; ***P* < 0.01; ****P* < 0.001.

selected genes, such as *IL36G*, *S100A9*, *LCN2*, *DEFB4*, *CXCL8*, and *CCL20* (Fig. 3E). Importantly, regulation of these IκBζ target genes was conserved in IL-17A- and TNFα-treated primary KCs as well as in IL-36α-, IL-36γ-, and IL-1β-treated HaCaT cells (Fig. 3F and *SI Appendix*, Fig. S4 A–C). These findings thus implicate IκBζ as a master regulator of proinflammatory gene expression not only in IL-36-stimulated but also in IL-17A-, TNFα-, or IL-1β-treated KCs.

IκBζ Promotes IL-36-Driven Psoriasis-Like Disease in Vivo. Global *Nfkbi*-KO mice are protected against IMQ-induced psoriasis-like skin inflammation (11). Since the TLR7 agonist IMQ directly activates the innate immune response, it is difficult to discriminate between the contribution of IL-17 and IL-36 to the disease onset. Moreover, global *Nfkbi*-KO mice develop an autoinflammatory phenotype in adulthood (36, 37), which could

influence the skin inflammation of IMQ-treated mice. We therefore generated a mouse model using tamoxifen-inducible *Nfkbi*-KO mice that received intradermal injections of active IL-36α into the ears. Tamoxifen-induced Cre recombinase activation just before IL-36α application led to an effective KO of IκBζ, thereby preventing potential congenital off-target effects (Fig. 4A). Intradermal injection of IL-36α into the ears of control animals induced *Nfkbi* transcription (Fig. 4A) and, moreover, triggered ear swelling, scaling, epidermal thickening, KC hyperproliferation and increased infiltration of immune cells (Fig. 4B




Fig. 4. Characterization of the IL-36/IκBζ axis in vivo. (A, Upper) Scheme of tamoxifen and IL-36α treatment of control and inducible *Nfkbi*-KO mice. (Lower) Verification of *Nfkbi* deletion at the protein and mRNA level. For induction of IκBζ KO, *Nfkbi* flox/flox (Ctrl) and Rosa-creERT2 *Nfkbi* flox/flox (KO) mice received i.p. injections of tamoxifen (75 mg/kg) for four consecutive days to induce activation of Cre recombinase. Afterward, 1 μg murine IL-36α or PBS control was intradermally injected into one ear of the mice for five consecutive days. (B) Ear and epidermal thickness (± SEM) of PBS- and IL-36α-treated mice at day 5 from two (for PBS) or six (for IL-36α) animals per group. Pictures were taken at day 5 to show scaling at the treatment area. (C) H&E staining of ears from PBS- and IL-36α-treated control and KO mice. (Scale bars: 140 μm.) (D) Immunohistochemistry for the macrophage marker F4/80 and the neutrophil marker MPO. (Scale bars: 80 μm.) (E) Characterization of CD45+ immune cell infiltrates by flow cytometry. Neutrophils were characterized as CD45+ Ly6G+ and macrophages as CD45+, CD11b^{hi} and F4/80+. Error bars indicate results from two independent experiments. (F) Psoriasis-related gene expression in ears from IL-36α-treated mice. Results are shown as means ± SEM; *n* = 6 animals per group. (G) Expression data from skin biopsies of 64 healthy individuals and 58 psoriasis patients were analyzed from the Gene Expression Omnibus profile dataset GDS4602. Shown are normalized expression values of *NFKBIZ* and *IL17A* or *NFKBIZ* and *IL36G*, which were plotted against each other in every single nonlesional and lesional biopsy. Depicted is the regression coefficient (R^2) from the expression values of the psoriatic skin biopsies. **P* < 0.05; ***P* < 0.01; ****P* < 0.001.

and *C* and *SI Appendix, Fig. S5A*). These alterations were nearly absent in the IL-36 α -treated KO mice. Histological and flow cytometric analyses revealed a marked increase in infiltrating CD45⁺ immune cells, macrophages, and neutrophils in the IL-36 α -treated control animals, which was significantly blocked in the KO mice (Fig. 4 *D* and *E*). T cell infiltration was reduced in the KO animals as well, although the degree of T cell infiltration was generally low in the ears of IL-36 α -treated mice (*SI Appendix, Fig. S5B*). Importantly, expression of several psoriasis-associated target genes, similar to those identified by transcriptome analysis of IL-36 α -treated KCs (Fig. 3), was up-regulated in the ears of IL-36 α -treated control but not in IL-36 α -treated KO mice (Fig. 4*E*). Likewise, the expression of I κ B ζ -dependent proteins involved in granulocyte and leukocyte chemotaxis was also decreased in the KO mice (*SI Appendix, Fig. S5C*). Thus, I κ B ζ KO strongly protected against IL-36-driven psoriasis-like disease in vivo, which could be mediated by effects of *Nfkbiz* deficiency in KCs as well as in immune cells.

As previously reported (11, 13), we validated increased *NFKBIZ* expression in lesions from psoriasis patients, as compared with nonlesional skin areas or unaffected individuals (*SI Appendix, Fig. S5D*). Expression of *IL17A* and especially *IL36G* was elevated in psoriatic lesions. We then correlated the expression of *NFKBIZ*, *IL36G*, and *IL17A* in nonlesional and lesional samples in the individual patients to obtain an idea of the relevance of the two cytokines in driving *NFKBIZ* expression in psoriatic tissue. The correlation of *IL36G* and *NFKBIZ* was stronger than the link between *IL17A* and *NFKBIZ*, implicating IL-36 as an important driver of *NFKBIZ* expression in psoriasis (Fig. 4*G*). Moreover, as IL-36-mediated *NFKBIZ* induction could account for increased expression of psoriasis-related cytokines, we correlated the expression of *LCN2*, a bona fide I κ B ζ target gene (38), to *IL36G*, *NFKBIZ*, and *IL17A* expression. Indeed, the expression level of *LCN2* matched strongly *IL36G* and *NFKBIZ* expression, whereas it was only weakly correlated to *IL17A* expression patterns in psoriatic lesions (*SI Appendix, Fig. S5E*). These findings support a major role of I κ B ζ in IL-36 signaling in KCs and psoriasis and suggest I κ B ζ as an attractive therapeutic target which mediates proinflammatory signaling downstream of IL-17A and IL-36.

Discussion

Previous studies by us and others found that I κ B ζ is overexpressed in psoriatic lesions, whereas *Nfkbiz* KO mice are protected against IMQ-induced psoriatic skin inflammation (11, 13). In these and follow-up studies, I κ B ζ was identified as a major mediator of IL-17A signaling, leading to the induction of proinflammatory signaling in KCs (11, 39). Interestingly, in *Il17a*- or *Il17ra*- KO mice neither induction of *Nfkbiz* nor skin inflammation were fully blocked after IMQ treatment (11, 40), implying additional pathways of *NFKBIZ* induction and promotion of psoriasis.

Recently, IL-36 α and IL-36 γ have been identified as being overexpressed in psoriatic lesions (22, 23). In agreement, IL-36 treatment of KCs induced proinflammatory signaling (14), whereas KO of the IL-36 receptor inhibited IMQ-induced skin inflammation in mice. Our results show that I κ B ζ provides an important link between IL-36 signaling and psoriasis-associated inflammatory gene expression. We revealed that IL-36 mediates I κ B ζ expression in HaCaT cells and primary KCs, which followed kinetics similar to those seen with IL-17A/TNF α treatment, implying similar signaling pathways in I κ B ζ induction.

By ChIP-seq data and our own RNA-seq analyses we identified that KCs induce transcription of *NFKBIZ* from the yet uncharacterized proximal promoter 2, which contains several conserved binding sites for proinflammatory transcription factors. Indeed, IL-36 and IL-17A stimulation led to the activation

of NF- κ B, whereas knockdown of the NF- κ B subunit p65 prevented I κ B ζ induction.

Besides NF- κ B, we identified STAT3 as a regulator of I κ B ζ expression, as its depletion was sufficient to block IL-36- and IL-17A-mediated induction of I κ B ζ . These findings are intriguing, as STAT3 itself can drive proinflammatory gene expression in psoriasis (41). Constitutively active STAT3 in the epidermis of psoriatic lesions is often detectable, whereas pharmacological inhibition of STAT3 ameliorated psoriasis-like skin lesions in mice (42, 43). Moreover, STAT3 was proposed to control I κ B ζ expression in T cells (44). As STAT3 is especially involved in IL-36-driven induction of I κ B ζ expression, STAT3 inhibitors could be promising agents for the effective treatment of general pustular psoriasis, which is caused by mutations of IL36RN and hyperactivation of the IL-36 pathway (27, 28).

Our gene-expression profiling revealed that IL-36 affected the expression of hundreds of genes at early and late stimulation time points. As early effects of IL-36 stimulation on gene expression have not been investigated before in KCs, we could not only validate defined IL-36 target genes (14) but also identify previously unknown IL-36 dependent genes (e.g., *IL17C*, *CSF2*, *CSF3*) that encode important psoriasis-promoting cytokines (45–46). Of note, *NFKBIZ* knockdown led to the deregulation of a specific subset of IL-36 target genes at early and late stimulation time points. Most of these I κ B ζ -dependent IL-36 target genes regulate antimicrobial and proinflammatory responses, neutrophil chemotaxis, and leukocyte activation and hence have been implicated in the pathogenesis of psoriasis. Moreover, I κ B ζ -dependent gene expression seems to be highly conserved, as we found similar changes in the expression of I κ B ζ -dependent genes (e.g., *DEFB4*, *CCL20*, *S100A7*, *S100A9*, and *LCN2*) in HaCaT cells and primary KCs as well as upon IL-36 α , IL-36 γ , or IL-17A/TNF α stimulation.

Employing an inducible *Nfkbiz*-KO model, we further demonstrate that the absence of I κ B ζ also impaired psoriasis-related gene expression under in vivo conditions of IL-36 α stimulation. *Nfkbiz*-KO mice exhibited significantly reduced skin pathology, including less ear swelling and KC proliferation, and a strongly reduced infiltration of immune cells, in particular neutrophils. The results are consistent with findings in *Il36r*-deficient mice that are also protected in the IMQ psoriasis model (30). Notably, our previous study demonstrated that *Nfkbiz*-KO mice were even more protected than *Il17a*-deficient mice (11), supporting the idea that I κ B ζ might also be involved in IL-17-independent effects of psoriasis development. Of note, *Il36r*-deficient mice also showed stronger protection in the IMQ model than *Il17a*-KO mice (30).

In agreement with our findings in cultured KCs and *Nfkbiz*-KO mice, expression data from psoriasis patients validated elevated *NFKBIZ* and *IL36G* levels in psoriatic lesions as compared with nonaffected skin areas or skin from unaffected healthy individuals. Moreover, the expression of *IL36G* and *NFKBIZ* was strongly correlated with the I κ B ζ target gene *LCN2* compared with a much weaker correlation of *IL17A* with *NFKBIZ* and *LCN2* expression levels. These data strengthen the hypothesis that IL-36 is an initial driver for *NFKBIZ* expression in psoriasis.

While our results clearly position I κ B ζ downstream of IL-36, IL-17A and IL-36, in turn, are also transcriptional targets downstream of I κ B ζ (*SI Appendix, Fig. S4D*). Likewise, IL-17A, especially in combination with TNF, is a strong inducer of I κ B ζ expression but can be also induced downstream of I κ B ζ . Thus, the strong expression of *NFKBIZ* in psoriasis patients might be caused not only by elevated IL-36 expression but also by increased IL-17-type responses. The exact contribution of each cytokine is complicated by the existence of multiple members of the IL-17 and IL-36 families. Because IL-17 and IL-36 can mutually reinforce each other (25, 26), the two cytokines drive

complex autoamplification loops in which I κ B ζ seems to have an integral role in promoting skin inflammation (for a scheme see [SI Appendix, Fig. S6](#)). In fact, our present and previous results (11) suggest a dual requirement for I κ B ζ in IL-36 signaling of innate epithelial cells, such as KCs, as well as in IL-17A signaling of T cells, both of which might be necessary to drive full-blown psoriasis.

In conclusion, our findings reveal that two major cytokines, IL-36 and IL-17A, promote psoriasis by inducing I κ B ζ expression. While IL-17A antibodies have proven therapeutic efficacy, blocking of IL-36 might represent an alternative for patients resistant to anti-IL-17A therapy. Moreover, targeting their common mediator I κ B ζ might lead to future approaches for efficient long-term treatment of psoriasis patients.

- Vallabhapurapu S, Karin M (2009) Regulation and function of NF-kappaB transcription factors in the immune system. *Annu Rev Immunol* 27:693–733.
- Renner F, Schmitz ML (2009) Autoregulatory feedback loops terminating the NF-kappaB response. *Trends Biochem Sci* 34:128–135.
- Smale ST (2011) Hierarchies of NF-kB target-gene regulation. *Nat Immunol* 12: 689–694.
- Annemann M, et al. (2016) Atypical I κ B proteins in immune cell differentiation and function. *Immunol Lett* 171:26–35.
- Totzke G, et al. (2006) A novel member of the I κ appaB family, human I κ appaB-zeta, inhibits transactivation of p65 and its DNA binding. *J Biol Chem* 281:12645–12654.
- Hildebrand DG, et al. (2013) I κ B ζ is a transcriptional key regulator of CCL2/MCP-1. *J Immunol* 190:4812–4820.
- Yamamoto M, et al. (2004) Regulation of Toll/IL-1-receptor-mediated gene expression by the inducible nuclear protein I κ appaBzeta. *Nature* 430:218–222.
- Kitamura H, Kanehira K, Okita K, Morimatsu M, Saito M (2000) MAIL, a novel nuclear I kappa B protein that potentiates LPS-induced IL-6 production. *FEBS Lett* 485:53–56.
- Kayama H, et al. (2008) Class-specific regulation of pro-inflammatory genes by MyD88 pathways and I κ appaBzeta. *J Biol Chem* 283:12468–12477.
- Tartey S, et al. (2014) Akirin2 is critical for inducing inflammatory genes by bridging I κ B- ζ and the SWI/SNF complex. *EMBO J* 33:2332–2348.
- Johansen C, et al. (2015) I κ B ζ is a key driver in the development of psoriasis. *Proc Natl Acad Sci USA* 112:E5825–E5833.
- Lowe MA, Suárez-Fariñas M, Krueger JG (2014) Immunology of psoriasis. *Annu Rev Immunol* 32:227–255.
- Tsoi LC, et al. (2015) Enhanced meta-analysis and replication studies identify five new psoriasis susceptibility loci. *Nat Commun* 6:7001.
- Mahil SK, et al. (2017) An analysis of IL-36 signature genes and individuals with IL17L2 knockout mutations validates IL-36 as a psoriasis therapeutic target. *Sci Transl Med* 9: eaan2514, and erratum (2017) 9:eaar6600.
- Dunn E, Sims JE, Nicklin MJ, O'Neill LA (2001) Annotating genes with potential roles in the immune system: Six new members of the IL-1 family. *Trends Immunol* 22:533–536.
- Dinarello CA (2018) Overview of the IL-1 family in innate inflammation and acquired immunity. *Immunol Rev* 281:8–27.
- Bassoy EY, Towne JE, Gabay C (2018) Regulation and function of interleukin-36 cytokines. *Immunol Rev* 281:169–178.
- Towne JE, Garka KE, Renshaw BR, Virca GD, Sims JE (2004) Interleukin (IL)-1F6, IL-1F8, and IL-1F9 signal through IL-1Rrp2 and IL-1RAcP to activate the pathway leading to NF-kappaB and MAPKs. *J Biol Chem* 279:13677–13688.
- Towne JE, et al. (2011) Interleukin-36 (IL-36) ligands require processing for full agonist (IL-36 α , IL-36 β , and IL-36 γ) or antagonist (IL-36Ra) activity. *J Biol Chem* 286: 42594–42602.
- Blumberg H, et al. (2007) Opposing activities of two novel members of the IL-1 ligand family regulate skin inflammation. *J Exp Med* 204:2603–2614.
- Afonina IS, Müller C, Martin SJ, Beyaert R (2015) Proteolytic processing of interleukin-1 family cytokines: Variations on a common theme. *Immunity* 42:991–1004.
- D'Erme AM, et al. (2015) IL-36 γ (IL-1F9) is a biomarker for psoriasis skin lesions. *J Invest Dermatol* 135:1025–1032.
- Boutet MA, et al. (2016) Distinct expression of interleukin (IL)-36 α , β and γ , their antagonist IL-36Ra and IL-38 in psoriasis, rheumatoid arthritis and Crohn's disease. *Clin Exp Immunol* 184:159–173.
- Johnston A, et al. (2011) IL-1F5, -F6, -F8, and -F9: A novel IL-1 family signaling system that is active in psoriasis and promotes keratinocyte antimicrobial peptide expression. *J Immunol* 186:2613–2622.
- Carrier Y, et al. (2011) Inter-regulation of Th17 cytokines and the IL-36 cytokines in vitro and in vivo: Implications in psoriasis pathogenesis. *J Invest Dermatol* 131: 2428–2437.

Materials and Methods

Detailed information on cell culture experiments, generation of knockdown cells, luciferase reporter assays, ChIP, analyses of RNA and protein expression, RNA-seq, cytokine antibody arrays, generation of *Nfkbiz*-KO mice, flow cytometry, histology, and analysis of patient data is provided in [SI Appendix](#).

ACKNOWLEDGMENTS. We thank C. Schönfeld, C. Resch, and J. Loeffler for assistance and A. Witten and M. Stoll (University of Münster) for transcriptome profiling. Murine IL-36 α was generously provided by Amgen. This work was supported by the grants Sonderforschungsbereich/Transregio SFB/TR 156 (to S.H. and D.K.), SFB/TR 209 (to K.S.-O. and S.H.), the Emmy-Noether program of the Deutsche Forschungsgemeinschaft (to S.H.), and the German Ministry for Education and Research Grant of the Network for Autoinflammatory Disorders in Children and Adolescents 01FP09104B (to K.S.-O.).

- Pfaff CM, Marquardt Y, Fietkau K, Baron JM, Lüscher B (2017) The psoriasis-associated IL-17A induces and cooperates with IL-36 cytokines to control keratinocyte differentiation and function. *Sci Rep* 7:15631.
- Marrakchi S, et al. (2011) Interleukin-36-receptor antagonist deficiency and generalized pustular psoriasis. *N Engl J Med* 365:620–628.
- Onoufriadis A, et al. (2011) Mutations in IL36RN/IL1F5 are associated with the severe episodic inflammatory skin disease known as generalized pustular psoriasis. *Am J Hum Genet* 89:432–437.
- Blumberg H, et al. (2010) IL-1RL2 and its ligands contribute to the cytokine network in psoriasis. *J Immunol* 185:4354–4362.
- Tortola L, et al. (2012) Psoriasisform dermatitis is driven by IL-36-mediated DC-keratinocyte crosstalk. *J Clin Invest* 122:3965–3976.
- Gaffen SL (2009) Structure and signalling in the IL-17 receptor family. *Nat Rev Immunol* 9:556–567.
- Yamazaki S, Muta T, Matsuo S, Takeshige K (2005) Stimulus-specific induction of a novel nuclear factor-kappaB regulator, I κ appaB-zeta, via toll/interleukin-1 receptor is mediated by mRNA stabilization. *J Biol Chem* 280:1678–1687.
- Bernstein BE, et al. (2010) The NIH roadmap epigenomics mapping consortium. *Nat Biotechnol* 28:1045–1048.
- Shiina T, et al. (2001) Genomic organization, chromosomal localization, and promoter analysis of the mouse Mail gene. *Immunogenetics* 53:649–655.
- Ishiguro-Oonuma T, Ochiai K, Hashizume K, Morimatsu M (2015) The role of IFN- γ in regulating *Nfkbiz* expression in epidermal keratinocytes. *Biomed Res* 36:103–107.
- Okuma A, et al. (2013) Enhanced apoptosis by disruption of the STAT3-I κ B- ζ signaling pathway in epithelial cells induces Sjögren's syndrome-like autoimmune disease. *Immunity* 38:450–460.
- Ueta M, et al. (2008) Stat6-independent tissue inflammation occurs selectively on the ocular surface and perioral skin of I κ appaBzeta-/- mice. *Invest Ophthalmol Vis Sci* 49: 3387–3394.
- Karlsen JR, Borregaard N, Cowland JB (2010) Induction of neutrophil gelatinase-associated lipocalin expression by co-stimulation with interleukin-17 and tumor necrosis factor-alpha is controlled by I κ appaB-zeta but neither by C/EBP-beta nor C/EBP-delta. *J Biol Chem* 285:14088–14100.
- Muromoto R, et al. (2016) IL-17A plays a central role in the expression of psoriasis signature genes through the induction of I κ B- ζ in keratinocytes. *Int Immunol* 28: 443–452.
- El Malki K, et al. (2013) An alternative pathway of imiquimod-induced psoriasis-like skin inflammation in the absence of interleukin-17 receptor signaling. *J Invest Dermatol* 133:441–451.
- Sano S, et al. (2005) Stat3 links activated keratinocytes and immunocytes required for development of psoriasis in a novel transgenic mouse model. *Nat Med* 11:43–49.
- Andrés RM, Montesinos MC, Navalón P, Payá M, Terencio MC (2013) NF-kB and STAT3 inhibition as a therapeutic strategy in psoriasis: In vitro and in vivo effects of BTH. *J Invest Dermatol* 133:2362–2371.
- Miyoshi K, et al. (2011) Stat3 as a therapeutic target for the treatment of psoriasis: A clinical feasibility study with STA-21, a Stat3 inhibitor. *J Invest Dermatol* 131:108–117.
- Okamoto K, et al. (2010) I κ appaBzeta regulates T(H)17 development by cooperating with ROR nuclear receptors. *Nature* 464:1381–1385.
- Johnston A, et al. (2013) Keratinocyte overexpression of IL-17C promotes psoriasisform skin inflammation. *J Immunol* 190:2252–2262.
- Scholz T, et al. (2017) GM-CSF in murine psoriasisform dermatitis: Redundant and pathogenic roles uncovered by antibody-induced neutralization and genetic deficiency. *PLoS One* 12:e0182646.