


10-20-2020

Male Food Pantry Users Have Low Fruit, Vegetable, and Pulse Intakes Despite Interest in Food Nutrition Content

Donna M. Winham
Iowa State University, dwinham@iastate.edu

Elizabeth D. Davitt
Iowa State University

Follow this and additional works at: https://lib.dr.iastate.edu/fshn_ag_conf

 Part of the [Community-Based Research Commons](#), [Food Science Commons](#), [Gender and Sexuality Commons](#), and the [Human and Clinical Nutrition Commons](#)

The complete bibliographic information for this item can be found at https://lib.dr.iastate.edu/fshn_ag_conf/2. For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

This Poster is brought to you for free and open access by the Food Science and Human Nutrition at Iowa State University Digital Repository. It has been accepted for inclusion in Food Science and Human Nutrition Presentations, Posters and Proceedings by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

Male Food Pantry Users Have Low Fruit, Vegetable, and Pulse Intakes Despite Interest in Food Nutrition Content

Abstract

Background: Low-income men's use of food pantries, and dietary intakes including pulses is understudied. Food pantries can help, but resource restriction limits ability to prepare meals. Convenience foods may not be optimal for nutrition. Pulses (beans, peas, lentils, and chickpeas) count as a vegetable and provide fiber, iron, folate, and other shortfall micronutrients. Increased pulse consumption could improve diet quality. **Purpose:** The study evaluated low-income men's use dietary intakes including pulses. **Methods:** Men aged 25-64 years who utilized food pantry services were eligible. Men completed survey questions on demographics, nutrition assistance programs use, fruit/vegetable/fiber dietary intakes, and pulse consumption patterns. Men received \$5 for survey completion. **Results:** Forty-seven men (46 years \pm 11) from 10 pantries completed the survey. Seventy-three percent had low food security with 29% receiving SNAP benefits. Fifteen percent met fruit/vegetable recommendations and 35% met fiber goals. Although 32% never ate pulses, 15% ate them 4 times per week. Sixty percent expressed concern with nutrition content of food. **Conclusion:** Further inquiry on actual nutrition knowledge is needed to guide suggestions for improving dietary intakes of male food pantry users. Increased pulse consumption could improve dietary quality.

Disciplines

Community-Based Research | Food Science | Gender and Sexuality | Human and Clinical Nutrition

Comments

This accepted poster is published as Winham, D., Davitt, E., Male Food Pantry Users Have Low Fruit, Vegetable, and Pulse Intakes Despite Interest in Food Nutrition Content. Poster at FNCE 2020 (Food & Nutrition Conference & Expo) Indianapolis, IN. October, 17-20, 2020. 120(S9); A83. Doi: [10.1016/j.jand.2020.06.083](https://doi.org/10.1016/j.jand.2020.06.083). Posted with permission.

This poster is available at Iowa State University Digital Repository: https://lib.dr.iastate.edu/fshn_ag_conf/2

Abstract Proposal Title: Male food pantry users have low fruit, vegetable, and pulse intakes despite interest in food nutrition content

ABSTRACT PROPOSAL PREVIEW: MALE FOOD PANTRY USERS HAVE LOW FRUIT, VEGETABLE, AND PULSE INTAKES DESPITE INTEREST IN FOOD NUTRITION CONTENT

[Edit Male food pantry users have low fruit, vegetable, and pulse intakes despite interest in food nutrition content](#)

Abstract Proposal ID: 852538

Abstract Category: Research

Abstract Proposal Status: Active

Author(s)

DW

Donna M. Winham, DrPH, RD

Position:

Assistant Professor

Organization:

Iowa State University

Role:

Co-author

ED

Elizabeth D. Davitt, BSMS

Position:

Graduate Research Assistant

Organization:

Iowa State University

Role:

Presenting Author

Research Abstract

Learning Need Code

- 4070 Food security and hunger

Abstract

Background: Low-income men's use of food pantries, and dietary intakes including pulses is understudied. Food pantries can help, but resource restriction limits ability to prepare meals. Convenience foods may not be optimal for nutrition. Pulses (beans, peas, lentils, and chickpeas) count as a vegetable and provide fiber, iron, folate, and other shortfall micronutrients. Increased pulse consumption could improve diet quality.

Purpose: The study evaluated low-income men's use dietary intakes including pulses.

Methods: Men aged 25-64 years who utilized food pantry services were eligible. Men completed survey questions on demographics, nutrition assistance programs use, fruit/vegetable/fiber dietary intakes, and pulse consumption patterns. Men received \$5 for survey completion.

Results: Forty-seven men (46 years \pm 11) from 10 pantries completed the survey. Seventy-three percent had low food security with 29% receiving SNAP benefits. Fifteen percent met fruit/vegetable recommendations and 35% met fiber goals. Although 32% never ate pulses, 15% ate them 4 times per week. Sixty percent expressed concern with nutrition content of food.

Conclusion: Further inquiry on actual nutrition knowledge is needed to guide suggestions for improving dietary intakes of male food pantry users. Increased pulse consumption could improve dietary quality.

Funding Source

Research support was provided by a grant from the University Translational Network, and the Agriculture and Home Economics Experiment Station, Ames, Iowa, Project No. IOW05462.

Abstract Learning Objective

1. describe low-income men's dietary patterns and use of pulses

Advisor Information-Students Only

Advisor Name

Advisor Email

Advisor Phone