

## Original article

# Mapping the ecoepidemiology of Zika virus infection in urban and rural areas of Pereira, Risaralda, Colombia, 2015–2016: Implications for public health and travel medicine<sup>☆</sup>



Alfonso J. Rodriguez-Morales<sup>a, b, c, \*</sup>, Pablo Ruiz<sup>c, d</sup>, Javier Tabares<sup>c, d</sup>, Carlos Augusto Ossa<sup>d</sup>, Maria Camila Yepes-Echeverry<sup>a</sup>, Valeria Ramirez-Jaramillo<sup>a</sup>, Maria Leonor Galindo-Marquez<sup>a</sup>, Carlos Julian Garcia-Loaiza<sup>a</sup>, Juan Alejandro Sabogal-Roman<sup>a</sup>, Esteban Parra-Valencia<sup>c, e</sup>, Guillermo J. Lagos-Grisales<sup>a, c</sup>, Carlos O. Lozada-Riascos<sup>a, c, f</sup>, Cornelis A. de Pijper<sup>g</sup>, Martin P. Grobusch<sup>c, g</sup>

<sup>a</sup> Public Health and Infection Research Group, Faculty of Health Sciences, Universidad Tecnologica de Pereira, Pereira, Risaralda, Colombia

<sup>b</sup> Committee on Travel Medicine, Asociación Panamericana de Infectología, Quito, Ecuador

<sup>c</sup> Colombian Collaborative Network of Zika (RECOLZIKA), Pereira, Risaralda, Colombia

<sup>d</sup> Secretary of Health and Social Security of Pereira, Pereira, Risaralda, Colombia

<sup>e</sup> Facultad de Medicina, Universidad Católica de la Santísima Concepción, Concepción, Chile

<sup>f</sup> Regional Information System, Universidad Tecnológica de Pereira, Pereira, Risaralda, Colombia

<sup>g</sup> Center of Tropical Medicine and Travel Medicine, Department of Infectious Diseases, Division of Internal Medicine, Academic Medical Center, University of Amsterdam, The Netherlands

## ARTICLE INFO

## Article history:

Received 2 February 2017

Received in revised form

2 May 2017

Accepted 5 May 2017

Available online 6 May 2017

## Keywords:

Zika virus (ZIKV)

Geographical information systems (GIS)

Public health

Travelers

Colombia

Latin America

## ABSTRACT

**Objective:** Geographical information systems (GIS) have been demonstrated earlier to be of great use to inform public health action against vector-borne infectious diseases.

**Methods:** Using surveillance data on the ongoing ZIKV outbreak from Pereira, Colombia (2015–2016), we estimated incidence rates (cases/100,000 population), and developed maps correlating with the ecoepidemiology of the area.

**Results:** Up to October 8, 2016, 439 cases of ZIKV were reported in Pereira (93 cases/100,000 pop.), with highest rates in the South-West area. At the corregimientos (sub-municipalities) of Pereira, Caimalito presented the highest rate. An urban area, Cuba, has 169 cases/100,000 pop., with a low economical level and the highest Aedic index (9.1%). Entomological indexes were associated with ZIKV incidence at simple and multiple non-linear regressions ( $r^2 > 0.25$ ;  $p < 0.05$ ).

**Conclusions:** Combining entomological, environmental, human population density, travel patterns and case data of vector-borne infections, such as ZIKV, leads to a valuable tool that can be used to pinpoint hotspots also for infections such as dengue, chikungunya and malaria. Such a tool is key to planning mosquito control and the prevention of mosquito-borne diseases in local populations. Such data also enable microepidemiology and the prediction of risk for travelers who visit specific areas in a destination country.

© 2017 Elsevier Ltd. All rights reserved.

## 1. Introduction

Zika virus (ZIKV) has caused epidemics during 2015–2017 in different countries of the Americas, with more than 50 countries/territories affected in this region, from 148 globally affected in any form (including imported cases) [1,2]. After Brazil, probably Colombia is the second most affected country in the region [3–5],

<sup>☆</sup> This study was partially presented at the XVIII Pan-American Congress of Infectious Diseases (API), Panama City, Panama, May 2017 (Poster H-16).

\* Corresponding author. Public Health and Infection Research Group, Faculty of Health Sciences, Universidad Tecnologica de Pereira, Pereira, Risaralda, Colombia.

E-mail address: [arodriguezm@utp.edu.co](mailto:arodriguezm@utp.edu.co) (A.J. Rodriguez-Morales).

with over 100,000 cases having been reported from this Northern South American country, reflecting overall incidence rates above 150 cases/100,000 [3]. However, as previously demonstrated; when detailed analyses, at primary, secondary and tertiary territory levels are conducted, incidence rates found tend to be highly variable, and higher [3,6–8].

Early-developed ZIKV epidemiological maps for Colombia using geographical information systems (GIS) at high incidence departments, such as La Guajira [8] and Sucre [7] (Caribbean region), Tolima [6] and Valle del Cauca [3] (Central and Western region), Santander and Norte de Santander (Northeastern area) were published, but these did not include ecoepidemiological analyses that would allow to a better understand the pattern of temporo-spatial distribution of ZIKV infection in these newly-endemic areas. This has been shown to be significant in other vector-borne and arboviral diseases, such as dengue [9–12], where multiple studies have documented the influence of climate variables on its epidemiology, including reports from Colombia [9–12]. Even more, there is a clear need from public health authorities to develop and use GIS-based maps not just for ZIKV, but also for dengue and chikungunya, which are co-circulating in Pereira, in Risaralda, as well Colombia and Latin America at the same time, even co-infecting [1,13,14].

ZIKV and other arboviral diseases could be associated with multiple factors including changes in environmental and social conditions that may influence their epidemiology [15–18]. Up to date, there is a lack of studies assessing eco-epidemiological factors that would be related to ZIKV in Latin America, including Colombia, during ongoing epidemics [19].


Eco-epidemiological information is of utmost importance, which should include the availability of risk maps to address recommendations related to prioritize interventions as well for identification of areas of risk by visitors or people returning from

visiting specific places [8,11,20,21]. Linked to educational programs and other interventional measures [10,20–23], eco-epidemiological data should be considered in an integrated and systematic approach for ZIKV and other arboviral diseases control by national and local health authorities, as well at the community level in order to reduce and mitigate the disease impact.

In the current study, we set out to obtain estimates of ZIKV infection incidence rates in the municipality of Pereira, the capital of Risaralda department and the major city of the Coffee-Triangle region of Colombia; which included the development of GIS-based maps for disease, as well as its ecosystems, rainfall, economical classification as well as entomological indicators, among other ecoepidemiological aspects, in 2015–2016.

**2. Methods**

Scientific publications using GIS for development of epidemiological maps in ZIKV in Latin America are still lacking [6–8,20,24]. Pereira is the major city of the Coffee-Triangle region, which includes three departments (first administrative territory level) and 53 municipalities (second administrative territory level). This is the capital of Risaralda (957,250 habitants in 2016), a department surrounded by six other western departments (Antioquia, Caldas, Tolima, Quindio, Valle del Cauca and Choco) [10,25]. Pereira is constituted by two major areas, urban and rural; the first consisting of 20 communities (the city) and the second by 12 corregimientos (sub-municipalities) (both, tertiary administrative territory level) (Fig. 1). There is a total population of 472,023 habitants in 2016, per the national statistics department ([www.dane.gov.co](http://www.dane.gov.co)). The communities of Rio Otun, Centro, San Joaquin, Del Café, Boston, El Oso, Consota and Cuba, are the most populated areas of the municipality, making up for 51% of its population (Fig. 2).



**Fig. 1.** Geographic distribution of ZIKV incidence rates (cases/100,000pop) in Pereira municipality, Risaralda department, Colombia, 2015–2016. Ecosystems and rainfall patterns for the municipality are shown. Cuba community within the municipality is shown from an aerial photography. (\*Up to epidemiological week 40th, October 8, 2016).





Institute Agustin Codazzi, Colombia. Data of forest density, economical classification and economical activities for Pereira, was provided by the GIS system of the municipality of Pereira. Monthly satellite images for Total Rainfall and Surface Temperature were obtained from the Tropical Rainfall Measuring Mission (1 month - TRMM) imagery database NASA Earth Observations (NEO, NASA, USA) (<http://neo.sci.gsfc.nasa.gov/>) and analyzed for Pereira with the software Google Earth<sup>®</sup>, as described before [10].

Non-linear Poisson simple and multiple regression models were used for determining potential associations between the 2013–2015 mean entomological indexes and the incidence rates of ZIKV infection. Statistical significance was defined as  $p < 0.05$ . Statistical analyses were performed on Stata 14.0<sup>®</sup>.

### 3. Results

Up to October 8, 2016, 439 cases of ZIKV were reported in Pereira (33.85% of Risaralda department), for cumulated rates of 93/100,000 (Table 1). Of the total number of cases, 50 (11.4%) were laboratory confirmed by RT-PCR.

From the total number of cases, 77.7% occurred in the urban area, with a global incidence rate of 85.5 cases/100,000 (Table 1) (Fig. 1), and 94.1/100,000 in the rural area (1.1 times higher). Rates differed from 0 to 460/100,000 (Caimalito, rural, 7.3% of the

municipality cases), followed by Cerritos (rural, 184.2 cases/100,000; 3.4%) (Fig. 1), El Rocio (urban, 170.5 cases/100,000; 0.5%) (Fig. 2), Cuba (urban, 169.0 cases/100,000; 8.0%) and San Nicolas (146/100,000; 3.2%) (Table 1, Fig. 1). These five territories (out of 32), reported 22.3% of cases of the municipality (Table 1) (Fig. 2).

Burden of ZIKV infection in the city has been concentrated in its South-West area, where communes have reported >100 cases/100,000 (Fig. 2). In the case of the corregimientos of Pereira, Caimalito, which borders with La Virginia, a municipality of Risaralda with high transmission of ZIKV infection (1043 cases/100,000) and Cerritos, which borders with the urban area of Pereira, concentrated 10.7% of the cases (Fig. 1) (Fig. 2). This rural area had a higher rainfall pattern with very dense vegetation ecosystems (pastures and cultures) (Fig. 1). The densely populated, low economic level (level 3) (Fig. 3) and mainly residential area of Cuba and it has 169 cases/100,000 (Fig. 4); this is close to the airport, the military field of the city and the major soccer stadium (Fig. 1), as well with the highest Aedic index of the urban area (9.1%) (Fig. 5).

Eight of the communities and two of the corregimientos have rates that are higher than the incidence rate of the municipality (incidence rates ratio) (Table 1). Among these, three communes and the two corregimientos have rates higher than the incidence rate of the department (Table 1). The incidence rate ratio between Pereira and Risaralda is 0.69.

**Table 1**  
ZIKV incidence rates (cases/100,000pop) by communes and corregimientos in the Pereira municipality, Risaralda department, Colombia, 2015–2016.<sup>a</sup>



Communes or corregimientos	Cases	%	Population	%	Incidence Rates (/100,000pop)	Rates compared to the Department	Rates compared to the Municipality
Whole Department	1297	–	957,250	–	135.5	–	–
Whole municipality	439	100.0	472,023	100.0	93.0	0.69	–
<i>Urban Area (Communes)</i>							
El Rocio	2	0.5	1173	0.2	170.5	1.26	1.83
Cuba	35	8.0	20,714	4.4	169.0	1.25	1.82
San Nicolas	14	3.2	9599	2.0	145.8	1.08	1.57
El Oso	34	7.7	26,311	5.6	129.2	0.95	1.39
Perla del Otun	19	4.3	15,381	3.3	123.5	0.91	1.33
San Joaquin	38	8.7	33,420	7.1	113.7	0.84	1.22
Olimpica	17	3.9	15,775	3.3	107.8	0.80	1.16
Rio Otun	42	9.6	43,107	9.1	97.4	0.72	1.05
Consota	22	5.0	24,205	5.1	90.9	0.67	0.98
Centro	34	7.7	37,927	8.0	89.6	0.66	0.96
Del Café	22	5.0	28,016	5.9	78.5	0.58	0.84
El Poblado	14	3.2	19,723	4.2	71.0	0.52	0.76
Villavicencio	10	2.3	14,595	3.1	68.5	0.51	0.74
Boston	12	2.7	26,787	5.7	44.8	0.33	0.48
Ferrocarril	5	1.1	11,636	2.5	43.0	0.32	0.46
Universidad	8	1.8	18,897	4.0	42.3	0.31	0.46
El Jardin	5	1.1	11,991	2.5	41.7	0.31	0.45
Oriente	6	1.4	18,300	3.9	32.8	0.24	0.35
Villasantana	2	0.5	18,424	3.9	10.9	0.08	0.12
Aeropuerto	0	0.0	2732	0.6	0.0	0.00	0.00
Subtotal urban area	341	77.7	398,713	84.5	85.5	0.63	0.92
<i>Rural Area (Corregimientos)</i>							
Caimalito	32	7.3	6952	1.5	460.3	3.40	4.95
Cerritos	15	3.4	8143	1.7	184.2	1.36	1.98
Tribunas	10	2.3	11,374	2.4	87.9	0.65	0.95
Puerto Caldas	5	1.1	6451	1.4	77.5	0.57	0.83
Combia Alta	2	0.5	4644	1.0	43.1	0.32	0.46
Morelia	2	0.5	5588	1.2	35.8	0.26	0.38
Combia Baja	1	0.2	4680	1.0	21.4	0.16	0.23
Arabia	1	0.2	5527	1.2	18.1	0.13	0.19
Altagracia	1	0.2	7481	1.6	13.4	0.10	0.14
La Bella	0	0.0	6240	1.3	0.0	0.00	0.00
La Florida	0	0.0	3505	0.7	0.0	0.00	0.00
La Estrella	0	0.0	2726	0.6	0.0	0.00	0.00
Subtotal rural area	69	15.7	73,310	15.5	94.1	0.69	1.01
Unknown	29	6.6	–	–	–	–	–

<sup>a</sup> Up to epidemiological week 40<sup>th</sup>, October 8, 2016.





Hydrographical map of Pereira



<http://sigper.pereira.gov.co/visor/snpot2015/>

Fig. 4. GIS-based map of hydrography and activities classification of the communes (urban area) of Pereira, Risaralda, Colombia, 2015–2016.

infectious diseases such as tuberculosis and HIV [31,32]. As part of this study, activities for education and control will be focused in the commune of Cuba (for the urban area) and at Caimalito (for the rural area). Pereira is also part of the Central West Metropolitan

Area, which has over 700,000 residents and is composed of Pereira and the neighboring cities of Dosquebradas (200,829 inhabitants in 2016) and La Virginia (32,114 inhabitants in 2016), also the implications would be extended to these other cities and municipalities,




Fig. 5. GIS-based map of Aedic index (%) in the urban area (communes) of Pereira, Risaralda, Colombia, 2013–2015.

Table 2

Entomological indicators, Aedic index, Deposit index, Breteau Index, for the communes and corregimientos in the Pereira municipality, Risaralda department, Colombia, 2013–2015.

Area	Level	Commune or corregimiento	2013			2014			2015			Mean		
			Aedic Index	Deposit Index	Breteau Index	Aedic Index	Deposit Index	Breteau Index	Aedic Index	Deposit Index	Breteau Index	Aedic Index	Deposit Index	Breteau Index
		Mean municipality	10.1	10.0	13.1	7.2	6.9	8.3	6.1	6.6	7.6	8.1	8.2	10.2
		Mean urban area	8.4	8.5	11.6	6.6	6.2	7.6	4.5	4.7	5.4	6.3	6.2	7.9
Urban	Commune	Cuba	10.0	11.6	14.2	SD	SD	SD	8.2	8.1	10.0	9.1	9.9	12.1
		Ferrocarril	10.1	11.2	15.1	10.2	10.5	12.2	6.4	6.6	7.8	8.9	9.4	11.7
		Del Café	7.6	9.0	10.7	13.9	12.2	17.5	2.3	2.2	2.5	7.9	7.8	10.2
		San Nicolás	7.8	10.6	11.8	8.0	6.3	7.4	7.6	8.1	9.5	7.8	8.3	9.6
		Perla del Otun	10.8	9.7	13.7	10.2	7.8	10.2	1.8	1.5	1.8	7.6	6.3	8.6
		San Joaquín	12.6	11.8	16.1	5.0	4.6	5.7	4.3	4.5	4.9	7.3	7.0	8.9
		Villasantana	11.6	10.6	13.5	7.1	7.0	8.8	3.0	1.8	4.7	7.2	6.5	9.0
		El Poblado	12.9	8.1	24.7	5.8	6.6	7.8	2.1	2.3	2.6	7.0	5.7	11.7
		El Oso	8.3	8.3	10.4	4.0	4.6	5.1	8.0	8.5	9.8	6.8	7.1	8.5
		Villavicencio	10.8	11.3	15.4	2.1	2.8	3.1	7.0	6.7	7.3	6.6	6.9	8.6
		Boston	6.2	6.0	7.7	7.4	7.3	7.4	6.2	5.0	6.2	6.6	6.1	7.1
		Oriente	7.6	7.8	9.8	8.4	8.3	9.0	3.7	5.7	4.5	6.5	7.3	7.8
		Consota	9.8	9.2	12.4	6.5	6.1	7.5	3.0	3.1	3.5	6.4	6.1	7.8
		Rio Otun	7.3	8.8	11.1	6.8	7.0	9.2	4.4	6.5	6.8	6.2	7.5	9.0
		Centro	5.7	6.3	8.6	5.9	5.6	5.9	SD	SD	SD	5.8	5.9	7.2
		El Rocio	4.6	4.7	5.6	7.5	6.6	9.2	4.8	4.1	4.8	5.6	5.1	6.5
		Olimpica	SD	SD	SD	2.8	2.4	3.2	SD	SD	SD	2.8	2.4	3.2
		Universidad	7.5	7.1	7.5	0.3	0.4	0.3	0.0	0.0	0.0	2.6	2.5	2.6
		El Jardín	0.0	0.0	0.0	SD	SD	SD	SD	SD	SD	0.0	0.0	0.0
		Aeropuerto	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
		Mean rural area	13.8	13.5	16.6	10.0	9.5	11.3	10.2	11.6	13.5	11.9	12.5	14.9
Rural	Corregimiento	La Estrella	23.4	21.5	31.1	N/A	N/A	N/A	N/A	N/A	N/A	23.4	21.5	31.1
		Morelia	14.1	15.6	16.9	N/A	N/A	N/A	20.0	19.0	26.7	17.0	17.3	21.8
		Caimalito	19.8	17.7	21.8	16.5	15.4	18.3	14.4	13.7	15.9	16.9	15.6	18.6
		Puerto Caldas	13.7	13.0	17.4	13.1	12.2	14.9	7.8	8.5	10.6	11.5	11.2	14.3
		Altagracia	14.4	13.7	18.4	N/A	N/A	N/A	6.4	7.6	9.8	10.4	10.6	14.1
		Arabia	8.4	13.8	8.4	N/A	N/A	N/A	N/A	N/A	N/A	8.4	13.8	8.4
		Cerritos	10.1	9.2	12.1	8.5	8.7	10.2	5.8	10.5	5.8	8.1	9.5	9.4
		Combia Baja	N/A	N/A	N/A	N/A	N/A	N/A	6.8	10.1	12.3	6.8	10.1	12.3
		Tribunales	6.2	3.6	7.1	1.8	1.8	1.8	N/A	N/A	N/A	4.0	2.7	4.4
		Combia Alta	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
		La Bella	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
		La Florida	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A

N/A = Not available/collected.

especially La Virginia, which has been also studied for ZIKV and chikungunya [29]. Pereira is a city with a high population mobility

and located in an area of the country which serves for transit to multiple destinations by road in all directions, including also an






Fig. 6. Non-linear regression models between entomological indexes (mean, 2013–2015) and the ZIKV infection incidence in the Pereira municipality, Risaralda, Colombia, 2015–2016.




Fig. 7. Map of sites of entomological collection for potential vectors of ZIKV in the Corregimiento of Caimalito, Pereira municipality, Risaralda, Colombia, March 2017.

international airport located in its own urban commune, the International Airport Matecaña (which moved 1,890,088 travelers in 2015, 11% directly from other countries), serving Pereira, but also being important for people with final destinations in the North of Valle del Cauca (department), such as Cartago (highly endemic for ZIKV infection) (>500 cases/100,000) or Toro, La Victoria, Roldanillo, Zarzal (all with >1000 cases/100,000) [3], and Armenia (capital of Quindío department), Manizales (capital of Caldas department) and south Antioquia and north Tolima [6]. This airport

is the main in the Coffee-Triangle area of Colombia, including international flights to New York (JFK) (USA) (with a stop in Cartagena, Colombia) and Panama City (Panama). It has been officially announced by a US-based airline, that in the next few months there will be a new route, to Miami. Domestic flights include Bogota (capital of Colombia), Cartagena, Medellín, Quibdó and San Andrés and Providencia islands. Implications for travel medicine should be considered when assessing the risk of international spread and measures related [1,8,21,30].



From this travel medicine point of view, it is important to mention that Pereira is in the center of the Golden Triangle (which consists of the cities of Bogotá, Medellín and Cali), and for that it has become especially important in the fields of trade and commerce, but also of tourism. Touristic places in the region (few kilometers away from Pereira) include the National Park of the Coffee, the National Natural Park of the Snow Mountains (*Los Nevados*), the Hot Springs of Santa Rosa de Cabal, Cocora Valley, and the BioPark Ukumari (former Matecaña Zoo), among others. For example, BioPark Ukumari is in the Cerritos corregimiento, the second most important area in terms of transmission of Zika (incidence rate) at the rural area of Pereira, and located just 8 km far away from the International Airport Matecaña. This 110.5 acres park, since its inauguration in September 2015 considered the largest biopark in Latin America, received 80,000 visitors during its first year. In 2011, Pereira was one of the cities in Colombia for a mass gathering event, the FIFA Sub-20 Soccer World Cup (Stadium Hernan Ramirez Villegas, 30,297 seats), which brought thousands of international travelers and fans as well international sport teams to this city, from Ecuador, Costa Rica, Saudi Arabia, Nigeria, Cameroon, Mexico, Brazil and Spain (all these teams played in Pereira), with all the related implications [33]. In addition, a proportion of travelers (teams from Australia, Guatemala, Croatia, England and South Korea) going to Armenia and Manizales (other cities of the Cup) arrived via Pereira due to its airport's larger capacity. Then, public health policies and strategies for integral control of ZIKV in people living and visiting those areas, should be considered and urgently implemented. Particularly in Caimalito, where a free-trade commerce zone is being implemented, it is a neighboring town with La Virginia (just separated by the Cauca river, with a road bridge), which is highly endemic for ZIKV infection now (more than 1000/100,000), an effective control is needed. In Pereira, as well as Risaralda, Colombia and Latin America as a whole, other arboviruses, such as dengue and chikungunya are also co-circulating, then those measures will mitigate and reduce not just ZIKV [1]. In this setting, co-infections are possible and have reported in Colombia in such epidemiological scenarios [13,14], as well in other countries of Latin America & the Caribbean (e.g. Brazil, Ecuador, Haiti) [34–36].

Although ZIKV was isolated in 1947, only significant research has been done during the past months (ending 2015-beginning 2016) [24], in countries particularly such as Brazil and in Colombia, due to multiple negative potentially linked outcomes, including not just birth and adult neurological compromise, but even deaths [4,37–40].

In this setting, the use of GIS would be important to provide models on the impact and suitability of ZIKV containment measures in the context of environmental and social factors. Recent evidence indicates that the 2015 El Niño caused extreme climatic conditions in Northeastern South America during winter and spring in the Southern hemisphere [41] that might have contributed to the rapid dispersal of the Zika virus in Brazil and other countries in the region, possibly happening in Pereira, as we observed, but in other areas of Colombia, as has been evidenced for dengue in different regions of this country [9,10]. Also, entomological surveillance, in airports, such the one in Pereira, should be enhanced, given the associated risks for exporting infected mosquitoes to other regions and countries, but also importation of infected and non-infected mosquito vectors; particularly in this city, where still there a mean Aedic index >5% for most areas (Fig. 5).

Finally, maps provide relevant information to assess the risk of travelers to specific destinations in highly transmission areas allowing detailed prevention advices [3]. Use of GIS-based epidemiological maps allows to integrate prevention and control strategies, as well as public health policies, for joint control of this

vector-borne disease in this and other areas of the country [21]. As other arboviruses are co-circulating (DENV, CHIKV) [1], maps of them as well for co-infections will be also needed [1,13,14,36]. Simultaneous or sequential arboviral infections occur and should be also assessed [42]. Preparedness in this setting should also consider the potential arrival of Mayaro and Yellow Fever in *Aedes* infested areas [43–46].

Combining entomological, environmental, human population density, travel patterns and case data of vector-borne infections, such as ZIKV, leads to a valuable tool that can be used to pinpoint hotspots also for infections such as dengue, chikungunya and malaria, among others [3,8,21,30]. Such a tool is key to planning mosquito control and the prevention of mosquito-borne diseases in local populations. Such data also enable microepidemiology and the prediction of risk for travelers who visit specific areas in a destination country.

In conclusion, GIS mapping appears as so far under-utilized tool to characterize (arboviral) disease outbreaks down to a detail which allows to interrupt the further spread early, targeted and hence cost-effectively.

### Ethical approval

Not required.

### Funding

Universidad Tecnológica de Pereira, Pereira, Risaralda, Colombia. This study is part of the project “Desarrollo de Mapas Epidemiológicos a través de Sistemas de Información Geográfica para la Caracterización Geográfica de Enfermedades Infecciosas y Tropicales en el Eje Cafetero de Colombia” (Code 5-15-5 [2015–2017]), Universidad Tecnológica de Pereira, Pereira, Risaralda, Colombia.

### Author contributions

Study design: AJRM, Data collection: PR, JT, CAO, Data analysis: AJRM, COLR, PR, JT, Writing: All authors. All authors contributed to, and approved the final version submitted.

### Data availability

Raw data for is available and will be provided on request.

### Grant information

This study was funded by the Universidad Tecnológica de Pereira, Pereira, Risaralda, Colombia (Code 5-15-5 [2015–2017]). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

### Conflicts of interest

The authors have no conflict of interest to disclose.

### Acknowledgments

To Yuan Lian (USA) and Bao Cai (USA), for their critical review and English improvement of the manuscript. This manuscript has been previously presented in part at the XVIII Pan-American Congress of Infectious Diseases (Pan-American Association of Infectious Diseases, API), Panama City, Panama, May 2017 (Poster H-16). The Faculty of Health Sciences and Vice-Rectorship of Research of the Universidad Tecnológica de Pereira, the Colombian Association of Infectious Diseases (ACIN) (Coffee-Triangle Region chapter),

API and the Kansas State University, USA; supported A. J. Rodriguez-Morales, G. J. Lagos-Grisales, V. Ramírez-Jaramillo, C. J. García-Loaiza and J. A. Sabogal-Roman, for their participation at the API meeting where this study was presented.

## References

- [1] Rodriguez-Morales AJ, Villamil-Gomez WE, Franco-Paredes C. The arboviral burden of disease caused by co-circulation and co-infection of dengue, chikungunya and Zika in the Americas. *Travel Med Infect Dis* 2016;14:177–9.
- [2] Rodriguez-Morales AJ. Zika: the new arbovirus threat for Latin America. *J Infect Dev Ctries* 2015;9:684–5.
- [3] Rodriguez-Morales AJ, Galindo-Marquez ML, Garcia-Loaiza CJ, Sabogal-Roman JA, Marin-Loaiza S, Ayala AF, et al. Mapping Zika virus disease incidence in Valle del Cauca. *Infection* 2017;45:93–102.
- [4] Sarmiento-Ospina A, Vasquez-Serna H, Jimenez-Canizales CE, Villamil-Gomez WE, Rodriguez-Morales AJ. Zika virus associated deaths in Colombia. *Lancet Infect Dis* 2016;16:523–4.
- [5] Cardona-Cardona AF, Rodriguez Morales AJ. Severe abdominal pain in a patient with Zika infection: a case in Risaralda. *Colomb J Infect Public Health* 2016;9:372–3.
- [6] Rodriguez-Morales AJ, Galindo-Marquez ML, Garcia-Loaiza CJ, Sabogal-Roman JA, Marin-Loaiza S, Ayala AF, et al. Mapping Zika virus infection using geographical information systems in Tolima, Colombia, 2015–2016. *F1000Res* 2016;5:568.
- [7] Rodriguez-Morales AJ, Patino-Cadavid LJ, Lozada-Riascos CO, Villamil-Gomez WE. Mapping Zika in municipalities of one coastal department of Colombia (Sucre) using geographic information systems during the 2015–2016 outbreak: implications for public health and travel advice. *Int J Infect Dis* 2016;48:70–2.
- [8] Rodriguez-Morales AJ, Garcia-Loaiza CJ, Galindo-Marquez ML, Sabogal-Roman JA, Marin-Loaiza S, Lozada-Riascos CO, et al. Zika infection GIS-based mapping suggest high transmission activity in the border area of La Guajira, Colombia, a northeastern coast Caribbean department, 2015–2016: implications for public health, migration and travel. *Travel Med Infect Dis* 2016;14:286–8.
- [9] Mattar S, Morales V, Cassab A, Rodriguez-Morales AJ. Effect of climate variables on dengue incidence in a tropical Caribbean municipality of Colombia, Cerete, 2003–2008. *Int J Infect Dis* 2013;17:e358–9.
- [10] Quintero-Herrera LL, Ramirez-Jaramillo V, Bernal-Gutierrez S, Cardenas-Giraldo EV, Guerrero-Matituy EA, Molina-Delgado AH, et al. Potential impact of climatic variability on the epidemiology of dengue in Risaralda, Colombia, 2010–2011. *J Infect Public Health* 2015;8:291–7.
- [11] Herrera-Martinez AD, Rodriguez-Morales AJ. Potential influence of climate variability on dengue incidence registered in a western pediatric hospital of Venezuela. *Trop Biomed* 2010;27:280–6.
- [12] Zambrano LI, Sevilla C, Reyes-Garcia SZ, Sierra M, Kafati R, Rodriguez-Morales AJ, et al. Potential impacts of climate variability on dengue hemorrhagic fever in Honduras. *Trop Biomed* 2010;2012(29):499–507.
- [13] Villamil-Gomez WE, Rodriguez-Morales AJ, Uribe-Garcia AM, Gonzalez-Arismendy E, Castellanos JE, Calvo EP, et al. Zika, dengue, and chikungunya co-infection in a pregnant woman from Colombia. *Int J Infect Dis* 2016;51:135–8.
- [14] Villamil-Gomez WE, Gonzalez-Camargo O, Rodriguez-Ayubi J, Zapata-Serpa D, Rodriguez-Morales AJ. Dengue, chikungunya and Zika co-infection in a patient from Colombia. *J Infect Public Health* 2016;9:684–6.
- [15] Singer M. The spread of Zika and the potential for global arbovirus syndemics. *Glob Public Health* 2016;1–18.
- [16] Bogoch II, Brady OJ, Kraemer MU, German M, Creatore MI, Brent S, et al. Potential for Zika virus introduction and transmission in resource-limited countries in Africa and the Asia-Pacific region: a modelling study. *Lancet Infect Dis* 2016;16(11):1237–45.
- [17] Chowell G, Hincapie-Palacio D, Ospina J, Pell B, Tariq A, Dahal S, et al. Using phenomenological models to characterize transmissibility and forecast patterns and final burden of Zika epidemics. *PLoS Curr* 2016;8.
- [18] Lysaght T, Lee TL, Watson S, Lederman Z, Bailey M, Tambyah PA. Zika in Singapore: insights from one health and social medicine. *Singap Med J* 2016;57(10):528–9.
- [19] Alfaro-Murillo JA, Parpia AS, Fitzpatrick MC, Tamagnan JA, Medlock J, Ndeffo-Mbah ML, et al. A cost-effectiveness tool for informing policies on Zika virus control. *PLoS Negl Trop Dis* 2016;10:e0004743.
- [20] Zambrano LI, Sierra M, Lara B, Rodriguez-Nunez I, Medina MT, Lozada-Riascos CO, et al. Estimating and mapping the incidence of dengue and chikungunya in Honduras during 2015 using geographic information systems (GIS). *J Infect Public Health* 2016. <http://dx.doi.org/10.1016/j.jiph.2016.08.003>.
- [21] Tami A, Grillet ME, Grobusch MP. Applying geographical information systems (GIS) to arboviral disease surveillance and control: a powerful tool. *Travel Med Infect Dis* 2016;14:9–10.
- [22] Salomon OD, Quintana MG, Mastrangelo AV, Fernandez MS. Leishmaniasis and climate change-case study: Argentina. *J Trop Med* 2012;2012:601242.
- [23] Coura JR, Junqueira AC. Ecological diversity of *Trypanosoma cruzi* transmission in the Amazon basin. The main scenarios in the Brazilian Amazon. *Acta Trop* 2015;151:51–7.
- [24] Martinez-Pulgarin DF, Acevedo-Mendoza WF, Cardona-Ospina JA, Rodriguez-Morales AJ, Paniz-Mondolfi AE. A bibliometric analysis of global Zika research. *Travel Med Infect Dis* 2016;14:55–7.
- [25] Rodriguez-Morales AJ, Orrego-Acevedo CA, Zambrano-Munoz Y, Garcia-Follecio FJ, Herrera-Giraldo AC, Lozada-Riascos CO. Mapping malaria in municipalities of the coffee triangle region of Colombia using geographic information systems (GIS). *J Infect Public Health* 2015;8:603–11.
- [26] Pena-Garcia VH, Triana-Chavez O, Mejia-Jaramillo AM, Diaz FJ, Gomez-Palacio A, Arboleda-Sanchez S. Infection rates by dengue virus in mosquitoes and the influence of temperature may be related to different endemicity patterns in three Colombian cities. *Int J Environ Res Public Health* 2016;13.
- [27] INS. Enfermedad por virus Zika. *Bol epidemiológico Sem* 2016;2016:103–9.
- [28] Pacheco O, Beltran M, Nelson CA, Valencia D, Tolosa N, Farr SL, et al. Zika virus disease in Colombia - preliminary report. *N Engl J Med* 2016. <http://dx.doi.org/10.1056/NEJMoa1604037>.
- [29] Rodriguez-Morales AJ, Gil-Restrepo AF, Ramirez-Jaramillo V, Montoya-Arias CP, Acevedo-Mendoza WF, Bedoya-Arias JE, et al. Post-chikungunya chronic inflammatory rheumatism: results from a retrospective follow-up study of 283 adult and child cases in La Virginia, Risaralda, Colombia. *F1000Res* 2016;5:360.
- [30] Rodriguez-Morales AJ, Bedoya-Arias JE, Ramirez-Jaramillo V, Montoya-Arias CP, Guerrero-Matituy EA, Cardenas-Giraldo EV. Using geographic information system (GIS) to mapping and assess changes in transmission patterns of chikungunya fever in municipalities of the Coffee-Triangle region of Colombia during 2014–2015 outbreak: implications for travel advice. *Travel Med Infect Dis* 2016;14:62–5.
- [31] Castaneda-Hernandez DM, Martinez-Ramirez JE, Bolivar-Mejia A, Rodriguez-Morales AJ. Differences in TB incidence between prison and general populations, Pereira, Colombia, 2010–2011. *Tuberc (Edinb)* 2013;93:275–6.
- [32] Saldarriaga-Arenas PA, Rodriguez-Morales AJ. Epidemiology of opportunistic diseases in AIDS patients from Pereira municipality, Colombia, 2010–2011. *J Infect Public Health* 2013;6:496–8.
- [33] Gallego V, Berberian G, Lloveras S, Chaves TS, Orduña T, et al. The 2014 FIFA World Cup: communicable disease risks and advice for visitors to Brazil—a review from the Latin American society for travel medicine (SLAMVI). *Travel Med Infect Dis* 2014;12:208–18.
- [34] Sardi SI, Somasekar S, Naccache SN, Bandeira AC, Tauro LB, Campos GS, et al. Coinfections of Zika and chikungunya viruses in Bahia, Brazil, identified by metagenomic next-generation sequencing. *J Clin Microbiol* 2016;54:2348–53.
- [35] Zambrano H, Waggoner JJ, Almeida C, Rivera L, Benjamin JQ, Pinsky BA. Zika virus and chikungunya virus coinfections: a series of three cases from a single center in Ecuador. *Am J Trop Med Hyg* 2016;95:894–6.
- [36] Iovine N, Lednický J, Cherabuddi K, Crooke H, White SK, Loeb JC, et al. Coinfection with Zika and Dengue-2 viruses in a traveler returning from Haiti, 2016: clinical presentation and genetic analysis. *Clin Infect Dis* 2017;64(1):72–5.
- [37] Arzuza-Ortega L, Polo A, Perez-Tatis G, Lopez-Garcia H, Parra E, Pardo-Herrera LC, et al. Fatal sickle cell disease and Zika virus infection in girl from Colombia. *Emerg Infect Dis* 2016;22:925–7.
- [38] Villamil-Gomez WE, Mendoza-Guete A, Villalobos E, Gonzalez-Arismendy E, Uribe-Garcia AM, Castellanos JE, et al. Diagnosis, management and follow-up of pregnant women with Zika virus infection: a preliminary report of the ZIKERNOL cohort study on Sincelejo. *Colomb Travel Med Infect Dis* 2016;14:155–8.
- [39] Parra B, Lizarazo J, Jimenez-Arango JA, Zea-Vera AF, Gonzalez-Manrique G, Vargas J, et al. Guillain-Barre syndrome associated with Zika virus infection in Colombia. *N Engl J Med* 2016;375(16):1513–23.
- [40] Butler D. First Zika-linked birth defects detected in Colombia. *Nature* 2016;531:153.
- [41] Paz S, Semenza JC. El Niño and climate change—contributing factors in the dispersal of Zika virus in the Americas? *Lancet* 2016;387:745.
- [42] Norman FF, Chamorro S, Vazquez A, Sanchez-Seco MP, Perez-Molina JA, Monge-Maillo B, et al. Lopez-velez R. Sequential chikungunya and Zika virus infections in a traveler from Honduras. *Am J Trop Med Hyg* 2016;95(5):1166–8.
- [43] Lednický J, De Rochars VM, Elbadry M, Loeb J, Telisma T, Chavannes S, et al. Mayaro virus in child with acute febrile illness, Haiti, 2015. *Emerg Infect Dis* 2016;22:2000–2.
- [44] Patino-Barbosa AM, Bedoya-Arias JE, Cardona-Ospina JA, Rodriguez-Morales AJ. Bibliometric assessment of the scientific production of literature regarding Mayaro. *J Infect Public Health* 2016;9:532–4.
- [45] Paniz-Mondolfi AE, Rodriguez-Morales AJ, Blohm G, Marquez M, Villamil-Gomez WE. ChikDenMaZika Syndrome: the challenge of diagnosing arboviral infections in the midst of concurrent epidemics. *Ann Clin Microbiol Antimicrob* 2016;15:42.
- [46] Woodall JP, Yuill TM. Why is the yellow fever outbreak in Angola a 'threat to the entire world'? *Int J Infect Dis* 2016;48:96–7.

Reproduced with permission of copyright owner. Further reproduction prohibited without permission.