

Na,K-ATPase $\alpha 4$ isoform is essential for sperm fertility

Tamara Jimenez¹, Jeffrey P. McDermott¹, Gladis Sánchez, and Gustavo Blanco²

Department of Molecular and Integrative Physiology, University of Kansas Medical Center, Kansas City, KS 66160

Edited by David E. Clapham, Children's Hospital Boston, The Howard Hughes Medical Institute, Boston, MA, and approved December 7, 2010 (received for review November 9, 2010)

Regulation of ion balance in spermatozoa has been shown to be essential for sperm motility and fertility. Control of intracellular ion levels requires the function of distinct ion-transport mechanisms at the cell plasma membrane. Active Na⁺ and K⁺ exchange in sperm is under the control of the Na,K-ATPase. Two molecular variants of the catalytic subunit of the Na,K-ATPase, $\alpha 1$ and $\alpha 4$, coexist in sperm. These isoforms exhibit different biochemical properties; however, their function in sperm fertility is unknown. In this work, we show that Na,K-ATPase $\alpha 4$ is essential for sperm fertility. Knockout male mice lacking $\alpha 4$ are completely sterile and spermatozoa from these mice are unable of fertilizing eggs in vitro. Furthermore, $\alpha 4$ deletion results in severe reduction in sperm motility and hyperactivation typical of sperm capacitation. In addition, absence of $\alpha 4$ causes a characteristic bend in the sperm flagellum, indicative of abnormal sperm ion regulation. Accordingly, $\alpha 4$ -null sperm present increased intracellular Na⁺ and cell plasma membrane depolarization. These results are unique in demonstrating the absolute requirement of $\alpha 4$ for sperm fertility. Moreover, the inability of $\alpha 1$ to compensate for $\alpha 4$ suggests that $\alpha 4$ is the Na,K-ATPase- α isoform directly involved in sperm fertility. Our findings show $\alpha 4$ as an attractive target for male contraception and open the possibility for the potential use of this Na,K-ATPase isoform as a biomarker for male fertility.

Na,K-ATPase is an ion transporter of the plasma membrane involved in the active exchange of intracellular Na⁺ for extracellular K⁺ in most cells (1). The Na⁺ and K⁺ gradients generated by Na,K-ATPase are essential in maintaining cell ion homeostasis, cell membrane resting potential, and the transport of a variety of solutes and water across the cell surface (2). Two major polypeptides, named the α and β subunits, constitute Na,K-ATPase (3). The catalytic α polypeptide is the subunit involved in the ATP hydrolysis and ion-translocation functions of Na,K-ATPase (1).

Three molecular variants of the Na,K-ATPase α polypeptide, the $\alpha 1$, $\alpha 2$, and $\alpha 3$ isoforms, have been found in somatic cells of mammals (4–6). Na,K-ATPase is also present in male germ cells and in differentiated spermatozoa (7). Although Na,K-ATPase activity in sperm has been known for some time (8), it was not until recently that the existence of a distinct α isoform of this transporter was reported in the male gamete (9). This polypeptide, named the Na,K-ATPase $\alpha 4$ isoform, is present only in male germ cells, it is expressed after meiosis of these cells (7, 10), and is abundant in the mid-piece of the sperm flagellum (7, 11, 12). In addition to $\alpha 4$, another Na,K-ATPase isoform, the $\alpha 1$ polypeptide, which is ubiquitously present in all tissues, is also expressed in spermatozoa (7). We have previously shown that $\alpha 4$ exhibits unique biochemical properties, including specific affinities for the physiological ligands, Na⁺, K⁺, and ATP, and a particular high sensitivity for the inhibitor ouabain (13). Selective inhibition of $\alpha 4$ with ouabain has been shown to affect rat sperm motility in vitro, reducing total, progressive, and different parameters of sperm movement (11, 12, 14). In addition, increased expression of $\alpha 4$ enhances sperm motility in transgenic mice (15). Although these observations suggest the involvement of $\alpha 4$ in flagellar beating, the role and mechanisms of action of $\alpha 4$ in sperm fertility still remain unknown.

In this work, we have studied the role of the Na,K-ATPase $\alpha 4$ isoform by directly disrupting the *Atp1a4* gene that encodes for the $\alpha 4$ polypeptide. Our results show that the $\alpha 4$ isoform is essential for the fertility of male mice and for the ability of mouse spermatozoa to fertilize eggs in vitro. Furthermore, we demonstrate that $\alpha 4$ activity is necessary for sperm motility and hyperactivation, a particular pattern of motility acquired by sperm during capacitation and required for fertilization (16, 17). Our data also show that sperm lacking $\alpha 4$ exhibit ion balance changes, high intracellular Na⁺ levels, and membrane depolarization, all parameters that are critical for sperm motility and fertility.

Results

Sperm from Mice in Which the *Atp1a4* Gene Is Disrupted Lack Na,K-ATPase $\alpha 4$ Expression and Activity. To determine the function of $\alpha 4$ in male fertility, we have used a genetic approach, targeting the *Atp1a4* gene to suppress expression of the $\alpha 4$ polypeptide. We have disrupted the $\alpha 4$ locus in ES cells by removing a region spanning exons 5 to 8 of the *Atp1a4* gene (Fig. 1A), which encodes for the ATP binding and phosphorylation sites of the catalytic domain of Na,K-ATPase (18). Positive ES cell clones were confirmed by Southern blot analysis (Fig. 1B). Once the chimeric mouse was obtained, a colony was established and the desired genotype of offspring was verified by PCR (Fig. 1C). The absence of the $\alpha 4$ isoform in the knockout mice was further validated by RT-PCR of testis RNA (Fig. 1D) and by immunoblotting of sperm protein samples (Fig. 1E). Heterozygous mice expressed approximately half the amount of $\alpha 4$ than wild-type mice. The null allele was inherited with the offspring of mated heterozygous F1 mice (27 litters) to the approximate expected Mendelian ratio (62 wild-type, 97 heterozygous, and 53 homozygous $\alpha 4$ mice).

Determination of Na,K-ATPase activity and ouabain inhibition in sperm homogenates showed a biphasic dose-response curve in wild-type animals, with inhibition IC₅₀ values in the nanomolar ($1.7 \pm 1.8 \times 10^{-9}$ M) and micromolar ($4.2 \pm 1.4 \times 10^{-5}$ M) range. As previously shown, these kinetic parameters represent the activity of the $\alpha 4$ and $\alpha 1$ isoforms, respectively (4) (Fig. 2A). In contrast to wild-type, Na,K-ATPase activity from $\alpha 4$ knockout-mice sperm responded to ouabain with a monophasic profile, lacking the enzyme component corresponding to $\alpha 4$, and showing a single IC₅₀ value ($1.6 \pm 0.5 \times 10^{-5}$ M) that is characteristic of $\alpha 1$ (Fig. 2A). In addition, the total Na,K-ATPase activity in spermatozoa from $\alpha 4$ null mice was significantly lower than in wild-type sperm (Fig. 2B). This decrease in activity did not depend on changes in the ATP hydrolysis catalyzed by the $\alpha 1$ isoform, but on the absence of $\alpha 4$. Further indication for the lack of $\alpha 4$ activity was obtained through ouabain-binding studies using the fluorescent derivative bodipy-ouabain. To favor binding to $\alpha 4$, we used concentrations of bodipy-ouabain that are rela-

Author contributions: G.B. designed research; T.J., J.P.M., and G.S. performed research; T.J., J.P.M., G.S., and G.B. analyzed data; and G.B. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

¹T.J. and J.P.M. contributed equally to this work.

²To whom correspondence should be addressed. E-mail: gblanco@kumc.edu.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1016902108/-DCSupplemental.




Fig. 1. Disruption of the Na,K-ATPase *Atp1a4* gene. (A) Scheme showing the targeting strategy for the *Atp1a4* gene. (B) Southern blot analysis strategy for screening ES cell clones. Identification of positive ES cell clones (hybrid 129/SvEv-C57BL/6 lines) was performed using probes that specifically hybridized to the indicated regions of the $\alpha 4$ DNA. Wild-type (WT) and $\alpha 4$ -null (null) mice exhibited the predicted DNA bands of 18.9 and 13.6 kb for the 5' end, and 8.1 and 4.8 kb for the 3' end. As negative controls, ES cells from the parent 129/SvEv (129), C57BL/6 (B6), and hybrid (Hyb) 129/SvEv-C57BL/6 lines were used. (C) PCR genotyping of mice. PCR of mouse tail clips produced distinct products of 465 and 531 bp in wild-type and $\alpha 4$ -null mice, respectively, and both DNA fragments in the heterozygous mice. Samples with no DNA were used as a negative control, whereas wild-type genomic DNA plus vector DNA were used as a positive control. (D) RNA analysis of wild-type and knockout mice. RT-PCR was performed on mouse testis RNA, using primers specific for $\alpha 4$ and GAPDH, used as a loading control. This process yielded expected bands of 478 and 430 bp for $\alpha 4$ and GAPDH, respectively (lines marked +). Samples in which reverse transcriptase was omitted (-) were used as a negative control. (E) Immunoblot analysis of $\alpha 4$ expression. Sperm lysates were used to determine expression of the $\alpha 4$ polypeptide. Anti- $\alpha 4$ antiserum generated in chicken and anti- $\alpha 1$ antiserum generated in rabbit, followed by horseradish peroxidase-conjugated secondary antibodies and chemiluminescence were used for detection. Tubulin was used as a loading control.

tively low (10^{-8} M). At this amount, bodipy-ouabain predominantly binds to the $\alpha 4$ isoform, which has a much higher affinity for ouabain than $\alpha 1$ (13). Sperm from $\alpha 4$ -null mice showed negligible binding of bodipy-ouabain compared with sperm from wild-type mice (Fig. 2C). The minimal amount of fluorescence detected in $\alpha 4$ -null sperm most likely represents some unspecific binding of bodipy-ouabain, which is unavoidable in this assay. Taken together, these results showed the functional absence of $\alpha 4$ in knockout sperm and that $\alpha 4$ deficiency was not compensated through up-regulation of $\alpha 1$ activity.

Mice Null in the Na,K-ATPase $\alpha 4$ Isoform Are Sterile. Both hetero- and homozygous $\alpha 4$ knockout mice were overall phenotypically normal, showing testis size, testis shape, and testis-to-body weight ratio indistinguishable from wild-type mice (Fig. 3A and B). Moreover, testis from wild-type, hetero- and homozygous mice exhibited no histological differences. As shown in Fig. 3C, partial or total deficit of $\alpha 4$ did not affect the general structure of the seminiferous tubules, which presented their typical composition and the male germ cell at different developmental stages. This finding shows that disruption of $\alpha 4$ is not followed by important changes in the testis seminiferous epithelium.

The most remarkable physiological consequence of disrupting $\alpha 4$ expression was that $\alpha 4$ -null male mice were completely infertile. Controlled matings of eight null male mice with multiple

wild-type females yielded no pregnancies after a period of 3 mo. This was not because of the inability of these mice to mate, as vaginal plugs were found in the females that had been with $\alpha 4$ homozygous mice with the same frequency as those mated with wild-type males. In contrast, $\alpha 4$ heterozygous male mice were fertile. In addition, all homozygous female mice were fertile and produced similar litter numbers as wild-type mice during the same time period.

Not only were the $\alpha 4$ -null mice sterile, but sperm from these mice was incapable of fertilizing oocytes in vitro (Fig. 4A). Thus, fertilization assays in which $\alpha 4$ -null sperm was incubated with ova, resulted in oocytes that did not develop to the two-cell stage (Fig. 4B). Taken together, these data indicate an essential requirement of $\alpha 4$ for sperm fertility not only in vivo, but also in vitro.

Spermatozoa from $\alpha 4$ -Null Mice Exhibit Reduced Motility. The abnormal fertility phenotype caused by $\alpha 4$ disruption was accompanied by a severe reduction in sperm motility. When measured in noncapacitated media, total motility of the $\alpha 4$ -null mice was decreased to $\approx 85\%$ of that of the wild-type mice (Fig. 5A and Movies S1 and S2). In addition, absence of $\alpha 4$ drastically affected other parameters of sperm motility, including progressive motility, straight line, curvilinear and average path velocities, beat cross frequency, amplitude of lateral head displacement, line-




Fig. 2. Spermatzoa from Na,K-ATPase $\alpha 4$ -null mice lack Na,K-ATPase $\alpha 4$ activity. (A) Dose-response curves for the inhibition of Na,K-ATPase by ouabain. Assays were performed on sperm homogenates (5 μ g total protein) at saturating concentrations of all Na,K-ATPase physiological ligands and the indicated doses of ouabain. The curves represent the best fit of the experimental data to highly ouabain-sensitive and ouabain-resistant $\alpha 4$ and $\alpha 1$ isoforms. Each value is the mean \pm SEM of three experiments. (B) Maximal Na,K-ATPase activity for total $\alpha 4$ and $\alpha 1$ isoforms in wild-type and $\alpha 4$ -null mice. Activity of the $\alpha 4$ and $\alpha 1$ isoforms was distinguished based on the difference in ouabain affinity that characterizes these isoforms (4). Total activity corresponded to that inhibited by 10^{-3} M ouabain. Hydrolysis of ATP by $\alpha 4$ was determined as that sensitive to 10^{-6} M ouabain. Activity of $\alpha 1$ corresponded to the difference in ATP hydrolysis between 10^{-6} and 10^{-3} M. (C) Ouabain binding assays. Sperm from wild-type and knockout mice were labeled with the fluorescent ouabain derivative, bodipy-ouabain. Samples were then subjected to flow cytometry and levels of fluorescence were determined. In B and C, bars represent the mean \pm SEM of three experiments. Values significantly different from the wild-type are indicated: * $P < 0.001$.

arity, and straightness (Fig. 5 B–I). Furthermore, when sperm was capacitated in vitro in media with BSA and bicarbonate, hyperactivation, a pattern of motility typical of sperm capacitation (17), was almost completely abolished in the $\alpha 4$ knockout mice (Fig. 5J). Cell viability, determined using the Live/Dead Sperm Viability Kit (Invitrogen), was similar for both $\alpha 4$ -null and wild-type sperm (97.0 ± 1.2 vs. 98.1 ± 0.1). This finding indicated that the lower sperm motility found in the $\alpha 4$ -null mouse was not a consequence of increased cell death. In contrast to the homozygous mice, heterozygous $\alpha 4$ mice showed only a partial reduction in sperm motility parameters (Fig. 5). Overall, these

results demonstrate that the $\alpha 4$ isoform is directly involved in multiple aspects of sperm flagellar movement, including the hyperactive motility that accompanies sperm capacitation.

Spermatzoa from $\alpha 4$ -Null Mice Show Several Other Abnormalities. Strikingly, the majority of spermatozoa from the $\alpha 4$ homozygous mice exhibited a distinct bend between the mid- and the principal piece of the flagellum that was not present in heterozygous or wild-type mice. This bend showed different degrees of angularity,




Fig. 3. Na,K-ATPase $\alpha 4$ -null mice testis exhibit normal morphology. (A) Macroscopic view of testis dissected from wild-type, heterozygous, and homozygous mouse testis. (B) Relationship between testis and body weight in wild-type, heterozygous, and homozygous mouse testis. Bars represent the mean \pm SEM of three experiments. (C) Microscopic analysis of wild-type, heterozygous, and homozygous mouse testis. Testis were dissected and fixed with Bouin's fixative (Ricca Chemical Co.), embedded in paraffin, and sectioned in 10- μ m sections. Tissue was treated with xylene and ethanol to remove the paraffin, and stained with H&E. (Upper) 20 \times magnification; (Lower) 40 \times magnification. (Scale bars, 100 μ m.)




Fig. 4. Sperm from $\alpha 4$ -null mice are unable to fertilize eggs. Sperm from wild-type or $\alpha 4$ -null sperm were incubated with oocytes in Cook's medium to allow in vitro fertilization. After removing the oocytes, they were washed and cultured overnight. The development of two-cell stage embryos was determined. (A) Fertilization rate from the in vitro fertilization assays. Bars represent the mean \pm SEM. Values significantly different from the wild-type are indicated with an asterisk, with $P < 0.001$. (B) Light microscopy images of oocytes and two cell embryos.




Fig. 5. Sperm from $\alpha 4$ -null mice show low motility. Spermatozoa from the cauda epididymis were isolated in modified Tyrode's medium. Sperm motility was determined using CASA and different parameters of sperm movement were analyzed. Total percent motility (A), progressive motility (B), straight line velocity (C), curvilinear velocity (D), average path velocity (E), beat cross frequency (F), amplitude of lateral head displacement (G), linearity (H), and straightness (I) were determined. In addition, hyperactive motility (J) was measured after capacitation of the sperm for 1 h. Bars represent the mean \pm SEM of three experiments. Values significantly different from the control are indicated with an asterisk, with P values ranging between 0.05 and 0.001.

ranging from obtuse and acute angles to serious retroflexion into a complete 180° folding of the head over the principal piece of the flagellum (Fig. 6A). Defects similar to those have been associated with deficiencies in sperm osmoregulation and volume control (19), suggesting that one of the functions of $\alpha 4$ is to maintain sperm osmotic balance.

Na,K-ATPase is responsible for maintaining the low intracellular Na^+ concentration ($[\text{Na}^+]_i$), which is crucial to many general and cell-specific processes (2). Determination of $[\text{Na}^+]_i$ in the sperm cytosol using the fluorescent indicator, Sodium Green Tetraacetate, showed that $\alpha 4$ -null mice exhibited significantly higher $[\text{Na}^+]_i$ levels, with an increase of $\approx 60\%$ compared with wild-type mice (Fig. 6B). This finding suggests that the $\alpha 4$ isoform plays an important role in maintaining sperm active transmembrane Na^+ transport.

The transmembrane distribution of electrical charges, resulting from the uneven Na^+ and K^+ transport catalyzed by Na,K-ATPase contributes to maintaining the resting cell-membrane potential (2). Membrane potential determinations, using the fluorescent indicator [DiSC₃(5)], showed that the plasma membrane from $\alpha 4$ knockout spermatozoa was highly depolarized compared with that of wild-type sperm (Fig. 6C). This finding suggests that the ion gradients created by the $\alpha 4$ isoform are important in maintaining sperm-membrane potential.

Discussion

Previous studies have explored the function of $\alpha 4$ using selective pharmacological inhibition of this isoform with ouabain (7, 11, 12, 20). More recently, we have investigated the function of $\alpha 4$ using a mouse model with increased expression of the rat $\alpha 4$ isoform in sperm (15). Although these approaches have helped us understand some functional properties of $\alpha 4$, they did not directly address the role of $\alpha 4$ in sperm physiology and, more importantly, could not assess the relevance of the Na,K-ATPase $\alpha 4$ isoform in male fertility. We have studied this topic herein, using targeted deletion of the *Atp1a4* gene in mice. Homozygous mice with the disrupted *Atp1a4* gene show complete abolishment

of $\alpha 4$ polypeptide expression and loss of the typical high ouabain-sensitive Na,K-ATPase activity of $\alpha 4$ (13, 21).

The $\alpha 4$ -null mice present normal overall testis structure and histology. The conserved pattern in male germ-cell developmental stages in seminiferous tubules from $\alpha 4$ knockout mice suggests that the $\alpha 4$ isoform is not essential for the process of spermatogenesis, but rather that $\alpha 4$ is important for the differentiated sperm. This is supported by previous findings, showing that $\alpha 4$ is expressed after meiosis of the male germ cells and during late stages of spermatogenesis (7, 11, 22).

In agreement with an essential role of $\alpha 4$ in sperm function is the complete infertility of the knockout mice. Only male $\alpha 4$ -null mice are sterile, and the normal fertility of the homozygous female mice further proves the gender specificity in expression of the $\alpha 4$ isoform (9). Different from homozygous male mice, the $\alpha 4$ heterozygous male mice are able to generate offspring, despite the reduced motility of their sperm. Therefore, it appears that partial expression of $\alpha 4$, although not sufficient to support full sperm motility, is still adequate to maintain male fertility.

Disruption of the *Atp1a4* gene resulted in a drastic reduction of sperm motility, indicating an important role of $\alpha 4$ in sperm flagellar beat. This finding is consistent with the localization of the $\alpha 4$ polypeptide, which is expressed in the sperm tail (7, 11). Inhibition of $\alpha 4$ with ouabain has been shown to reduce sperm movement (11, 14); however, this produced a decrease in motility that was less pronounced than that caused by disruption of the *Atp1a4* gene. This finding emphasizes the higher efficiency of our genetic approach in blocking $\alpha 4$ activity over the pharmacological approach. Heterozygous $\alpha 4$ -null mice also have diminished sperm motility; however, to a lesser extent than the $\alpha 4$ -homozygous mice, suggesting that a gene dose effect is associated with expression of the $\alpha 4$ isoform. The dependence of sperm motility on differential expression of the Na,K-ATPase $\alpha 4$ isoform is also supported by the increase in sperm movement observed when the rat $\alpha 4$ isoform is overexpressed in mice (15).

Lack of the $\alpha 4$ isoform affects not only total sperm motility, but also a variety of parameters of sperm motility, suggesting




Fig. 6. Spermatozoa from Na,K-ATPase $\alpha 4$ -null mice display abnormal morphology and show altered $[Na^+]_i$ and membrane potential. (A) Sperm morphology. Sperm was plated onto glass coverslips, fixed using buffered formalin phosphate, and stained with Trypan Blue. Images were taken under bright field using a 60x objective in a Nikon Eclipse 80i scope (Nikon Instruments, Inc.). (Scale bar, 100 μm .) (B) $[Na^+]_i$ determinations in sperm using the fluorescent indicator Sodium Green Tetraacetate. (C) Sperm membrane-potential determinations using [DiSC₃(5)]. Bars represent the mean \pm SEM of three experiments. Values significantly different from the wild-type: * $P < 0.001$.

a general role of $\alpha 4$ in multiple aspects of sperm flagellar beat. In addition, $\alpha 4$ -null mice exhibit severe decrease in sperm hypermotility, which is required for sperm penetration through the egg vestments (17). Because $\alpha 4$ is not directly involved in the development of the sperm acrosomal reaction (15), the profound reduction in sperm motility and hypermotility represent central causes leading to the lack of fertilizing capacity of $\alpha 4$ -null sperm in vitro fertility assays, and to the infertility presented by the $\alpha 4$ -null mice. Therefore, sperm general motility and hyperactivation are main factors through which the Na,K-ATPase $\alpha 4$ isoform controls sperm fertility.

An important question regards the mechanisms by which the $\alpha 4$ isoform influences sperm motility. The $\alpha 4$ -null mice present a characteristic bend in the sperm flagellum. This abnormality in cell shape has been previously noted in wild-type sperm after treatment with ion channel blockers or after hypotonic shock (19), and in knockout mice in which the SLO3 potassium channel had been deleted (23). These changes in sperm shape have been ascribed to alterations in osmolarity and ion balance (19). The similar changes in sperm morphology we observe suggest that the ion gradients maintained by $\alpha 4$ are important in controlling sperm cytoplasmic ion homeostasis. Accordingly, the $[Na^+]_i$ in spermatozoa from $\alpha 4$ -null mice is increased compared with wild-type mice. Another expression of ion imbalance in the $\alpha 4$ -null mouse is the depolarization of the sperm plasma membrane. Sperm motility has been shown to depend on ion balance and specifically $[Na^+]_i$ levels (24). In addition, appropriate values of plasma membrane potential are required for sperm motility and during sperm capacitation (24). Moreover, cell membrane depolarization is associated with infertility in patients with asthenozoospermia (25). Our results highlight the importance of ion balance and membrane excitability as the mechanisms by which the $\alpha 4$ isoform supports sperm motility.

Our results also show that absence of $\alpha 4$ did not produce changes in expression of the ubiquitous $\alpha 1$ isoform. This absence of compensatory up-regulation of $\alpha 1$ in the $\alpha 4$ -null mice, suggests that regulation of $\alpha 1$ and $\alpha 4$ activity is independent from each other. This idea is also supported by our previous study, which found no reciprocal modulation of $\alpha 1$ expression and activity after expression of the rat $\alpha 4$ isoform in mice (15). The inability of $\alpha 1$ to substitute for $\alpha 4$ also suggests that these isoforms perform different roles in male fertility. In support of this, the $\alpha 1$ and $\alpha 4$ polypeptides have been shown to have distinct enzymatic properties (13). Although additional experiments will be necessary to define the function of $\alpha 1$ in sperm, it is clear that the $\alpha 4$ isoform plays an essential role in controlling Na^+ and K^+ exchange in sperm and in supporting fertility of the male gamete.

Taken together, our results are unique in demonstrating that the Na,K-ATPase $\alpha 4$ isoform is essential for sperm fertility both in vivo and in vitro and that the $\alpha 1$ subunit is not able to substitute for this role. If alterations in the *Atp1a4* gene produce the same defects in human sperm, our results open the possibility for the future use of the $\alpha 4$ isoform as a biomarker for male fertility. Finally, our data also provide compelling evidence for the use of the Na,K-ATPase $\alpha 4$ isoform as a target for male contraception.

Materials and Methods

Preparation of Knockout Animals. A mouse 129 BAC containing the *Atp1a4* genomic locus was used to generate the *Atp1a4* knockout gene-targeting vector. The BAC was subcloned into pSP72 vector (Promega Corporation) and a region of 6.8 Kb of the *Atp1a4* gene, including exons 5 through 8, was deleted and replaced by a neomycin expression cassette (Fig. 1A). The targeting vector was linearized with NotI and electroporated into BA1 hybrid (C57BL/6 \times 129/SvEv) ES cells for homologous recombination. After neomycin selection, isolated ES cell clones were screened by PCR. Clones were confirmed by Southern blotting following the scheme of Fig. 1B, and positive clones were further characterized by karyotyping. Targeted ES clones were microinjected into C57BL/6 blastocysts and these were implanted into pseudopregnant females. Resulting chimeras were mated to wild-type C57BL/6 mice to generate F1 heterozygous offspring and breeding continued to establish the founder line. Preparation of the knockout mice were performed at Ingenious Targeting Laboratory, Inc. (Stony Brook, NY). Genotype analysis was performed on tail biopsies by PCR on isolated genomic DNA using the REExtract-N-Amp Tissue PCR Kit (Sigma).

Sperm Isolation. All experimental protocols involving animals in this work were approved by the University of Kansas Medical Center Institutional Animal Care and Use Committee. Spermatozoa from wild-type, $\alpha 4$ heterozygous and null mice were obtained from the cauda of adult mice epididymides after swim-up of the cells, as previously described (14). Sperm was resuspended in modified Tyrode's medium, containing: 100 mM NaCl, 4.7 mM KCl, 1.2 mM KH_2PO_4 , 1.2 mM $MgSO_4$, 5.5 mM Glucose, 0.8 mM pyruvic acid, 4.8 mM lactic acid, 20 mM Hepes (pH 7.4), counted, and used for the different assays. For some experiments, sperm was capacitated in modified Tyrode's medium, supplemented with 1.7 mM $CaCl_2$, 25 mM sodium bicarbonate and 0.5% BSA.

Southern Blot. Southern blot was used to confirm the ES cell clones containing the $\alpha 4$ construct. DNA was first digested with either NheI or PstI and electrophoretically separated on a 0.8% agarose gel. After transfer to a nylon membrane, DNA was hybridized with probes targeted to the 5' external region of the NheI digested DNA, or to the 3' internal region of the PstI digested DNA. After washing off the nonhybridized probe, membranes were processed for autoradiography.

PCR. PCR was used for analysis of genomic DNA. Reactions were performed using primers directed to Na,K-ATPase $\alpha 4$ isoform sequences (sense 5'-CAG-GAGAGCAGGATCGTGGAC-3') and antisense (5'-GGTGATACATGCTAGTCACTG-AGGTGCC-3') and to the neomycin cassette sequence (5'-GCATCGCCTTCTA-TCCGCTTCTTG-3'). Procedures were as described (7), using a first cycle of 30 s at 94 $^{\circ}C$, followed by 30 cycles at 94 $^{\circ}C$ for 15 s, 60 $^{\circ}C$ for 15 s, and 72 $^{\circ}C$ for 30 s.

RT-PCR. Total testis RNA was prepared using TRIzol reagent according to the supplier specifications (Invitrogen). A total of 10.0 μg RNA was incubated in 100 μl of 1 \times DNase buffer, 1 U RNase-free DNase I (Roche) at 37 $^{\circ}C$ for 20 min. Complementary DNA was prepared by reverse transcription of 1.0 μg of total

RNA using SuperScript II Reverse Transcriptase (Invitrogen), as described (7). The resulting first-strand cDNA was subjected to amplification using primers specific for the mouse Na,K-ATPase $\alpha 4$ isoform (sense 5'-GAGGAGCAAC-CACGGGAAAACG-3') and (antisense 5'-GCTAGGCAAGTCAAGAGCAGAA-CC-3'), and for the glyceraldehyde phosphate dehydrogenase (GAPDH) sense (5'-GGAGATTGTGCCATCAACG-3') and antisense (5'-CACAAATGCCAAAGTTG-TCATGG-3') primers. Amplified DNA fragments were identified by electrophoresis in 1% agarose gels stained with ethidium bromide.

Immunoblot. Protein expression was analyzed by SDS/PAGE (7.5% gel) and immunoblotting, as previously described (7). The Na,K-ATPase $\alpha 1$ and $\alpha 4$ isoforms were identified using antisera generated against specific regions of the polypeptides in rabbit and chicken respectively. Horseradish peroxidase-conjugated secondary antibodies (Jackson ImmunoResearch Laboratories, Inc.) and chemiluminescence were used for detection.

Na,K-ATPase and Ouabain Binding Assays. Na,K-ATPase activity was assayed on sperm homogenates, through determination of the initial rate of release of $^{32}\text{P}_i$ from γ [^{32}P]-ATP, as previously described (14). The ATPase activity of 10 μg of total protein per sample was measured in a final volume of 0.25 mL in medium containing 120 mM NaCl, 30 mM KCl, 3 mM MgCl_2 , 0.2 mM EGTA, 30 mM Tris-HCl (pH 7.4), 3 mM ATP with 0.2 μCi γ [^{32}P]-ATP in the presence and absence of the indicated ouabain concentrations. Curve fitting of the experimental data were performed using a Marquardt least-squares non-linear regression computing program (Sigma Plot; Jandel Scientific). Ouabain binding capacity of spermatozoa from wild-type and null mice was measured using bodipy-ouabain (Invitrogen). Sperm was placed in modified Tyrode's medium at a concentration of 2×10^6 cells/mL, and 10^{-8} M bodipy-ouabain was added. Samples in which 1 mM unlabeled ouabain was also added were used as a control for specific bodipy-ouabain binding. After incubation for 20 min at 37 °C, cells were washed twice for 5 min in modified Tyrode's medium. Finally, fluorescence was measured at 488 nm, using a LSRII flow cytometer (BD Biosciences).

Membrane-Potential Assays. Approximately 2×10^6 cells/mL were suspended in modified Tyrode's medium and membrane potential was determined using the fluorescent indicator [DiSC₃(5)]. Calibration of the fluorescence

changes into millivolts and determination of the cell membrane potential was calculated following protocols previously described (14).

Intracellular Na⁺ Measurements. The $[\text{Na}^+]_i$ was measured in sperm (20×10^6 cells/mL) in modified Tyrode's medium with Sodium Green Tetraacetate, as described previously (14).

Sperm-Motility Assays. Approximately 3×10^6 cells from wild-type, $\alpha 4$ heterozygous and null mice were used to determine sperm motility, as previously described (14). Samples were analyzed by CASA, using the Minitube Sperm-Vision Digital Semen Evaluation system (version 3.5, Penetrating Innovations). Total sperm motility and different parameters of sperm movement were analyzed using analytical setup parameters defined elsewhere (14).

In Vitro Fertilization Assays. Cumulus-oocyte masses from superovulated CD-1 female mice were recovered after stimulation with 5 IU of PG 600 (Intervet) and 5 IU of human chorionic gonadotrophin. Spermatozoa were collected from the cauda epididymis after swim-up. In vitro fertilization assays were performed following standard procedures with some modifications (14). Approximately 10^5 sperm/mL from wild-type or $\alpha 4$ -null sperm were incubated with oocytes in Cook's medium (Research Vitro Fert K-RVFE-50; Cook Medical) for 6 h at 37 °C, with 5% CO₂ and 5% O₂ under oil. After removing and washing the oocytes in Cook's medium, they were cultured overnight. Fertilization success was determined as the development of two-cell stage embryos.

Statistical Analysis. Statistical significance of differences between controls and ouabain treated samples was determined by the Student's *t* test, using Sigma Plot software (Jandel Scientific). Statistical significance was defined as $P < 0.05$.

ACKNOWLEDGMENTS. We thank Drs. V. Chennathuzuzhi and Susana Gonzalo for critical discussions and reading the manuscript, Ingenious Targeting Laboratory (Stony Brook, NY) for developing the knockout mice used in this study, and Dr. Melissa Larson for her help with the in vitro fertilization assays. This work was supported by Grants HD043044 and HD055763 from the National Institutes of Health.

- Kaplan JH (2002) Biochemistry of Na,K-ATPase. *Annu Rev Biochem* 71:511–535.
- Féraille E, Doucet A (2001) Sodium-potassium-adenosinetriphosphatase-dependent sodium transport in the kidney: hormonal control. *Physiol Rev* 81:345–418.
- Morth JP, et al. (2007) Crystal structure of the sodium-potassium pump. *Nature* 450: 1043–1049.
- Blanco G, Mercer RW (1998) Isozymes of the Na-K-ATPase: Heterogeneity in structure, diversity in function. *Am J Physiol* 275:F633–F650.
- Blanco G (2005) Na,K-ATPase subunit heterogeneity as a mechanism for tissue-specific ion regulation. *Semin Nephrol* 25:292–303.
- Mobasheri A, et al. (2000) Na⁺, K⁺-ATPase isozyme diversity; comparative biochemistry and physiological implications of novel functional interactions. *Biosci Rep* 20: 51–91.
- Wagoner K, Sanchez G, Nguyen AN, Enders GC, Blanco G (2005) Different expression and activity of the alpha1 and alpha4 isoforms of the Na,K-ATPase during rat male germ cell ontogeny. *Reproduction* 130:627–641.
- Uesugi S, Yamazoe S (1966) Presence of sodium-potassium-stimulated ATPase in boar epididymal spermatozoon. *Nature* 209:403.
- Shamraj OI, Lingrel JB (1994) A putative fourth Na⁺,K⁽⁺⁾-ATPase alpha-subunit gene is expressed in testis. *Proc Natl Acad Sci USA* 91:12952–12956.
- Blanco G, Sánchez G, Melton RJ, Tourtellotte WG, Mercer RW (2000) The alpha4 isoform of the Na,K-ATPase is expressed in the germ cells of the testes. *J Histochem Cytochem* 48:1023–1032.
- Woo AL, James PF, Lingrel JB (2000) Sperm motility is dependent on a unique isoform of the Na,K-ATPase. *J Biol Chem* 275:20693–20699.
- Sanchez G, Nguyen AN, Timmerberg B, Tash JS, Blanco G (2006) The Na,K-ATPase alpha4 isoform from humans has distinct enzymatic properties and is important for sperm motility. *Mol Hum Reprod* 12:565–576.
- Blanco G, Melton RJ, Sánchez G, Mercer RW (1999) Functional characterization of a testes-specific alpha-subunit isoform of the sodium/potassium adenosinetriphosphatase. *Biochemistry* 38:13661–13669.
- Jimenez T, Sanchez G, Wertheimer E, Blanco G (2006) Activity of the Na,K-ATPase alpha4 isoform is important for membrane potential, intracellular Ca²⁺ and pH, to maintain motility in rat spermatozoa. *Reproduction* 139:835–845.
- Jimenez T, et al. Increased expression of the Na,K-ATPase alpha4 isoform enhances sperm motility in transgenic mice. *Biol Reprod*, in press.
- Visconti PE, et al. (1998) The molecular basis of sperm capacitation. *J Androl* 19: 242–248.
- Suarez SS (2008) Control of hyperactivation in sperm. *Hum Reprod Update* 14: 647–657.
- Pedersen BP, Buch-Pedersen MJ, Morth JP, Palmgren MG, Nissen P (2007) Crystal structure of the plasma membrane proton pump. *Nature* 450:1111–1114.
- Cooper TG, et al. (2004) Mouse models of infertility due to swollen spermatozoa. *Mol Cell Endocrinol* 216:55–63.
- Woo AL, James PF, Lingrel JB (2002) Roles of the Na,K-ATPase alpha4 isoform and the Na⁺/H⁺ exchanger in sperm motility. *Mol Reprod Dev* 62:348–356.
- Woo AL, James PF, Lingrel JB (1999) Characterization of the fourth alpha isoform of the Na,K-ATPase. *J Membr Biol* 169:39–44.
- Rodova M, Nguyen AN, Blanco G (2006) The transcription factor CREMtau and cAMP regulate promoter activity of the Na,K-ATPase alpha4 isoform. *Mol Reprod Dev* 73: 1435–1447.
- Santi CM, et al. (2010) The SLO3 sperm-specific potassium channel plays a vital role in male fertility. *FEBS Lett* 584:1041–1046.
- Darszon A, et al. (2006) Sperm channel diversity and functional multiplicity. *Reproduction* 131:977–988.
- Calzada L, Tellez J (1997) Defective function of membrane potential (psi) on sperm of infertile men. *Arch Androl* 38:151–155.