

# Editorial Expression of Concern

## CELL BIOLOGY

PNAS is publishing an Editorial Expression of Concern regarding the following article: “New class of transcription factors controls flagellar assembly by recruiting RNA polymerase II in *Chlamydomonas*,” by Lili Li, Guangmei Tian, Hai Peng, Dan Meng, Liang Wang, Xiao Hu, Cheng Tian, Miao He, Junfei Zhou, Lihong Chen, Cheng Fu, Weixiong Zhang, and Zhangfeng Hu, which was first published April 9, 2018, and appeared in issue 17, April 24, 2018; 10.1073/pnas.1719206115 (*Proc. Natl. Acad. Sci. U.S.A.* **115**, 4435–4440). We have recently learned that, since 2019, the authors have not shared the mutant strains described in the above-noted article to qualified investigators due to ongoing research. This is a violation of PNAS editorial policy, and we are publishing this statement to alert readers. PNAS authors must ensure that all unique reagents described in their published papers are available to qualified researchers, and any restrictions on the sharing of materials must be disclosed to the editorial office at the time of submission (<https://www.pnas.org/authors/editorial-and-journal-policies#materials-and-data-availability>).

May R. Berenbaum  
*Editor-in-Chief*

Published under the [PNAS license](#).

Published June 7, 2021.

[www.pnas.org/cgi/doi/10.1073/pnas.2108930118](http://www.pnas.org/cgi/doi/10.1073/pnas.2108930118)



# New class of transcription factors controls flagellar assembly by recruiting RNA polymerase II in *Chlamydomonas*

Lili Li<sup>a,1</sup>, Guangmei Tian<sup>a,1</sup>, Hai Peng<sup>a,1</sup>, Dan Meng<sup>b</sup>, Liang Wang<sup>c</sup>, Xiao Hu<sup>a</sup>, Cheng Tian<sup>a</sup>, Miao He<sup>a</sup>, Junfei Zhou<sup>a</sup>, Lihong Chen<sup>a</sup>, Cheng Fu<sup>d</sup>, Weixiong Zhang<sup>a,e,f,2</sup>, and Zhangfeng Hu<sup>a,2</sup>

<sup>a</sup>Institute for Systems Biology, Jiangnan University, Wuhan, 430056 Hubei, China; <sup>b</sup>Tianjin Key Laboratory of Food and Biotechnology, School of Biotechnology and Food Science, Tianjin University of Commerce, 300134 Tianjin, China; <sup>c</sup>School of Life Sciences, Jiangsu Normal University, Xuzhou, 221116 Jiangsu, China; <sup>d</sup>Institute for Interdisciplinary Research, Jiangnan University, Wuhan, 430056 Hubei, China; <sup>e</sup>Department of Computer Science and Engineering, Washington University in St. Louis, St. Louis, MO 63130; and <sup>f</sup>Department of Genetics, Washington University in St. Louis, St. Louis, MO 63130

Edited by Chris Kintner, Salk Institute, and accepted by Editorial Board Member Kathryn V. Anderson March 6, 2018 (received for review November 3, 2017)

Cells have developed regulatory mechanisms that underlie flagellar assembly and maintenance, including the transcriptional regulation of flagellar genes, an initial step for making flagella. Although transcriptional regulation of flagellar gene expression is required for flagellar assembly in *Chlamydomonas*, no transcription factor that regulates the transcription of flagellar genes has been identified. We report that X chromosome-associated protein 5 (XAP5) acts as a transcription factor to regulate flagellar assembly in *Chlamydomonas*. While XAP5 proteins are evolutionarily conserved across diverse organisms and play vital roles in diverse biological processes, nothing is known about the biochemical function of any member of this important protein family. Our data show that loss of XAP5 leads to defects in flagellar assembly. Post-translational modifications of XAP5 track flagellar length during flagellar assembly, suggesting that cells possess a feedback system that modulates modifications to XAP5. Notably, XAP5 regulates flagellar gene expression via directly binding to a motif containing a CTGGGGTG-core. Furthermore, recruitment of RNA polymerase II (Pol II) machinery for transcriptional activation depends on the activities of XAP5. Our data demonstrate that, through recruitment of Pol II, XAP5 defines a class of transcription factors for transcriptional regulation of ciliary genes. This work provides insights into the biochemical function of the XAP5 family and the fundamental biology of the flagellar assembly, which enhance our understanding of the signaling and functions of flagella.

cilia | XAP5 | transcription factor | transcriptional regulation | *Chlamydomonas*

Cilia and flagella are subcellular organelles that protrude from the surface of cells (1). They have evolved to play crucial roles in cell motility, sensory reception, and signal transduction in various organisms (2). Cilia are highly conserved throughout evolution and are structurally dynamic, undergoing growth and resorption during the cell cycle (3). Ciliary dysfunction can cause a substantial number of human genetic disorders, including polycystic kidney disease, Joubert syndrome, and Bardet-Biedl syndrome, which are collectively referred to as “ciliopathies” (4).

Previous studies have shown that cilia are complex structures with hundreds of components involved in structure and function (5–7). These studies have also identified a number of human ciliopathy disease genes. They are involved in intraflagellar transport (IFT) and Bardet-Biedl syndrome (BBS) and are widely distributed and highly conserved among organisms with cilia and flagella (8, 9). IFT is a complex transport mechanism that regulates the trafficking of ciliary proteins along the axonemal doublet microtubules and is required for assembly and maintenance of cilia (8). The BBS subunits form a complex to transport ciliary proteins into the ciliary compartment (9). The expression of ciliary components is regulated at the transcriptional level, an important step during cilia formation (7, 10). Transcriptional regulation of ciliary gene expression was first recorded via studies of flagellar regeneration

after experimentally inducing deflagellation in *Chlamydomonas* (11). Cells regenerate half-length flagella after flagellar amputation by using a preexisting pool of cytoplasmic proteins (12). Nevertheless, the synthesis of mRNA and proteins of the flagellar components must be switched on to complete reassembly of flagella (11, 12). Although the specific transcriptional regulation of ciliary genes' expression during cilia assembly has been widely studied in *Chlamydomonas*, the mechanisms by which cilia assembly is programmed at the transcriptional level remain elusive.

Emerging evidence suggests that transcription regulation may be critically important in the control of different functional and structural modules to generate ciliary diversity in various cell types (10, 13–15). The regulatory factor X (RFX) family proteins share a highly conserved winged-helix DNA-binding domain and are required for the transcriptional regulation of ciliary genes in primary cilia (13). FOXJ1 is a forkhead/winged-helix family transcription factor that is necessary for motile ciliogenesis in

## Significance

Transcriptional regulation of flagellar genes controls an initial step in flagellar assembly. In this study, we show that XAP5, a conserved protein of unknown function, defines a class of transcription factor for transcriptional regulation of genes involved in flagellar assembly. Phosphorylation of X chromosome-associated protein 5 (XAP5) during flagellar regeneration tracks flagellar length. Remarkably, recruitment of RNA polymerase II (Pol II) machinery for transcriptional activation depends on the activities of XAP5. Our data demonstrate that XAP5 functions as a transcription factor for transcriptional regulation of flagellar genes through recruitment of RNA Pol II. Our results enhance our understanding of the biochemical function of the XAP5 family and the transcriptional regulation of flagellar assembly.

Author contributions: L.L., G.T., H.P., W.Z., and Z.H. designed research; L.L., G.T., H.P., D.M., L.W., X.H., C.T., M.H., J.Z., L.C., and C.F. performed research; Z.H. contributed new reagents/analytic tools; L.L., G.T., H.P., D.M., L.W., W.Z., and Z.H. analyzed data; and W.Z. and Z.H. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission. C.K. is a guest editor invited by the Editorial Board.

Published under the PNAS license.

Data deposition: The *Chlamydomonas* XAP5 cDNA and protein sequences have been deposited in the National Center for Biotechnology Information (NCBI) GenBank data library (<https://www.ncbi.nlm.nih.gov/nuccore/KU361202>) (ID codes KU361202.1 and AMY57988.1). The RNA-sequencing data are available at the NCBI Sequence Read Archive (SRA) database (<https://www.ncbi.nlm.nih.gov/sra/SRP131193>) (accession no. SRP131193).

<sup>1</sup>L.L., G.T., and H.P. contributed equally to this work.

<sup>2</sup>To whom correspondence may be addressed. Email: weixiong.zhang@wustl.edu or huzhangfengzhf@163.com.

This article contains supporting information online at [www.pnas.org/lookup/suppl/doi:10.1073/pnas.1719206115/-DCSupplemental](http://www.pnas.org/lookup/suppl/doi:10.1073/pnas.1719206115/-DCSupplemental).

Published online April 9, 2018.




XAP5 is not required by processes essential for cell size or growth and supporting the idea that there is no necessary association between flagella phenotypes and cell size.

**Posttranslational Modification of XAP5 During Flagellar Assembly.** XAP5 proteins resided preferentially in the nucleus and seldom existed in the cytoplasm (Fig. S2 A–C). XAP5 contains a predicted nuclear-localization signal (NLS), KRKR, at amino acids 106–109 (Fig. S2D). In contrast to the wild-type gene, both *XAP5-ΔNLS-HA* and *XAP5-K106A-HA* failed to rescue the flagellar-assembly defects (Fig. S2 E and F). Deletion or mutation of the NLS KRKR disrupted the nuclear localization of XAP5 and led to a diffused cytoplasmic distribution (Fig. S2 E and G). Thus, the appropriate nuclear localization of XAP5 mediated by the NLS is required for flagellar assembly.

To understand the role that XAP5 plays in flagellar assembly, we examined its expression and subcellular localization. The expression and subcellular location of XAP5 had no significant change during flagellar regeneration (Fig. 2A and Fig. S2H). Unexpectedly, a posttranslational modification in XAP5 was reflected by a change in its electrophoretic mobility (Fig. 2A and Fig. S3 A and B). Importantly, XAP5 was modified immediately after deflagellation, and the modification of XAP5 disappeared as the flagella reached nearly their full length (Fig. 2A and Fig. S3 A and B). After lysates of *xap5::XAP5-HA* cells were incubated with a phosphatase, XAP5 was not modified (Fig. 2B), suggesting that XAP5 was phosphorylated during flagellar regeneration. To determine whether the phosphorylation of XAP5 occurred under the physiological conditions, we analyzed the status of XAP5 during the cell cycle. As expected, XAP5 was also phosphorylated during flagellar assembly after mitosis (Fig. S3C) (3). Therefore, XAP5 is an important regulator of flagellar assembly that occurs under physiological and nonphysiological conditions, and the phosphorylation of XAP5 may play a vital role in flagellar assembly. Nuclear XAP5, rather than cytoplasmic XAP5, underwent phosphorylation in response to pH shock-induced deflagellation (Fig. 2C). Correspondingly, the cytoplasmic localized *XAP5-ΔNLS-HA* and *XAP5-K106A-HA* mutants failed to undergo a mobility shift-associated phosphorylation after deflagellation (Fig. 2D). Thus, nuclear localization of XAP5 is necessary for its phosphorylation during flagellar regeneration.

Deflagellated cells regenerated to only half the original length after treatment with staurosporine (Fig. 2E), a protein kinase inhibitor which has been reported to block the phosphorylation of nuclear protein that may play a crucial role in transcriptional activation (23). Consistent with prior studies, blocking protein synthesis by cycloheximide resulted in regeneration of half-length flagella and a slow regeneration of flagella by rapamycin after deflagellation (Fig. 2E) (24), implying that cells have developed diverse regulatory mechanisms to ensure the formation of functional cilia. XAP5 did not undergo an obvious mobility shift in cells treated with staurosporine (Fig. S2 D and E). These data raise the possibility that staurosporine affects flagellar gene transcription, probably through the regulation of XAP5. Furthermore, in cells treated with actinomycin D and deflagellated by two successive pH shocks, flagellar assembly was completely inhibited after the second amputation (Fig. 2F), showing that inhibition of transcription impairs flagellar assembly.


To understand whether phosphorylation of XAP5 is required for flagellar assembly, we expressed in *xap5* HA-tagged XAP5 constructs with multipoint mutations or a single-point mutation at serine/threonine/tyrosine residue(s). We found that S119, a consensus phosphorylation site in the coiled-coil region of XAP5, is required for flagellar assembly (Fig. S4 A–C). The wild-type XAP5 and a phosphomimetic (S119D) mutant fully complemented the flagellar type, whereas phosphodeficient (S119A) mutant cells were aflagellate (Fig. S4 B and C). The S119D and S119A mutant proteins did not undergo a mobility shift-associated phosphorylation after flagellar amputation (Fig. S4D), and these mutations did not affect the nuclear localization of XAP5 (Fig. S4E). Thus, S119 phosphorylation of XAP5 is essential for flagellar assembly. Furthermore, the



**Fig. 2.** XAP5 is phosphorylated in response to signals that induce flagellar assembly. (A) Immunoblot analysis of XAP5 during flagellar regeneration using an antibody against XAP5. Kinetics of flagellar regeneration upon shock-induced deflagellation is shown above the blot. Coomassie Brilliant Blue (CBB) showed an equal loading of all of the samples. H3, histone H3 antibody; pdf, cell sample before deflagellation. (B) Western blot analysis of whole-cell lysates from control and deflagellated *xap5::XAP5-HA* cells that were hatched with or without phosphatase (Ptase). (C) Control and deflagellated *xap5::XAP5-HA* cells were fractionated into whole-cell (WC), cytoplasm (Cyt), and nuclei (Nuc) fractions and were analyzed by immunoblotting. Antibodies against NAB1 and histone H3 were used to mark the cytoplasm and nuclei, respectively. (D) Immunoblot analysis of cell lysates from control and deflagellated *xap5* cells expressing the HA-tagged wild-type,  $\Delta$ NLS, and K106A-XAP5 protein. (E) Inhibition of flagellar regeneration by staurosporine. Cells were triggered to regenerate flagella by pH shock in the presence of actinomycin D (AD) (100  $\mu$ g/mL), cycloheximide (CHX) (10  $\mu$ g/mL), rapamycin (Rap) (50  $\mu$ M), or staurosporine (Stau) (1  $\mu$ M). Con, control. Data in A and E represent mean  $\pm$  SD. One flagellum from at least 50 cells was measured at each time point in three independent experiments.

phosphomimetic (S119D) mutant cells subjected to staurosporine treatment regrew approximately half-length flagella after amputation (Fig. S4 F and G), suggesting that staurosporine also suppresses other processes involved in flagellar assembly.

**XAP5 Modulates the Transcription of Ciliary Genes.** To elucidate the mechanism by which XAP5 regulates flagellar assembly, we studied the possible function of XAP5 in modulating the expression of conserved genes necessary for ciliary assembly. We analyzed some of IFT- and BBS-related genes that broadly exist and are highly conserved in ciliated organisms. We found that *IFT144*, *IFT70*, and *BBS8* were down-regulated in *xap5* cells (Fig. 3A), whereas the expression levels of *IFT139*, *IFT20*, *BBS4*, and  $\alpha$ -tubulin in *xap5* cells were not changed significantly (Fig. 3A). However, the expression of *IFT144*, *IFT70*, and *BBS8* appeared to be normal in other flagellaless mutants (Fig. 3B), suggesting



**Fig. 3.** XAP5 modulates the transcription of ciliary genes via directly binding to the CTGGGGTG motif. (A) Analysis of the transcript abundance of flagellar-associated genes in wild-type and *xap5* cells by real-time qPCR. (B) Quantification of the transcript abundance of ciliary genes in mutants defective in flagellar assembly. (C) Changes in the relative expression level of genes in wild-type and *xap5* cells after pH shock-induced deflagellation. (D) DIC images of wild-type, *ift70*, and rescued (*ift70::IFT70*) cells. Palmelloid *ift70* mutant cells were incubated with autolysin and released from the mother cell wall. (Scale bar, 5  $\mu$ m.) (E) Schematic drawing representing the potential XAP5 target sites in the promoter regions of the ciliary genes. Data in A–C represent the mean  $\pm$  SD of three independent experiments. \* $P < 0.05$ . N.S., not significant ( $P > 0.05$ ).

that the lowered transcription levels of *IFT144*, *IFT70*, and *BBS8* in *xap5* cells were unrelated to defects in flagellar assembly.

*IFT144*, *IFT70*, and *BBS8* were up-regulated in wild-type cells during flagellar regeneration while their abundances in *xap5* cells were comparable (Fig. 3C). However, the *BBS4* gene was up-regulated in *xap5* compared with wild-type cells after flagellar amputation (Fig. 3C). These data suggest that the transcriptional regulation of flagellar genes might be classified into two types: XAP5-dependent and XAP5-independent. In addition, the expression abundances of *IFT144*, *IFT70*, and *BBS8* were significantly reduced in *xap5::XAP5- $\Delta$ NLS-HA* and *xap5::XAP5-K106A-HA* cells compared with the wild-type cells (Fig. S5A), indicating that nuclear localization of XAP5 is essential for ciliary gene transcription. Moreover, *IFT70* proteins in *xap5* cells exhibited distinctly decreased levels compared with the levels in wild-type cells and flagellaless mutants (Fig. S5B–G). Consistent with the previous findings (25), the *IFT70*-null mutation impaired flagellar assembly (Fig. 3D and Fig. S5G). Collectively, these results reveal that XAP5 regulates flagellar assembly via modulating the transcription of XAP5-dependent flagellar genes.

**Sequence-Specific Binding of XAP5 in the Promoters of Ciliogenic Genes.** To assess whether XAP5 directly regulates ciliogenic gene expression, we first performed ChIP assays with an antibody against HA. Remarkably, the promoters of the candidate XAP5-target genes *IFT144*, *IFT70*, and *BBS8* were preferentially enriched in the *xap5::XAP5-HA* ChIP samples, especially near the predicted transcription start site (TSS) (Fig. S6A). Furthermore, XAP5 occupancy in the promoter regions was significantly elevated after flagellar detachment (Fig. S6B). However, when cells were treated with staurosporine, XAP5 occupancy exhibited no significant change during flagellar regeneration (Fig. S6B), implying that posttranslational modification directed the localization of XAP5 in chromatin regions. Therefore, XAP5 is presumably a potential factor affecting the activation of *IFT144*, *IFT70*, and *BBS8* promoters.

We observed that the 298-bp, 254-bp, and 224-bp regions upstream of the TSSs of *IFT144*, *IFT70*, and *BBS8*, respectively, were required for their normal promoter activities (Fig. S6C). Intriguingly, a BLAST analysis of these regions revealed a conserved CTGGGGTG sequence located immediately upstream of the TSS (Fig. 3E). An analysis by EMSA showed that XAP5 could bind directly to the motif (Fig. S6D and E). Together, these results show that XAP5 regulates ciliary gene expression through direct binding to the CTGGGGTG core sequence motif in the ciliary gene promoters.

To investigate the potential scope of involvement of XAP5 in the transcription of flagellar and nonflagellar genes, we carried out genome-wide profiling of gene expression in *xap5* and in wild-type cells using deep sequencing (RNA-seq). The results showed that 2,126 genes, including 267 flagellar-associated genes, were down-regulated by at least twofold in *xap5* mutants (Fig. S7A). A gene ontology (GO) analysis revealed that these differentially expressed genes were enriched with functions for signal transduction-, cilium-dependent cell motility-, and cilium assembly-related biological processes (Fig. S7B), in motile cilium-, microtubule-, and intraciliary transport particle B (IFT B)-involved cellular components (Fig. S7C), as well as motor activity, structural constituent of cytoskeleton, and calcium ion binding in molecular functions (Fig. S7D).

We identified 111 known flagellar genes that were down-regulated in *xap5* cells (Fig. S7E and Table S1). Importantly, 106 of these genes contained the putative XAP5-binding motifs and thus may be directly regulated by XAP5 (Fig. S7F and G and Table S1). The remaining five flagellar genes did not contain the putative binding motif (Fig. S7G). Their decreased expression might be due to changes in the abundance of other flagellar gene transcripts. We also examined the expression of three XAP5-dependent flagellar genes (*ODAI3*, *FAP147*, and *FAP5*) and one XAP5-independent flagellar gene (*IFT20*) after amputation induced by pH shock in the wild-type and *xap5* cells. The expression of *IFT20* was elevated after pH shock in both wild-type and *xap5* cells. In contrast, the three XAP5-dependent


flagellar genes were up-regulated only in wild-type cells after flagellar amputation but were not significantly changed in *xap5* cells (Fig. S7H). Additionally, the abundance of XAP5-dependent flagellar genes was significantly decreased in the phosphodeficient (*S119A*) mutant cells compared with the wild-type XAP5 and the phosphomimetic (*S119D*) mutant cells (Fig. S7I). Interestingly, the expression of the XAP5-dependent flagellar genes increased slightly in the *S119D* mutant compared with wild-type XAP5 cells (Fig. S7I). However, no change in the expression levels of two XAP5-independent flagellar genes (*IFT20* and *BBS4*) was observed in these cells (Fig. S7I). Thus, proper phosphorylation of XAP5 is required for normal transcriptional control of XAP5-dependent flagellar genes.

**XAP5 Facilitates Recruitment of the Polymerase II Apparatus to Promoters.** To directly demonstrate that XAP5 functions as a positive transcriptional activator of *IFT144*, *IFT70*, and *BBS8*, fragments of a putative promoter region containing the CTGGGGTG motif or a mutant form, AGTTTGTG, were assessed for their ability to activate the expression of a luciferase reporter gene. The result of luciferase activities showed that the CTGGGGTG motif in the promoter regions of *IFT144*, *IFT70*, and *BBS8* was critical for the expression of these genes (Fig. S8A and B). Importantly, the *ift70* mutant phenotype was rescued by a transformation with the wild-type promoter driving *IFT70* expression. In addition, about 28% of transformants rescued the mutant flagellar phenotype (Fig. S8C). None of the transformants recovered the flagellar phenotype when *ift70* mutants were transformed with a mutated promoter driving *IFT70* expression (Fig. S8C). Therefore, XAP5 is a potent transactivator of *IFT144*, *IFT70*, and *BBS8* genes. Furthermore, we

investigated the capacity of the promoters of wild-type ciliary genes to activate the expression of the luciferase (*Luc*) reporter gene in the wild type and *xap5* mutant (Fig. S8D). The *xap5* mutant showed a lower relative luminescence unit (RLU) average compared with the wild type (Fig. S8D). Additionally, the RLU value of the wild-type *IFT70* promoter directing the *Luc* gene was comparable in the wild type and flagellaless mutants (Fig. S8E), indicating that ciliary gene promoter activities are unrelated to ciliary defects.

A hallmark function of XAP5 is its direct regulation of ciliary gene expression, indicating the possibility that XAP5 could recruit the polymerase II (Pol II) machinery to the ciliary gene promoter for transcriptional activation. To further verify that loss of XAP5 reduces the levels of transcription of target genes, a ChIP analysis revealed that the occupancy of Pol II at the promoter regions of *IFT144*, *IFT70*, and *BBS8* genes with an antibody recognized the tandem heptapeptide repeats in the C-terminal domain (CTD) of Pol II's largest subunit (RBP1) (Fig. S8F and G). An enrichment of Pol II at the target genes was observed in wild-type cells (Fig. 4A). In contrast, the occupancy of the Pol II apparatus at the *IFT144*, *IFT70*, and *BBS8* genes was reduced significantly in *xap5* cells (Fig. 4A).

We further reasoned that the ciliary genes were up-regulated during ciliogenesis as a result of the accumulation of Pol II at the genes. Strikingly, increased Pol II occupancy at ciliary genes was accompanied by a simultaneously increased presence of XAP5 at the proximal promoters after pH shock in *xap5::XAP5-HA* cells (Fig. 4B). In contrast, the occupancy of Pol II was not significantly changed in *xap5* mutants during flagellar regeneration (Fig. 4B). The component of the Pol II machinery was present in the immunoprecipitates prepared with an antibody against HA



**Fig. 4.** XAP5-mediated recruitment of the Pol II machinery to ciliary gene promoters. (A) ChIP-qPCR analysis of the indicated genes using three independently prepared samples. ChIP was performed in wild-type and *xap5* cells with or without (Control) an antibody against RPB1. The regions amplified from chromatin immunoprecipitates by qPCR are indicated below each gene. (B) Detection of Pol II occupancy at ciliary genes in *xap5::XAP5-HA* and *xap5* cells after pH shock-induced deflagellation (Defla). (C) Coimmunoprecipitation of XAP5 and Pol II machinery. Isolated nuclei from cells in the presence or absence of staurosporine were immunoprecipitated with antibody against HA or with preimmune IgG, followed by Western blotting using antibodies against HA and RPB1, respectively. (D) Detection of Pol II occupancy at ciliary genes by ChIP-qPCR in control and deflagellated wild-type cells in the presence or absence of staurosporine. Data in A, B, and D represent the mean  $\pm$  SD of three independent experiments. \* $P < 0.05$ ; N.S., not significant ( $P > 0.05$ ).

and was absent in the control group (preimmune IgG) (Fig. 4C). Importantly, phosphorylated XAP5 had a stronger affinity to the Pol II machinery than did nonphosphorylated XAP5 (Fig. 4C). Furthermore, *xap5::XAP5-HA* cells treated with staurosporine failed to accumulate Pol II at target genes after deflagellation (Fig. 4C and D). Combined, these results reveal the importance of phosphorylated XAP5-mediated recruitment of the Pol II machinery for transcriptional activation, suggesting that XAP5 acts as a transcription factor for transcriptional regulation of flagellar genes.

## Discussion

The specific transcriptional induction of ciliary genes is precisely controlled and coordinated during cilia formation, and inhibition of transcription impairs ciliogenesis (10). Although the significance of the transcriptional regulation of ciliary gene expression was initially documented in *Chlamydomonas* (11), the mechanisms underlying this regulation remain poorly understood. To our knowledge, the mechanisms and genes that regulate the transcription and expression of ciliary genes have not been reported in *Chlamydomonas* or other unicellular organisms.

It has been generally acknowledged that the last eukaryotic common ancestor is a flagellated unicellular organism (26). Despite flagellar genes being under direct transcriptional regulation in both unicellular and multicellular organisms, the evolution of the transcriptional program controlling flagellar assembly is largely unknown (26). Previous studies have shown that RFX transcription factors, FOXJ1 transcription factors, and ciliary genes all evolved independently (26–28). The transcriptional control of ciliary genes generally regulates the formation of one type of cilium in unicellular organisms. Nevertheless, in multicellular organisms, many ciliary genes are differentially regulated with cell type-specific patterns of expression to generate ciliary diversity (10). RFX and FOXJ1 are key regulators of ciliary gene expression in animals (10). However, both are absent from many unicellular organisms, including *Chlamydomonas*, indicating that the transcriptional control of the ciliogenesis is fundamentally different in unicellular and multicellular organisms.

Previous data suggest that XAP5 proteins play vital roles in many biological processes (17–20). However, little is known about the precise molecular function of the conserved nucleus-localized protein XAP5. In the current study, an in vitro binding analysis with predicted transcription regulatory elements revealed that XAP5 could specifically recognize and bind to a motif with a CTGGGGTG core in the promoter regions of ciliary genes (Fig. 3). Moreover, we demonstrated that XAP5 induced the promoter activities of targeted ciliary genes (Fig. 4). Therefore, our results show that XAP5 functions as a transcription factor to regulate flagellar assembly in *Chlamydomonas*. The transcriptional control of more than 100 flagellar genes was XAP5 dependent, whereas the expression of several flagellar genes was XAP5 independent, implying that the regulation of flagellar genes at the transcriptional level is a highly complex process and that XAP5-independent flagellar gene expression can be regulated by other, undiscovered transcriptional mechanisms. In addition, the promoter regions of almost all the XAP5-dependent flagellar genes contained the putative XAP5-binding motif. Thus, it is possible to use the XAP5-binding motif to predict XAP5 targets, particularly those among flagellar genes, which provides a way to identify genes potentially involved in ciliary assembly and function.

## Materials and Methods

The experimental materials and detailed experimental procedures can be found in *SI Materials and Methods*, which includes strains and cell cultures, transformation, screening for flagellar-assembly mutants, complementation of *xap5* and *ift70*, flagellar regeneration, flagellar length and cell size measurements, data availability, alignment and phylogenetic analysis, live-cell imaging and movies, real-time qPCR and RNA-seq analysis, ChIP, EMSA, luciferase activity assay, isolation of nuclei, and anti-IFT70 and anti-XAP5 antibodies.

**ACKNOWLEDGMENTS.** We thank Fangzhi Mou (Wuhan University of Technology) and Zemao Gu (Huazhong Agricultural University) for technical assistance and Kevin Zhang for a critical review of the manuscript and helpful discussions. This work was supported by National Natural Science Foundation of China Grant 31601090 (to Z.H.), NIH Grant R01GM100364 (to W.Z.), and Jiangnan University Internal Research Grant 040 (to Z.H.).

- Goetz SC, Anderson KV (2010) The primary cilium: A signalling centre during vertebrate development. *Nat Rev Genet* 11:331–344.
- Ishikawa H, Marshall WF (2011) Ciliogenesis: Building the cell's antenna. *Nat Rev Mol Cell Biol* 12:222–234.
- Hu Z, Liang Y, He W, Pan J (2015) Cilia disassembly with two distinct phases of regulation. *Cell Rep* 10:1803–1810.
- Reiter JF, Leroux MR (2017) Genes and molecular pathways underpinning ciliopathies. *Nat Rev Mol Cell Biol* 18:533–547.
- Ishikawa H, Thompson J, Yates JR, 3rd, Marshall WF (2012) Proteomic analysis of mammalian primary cilia. *Curr Biol* 22:414–419.
- Stolc V, Samanta MP, Tongprasit W, Marshall WF (2005) Genome-wide transcriptional analysis of flagellar regeneration in *Chlamydomonas reinhardtii* identifies orthologs of ciliary disease genes. *Proc Natl Acad Sci USA* 102:3703–3707.
- Pazour GJ, Agrin N, Leszyk J, Witman GB (2005) Proteomic analysis of a eukaryotic cilium. *J Cell Biol* 170:103–113.
- Davis EE, Katsanis N (2014) Dissecting intraflagellar transport, one molecule at a time. *Dev Cell* 31:263–264.
- Jin H, et al. (2010) The conserved Bardet-Biedl syndrome proteins assemble a coat that traffics membrane proteins to cilia. *Cell* 141:1208–1219.
- Choksi SP, Lauter G, Swoboda P, Roy S (2014) Switching on cilia: Transcriptional networks regulating ciliogenesis. *Development* 141:1427–1441.
- Lefebvre PA, Silflow CD, Wieben ED, Rosenbaum JL (1980) Increased levels of mRNAs for tubulin and other flagellar proteins after amputation or shortening of *Chlamydomonas* flagella. *Cell* 20:469–477.
- Rosenbaum JL, Moulder JE, Ringo DL (1969) Flagellar elongation and shortening in *Chlamydomonas*. The use of cycloheximide and colchicine to study the synthesis and assembly of flagellar proteins. *J Cell Biol* 41:600–619.
- Swoboda P, Adler HT, Thomas JH (2000) The RFX-type transcription factor DAF-19 regulates sensory neuron cilium formation in *C. elegans*. *Mol Cell* 5:411–421.
- Yu X, Ng CP, Habacher H, Roy S (2008) Foxj1 transcription factors are master regulators of the motile ciliogenic program. *Nat Genet* 40:1445–1453.
- Stubbs JL, Oishi I, Izpisua Belmonte JC, Kintner C (2008) The forkhead protein Foxj1 specifies node-like cilia in Xenopus and zebrafish embryos. *Nat Genet* 40:1454–1460.
- Mazzarella R, Pengue G, Yoon J, Jones J, Schlessinger D (1997) Differential expression of XAP5, a candidate disease gene. *Genomics* 45:216–219.
- Zhang A, et al. (2011) Novel retrotransposed imprinted locus identified at human 6p25. *Nucleic Acids Res* 39:5388–5400.
- Martin-Tryon EL, Harmer SL (2008) XAP5 CIRCADIAN TIMEKEEPER coordinates light signals for proper timing of photomorphogenesis and the circadian clock in Arabidopsis. *Plant Cell* 20:1244–1259.
- Fang X, Shi Y, Lu X, Chen Z, Qi Y (2015) CMA33/XCT regulates small RNA production through modulating the transcription of Dicer-like genes in Arabidopsis. *Mol Plant* 8:1227–1236.
- Anver S, et al. (2014) Yeast X-chromosome-associated protein 5 (Xap5) functions with H2A.Z to suppress aberrant transcripts. *EMBO Rep* 15:894–902.
- Dutcher SK, Morrisette NS, Preble AM, Rackley C, Stanga J (2002) Epsilon-tubulin is an essential component of the centriole. *Mol Biol Cell* 13:3859–3869.
- Matsuura K, Lefebvre PA, Kamiya R, Hirono M (2002) Kinesin-II is not essential for mitosis and cell growth in *Chlamydomonas*. *Cell Motil Cytoskeleton* 52:195–201.
- Harper JD, Sanders MA, Salisbury JL (1993) Phosphorylation of nuclear and flagellar basal apparatus proteins during flagellar regeneration in *Chlamydomonas reinhardtii*. *J Cell Biol* 122:877–886.
- Yuan S, et al. (2012) Target-of-rapamycin complex 1 (Torc1) signaling modulates cilia size and function through protein synthesis regulation. *Proc Natl Acad Sci USA* 109:2021–2026.
- Fan ZC, et al. (2010) *Chlamydomonas* IFT70/CrDYF-1 is a core component of IFT particle complex B and is required for flagellar assembly. *Mol Biol Cell* 21:2696–2706.
- Piasecki BP, Burghoorn J, Swoboda P (2010) Regulatory Factor X (RFX)-mediated transcriptional rewiring of ciliary genes in animals. *Proc Natl Acad Sci USA* 107:12969–12974.
- Chu JS, Baillie DL, Chen N (2010) Convergent evolution of RFX transcription factors and ciliary genes predated the origin of metazoans. *BMC Evol Biol* 10:130.
- Vij S, et al. (2012) Evolutionarily ancient association of the FoxJ1 transcription factor with the motile ciliogenic program. *PLoS Genet* 8:e1003019.