

Order Statistics from Beta Inverse Weibull Distribution and Characterizations	العنوان:
المجلة العلمية للدراسات التجارية والبيئية	المصدر:
جامعة قناة السويس - كلية التجارة بالاسماعيلية	الناشر:
Mohamed, Abeer S.	المؤلف الرئيسي:
مج 6, ع 3	المجلد/العدد:
نعم	محكمة:
2015	التاريخ الميلادي:
92 - 79	الصفحات:
932498	رقم MD:
بحوث ومقالات	نوع المحتوى:
English	اللغة:
EcoLink	قواعد المعلومات:
التوزيعات الاحتمالية، توزيع معكوس وايبول، توزيع بيتا معكوس ويبل	مواضيع:
http://search.mandumah.com/Record/932498	رابط:

Order statistics from Beta inverse Weibull distribution and characterizations

Dr/Abeer S.Mohamed

Department of Statistics, Faculty of Commerce (Girls' Branch), Al-Azhar University, Cairo, Egypt

المخلص

يعتبر توزيع معكوس وايبل من اكثر التوزيعات الاحتمالية استخداما في دراسة نظريات الصلاحية والتي تستخدم بكثرة في العلوم التطبيقية والفيزيائية وبداية ظهوره علي يد كيللر وكاماث ١٩٨٢ لاعتبارات فيزيقية خاصه بدراسة اعطال محركات الديزل وكتعميم لهذا التوزيع لاعطائه مرونة اكثر تم اشتقاق توزيع بيتا - معكوس وايبل باستخدام لوغاريتم لتوزيع بيتا علي يد خان ٢٠١٠ وانضم الي التوزيعات التي تستخدم لدراسة اعطال الاجزاء الساكنه في الأجهزة والمحركات والفشل الحيوي. الاحصاءات الترتيبية هي عناصر عينه عشوائية مرتبة من الاصغر الي الاكبر يكثر استخدامها للحصول علي احصاءات وصفية كوسيط ومدى العينة وتعطي مقدرات ذات مأمونية عالية للوسط والوسيط. ولان توزيع بيتا معكوس وايبل من احدث التوزيعات تم استخدامه في الدراسة للحصول علي الاحصاءات الترتيبية له مثل العزوم والمدى والدوال المشتركة وكذلك الاحصاءات الترتيبية للتوزيع عند قطعه بشكل مزدوج (الطرف الايمن والايسر).

Abstract

Inverse Weibull distribution (IW) referred to as the Beta Inverse Weibull distribution(BIW) is generated from the logit of a beta random variable to be considered as a new reliability model. The basic distribution theory of order statistics (ordered random variables) is developed for both finite and asymptotic random samples. Order statistics provide simple robust estimators of location such as the median or trimmed means. The smallest, largest, or middle observations for a given random sample can be computed using order statistic.

In this paper the order statistics from Beta inverse (BIW) distribution is studied. The cumulative distribution and the density for maximum value and minimum are also presented, the exact expression for the single and double moments of order statistics from BIW distribution are derived , finally the order statistic for doubly truncated of the BIW, its r^{th} single and joint r^{th} moments are presented.

Keywords: Beta inverse Weibull, order statistics, single moment of order statistic, double moment of order statistic, doubly truncated.

Research Methodology

- 1. Selection of the research topic:** distribution beta inverse Weibull distribution has been chosen as a modern distribution to study its properties using order statistics has not been studied before.
- 2. The importance of research:** the most continues distributions related to that used in the distributions of life for estimating the reliability and failure in the industrial, medical and engineering field distribution are Weibull and inverse Weibull. beta inverse Weibull distribution has been derived recently to give the prior distributions more flexibility in the presentation of the data to the medical, engineering and industrial study.
- 3. The research plan:** research plan adopted to study the references that were used order statistics with continuous distributions. Then we using the theorems of the order statistic to put the characteristic to the new distribution beta inverse Weibull
- 4. Search Results:** we drive the cumulative probability density function and probability density function to the greatest value and the lowest value of the distribution. the exact expression for the single and double moments of order statistics from BIW distribution are derived , finally the order statistic for doubly truncated of the BIW was presented.

1. Introduction

Order statistics has been used in many real-life applications involving data relating to life testing studies and the BIW distribution was introduced by Khan (2010) as a new reliability model which is generated from the logit of a beta random variable. This model has been introduced as a life time distribution giving more flexibility than the inverse Weibull distribution which was introduced by Keller and Kamath (1982) and was based on the IW distribution. Many authors have studied order statistics and the associated inference for many distributions. Shawky and Bakoban (2009) derived the order statistics from an exponentiated gamma distribution with exact expression for the single and double moments. Raqab and Ahsanullah (2001) obtained the BLUE's of the location and scale parameters of generalized exponential distribution based on order statistics. Balakrishnan and Chan (1998) have discussed linear estimation for log-gamma parameters based on order statistics.

Mahmoud et al. (2003) introduced the exact expression for the single and double moments of order statistics from the inverse Weibull distribution and calculated the variances and covariance's and they used these moments to obtain the BLUE's of the location and scale parameters of inverse Weibull distribution under Type-II censored sample. Mohamed (2013) introduced the Bayesian estimation to obtain the estimators of the scale parameter α for BIW distribution. The Bayes estimators are derived by considering non-informative and informative prior distributions based on LINEX, general

entropy and binary loss functions. Mohamed (2014) presented a comparison between Bayesian estimation and maximum likelihood estimation for BIW.

Aboutahoun and Al-Otaibi (2009) are derive the recurrence relations between the single and the product moments for order statistics from doubly truncated Makeham distribution.

Let Y_1, Y_2, \dots, Y_n be a random sample of size n from BIW distribution with the probability density function (p.d.f) is given by

$$f(y) = \frac{\beta\alpha^{-\beta}}{\beta(a,b)} y^{-(\beta+1)} e^{-a(\alpha y)^{-\beta}} \left[1 - e^{-(\alpha y)^{-\beta}}\right]^{b-1} \quad (1.1)$$

for $y \geq 0, \alpha > 0, \beta > 0, a > 0$ and $b > 0$.

and the cumulative distribution function is given by 1.2.

$$F(y) = \frac{\Gamma(b)}{\beta(a,b)} \sum_{i=0}^{\infty} \frac{(-1)^i}{\Gamma(b-i) i! (a+i)} e^{-(a+i)(\alpha y)^{-\beta}} \quad (1.2)$$

This paper is organized as it follows. Section 2 introduces the order statistic, the minimum, the maximum of density and the cumulative density function of BIW. We derive exact expression for the single moments of order distribution to the BIW in section 3.

2. The Order statistic of BIW

Let $Y_{1:n} \leq Y_{2:n} \leq \dots \leq Y_{n:n}$ denote the order statistics for the sample of size n from the BIW distribution and $Y_{j:n}, j = 1, 2, \dots, n$ is the j^{th} order statistic with probability density function given by David (1981).

$$f_{j:n}(y) = C_{j:n} f(y) [F(y)]^{j-1} [1 - F(y)]^{n-j}, \quad -\infty < y < \infty \quad (2.1)$$

where $f(\cdot)$ and $F(\cdot)$ are given in (1.1) and (1.2), respectively, and

$$C_{j:n} = \frac{n!}{(n-j)!(j-1)!}, \quad (2.2)$$

And the joint p.d.f of $Y_{j:n}$ and $U_{k:n}$ with integer j and k $1 \leq j \leq k \leq n$ is given by David (1981) as follows:

$$f_{j;k:n}(y,u) = C_{j;k:n} (F(u))^{j-1} (F(y)-F(u))^{k-j-1} (1-F(y))^{n-k} f(y) f(u) \quad (2.3)$$

$$\text{where } C_{j;k:n} = \frac{n!}{(j-1)!(k-j-1)!(n-k)!} \quad (2.4)$$

and $f(\cdot)$ and $F(\cdot)$ are given in (1.1) and (1.2), respectively

2.1 The Cumulative distribution and the density for the smallest order statistic

Theorem:

For $Y_{1:n} \leq Y_{2:n} \leq \dots \leq Y_{n:n}$ i id continuous random sample of size n with p.d.f and c.d.f with continuous distribution then the smallest order statistic is given as:

$$\begin{aligned} f_{(1)}(y) &= \frac{d}{dy} (1 - (1 - F(y))^n) = n (1 - F(y))^{n-1} = n (1 - F(y))^{n-1} \frac{d}{dy} F(y) \\ &= n f(y) (1 - F(y))^{n-1} \end{aligned} \quad (2.5)$$

And

$$\begin{aligned} F_{(1)}(y) &= P(Y_{(1)} < y) = 1 - P(Y_{(1)} > y) = 1 - P(Y_1 > y) \dots P(Y_n > y) \\ &= 1 - (1 - F(y))^n \end{aligned} \quad (2.6)$$

From (2.5) and (2.6) and using (1.1) and (1.2) (for BIW distribution), then

$$f_{(1)}(y) = My^{-(\beta+1)} e^{-a(\alpha y)^{-\beta}} \left[1 - e^{-(\alpha y)^{-\beta}} \right]^{b-1} \\ * \left(1 - \frac{\Gamma(b)}{\beta(a,b)} \sum_{i=0}^{\infty} \frac{(-1)^i}{\Gamma(b-i)! (a+i)} e^{-(a+i)(\alpha y)^{-\beta}} \right)^{n-1} \quad (2.7)$$

Where $M = n \frac{\beta \alpha^{-\beta}}{\beta(a,b)}$

and

$$F_{(1)}(y) = 1 - \left(1 - \frac{\Gamma(b)}{\beta(a,b)} \sum_{i=0}^{\infty} \frac{(-1)^i}{\Gamma(b-i)! (a+i)} e^{-(a+i)(\alpha y)^{-\beta}} \right)^n \quad (2.8)$$

2.2 The Cumulative distribution and the density the largest order statistics:

Theorem:

If Y_1, Y_2, \dots, Y_n be a random sample of size n from a population with *continuous* p.d.f $f(y)$, then the p.d.f. and c.d.f of the largest order statistics $Y_{(n)}$ is given as:

$$f_{(n)}(y) = \frac{d}{dy} F(y)^n = n F(y)^{n-1} \frac{d}{dy} F(y) \\ = n f(y) F(y)^{n-1} \quad (2.9)$$

And

$$F_{(n)}(y) = P(Y_{(n)} < y) = 1 - P(Y_{(n)} > y) = P(Y_1 < y \dots P(Y_n < y) \\ = F(y)^n, \quad (2.10)$$

Then., for the BIW in (1.1) and (1.2)

$$f_{(n)}(y) = My^{-(\beta+1)} e^{-a(\alpha y)^{-\beta}} \left[1 - e^{-(\alpha y)^{-\beta}} \right]^{b-1} \\ * \left(\frac{\Gamma(b)}{\beta(a,b)} \sum_{i=0}^{\infty} \frac{(-1)^i}{\Gamma(b-i)! (a+i)} e^{-(a+i)(\alpha y)^{-\beta}} \right)^{n-1} \quad (2.11)$$

and

$$F_{(n)}(y) = \left(\frac{\Gamma(b)}{\beta(a,b)} \sum_{i=0}^{\infty} \frac{(-1)^i}{\Gamma(b-i)j!(a+i)} e^{-(a+i)(\alpha y)^{-\beta}} \right)^n \quad (2.12)$$

3. The moments

The s^{th} moments of the j^{th} order statistics from BIW is given by

$$\mu_{j:n}^{(s)} = E(y_{j:n}^s) = C_{j:n} \int_{-\infty}^{\infty} y^s f(y) (F(y))^{j-1} (1-F(y))^{n-j} dy \quad (3.1)$$

Using the equations (1.1), (1.2) and (3.1) , then the single moment of order statistics from the BIW distribution can be written as follows:

$$\begin{aligned} \mu_{j:n}^{(s)} = E(y_{j:n}^s) &= C_{j:n} A \int_{-\infty}^{\infty} y^{-(\beta+1-s)} e^{-a(\alpha y)^{\beta}} (1-e^{-(\alpha y)^{\beta}})^{b-1} \\ &\left(\sum_{i=0}^{\infty} \frac{(-1)^i}{\Gamma(a-i) i!(a+i)} e^{-(a+i)(\alpha y)^{-\beta}} \right)^{j-1} \left(1 - \frac{\Gamma b}{\beta(a,b)} \sum_{i=0}^{\infty} \frac{(-1)^i}{\Gamma(a-i) i!(a+i)} e^{-(a+i)(\alpha y)^{-\beta}} \right)^{n-j} dy \end{aligned} \quad (3.2)$$

Where

$$A = \left(\frac{\Gamma b}{\beta(a,b)} \right)^i \quad (3.3)$$

The equation (3.2) can be solved numerically

The double (h,s)th moment of $Y_{j:n}$ and $U_{k:n}$, $1 \leq j \leq k \leq n$ is given by

$$\begin{aligned} \mu_{j,k:n}^{(h,s)} = E(y_{j:n}^s u_{k:n}^h) &= C_{j,k:n} \int_{-\infty}^{\infty} \int_u^{\infty} y^s u^h f(y) f(u) (F(y))^{j-1} (1-F(y))^{n-k} \\ &(F(y) - F(u))^{j-k-1} dy du \end{aligned} \quad (3.4)$$

where $f(\cdot)$ and $F(\cdot)$ are given in (1.1) and (1.2), respectively and $C_{j;kn}$ is given in (2.4)

if we would like to derive the c.d.f of the sample range R_n of a random sample of size n form a BIW distribution,

Theorem :

Let $R_n := Y_{n:n} - Y_{1:n}$ be the range of a random sample of size n from a continuous distribution with $(-\infty, \infty)$. Then, for any real number $r > 0$, we have

$$F(R_n) = n \int_{-\infty}^{\infty} (F(y+r) - F(y))^{n-1} f(y) dy \tag{3.5}$$

Using (1.1) and (1.2) for BIW,

$$F(R_n) = \int_{-\infty}^{\infty} \frac{\beta \alpha^{-\beta}}{\beta(a,b)} y^{-(\beta+1)} e^{-a(\alpha y)^\beta} (1 - e^{-(\alpha y)^\beta})^{b-1} \left(I - \frac{\Gamma b}{\beta(a,b)} \sum_{i=0}^{\infty} \frac{(-1)^i}{\Gamma(a-i) i! (a+i)} \right) e^{-(a+i)(\alpha y)^{-\beta}})^{n-1} dy \tag{3.6}$$

where

$$I = F(y+r) = \frac{\Gamma(b)}{\beta(a,b)} \sum_{i=0}^{\infty} \frac{(-1)^i}{\Gamma(b-i) i! (a+i)} e^{-(a+i)(\alpha(y+r))^{-\beta}}$$

3.1 Doubly truncated

We can derive The doubly truncated p.d.f of continuous random variable from the BIW from (1.1) as follows:

$$g(y) = \frac{f(y)}{P-Q} = \frac{\beta \alpha^{-\beta}}{P-O(\beta(a,b))} y^{-(\beta+1)} e^{-a(\alpha y)^\beta} (1 - e^{-(\alpha y)^\beta})^{b-1}, Q_1 \leq y \leq P_1 \tag{3.7}$$

Where

$$P = \frac{\Gamma b}{\beta(a,b)} \sum_{i=0}^{\infty} \frac{(-1)^i}{\Gamma(a-i) i! (a+i)} e^{-(a+i)(\alpha P_1)^{-\beta}} \tag{3.8}$$

$$Q = \frac{\Gamma b}{\beta(a,b)} \sum_{i=0}^{\infty} \frac{(-1)^i}{\Gamma(a-i) \Gamma(a+i)} e^{-(a+i)(\alpha Q_1)^{-\beta}} \tag{3.9}$$

and Q, P ($0 < Q < P < 1$) are respectively the proportions of the truncation on the left and the right of BIW distribution in (1.1),

where the doubly truncated BIW can be used in some situation that depend on the time start $t_0 \neq 0$

The cumulative distribution function c.d.f of the doubly truncated BIW is given by

$$G(x) = Q_2 - \left[\frac{\beta \alpha^{-\beta}}{P - O(\beta(a,b))} y^{-(\beta+1)} e^{-a(\alpha y)^\beta} (1 - e^{-(\alpha y)^\beta})^{b-1} \right]$$

where $Q_2 = \frac{Q}{P - Q}$

$$\tag{3.10}$$

Let $Y_{1:n} \leq Y_{2:n} \leq \dots \leq Y_{n:n}$ is a sample of order statistics of size n from the doubly truncated BIW distribution which given in (3.7) and the c.d.f in (3.10) then we can derive the p.d.f for the j^{th} order statistic as follows:

$$g_{j:n}(y) = C_{j:n} g(y) [G(y)]^{j-1} [1 - G(y)]^{n-j} \quad Q < y < P_1 \tag{3.11}$$

where $g(\cdot)$ and $G(\cdot)$ are given in (3.7) and (3.10), respectively, and

$$C_{j:n} = \frac{n!}{(n-j)!(j-1)!}$$

And the joint p.d.f of $Y_{j:n}$ and $U_{k:n}$ with integer j and k $1 \leq j \leq k \leq n$ is given by David (1981) as follows:

$$g_{j;k:n}(y,u) = C_{j;k:n} (G(u))^{j-1} (G(y)-G(u))^{k-j-1} (1-G(y))^{n-k} g(y) g(u) \quad (3.12)$$

$$C_{j;k:n} = \frac{n!}{(j-1)!(k-j-1)!(n-k)!} \quad (3.13)$$

and $g(\cdot)$ and $G(\cdot)$ are given in (3.7) and (3.10), respectively.

The s^{th} moments of the j^{th} order statistics from BIW (the single moments) is given by

$$\mu_{j:n}^{(s)} = E(y_{j:n}^s) = C_{j:n} \int_{-Q_1}^{P_1} y^s g(y) (G(y))^{j-1} (1-G(y))^{n-j} dy, \quad 1 \leq r \leq n \quad (3.14)$$

The equation (3.14) can be solved numerically

The double $(h,s)^{\text{th}}$ moment of $Y_{j:n}$ and $U_{k:n}$, $1 \leq j \leq k \leq n$ is given by

$$\mu_{j;k:n}^{(h,s)} = E(y_{j:n}^s u_{k:n}^h) = C_{j;k:n} \int_0^\infty \int_y^\infty y^s u^h g(y) g(u) (G(y))^{j-1} (1-G(y))^{n-k} (G(y)-G(u))^{j-k-1} dy du \quad (3.1)$$

3.2 Special Cases

The single and double moments of order statistic to the many of distributions can be derived from (3.2) and (3.4) as a special cases from BIW.

for $a = 1$, $b = 1$, and $\beta = 2$ the distribution in (1.1) reduces to standard inverse Rayleigh distribution (SIR), i.e.,

$$f(y) = \frac{2}{\alpha^2 y^3} e^{-\frac{1}{(\alpha y)^2}}, \quad \text{for } y \geq 0, \alpha > 0. \quad (3.16)$$

$$F(x) = e^{-\frac{\alpha^2}{t^2}} \quad (3.17)$$

If $a = 1$ and $b = 1$, then (1.1) reduces to the two parameter Inverse Weibull distribution (IW), i.e.,

$$f(y) = \beta \alpha^{-\beta} y^{-(\beta+1)} e^{-(\alpha y)^{-\beta}}, \quad \text{for } y \geq 0, \alpha > 0, \beta > 0. \quad (3.1)$$

$$F(x) = e^{-(\alpha y)^{-\beta}} \quad (3.19)$$

4. The Conclusions

A new reliability model (Beta Inverse Weibull distribution) was introduced by Khan 2010 as the logit of a beta random variable. This model is based on BIW and it offers more advantages than the IW. The smallest, largest, or middle observations for a give random sample was computed. The basic distribution theory of order statistics (ordered random variables) was developed for both finite and asymptotic random samples. This research considers order statistics as a simple robust location estimator of location generated from Beta inverse Weibull (BIW) distribution. The cumulative distribution and the minimum and maximum density were also presented , the exact expression for the single and double moments of order statistics from BIW distribution have been derived and the doubly truncated BIW was derived.

References

- Aboutahoun, A. W. and Al-Otaibi, N. M. (2009). Recurrence relations between moments of order statistics from doubly truncated Makeham distribution. *Computational and applied mathematics* Volume 28, N.3, pp. 277–290
- Balakrishnan N. and Chan P. S., (1998). Log-gamma order statistics and linear estimation of parameters, In: Balakrishnan, N. and Rao, C. R. (Eds.), *Handbook of Statistics*, Vol.17. Elsevier Science, Amsterdam, (1998), 61-83.
- Keller, A. Z. & Kamath, A. R. R. (1982). Alternative reliability models for mechanical systems. In *Proceedings of the Third International Conference on Reliability and Maintainability*, pp. 411–415.
- Khan, M. S. (2010). The Beta inverse Weibull. *International Transactions in Mathematical Sciences and Computer*. Volume 3, No. 1, 2010, pp. 113-119
- Mahmoud M. A. Sultan W., K. S. and Amer S. M. (2003) Order statistics from inverse Weibull distribution and associated inference, *Computational Statistics & Data Analysis*, 42, 149-163.
- Mohamed, S. A. (2013). Bayesian Estimation of the Scale Parameter Under Asymmetric Loss Functions to the Beta Inverse Weibull Distribution. *Saleh Kamel center for Islamic economic*. no 49.437-456

Mohamed.S.A (2014).Comparison Study Between the Maximum Likelihood and Bayesian Estimation for the Beta Inverse Weibull Distribution

SalehKamel canter for Islamic economic. no 52381-396

RaqabM. Z. and M. Ahsanullah,(2001) Estimation of the location and scale parameters of generalized exponential distribution based on order statistics, J. Statist. Comput. Simul., 69, 109-123

ShawkyA. I. and Bakoban R. A.(2009). Order Statistics from Exponentiated Gamma Distribution and Associated Inference. Int. J. Contemp. Math. Sciences, Vol. 4, no. 2, 71 - 91