

2017

# Patterns Of Participation And Performance In Youth Baseball Players

Amanda Arnold  
*University of South Carolina*

Follow this and additional works at: <https://scholarcommons.sc.edu/etd>

 Part of the [Exercise Science Commons](#)

---

## Recommended Citation

Arnold, A. (2017). *Patterns Of Participation And Performance In Youth Baseball Players*. (Doctoral dissertation). Retrieved from <https://scholarcommons.sc.edu/etd/4451>

This Open Access Dissertation is brought to you by Scholar Commons. It has been accepted for inclusion in Theses and Dissertations by an authorized administrator of Scholar Commons. For more information, please contact [dillarda@mailbox.sc.edu](mailto:dillarda@mailbox.sc.edu).

PATTERNS OF PARTICIPATION AND PERFORMANCE IN YOUTH BASEBALL  
PLAYERS

by

Amanda Arnold

Bachelor of Science  
Texas Woman's University, 2006

Doctor of Physical Therapy  
Texas Woman's University, 2010

---

Submitted in Partial Fulfillment of the Requirements

For the Degree of Doctor of Philosophy in

Exercise Science

The Norman J. Arnold School of Public Health

University of South Carolina

2017

Accepted by:

Paul F. Beattie, Major Professor

Ellen Shanley, Committee Member

Charles A. Thigpen, Committee Member

Stacy L. Fritz, Committee Member

John M. Brooks, Committee Member

Cheryl L. Addy, Vice Provost and Dean of the Graduate School

© Copyright by Amanda Arnold, 2017  
All Rights Reserved.

## DEDICATION

For Jeff.

*“Baseball was, is and always will be to me the best game in the world.”*

- Babe Ruth

## ACKNOWLEDGEMENTS

There is an old proverb that states ‘It takes a village to raise a child.’ While this statement is undoubtedly true, I feel that the concept also applies to learning and teaching with students. Throughout my academic and professional careers, I have been fortunate to be consistently surrounded by supportive, intelligent and giving people who continue to mentor me along my path. This dissertation is an homage to them.

To my family and friends: Thank you. Thank you for your love, support and belief in me and for constantly reminding me of who I am especially on the days I was unable to see it.

To my wife: We did it! I am fairly certain you had no idea what you were signing up for when we left Texas and began this adventure but we made it! Thank you for your love, patience and positive outlook on life and for being the best research assistant a PhD student could ask for. Here’s to starting the next chapter in our lives together!

To my dissertation committee: Thank you to Dr. Ellen Shanley and Dr. Chuck Thigpen for challenging me to think critically at every level both as a researcher and as a clinician. Your incalculable time, efforts and dedication to me, our profession and our communities are greatly appreciated and have set the standard for excellence in clinical research and education. Thank you to Dr. Paul Beattie, Dr. Stacy Fritz and Dr. John Brooks for challenging me to question what I know and to think what could be? Your mentorship, perspectives and support throughout this process have positively influenced my life on both a personal and professional level.

To Dr. Richard Hawkins, Dr. Michael Kissenberth and Dr. JT Tokish: Thank you. Thank you for your curiosity and never ending desire to learn and teach with those around you. Each of you lead by example and for that I am grateful. Your intelligence, compassion and approachability have profoundly impacted my life as well as the lives of your patients and colleagues.

To the baseball community in the Upstate of South Carolina: Thank you. This research is for you and would not be possible without you. Your continued participation and support allow researchers like me to pursue the ultimate goal of keeping kids healthy and active in sport.

## ABSTRACT

Baseball is a popular sport to play in the United States, with approximately 13-17 million athletes participating across all levels of competition. Youth (9-12 years) and adolescent (13-18 years) players comprise the majority of this population playing at the club and high school levels, yet less than 10% of research studies include athletes <18 years old. Despite increased awareness of the risks surrounding sports participation, youth and adolescent baseball players continue to report overuse injuries at alarming rates.

The lack of high-quality research describing athletic performance and injury risk factors, such as sport specialization, in young athlete populations poses a significant knowledge gap in the literature. The current investigation sought to establish the incidence of upper extremity (UE) injuries while examining population-specific risk factors in a cohort of youth baseball players (**Aim 1**). The current study also examined the measurement properties of normalized isometric shoulder strength, by 5 separate methods, for use as a multi-faceted clinical assessment tool that was responsive to changes in physical growth and development over time (**Aim 2**).

Youth baseball players were examined for baseline participation and isometric shoulder strength data and then prospectively followed via coach and parent surveys. Athletic exposures (AE) and the presence of UE injuries were tracked for each player. Chi square analyses were used to compare the frequency of UE injuries based on position group, sports specialization status and participation in additional specialty training. Odds

ratios as well as absolute and absolute risk differences with 95% confidence intervals (CI) were calculated between groups for **Aim 1**. A subset of athletes (n = 58) was physically re-examined during the follow-up period to establish the test-retest reliability of each of the normalized isometric shoulder strength measures. Repeated measures analyses of variance (ANOVA) were conducted to compare changes in isometric shoulder strength at 2 time points after normalizing to 5 separate measures of body size. Linear regression models were used to examine the relationships between normalized isometric shoulder torque measures and ball velocity in youth baseball players for **Aim 2**.

Results showed that youth baseball players demonstrated an UE injury incidence rate of 16.3/1000 AEs. Specialized athletes, who comprised 83.0% of this cohort, demonstrated a 15.9% increase in absolute risk for developing an UE injury when compared to multi-sport counterparts. Following comparisons across 5 methods of normalization, only torque, defined as the measure of shoulder strength divided by the corresponding ulnar length, demonstrated excellent reliability and detected significant changes between shoulder strength in each of the 4 measures tested. Torque was the most stable and reliable normalization method evaluated in this study. Modest but significant correlations were observed between shoulder scaption torque, shoulder external rotation (ER) torque at 0° and ball velocity suggesting that these measures were the most useful predictors of throwing performance in 9-12 year old baseball players.



## TABLE OF CONTENTS

DEDICATION .....	iii
ACKNOWLEDGEMENTS.....	iv
ABSTRACT .....	vi
LIST OF TABLES .....	xi
LIST OF FIGURES .....	xii
LIST OF ABBREVIATIONS.....	xiii
CHAPTER 1: INTRODUCTION.....	1
1.1 UPPER EXTREMITY INJURIES IN YOUTH BASEBALL PLAYERS.....	1
1.2 NORMALIZATION METHODS FOR ISOMETRIC SHOULDER STRENGTH IN YOUTH BASEBALL PLAYERS.....	3
CHAPTER 2: REVIEW OF THE LITERATURE .....	6
2.1 ABSTRACT .....	6
2.2 INTRODUCTION.....	7
2.3 METHODS .....	8
2.4 RESULTS.....	9
2.5 DISCUSSION.....	14
2.6 CONCLUSION .....	17
2.7 CLINICAL RECOMMENDATIONS .....	18
CHAPTER 3: METHODS.....	28
3.1 RESEARCH DESIGN .....	28

3.2 STUDY SETTING.....	28
3.3 STUDY SUBJECTS.....	28
3.4 PROCEDURES .....	29
3.5 STATISTICAL ANALYSES.....	34
CHAPTER 4: SPORT SPECIALIZATION INCREASES UPPER EXTREMITY INJURY RISK IN YOUTH BASEBALL PLAYERS: A PROSPECTIVE COHORT STUDY .....	42
4.1 ABSTRACT .....	42
4.2 INTRODUCTION.....	44
4.3 METHODS .....	45
4.4 RESULTS.....	48
4.5 DISCUSSION .....	50
4.6 CONCLUSION .....	56
CHAPTER 5: NORMALIZATION METHODS IN ISOMETRIC SHOULDER STRENGTH IN YOUTH BASEBALL PLAYERS: A COMPARISON ACROSS 5 METHODS .....	59
5.1 ABSTRACT .....	59
5.2 INTRODUCTION.....	61
5.3 METHODS .....	63
5.4 RESULTS.....	66
5.5 DISCUSSION .....	68
5.6 CONCLUSION .....	72
CHAPTER 6: DISCUSSION.....	79
6.1 UPPER EXTREMITY INJURIES IN YOUTH BASEBALL PLAYERS.....	80
6.2 NORMALIZATION METHODS FOR ISOMETRIC SHOULDER STRENGTH IN YOUTH BASEBALL PLAYERS.....	82
6.3 LIMITATIONS .....	84

6.4 CLINICAL IMPLICATIONS .....	86
6.5 FUTURE STUDIES .....	87
REFERENCES .....	89
APPENDIX A – SAGE PUBLISHING PERMISSION TO REPRINT POLICY .....	105
APPENDIX B – REDCAP ONLINE SURVEY SAMPLE: YOUTH BASEBALL QUESTIONNAIRE..	106
APPENDIX C – DATA COLLECTION SHEET .....	108

## LIST OF TABLES

Table 2.1 Studies that Report Lower Extremity Physcal Injuries.....	25
Table 2.2 Studies that Report Upper Extremity Physcal Injuries .....	27
Table 3.1 Data Collection Measures.....	38
Table 3.2 Normalization Methods for Isometric Shoulder Strength.....	40
Table 3.3 Power Analyses Based on Expected Injury Rates .....	41
Table 3.4 Power Analyses Based on Risk Factor Comparisons .....	41
Table 4.1 Analysis of Risk Factors for Upper Extremity Injuries .....	58
Table 5.1 Baseline Characteristics of Youth Baseball Players .....	72
Table 5.2 Overview of Anthropometric & Isometric Shoulder Strength Measures .....	73
Table 5.3 Normalization Methods for Isometric Shoulder Strength Measures .....	74
Table 5.4 Reliability of Normalized Dominant Isometric Shoulder Strength Measures using Torque Method.....	75
Table 6.1 Injury Frequency Comparisons Across Multiple Youth Baseball Epidemiologic Studies.....	88

## LIST OF FIGURES

Figure 2.1 Preferred Reporting Items for Systematic Review & Meta-Analysis (PRISMA) Flow Diagram .....	20
Figure 2.2 Strength-of-Recommendation Taxonomy (SORT) Diagram .....	21
Figure 2.3 Radiograph of 13-year-old Male Football Player with Osgood-Schlatter Disease .....	22
Figure 2.4 Radiograph of 14-year-old Male Baseball Player with Little League Shoulder .....	23
Figure 2.5 Radiograph of 13-year-old Male Baseball Player with Little League Elbow ..	24
Figure 3.1 Prospective Cohort Study Design Flowchart .....	37
Figure 4.1 Sport Specialization – Perception versus Reality .....	57
Figure 5.1 Examination of Isometric Shoulder Scaption & Shoulder External Rotation at 0° Strength in Seated Position .....	76
Figure 5.2 Examination of Isometric Shoulder External Rotation at 90° and Shoulder Internal Rotation at 90° Strength in Prone Position.....	76
Figure 5.3 Changes in Normalized Dominant Shoulder Strength by Method over 6-month Period .....	77
Figure 5.4 Relationship Between Normalized Dominant Shoulder Scaption Torque, External Rotation at 0° Torque and Ball Velocity .....	78

## LIST OF ABBREVIATIONS

AE	Athletic Exposure
ANOVA	Analysis of Variance
AR	Absolute Risk
ARD	Absolute Risk Difference
BMI	Body Mass Index
CI	Confidence Interval
Cm	Centimeter
D	Dominant
ER	External Rotation
HHD	Hand-Held Dynamometry
HT	Humeral Torsion
ICC	Intraclass Correlation Coefficient
IR	Internal Rotation
IRB	Institutional Review Board
Kg	Kilogram
LE	Lower Extremity
MDC	Minimally Detectable Change
MHz	Megahertz
MPH	Miles Per Hour
ND	Non-Dominant
Nm	Newton Meter

NSAID ..... Non-Steroidal Anti-Inflammatory Drug  
OCEBM ..... Oxford Centre for Evidence-based Medicine  
OR ..... Odds Ratio  
PRISMA ..... Preferred Reporting Items for Systematic Review & Meta-Analysis  
PT ..... Physical Therapy  
ROM ..... Range of Motion  
RR ..... Rate Ratio  
SD ..... Standard Deviation  
SEM ..... Standard Error of Measure  
SORT ..... Strength-of-Recommendation Taxonomy  
UE ..... Upper Extremity  
Wks ..... Weeks

# CHAPTER 1

## INTRODUCTION

### 1.1 Upper Extremity Injuries in Youth Baseball Players

Baseball is a popular sport to play in the United States, with approximately 13-17 million athletes participating across all levels of competition.<sup>31,51,143</sup> Youth (9-12 years) and adolescent (13-18 years) players comprise the majority of this population playing nearly year-round with minimal rest at the club and high school levels.<sup>4,31,51,143</sup> Despite increased awareness surrounding the risks associated with overtraining, youth and adolescent baseball players continue to report overuse injuries at alarming rates.<sup>12,19,90,91,136,140,148,154,155</sup> The incidence of baseball-related overuse injuries in adolescent players was reported to be 1.3 – 4.0 injuries per 1,000 athletic exposures however this data is unknown in youth players.<sup>31,137</sup> The research does indicate that the majority of baseball-related overuse injuries are reported in the upper extremity (UE), specifically at the shoulder and elbow, however little is known about the etiology and development of these injuries in youth athletes.<sup>6,31,137</sup>

Despite evidence suggesting that sport specialization may be related to the development of overuse injuries in youth and adolescent athletes, its prevalence continues to increase in the U.S.<sup>72,107</sup> Research studies have previously defined sport specialization using a battery of criteria including year-round training in a single sport (>8 months/year), identification of a primary sport over additional sports and the cessation of additional sports to focus on a primary sport.<sup>8,71,72,84</sup> Established definitions exist in the



literature, however less is understood about the public perception of sports specialization with parents and coaches.<sup>71,72,84,107,108</sup> The effects of sport specialization on UE injury risk has not been previously examined in the baseball literature. Other risk factors, such as excessive pitch counts, varied pitch types and faulty throwing mechanics, have been linked to the development of shoulder and elbow pain in youth throwers using self-report and survey data.<sup>12,71,72,90,91,116,162</sup> The USA Baseball Medical & Safety Advisory Committee has used this research to establish age-appropriate guidelines for pitch counts, pitch types and throwing mechanics.<sup>12,19,90,91,136,140,148,154,155</sup> The effectiveness of these recommendations on the reduction of baseball-related overuse injuries is unknown, as the extent of the problem has not been previously established in the literature. The absence of epidemiologic studies describing overuse injury rates in youth athletes combined with the lack of population-specific risk factor data pose significant knowledge gaps in the evaluation and treatment of this population. To address these gaps in the literature, the first aim of the current project was:

**Aim 1:** Examine the effects of population-specific risk factors on UE injury risk in a cohort of 9-12 year old male baseball players.

1. The primary objective of this study was to determine the specific incidence of baseball-related UE injuries using athletic exposures (number of team-recorded practices and games) as the denominator.
2. The secondary objective of this study was to examine the effects of player position, sport specialization and participation in specialty training on baseball-related UE injury risk.

## 1.2 Normalization Methods for Isometric Shoulder Strength in Youth Baseball

### Players

Upper extremity muscle strength is an important component in the assessment of throwing performance and injury prevention in baseball players.<sup>17,28,44,50,52,61,97,105,148,156,160</sup> Strength is defined as the amount of force a muscle can maximally produce during a single repetition.<sup>67,68,70</sup> Clinicians and researchers routinely use batteries of strength measures in performance assessments, injury diagnostics and return to sport decisions following injury.<sup>17,33,61,148</sup>

A variety of methods, including isokinetic, isometric and functional testing, have been used to assess upper extremity strength in athletic populations.<sup>17,28,44,61,97,148,160</sup> While isokinetic testing is considered the gold standard in strength assessment; the high equipment costs and lack of portability make it impractical for use outside of laboratory settings.<sup>33,52,142</sup> Isometric testing using hand held dynamometry (HHD) has proven to be a reliable, low cost and portable alternative to isokinetics in assessing strength, particularly in the throwing shoulder.<sup>17,33,142,148</sup>

While the majority of upper extremity strength testing has been conducted at the collegiate and professional levels, few studies have sought to assess strength measures in younger athletes.<sup>17,44,61,97,105,148</sup> One potential reason for this gap in the literature may be related to the inherent variability of strength measures, especially when assessed in physically developing populations.<sup>67,68,70</sup> Studies have shown that anthropometric measures, such as height and weight, influence the body's ability to produce force, suggesting that changes during growth and development may impact a youth athlete's muscle strength and performance measures.<sup>44,67,68,97,105</sup> Relying solely on the measure to quantify changes over time, without accounting for alterations in body size, may not

adequately reflect how athletic performance and injury risk develop in youth populations.<sup>67,68,70</sup> Normalization is one option for assessing strength changes in physically developing populations however these methods have been inconsistently reported in the literature.<sup>68,70</sup>

Evaluating isometric shoulder strength in youth athletes is inherently different from collegiate and professional athletes.<sup>67,68,92</sup> Height, weight and neuromuscular control can fluctuate frequently in physically developing populations with the potential to rapidly change over short periods of time. Performance assessments that rely on absolute measures, without anthropometric normalization, may lack the ability to discern changes in muscle strength from changes in body size in youth populations.<sup>67,68,70</sup> Prior research studies suggest that normalization methods, which include body mass, body mass index (BMI), height, torque and percent of non-dominant shoulder strength described by Trakis<sup>148</sup> may be potential ways to assess muscle strength and changes in muscle strength over time in this population.<sup>63,67,68</sup> Accounting for these alterations in growth and development through normalization is critical to accurately assessing muscle function, throwing performance and injury risk in youth athletes.<sup>67,68,92</sup> Establishing an objective and reliable method for evaluating strength is an important step in understanding shoulder function in youth baseball players.<sup>33,148</sup> Once a reliable method has been established, the next steps are to examine the relationships between shoulder strength and ball velocity, a performance variable of interest in this population, and shoulder strength and UE injury risk in youth players. To address the lack of population-specific strength measures with related data in the youth athlete literature, the second aim of the current project was:

**Aim 2:** Compare the measurement properties of normalized isometric shoulder strength, using 5 separate methods, for use as a multi-faceted clinical assessment tool that was responsive to changes in physical growth and development over time in a cohort of 9-12 year old male baseball players.

1. The primary objective of this aim was to assess the test-retest reliability of 4 isometric shoulder strength measures.
2. The secondary objective of this aim was to assess changes in normalized isometric shoulder strength over time using baseline and follow-up evaluation measures.
3. The tertiary objective of this aim was to examine the relationship between normalized isometric shoulder strength measures and ball velocity, a performance variable of interest in youth baseball players.

## CHAPTER 2

### REVIEW OF THE LITERATURE

#### 2.1 Abstract

Introduction: Despite rising awareness of the risks associated with sports participation, overuse injuries continue to increase in youth athlete populations. Physeal injuries are one type of overuse injury exclusive to pediatric populations that are often sustained during athletic practice or competition. Overuse physeal injuries are, in theory, preventable, however little consensus has been reached surrounding the risk factors, prevention and treatment strategies reported in the literature.

Objective: The purpose of this systematic review was to summarize the best available evidence concerning overuse physeal injuries in youth and adolescent athletes. The information can then be used to guide the development of prevention and treatment programs specific to this population as well as identify any knowledge gaps for future research.

Methods: PubMed and Academic Search Complete (EBSCOhost) were explored using *physeal injuries* from January 1950 through May 2015. Original research studies performed in athletic populations with mechanisms of injury related to sport were chosen. A total of 24 studies were included in this systematic review.<sup>1</sup>

---

<sup>1</sup>Arnold A, Thigpen CA, Beattie PF, Kissenberth MJ, Shanley E. Overuse Physeal Injuries in Youth Athletes. *Sports Health*. 2017;9(2):139-147. Reprinted here with permission of publisher.

Results: Risk factors for injury include periods of accelerated growth, chronological age, body size, training volume and history of previous injury. Injury prevention strategies currently emphasize participation limitations and sport-specific training programs in skeletally immature athletes. The most effective treatment following an overuse physal injury was an extended period of active rest and joint immobilization when necessary.

*Overall Strength-of-Recommendations Taxonomy (SORT): C.*

Conclusion: Overuse physal injuries are multi-factorial in nature. Muscular imbalances following accelerated growth periods are thought to predispose young athletes to overuse injuries. Modifiable risk factors such as flexibility, strength and training volume should be regularly monitored in an effort to prevent these injuries when possible.

Keywords: physis; physal injury; overuse; sports injuries; pediatric injuries

## **2.2 Introduction**

An estimated 30 million children in the U.S. are involved annually in organized sport.<sup>1,20</sup> Despite rising awareness of the risks associated with sports participation, overuse injuries continue to increase in youth athlete populations.<sup>1,19-22,24,59</sup> Physal injuries are one type of overuse injury exclusive to pediatric populations that are most often sustained during athletic practice or competition.<sup>14,15,18-20,22-24,26,34,41,48,55,56,76,81,87,93,102,119,141,144,145,152</sup> While specific mechanisms of injury are heterogeneous and differ by sport, the physis, as the weakest part of the bone, is a site highly prone to injury in youth athletes.<sup>18,19,21,38</sup>

Overuse physal injuries develop in response to excess stress placed on immature bony and soft tissue structures.<sup>19-24,34,35,41,42,45,47,48,56,59,74,77,79,87,102,106,114,131</sup> Rapid physical

changes combined with repetitive sport-related tasks such as running and overhead throwing are frequently associated with the development of physeal injuries in youth athletes.<sup>19,21,26</sup> The gradual nature of this injury progression provides clinicians with multiple opportunities for effective intervention. Overuse physeal injuries are, in theory, preventable. Prevention and treatment strategies should be population-specific, taking into account previously established risk factors and clinical impairments observed in youth athletes.<sup>21</sup> The purpose of this work was to review and aggregate the best available evidence concerning recommended prevention and treatment strategies for overuse physeal injuries for application to clinical practice.

## **2.3 Methods**

### **2.3.a Literature Review Methods and Article Identification**

An electronic literature search was performed accessing papers published from January 1950 to May 2015 in the PubMed and all EBSCOhost databases. Search terms included *epiphyseal injury*, *epiphyseal plate injury*, *pediatric sports injury* and *physeal sports injury*. Additional searches in the aforementioned databases were performed using the terms *little league shoulder*, *gymnast wrist*, *little league elbow*, *lower extremity physeal injury*, *osgood schlatter disease*, *sever's disease* and *sinding-larsen-johansson disease* as they were the most commonly reported mechanisms of injury during the primary search. Only English language articles published in peer-reviewed journals with an emphasis on human participants were initially included. Articles were also required to meet *Level IV* standards or higher based on criteria developed by the Oxford Centre for Evidence-based Medicine (OCEBM). Abstracts and non-published works were not included. Based on these search criteria, 3,663 articles were located. Using the Preferred

Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines, studies were selected based on appropriateness of topic and full text options.<sup>104</sup> All clinical commentaries and review articles were omitted. A total of 24 original research studies were included in this systematic review (Figure 2.1).

### 2.3.b Eligibility Criteria

Article selection was based on repetitive stress as a mechanism of injury in young athletes. Case reports, case series and cohort studies that described non-sport related mechanisms of injury, such as falls, were not included in this review. Acute sport-related injuries were also excluded. The scope of this systematic review was limited to overuse physéal injuries sustained during athletic competition.

## 2.4 Results

Twenty-four studies were included in this systematic review (Tables 2.1 and 2.2).<sup>2,5,9,10,13,27,34,45,47,59,60,66,79,82,83,88,103,114,115,117,130,131,147</sup> Thirty-three percent of studies included descriptions of known physéal injury risk factors while only 8% percent of studies used those factors to outline effective prevention strategies.<sup>10,27,34,45,79,117,130,147</sup> Eighty-eight percent of studies included data describing treatment strategies following an overuse physéal injury.<sup>2,9,10,13,27,34,45,47,59,66,82,83,88,103,114,115,117,131,147,150</sup> Review of current evidence suggests that more emphasis has been placed on the treatment of overuse physéal injuries and that further research is needed to establish effective prevention strategies for these diagnoses.

Risk factors common to both lower extremity (LE) and upper extremity (UE) physéal injuries include age, physical characteristics, growth patterns and training volume.<sup>19,21,26</sup> While limited evidence was available describing effective prevention



strategies in this population, studies did emphasize that youth athletes should engage in minimum periods of active rest following their competition cycles.<sup>19,38</sup> Adequate physical training and variation in sport-specific tasks were also encouraged.<sup>19,39</sup> Treatment strategies following an overuse physal injury included varying periods of active rest and when necessary, immobilization of the affected joint.<sup>2,5,9,10,27,47,59,66,82,83,103,115,150</sup> Gradual return to physical training and conditioning tasks was recommended prior to full return to sport.<sup>5,9,79,82</sup>

#### 2.4.a Lower Extremity Injuries

Overuse physal injuries in the LE typically occur when excess stress is placed across areas with major tendons insertions.<sup>11,19</sup> Osgood-Schlatter Disease, Sever's Disease and Sinding-Larsen-Johansson Syndrome are 3 of the most common overuse physal injuries sustained during childhood.<sup>15,93,96</sup> The first two syndromes account for a staggering 18% of all pediatric overuse injuries reported in the literature.<sup>93</sup> Osgood-Schlatter Disease is described as chronic apophysitis of the patellar tendon where it inserts on the tibial tuberosity apophysis. It is typically observed in girls ages 8-13 years and boys ages 10-15 years (Figure 2.3).<sup>34</sup> The same inflammatory process occurs with Sever's Disease but at the Achilles tendon insertion into the vertical calcaneal apophysis.<sup>117</sup> This condition appears to present more often in young boys between the ages of 8 and 12 years old.<sup>74</sup> Sinding-Larsen-Johansson Syndrome has a similar etiology but develops at the junction of the inferior pole of the patella and the proximal portion of the patellar tendon.<sup>150</sup> While this syndrome appears less frequently in the literature than the previous two, Sinding-Larsen-Johansson Syndrome does occur in youth athletes

between the ages 10-15 years, limiting their function and participation levels (Table 2.1).<sup>150</sup>

Prevention strategies in the literature emphasize the correction of modifiable risk factors such as deficits in trunk and LE flexibility, which is often attributed to rapid changes in physical growth common during childhood and adolescence.<sup>19,21,26,39,150</sup> Programs designed to enhance cardiovascular endurance and correct physical training errors are also recommended to prevent these types of injuries.<sup>34,38,79</sup> Following an overuse physal injury in the LE, 50% of studies recommend a 3-5 month period of active rest with complete cessation of sport-specific activities.<sup>9,59,66,82,103,117,150</sup> Twenty-one percent of studies suggest activity modifications may be appropriate based on the symptom presentation of the athlete, thereby limiting their total time away from sport.<sup>9,82,117</sup> Lower extremity stretching and conditioning programs were also used in 21% of the studies as either a stand alone treatment or in conjunction with additional strategies.<sup>34,79,117</sup> Several studies reported joint immobilization and surgical intervention for long standing physal injuries related to overuse, however these strategies were only employed in severe cases.<sup>59,66,83,103,114,117</sup> Irrespective of the treatment strategy used, an athlete should not fully return to sport until symptom resolution has occurred. No studies to date have examined or compared the effectiveness of these treatments in youth athlete populations.<sup>134</sup>

#### 2.4.b Upper Extremity Injuries

Overuse physal injuries in the UE occur due to excess compression or traction forces placed across a joint during sport.<sup>19,41</sup> Gymnast Wrist, Little League Shoulder and Little League Elbow are 3 UE physal injuries that are highly prevalent and described

frequently in the pediatric sports literature.<sup>2,5,10,13,27,47,60,79,115,117,131,147</sup> Seventy nine percent of youth gymnasts report wrist pain during practice or competition while 32% of youth baseball pitchers report arm pain while throwing.<sup>41,90,91</sup>

Gymnast wrist occurs in response to the premature closure of the distal radial physis following excessive compression loads during UE weight bearing.<sup>24,88</sup> Gymnastics is one of the few sports that repeatedly performs closed chain weight bearing activities on both their upper and lower extremities.<sup>3,16,20,22,23,35,41,42</sup> This injury is typically seen in athletes between the ages of 10 and 14 years old (Table 2.2).<sup>41</sup> Little League Shoulder has been described in the literature as a widening of the proximal humeral epiphysis or epiphysiolysis (Figure 2.4). It is most often seen in the dominant shoulder and is thought to occur secondary to the repetitive rotational and traction stresses associated with overhead throwing.<sup>2,5,10,14,27,47,115,147</sup> Little League Elbow is a term often used to describe a variety of physeal and cartilaginous injuries at the pediatric elbow.<sup>18,19,21,55,60,77</sup> By definition, Little League Elbow is a repetitive traction injury to the medial epicondylar apophysis (Figure 2.5).<sup>55</sup> Diagnoses of Little League Shoulder and Little League Elbow are most often made following reports of persistent arm pain and loss of function in youth baseball pitchers between the ages of 11 and 15 years old (Table 2.2).<sup>27,77,115</sup>

Risk factors associated with the development of Gymnast Wrist include consistent UE loading and timing of growth spurts.<sup>40</sup> Studies suggest that participation in repetitive UE weight bearing tasks, especially during periods of rapid physical growth, is directly associated with this highly prevalent, population-specific injury.<sup>19,37,40-42</sup> Risk factors related to the development of Little League Shoulder and Little League Elbow are similar. Excessive game, season and yearly pitch counts and pitching while fatigued are

factors that have been associated with shoulder and elbow dysfunction in youth baseball players.<sup>90,91</sup> Pitch type and selection are also important for the health of this population. Youth baseball players who reported throwing breaking pitches such as curveballs or sliders over the course of the season were more at risk to develop shoulder and elbow pain when compared to those who did not.<sup>90</sup> Anthropometric measures such as increased height and weight also impacted injury risk but were more significant to the development of elbow pathology than shoulder pathology.<sup>91</sup>

Despite the lack of epidemiological data concerning Gymnast Wrist, multiple prevention strategies have been suggested in the literature.<sup>37,38,40</sup> The gradual progression and variation of training loads is imperative to limit the volume of compressive forces sustained through the distal radial physis.<sup>37,39,40</sup> Studies suggest that coaches and parents should be cognizant of rapid changes in growth, as the athlete is most at risk for overuse physal injuries during this period.<sup>37,39</sup> In an effort to prevent Little League Shoulder and Little League Elbow, USA Baseball implemented yearly, seasonal and game pitch count limitations based on an athlete's age at the time of competition.<sup>90,91,116</sup> These recommendations were designed to decrease an athlete's risk for injury by limiting excessive stress and fatigue during sports participation.<sup>89-91</sup>

Treatment strategies for all three overuse physal injuries center around an extended period of active rest. Following an injury, 50% of studies recommend active rest from sport-specific training to ensure adequate healing and symptom resolution.<sup>2,5,27,47,115,117</sup> Recommended periods of active rest range from 4-6 weeks for a diagnosis of Gymnast Wrist or Little League Elbow to 3-5 months for athletes with Little League Shoulder.<sup>2,27,40,115,140</sup> In severe cases of Little League Elbow, joint immobilization

and/or surgical intervention have been employed to ensure optimal functional outcomes.<sup>78,119,125</sup> However, an extended period of active rest remains the main treatment of choice for overuse physeal injuries in the UE.

## 2.5 Discussion

The main purposes of this systematic review were to identify known risk factors associated with overuse physeal injuries and to determine which prevention and treatment strategies were most effective and supported by the evidence. Physeal injuries represent approximately 15% of all pediatric sports injuries currently reported in the literature.<sup>19</sup>

The physis, as the weakest physiologic structure in a young athlete, is particularly susceptible to overuse injuries.<sup>14,15,18-26,34,40,48,55,56,76,81,87,93,102,119,141,144,145,152</sup> As participation in youth sports continues to increase, clinicians should become cognizant of the risk factors, prevention strategies and treatment options associated with overuse physeal injuries.<sup>14,15,19,21,22,55,144,145</sup>

### 2.5.a Risk Factors

Risk factors associated with participation in youth sports have been reported throughout the literature, however no research studies have examined injury risk with respect to physeal injuries.<sup>21</sup> Physeal injuries are exclusive to skeletally immature individuals suggesting that modifiable and non-modifiable risk factors are specific to this population.<sup>21,26,34,35,56,79,100,134</sup>

Non-modifiable risk factors for overuse injuries can include timing of accelerated growth spurts, chronological age, body size and history of previous injury.<sup>21,38,40</sup> Previous injury is the strongest predictor for the development of future injuries supported by the

literature.<sup>21,26,126,149</sup> Studies show that athletes who reported a prior injury were at a much higher risk to sustain an injury when compared to a previously healthy cohort.<sup>149</sup>

Modifiable risk factors such as flexibility, strength, training volume and coaching styles also impact overall injury risk in youth and adolescent athletes. Multiple studies suggest that excessive training loads often lead to physical fatigue in youth athletes. Continued participation in sport once fatigued can damage an athlete's physical development thereby illustrating the importance of responsible coaching, especially during the early years of sport.<sup>21,38,40,76</sup>

#### 2.5.b Prevention and Treatment Strategies

Injury prevention strategies for youth and adolescent athletes focus on limiting time spent participating in sport as well as encouraging 2-3 months of scheduled rest away from training and competition.<sup>38,121</sup> This is designed to mediate the effects of repetitive risk-prone activities on physically maturing bodies. Pitch count regulations, which are enforced by the governing bodies in youth baseball, is one notable attempt at preventing upper extremity overuse injuries at the policy level.<sup>90,91,135,139</sup> Multiple studies have also recommended that clinicians monitor known risk factors such as anthropometric (i.e. height and weight) and physical characteristics (i.e. range of motion and strength) as youth athletes mature over time.<sup>34,37,38,136,148</sup> Multiple programs designed to improve flexibility, strength and balance deficits have been shown to have protective effects against injuries in this population.<sup>21,25,94,95,120,138</sup>

The most widely accepted treatment strategy following any physical injury is an extended period of active rest.<sup>3,10,19,27,30,74,77,114,115,134</sup> Recommended durations of active rest vary from 4-6 weeks to 3-5 months depending on by diagnosis, sport and severity of

symptoms.<sup>26,27,77</sup> During this time, strategies can include changes in field position to limit throwing in the cases of Little League Shoulder and Little League Elbow or the recommendation of no running for a specified period of time in athletes with Osgood-Schlatter, Sinding-Larsen-Johansson's or Sever's Disease. In most cases, non-symptomatic activities such as hitting a baseball or footwork drills in soccer can be continued. This allows young athletes to continue training without prolonging their recovery by re-aggravating the affected joint.

During a period of active rest, conservative measures such as physical therapy can prove beneficial. Once the pain has subsided, clinicians can begin to restore the necessary flexibility, strength and neuromuscular control required to participate safely in sport.<sup>2,3,5,13,20,22-25,27,29,35,40,42,47,78,79,115,132,140</sup> Progressive strength training programs, lasting approximately 6-8 weeks, can be augmented with return to throwing or running programs when appropriate.<sup>77</sup> The rehabilitation programs reported in the literature were vague and lacked return to sport criteria. Future research should focus on the development of age- and injury-specific return to sport progressions designed to provide clinicians with evidence-based guidelines to return their athletes safely back to sport.

#### 2.5.c Limitations

The main limitation of this systematic review was the lack of experimental and epidemiological data concerning overuse injuries in youth sports. Review studies typically described pediatric sports injuries in general terms with little respect to injury type. The current evidence surrounding risk factors, prevention and treatment strategies for overuse injuries in youth sports was primarily limited to review studies and *Level III and IV* publications. The paucity of high quality evidence combined with strict inclusion

criteria appeared to impact the study selection process. A variety of search terms were used however a disproportionate number of studies featured Little League Shoulder as a diagnosis of interest. This selection bias towards overuse physeal injuries in the UE may have influenced the generalizability of the clinical recommendations made in this systematic review.

#### 2.5.d Knowledge Gaps

The lack of high quality, patient-oriented research in younger athlete populations and the absence of research describing physeal injuries pose notable gaps in the literature. These gaps include minimal data establishing the incidence, prevalence and severity of overuse injuries in youth athletes, especially with respect to physeal involvement.<sup>91,116</sup> No original research studies have clearly defined physeal injuries at this time. Also, little is known about the effects of population-specific risk factors, like growth-related changes and training volume, on the development of injuries in skeletally immature individuals. Future studies should 1) seek to establish a clear definition of physeal injuries in sport, and 2) understand the mechanisms and risk factors associated with their development. This will provide the foundation for more effective prevention and treatment strategies at the policy level, including the paradigm-shifting concept of scheduled periods of rest from sport. Scheduled rest provides youth athletes the time they need to physically and mentally recover from the rigors of competitive sport.

#### 2.6 Conclusion

Overuse physeal injuries are multi-factorial in nature.<sup>95</sup> Periods of accelerated growth, chronological age, skeletal maturity and history of previous injury can predispose young athletes to repetitive stress injuries.<sup>95</sup> Modifiable risk factors such as flexibility,



strength and training volume should be regularly monitored in an effort to limit risk-prone activities and prevent injuries when possible.<sup>21,26,34,35,56,79,134,149</sup>

The most effective treatment strategy following overuse physal injuries is an extended period of active rest. Following symptom resolution, clinicians can begin to restore function through improvements in flexibility, strength and neuromuscular control.<sup>2,3,5,13,20,22-24,26,27,29,35,41,42,47,78,79,132,140</sup> Progressive strength training programs should include gradual return to throwing or running programs when appropriate.<sup>77</sup> Return to sport timelines typically range from 4-6 weeks in most cases, but can extend to 3-5 months when symptoms persist.<sup>27,77,83</sup>

## **2.7 Clinical Recommendations**

The most widely accepted treatment option following any physal injury is an extended period of active rest and when necessary joint immobilization. Once the pain has subsided, emphasis on the restoration of flexibility, strength and sport-specific endurance is appropriate.<sup>3,10,19,27,30,74,76,114,115,134</sup>

### *Strength-of-Recommendation Taxonomy: C*

- 1) Modifications such as implementing sport-specific flexibility and strength programs as well as limiting training and competition volumes (i.e. pitch counts) are 2 strategies to avoid overuse and fatigue-related injuries. This is especially important during periods of rapid growth.<sup>21,25,26,77,90,91,95,116,120,127,135,138,140,148</sup>

### *Strength-of-Recommendation Taxonomy: B*

- 2) Regular monitoring of anthropometric (i.e. height and weight) and physical characteristics (i.e. range of motion and strength) in youth athletes may prove

preventative as deficits have been linked to both UE and LE injuries in multiple sporting events.<sup>30,77,116,148,152</sup>

*Strength-of-Recommendation Taxonomy: C*




Figure 2.1 PRISMA Flow Diagram<sup>104</sup>

Strength of Recommendations	Definition
A	Recommendation based on consistent, good quality patient-oriented evidence* (morbidity, mortality, exercise and cognitive performance, physiologic responses).
B	Recommendation based on inconsistent or limited quality patient-oriented* evidence.
C	Recommendation based on consensus, usual practice, opinion, disease-oriented evidence* case series or studies of diagnosis, treatment, prevention, screening, or extrapolations from quasi-experimental research.

\*Patient-oriented evidence measures outcomes that matter to patients: morbidity, mortality, symptom improvement, cost reduction, and quality of life. Disease-oriented evidence measures intermediate, physiologic, or surrogate end points that may or may not reflect improvements in patient outcomes (e.g.: blood pressure, blood chemistry, physiologic function, pathologic findings).

Figure 2.2 Strength-of-Recommendation Taxonomy (SORT) Diagram<sup>49</sup>



Figure 2.3 Radiograph of 13-year-old male football player with Osgood-Schlatter Disease



Figure 2.4 Radiograph of 14-year-old male baseball player with Little League Shoulder



Figure 2.5 Radiograph of 13-year-old male baseball player with Little League Elbow

**Table 2.1. Studies that Report Lower Extremity Physeal Injuries**

Author	Level of Evidence (OCEBM)	Sample Size	Age (yrs)	Injury Site	Sport	Treatment Options
Beovich <sup>9</sup> (1988)	III	22	9 – 18	Proximal tibial tubercle	Multiple	Activity modifications (20); Active rest (2)
de Lucena <sup>34</sup> (2010)	III	954	12 – 15	Proximal tibial tubercle	Multiple	Stretching program
Doral <sup>45</sup> (2005)	IV	1	16	Anterior superior iliac spine	Soccer	Surgical intervention
Hajdu <sup>59</sup> (2000)	IV	7	13 – 16	Proximal tibial tubercle	Ball games, skiing	Active rest (1); Surgical intervention (6)
Hussain <sup>66</sup> (1996)	III	261	11 – 18	Proximal tibial tubercle	Multiple	Active rest & NSAIDs (237); Surgical intervention (24)
Kolt <sup>79</sup> (1999)	III	43	11 – 19	Multiple sites	Gymnastics	Physical conditioning program
Kujala <sup>82</sup> (1985)	III	68	9 – 18	Proximal tibial tubercle	Multiple	Active rest 3 months; Activity modifications 7 months
Laor <sup>83</sup> (2006)	IV	6	8 – 15	Distal femur, proximal tibia, proximal fibula	Football, basketball, gymnastics, other	Joint immobilization 1-5 wks
Liebling <sup>88</sup> (1995)	IV	1	13	Distal femur, proximal tibia	Baseball	None
Mital <sup>103</sup> (1980)	III	118	9 – 18	Proximal tibial tubercle	Multiple	Active rest/Joint immobilization (104); Surgical intervention (14)
Nanni <sup>114</sup> (2005)	IV	1	15	Proximal tibia	Rugby	Surgical intervention



Orava <sup>117</sup> (1982)	III	185	9 – 26	Multiple sites	Multiple	Varied
Rossi <sup>130</sup> (2001)	III	203	11 – 18	Pelvic apophyses	Soccer, gymnastics, fencing, tennis	None
Valentino <sup>150</sup> (2012)	IV	1	13	Inferior patellar pole	Football	Active rest 5 months

**Table 2.2. Studies that Report Upper Extremity Physéal Injuries**

Author	Level of Evidence (OCEBM)	Sample Size	Age (yrs)	Injury Site	Sport	Treatment Options
Akgul <sup>2</sup> (2011)	IV	1	13	Proximal humerus	Non-athlete	Active rest 4 months
Anton <sup>5</sup> (2010)	IV	1	13	Proximal humerus	Baseball	Active rest, Physical therapy
Binder <sup>10</sup> (2011)	III	72	8 – 13	Proximal humerus	Unknown	Joint immobilization 1-4 wks (57); Surgical intervention (15)
Boyd <sup>13</sup> (1997)	IV	1	15	Proximal humerus	Badminton	None
Carson <sup>27</sup> (1998)	III	23	14	Proximal humerus	Baseball	Active rest 3 months
Drescher <sup>47</sup> (2004)	IV	1	12	Proximal humerus	Cricket	Joint immobilization 3 wks; Active rest 3 months
Hang <sup>60</sup> (2004)	III	343	8 – 12	Distal humerus	Baseball	None
Kolt <sup>79</sup> (1999)	III	43	11 – 19	Multiple sites	Gymnastics	Physical conditioning program
Obembe <sup>115</sup> (2007)	IV	4	11 – 15	Proximal humerus	Baseball, Tennis	Active rest 3 months
Orava <sup>117</sup> (1982)	III	185	9 – 26	Multiple sites	Multiple	Varied
Roy <sup>131</sup> (1985)	IV	21	11 – 18	Distal radius	Gymnastics	None
Torg <sup>147</sup> (1972)	IV	1	12	Proximal humerus	Baseball	None

## CHAPTER 3

### METHODS

#### **3.1 Research Design**

This study prospectively followed a cohort of competitive youth baseball players over the course of a 6-month season (Figure 3.1). Approval for this study was received from the University of South Carolina's Institutional Review Board (IRB). The research team complied with all rules, regulations and training requirements put forth by the IRB.

#### **3.2 Study Setting**

This study was conducted at local baseball clubs (Hit House, Elite Baseball and Southern Athletics), as well as the Greenville and Northwood Little Leagues. ATI Physical Therapy and Steadman-Hawkins Clinic of the Carolinas facilities were also used during this study.

#### **3.3 Study Subjects**

Two hundred and sixty-one competitive baseball players were recruited for this study based on sample size calculations with 80% power and an alpha level of 0.05. Inclusion criteria required that all participants be male, between 9 and 12 years of age and uninjured at the time of baseline data collection. Players were excluded from the study if they reported any current injuries that restricted their ability to participate in baseball activities or if they reported a shoulder or elbow injury that required medical attention during the 3 months prior to the start of the study. Recruitment strategies were based on well-established community relationships with the coaches and parents of several local

baseball clubs as well as the Greenville and Northwood Little League Baseball programs. To increase retention and response rates over the course of a 6-month follow up period, the primary investigator emphasized the importance of both team coaches' and parents' participation in the bi-monthly online surveys. The primary investigator also acted as the established point of entry into the medical community, for coaches and parents, when a baseball player suffered a baseball-related upper extremity injury.

### **3.4 Procedures**

An online survey (see Appendix A), in conjunction with a baseline physical examination (see Appendix B), was completed during the course of the competitive baseball season to identify potential risk factors for injury in this population (Table 3.1). Players' training and playing histories, as well as any injuries sustained, was be tracked every 2 weeks over the course of a 6-month period (Table 3.1). Team coaches and participants' parents were contacted every 2 weeks via online surveys, phone conversations and in person visits in an effort to improve response rates as well as corroborate data reported for each player. All baseball players who reported baseball-related upper extremity impairments underwent a subsequent physical examination performed by the primary investigator to confirm the presence of an injury. A second exam was also performed in a subset of youth players to assess any changes from the baseline physical data recorded earlier in the season. The subsequent examination included height, weight, shoulder and elbow range of motion (ROM) and isometric shoulder strength.

Height and weight measurements for each participant were measured using a portable stadiometer and digital weight scale, respectively. Humeral torsion, passive

shoulder and elbow ROM, isometric shoulder strength and ulnar length were assessed in both the dominant and non-dominant arms of each participant. Two values of each measure were recorded per arm.

Humeral torsion was measured using a Sonosite-Edge (Sonosite Inc, Bothell, WA, USA) ultrasonography unit with a 6 cm linear array transducer (6-15 MHz). Examiners used a previously validated indirect ultrasonography method to assess the differences in bony development between the dominant and non-dominant arms.<sup>57,110-112,118,153-155</sup> All measurements were taken in the supine position with the arm at 90° of abduction and the elbow at 90° of flexion. One examiner passively rotated the arm until the apices of the greater and lesser tuberosities were parallel in the coronal plane. A second examiner placed a digital inclinometer (Fabrication Enterprises Inc, White Plains, NY, USA) along the ulnar border and recorded the corresponding angle of the forearm. In this method, the larger angle indicated less humeral retrotorsion.<sup>7,111,153,161</sup> Acceptable intra-rater reliability for measurement of HT was established prior to the data collection (ICC<sub>2,1</sub> = 0.92-0.99; SEM = 1.7°-3.8°).

Shoulder external rotation (ER) and internal rotation (IR) ROM was assessed bilaterally in supine using a digital inclinometer and methods previously reported in the literature.<sup>7,136,158</sup> The scapula was stabilized at the corocoid process with the arm at 90° of abduction and elbow flexion. A towel roll was placed under the distal humerus to maintain the scapular plane. The arm was then passively rotated to end range for measurement. No overpressure was applied. Horizontal adduction ROM was assessed with the athlete in the supine position. Full scapular retraction was maintained with stabilization at the lateral scapula, while the examiner horizontally adducted the arm

maintaining neutral humeral rotation until resistance was felt. A digital inclinometer was used to assess the angle between the humerus and the horizontal plane of the body.<sup>7,136</sup> Elbow extension ROM was assessed with the athlete in the supine position with the arm in 90° of abduction and neutral rotation. A towel roll was placed under the distal humerus to maintain the scapular plane. A digital inclinometer was placed along the anterior surface of the forearm in the plane of the acromion using the radial styloid process as the primary landmark. The angle of the radial styloid process relative to the parallel line of the acromion was recorded. Positive values indicated elbow hyperextension. Acceptable intra-rater reliability was established for all ROM measurements prior to data collection (ICC<sub>2,1</sub> = 0.92-0.99; SEM = 1.3°-3.8°). An average of 2 trials was used for each measure in data analysis. Total arc of motion was calculated by adding mean ER ROM with IR ROM for the dominant and non-dominant shoulders, respectively. Side-to-side differences were calculated for each ROM variable by subtracting dominant values from the non-dominant values.

Isometric shoulder strength was assessed bilaterally using a Lafayette Manual Muscle Tester hand-held dynamometer (Lafayette Instrument Company, Lafayette, IN, USA) and previously reported methods.<sup>17</sup> Isometric shoulder strength measures included scaption at 90°, ER at 0°, ER at 90° and IR at 90° for the dominant and non-dominant arms. Make tests were used for each isometric strength measure based on higher reliability when compared to break tests in hand-held dynamometers.<sup>146</sup> Scaption strength was measured in the seated position. The arm was abducted to 90° and then horizontally adducted to 45° with neutral shoulder rotation. The hand-held dynamometer was placed 5 cm distal to the cubital fossa. The participant raised the arm perpendicular to the floor

using maximum effort.<sup>17</sup> External rotation at 0° was assessed with the arm in 0° of shoulder abduction with a towel roll placed under the axilla. The elbow was positioned in 90° of flexion with neutral forearm rotation. The hand-held dynamometer was placed on the dorsal aspect of the forearm, 2 cm proximal to ulnar styloid process. The participant then externally rotated the arm with maximum effort.<sup>17</sup> External rotation at 90° was measured with the shoulder in 90° of abduction, 90° of ER and 90° of elbow flexion. A towel roll was placed under the distal humerus to maintain the arm in the plane of the body. The dynamometer was placed on the dorsal aspect of the forearm, 2 cm proximal to ulnar styloid process. The participant then externally rotated the arm with maximum effort.<sup>17,46</sup> Internal rotation at 90° was assessed in a similar fashion to ER at 90°, however the shoulder was in a state of neutral rotation and the dynamometer was placed on the volar aspect of the forearm. The participant was asked to maximally internally rotate his arm.<sup>17</sup> Acceptable intra-rater reliability for all isometric shoulder strength measures was established prior to the data collection ( $ICC_{2,1} = 0.94-0.99$ ; SEM = 1.3-3.6 lbs). To ensure that minimal detectable change (MDC) exceeded the standard error of measure (SEM), intra-rate reliability was re-calculated using the first 10 participants' data. Ulnar length measurements were recorded for the dominant (D) and non-dominant (ND) arms and used to calculate both ER and IR torque values for each participant.<sup>68</sup> Isometric shoulder strength was then normalized prior to data analysis using 5 separate methods: body mass, body mass index (BMI), height, torque and the Trakis Method (Table 3.2).<sup>148</sup>

Throwing performance was assessed using ball velocity during an overhead throw. This measure was assessed within 10 days of baseline data collection in a subset of 80 participants. Following a warm up period during team practice, participants were

asked to throw 3 balls from a distance of 46 feet on flat ground to a specified target. A Stalker radar gun was used to record the velocity of each throw in miles per hour (mph). Only accurate ball velocities were recorded and then used in data analyses to determine if isometric shoulder strength and throwing performance were related.

Baseball-related upper extremity injuries will be defined as any shoulder or elbow impairment that resulted in either: 1) an athlete missing  $\geq 1$  practice(s) or game(s), or 2) an athlete experiencing a reduction in their performance (i.e. decreased ball control or velocity) or change in position (i.e. moving from pitcher to 1<sup>st</sup> base) related to the UE complaint.<sup>43,75,113,137</sup> Injuries were originally identified via self-report from the team coaches and the participant's parents using the RedCap online survey system, phone conversations and in person interviews. Players, parents and coaches were also encouraged to contact the research team if any concerns arose throughout the study. Once identified, all baseball-related upper extremity injuries were examined by the primary investigator, a licensed physical therapist and when necessary, referred to a board certified, fellowship-trained, sports medicine physician at the Steadman Hawkins Clinic of the Carolinas for further evaluation. All injured baseball players who were examined by a physician may have received x-rays as part of their evaluation per typical standard of care. Any damage to the physis or 'growth plate' at the shoulder or elbow was confirmed through physician evaluation and diagnosis. All costs related to physician visits, including diagnostic imaging, was billed to the players' insurance companies. This study was not responsible for any medical expenses incurred as a result of a baseball-related injury.



### 3.5 Statistical Analyses

#### 3.5.a Specific Aim 1

Determine the incidence of baseball-related upper extremity injuries in a cohort of 9-12 year old male baseball players.

Specific incidence was defined as the number of baseball-related upper extremity injuries recorded per 1,000 athletic exposures. As stated previously, an athletic exposure was defined as 1 organized team practice or game that an athlete participated in. Means and standard deviations were calculated for each variable based on injury status.

Injury status was the main outcome of interest used in the power calculations for Specific Aim 1. The number of baseball-related upper extremity injuries necessary to statistically compare injured and uninjured groups was approximately 50 baseball players. Based on previous research, a baseball-related injury rate of 0.18 was expected, which indicated that a sample size of 275 baseball players would have been adequate to capture the target population of youth players who sustain baseball-related injuries.<sup>31,137</sup> Those calculations assumed a 2-sided alpha level of 0.05. Based on that analysis, the current study had sufficient power (100%) to assess upper extremity injury risk in a cohort of 9-12 year old male baseball players.

#### 3.5.b Specific Aim 2

Compare the measurement properties of normalized isometric shoulder strength, by 5 separate methods, for use as a multi-faceted clinical assessment tool that was responsive to changes in physical growth and development over time in a cohort of 9-12 year old male baseball players.

The main outcomes of interest were the measurement properties of each of the 5 normalization methods (body mass, BMI, height, torque and the Trakis method<sup>148</sup>) which included test-retest reliability of each strength measure, the ability to detect changes in shoulder strength over time and the strength of the relationship between shoulder strength and ball velocity as a measure of throwing performance. Means and standard deviations were calculated for each normalized isometric shoulder strength measures by method. Humeral torsion and shoulder ROM measures were also assessed as they were thought to potentially influence isometric shoulder strength in youth baseball players. When analyzed, no significant relationships were observed between isometric shoulder strength and humeral torsion or isometric shoulder strength and shoulder ROM in this population.

Reliability was assessed for all baseline and follow-up strength measures using intraclass correlation coefficients (ICC) with corresponding 95% confidence intervals (CI).<sup>122</sup> Standard errors of measurement (SEM) were also calculated to determine the absolute reliability of each strength measure using the largest SD in the formula  $SD \times \sqrt{1 - ICC}$ .<sup>122</sup> Individual SEMs were then used to calculate corresponding minimal detectable change (MDC<sub>95</sub>) values for each of the normalized strength measures using the formula  $SEM \times 1.96 \times \sqrt{2}$ .<sup>122</sup> Repeated measures analyses of variance (ANOVA) were conducted to compare changes in isometric shoulder strength at 2 time points (baseline and follow-up) after co-varying for physical growth and body size. Effect sizes were calculated to identify the magnitude of change detected between the 2 time points for each of the normalized strength measures. Linear regression models were used to examine the relationships between the normalized isometric shoulder strength measures and ball velocity in youth baseball players. The method with the most consistent

measurement properties for normalizing isometric shoulder strength in youth baseball players was determined based on each measure's test-retest reliability, ability to detect changes over time and strength of association with ball velocity. Statistical significance was set a priori at  $\alpha=0.05$ . All statistical analyses were performed using SPSS Statistics 21.0 (SPSS Inc., Chicago, IL, USA) software.

The study team demonstrated good reliability for all the variables of interest including isometric shoulder strength with a standard error of measurement of less than 5% for each measure. Only isometric shoulder strength data was presented, as they were the main variables of interest and represented the most variable data in the analyses. The small to moderate effect sizes used in the power calculations below were estimated using data from previous studies. These calculations assumed a 2-sided alpha level of 0.05 and showed that the current study was sufficiently powered (>91% for the isometric force production variables of interest).




Figure 3.1. Prospective Cohort Study Design Flowchart

**Table 3.1. Data Collection Measures**

<b>Dependent Variable Name</b>	<b>Definition</b>	<b>Collection Method</b>
<b>Upper Extremity Injury</b>	Any shoulder or elbow impairment that resulted in either: 1) athlete missing $\geq 1$ practice(s) or game(s), or 2) athlete experiencing a reduction in their performance (i.e. decreased ball control or velocity) or change in position (i.e. moving from pitcher to 1 <sup>st</sup> base) related to the UE complaint. <sup>43,75,113,137</sup>	Self-report followed by PT examination and physician examination when necessary
<b>Physical Injury</b>	An injury at the shoulder or elbow joint with physician confirmed damage to the corresponding physis <sup>18</sup>	Physician evaluation and diagnosis
<b>Independent Variable Name</b>	<b>Definition</b>	<b>Collection Method</b>
<b>Athletic Exposure</b>	1 organized team practice or game <sup>137</sup>	Bi-monthly self report from coaches and parents via online survey system and phone responses
<b>Age (years)</b>	= Date of data collection – date of birth	Date of birth
<b>Height (cm)</b>	Measured to the nearest 0.5 cm	Stadiometer
<b>Weight (lbs)</b>	Measured to the 1 <sup>st</sup> decimal place (ex: 119.1 lbs)	Digital scale
<b>Position: Primary and Additional Positions</b>	Position categories: 1 = pitcher 2 = position player	Self report
<b>Sport Specialization Status</b>	Self-Classification: Athletes were asked to identify as a specialized or multi-sport athlete Research-Based Classification: Athletes met at least 2/3 of the following research-based criteria: <ul style="list-style-type: none"> <li>• Participated in organized baseball activities <math>\geq 8</math> months/year</li> <li>• Participated on <math>\geq 1</math> organized baseball team during the year</li> <li>• Participated in additional baseball-specific specialty training (i.e. pitching lessons)</li> </ul>	Self report and Research-based criteria <sup>8,71,72,84</sup>

<b>Participation in Additional Specialty Training</b>	Training type categories: 1 = hitting lessons 2 = pitching lessons 3 = hitting or pitching lessons (baseball-specific) 4 = strength and agility training	Self report
<b>Pain Level</b>	0-10 with 10 being the highest level of pain	Pediatric Visual Analog Scale (VAS) score <sup>159</sup>
<b>Humeral Torsion</b>	The orientation of the humeral head relative to the transverse plan of the body <sup>57,110-112,118,153-155</sup>	Indirect method using diagnostic ultrasonography <sup>110,153</sup>
<b>ROM: ER at 90°</b>	Maximal shoulder ER with scapula stabilized and no overpressure <sup>64,65,73,101,118,129,139,148,149,163</sup>	Supine with digital inclinometer <sup>64,65,139</sup>
<b>ROM: IR at 90°</b>	Maximal shoulder IR with scapula stabilized and no overpressure <sup>64,65,73,101,118,129,139,148,149,163</sup>	
<b>ROM: Elbow Extension</b>	Maximal elbow extension with shoulder at 90° of abduction and no overpressure	
<b>ROM: Horizontal Adduction</b>	Maximal shoulder HA with scapula laterally stabilized and no overpressure <sup>64,65,73,101,118,129,139,148,149,163</sup>	
<b>Isometric Strength: Scaption at 90°</b>	Maximal force produced against manual resistance with upright posture and no trunk support <sup>1,17,44,73,97,105,128,129,148,149</sup>	
<b>Isometric Strength: ER at 0°</b>	Maximal force produced against manual resistance with upright posture and no trunk support <sup>1,17,44,73,97,105,128,129,148,149</sup>	
<b>Isometric Strength: ER at 90°</b>	Maximal force produced against manual resistance with shoulder at 90° abduction/90° ER <sup>1,17,44,73,97,105,128,129,148,149</sup>	
<b>Isometric Strength: IR at 90°</b>	Maximal force produced against manual resistance with shoulder at neutral rotation <sup>1,17,44,46,73,97,105,128,129,148,149</sup>	Prone make test with HHD <sup>17</sup>
<b>Isometric Strength: ER:IR Ratio</b>	= $\frac{\text{ER strength at 90°}}{\text{IR strength at 90°}}$	
<b>Ulnar Length</b>	Measurement between tip of olecranon process to most distal portion of ulnar styloid process <sup>67,68</sup>	Supine with shoulder in 90°/90° position with tape measure

<b>Torque</b>	Torque = isometric force production x lever arm <sup>67,68</sup>	= ER strength at 90° x ulnar length <sup>67,68</sup> = IR strength at 90° x ulnar length <sup>67,68</sup>
<b>Throwing Performance: Ball velocity</b>	Miles per hour (mph)	46 ft throw to a specified target with Stalker radar gun
<b>Exposure to Training: # of practices/time period</b>	Total # of practices played over the 6 month follow up period <sup>137</sup>	Bi-monthly self report from coaches and parents via online survey system and phone responses
<b>Exposure to Training: # of games/time period</b>	Total # of games played over the 6 month follow up period <sup>137</sup>	
<b>Exposure to Training: # of months played/year</b>	Total # of months where the athlete participated in organized baseball activities	
<b>Previous History of Injury</b>	Any previous injury that required medical attention or resulted in the athlete missing $\geq 1$ practice or game <sup>149</sup>	Self report

**Table 3.2. Normalized Isometric Shoulder Strength Calculations by Method**

<b>Method</b>	<b>Calculation</b>
<b>Body Mass (kg)</b>	= $\frac{\text{Shoulder Strength Measure}}{\text{Body Mass}}$
<b>BMI (kg/m<sup>2</sup>)</b>	= $\frac{\text{Shoulder Strength Measure}}{\text{BMI}}$
<b>Height (cm)</b>	= $\frac{\text{Shoulder Strength Measure}}{\text{Height}}$
<b>Torque (Nm)</b>	= Shoulder Strength Measure (N) x Ulnar Length (m)
<b>ND Strength (%)<sup>148</sup></b>	= $\frac{(\text{Dominant Shoulder Strength} - \text{Non-Dominant Shoulder Strength})}{(\text{Non-Dominant Shoulder Strength})}$

**Table 3.3. Aim 1 – A Priori Power Analyses Based on Expected Injury Rates<sup>a</sup>**

	<b>Sample Size</b>	<b>Total Injury Rate</b>	<b>Estimated # of Injuries</b>	<b>Estimated Power</b>
<b>Pilot Study</b>	275	0.16	44	100%
<b>Based on Previous Research<sup>31,137</sup></b>	275	0.18	50	100%

<sup>a</sup> Total injury rates determined using data from a pilot study and two previous research studies conducted in adolescent baseball players (13-18 years old).

**Table 3.4. Aim 2 – A Priori Power Analyses Based on Isometric Shoulder Strength Measures<sup>a</sup>**

<b>Independent Measure</b>	<b>Effect Size<sup>b</sup></b>	<b>Alpha Level</b>	<b>Estimated Power</b>
<b>Dominant Shoulder ER Strength</b>	0.41	0.05	99%
<b>Dominant Shoulder ER:IR Strength Ratio</b>	0.20	0.05	91%
<b>Normalized Dominant Shoulder ER Strength using %ND Strength Method<sup>148</sup></b>	0.22	0.05	95%

<sup>a</sup> The power analyses for Aim 2 were conducted using isometric shoulder strength data from a pilot study.

<sup>b</sup> Effect size was calculated to examine the expected difference between baseline and follow up measures using isometric shoulder strength data collected in a pilot study.



## CHAPTER 4

### SPORT SPECIALIZATION INCREASES UPPER EXTREMITY INJURY RISK IN YOUTH BASEBALL PLAYERS: A PROSPECTIVE COHORT STUDY

#### 4.1 Abstract

**Introduction:** Despite rising awareness of the risks associated with year-round sports specialization, athletes continue to specialize at increasing rates across the U.S. The effects of sport specialization on the development of upper extremity (UE) injuries in baseball players have not been previously studied in youth populations.

**Objective:** The purposes of this study were to 1) establish UE injury incidence, and 2) examine the association of sport specialization and specialty training as a pitcher on UE injury rates in a cohort of youth athletes.

**Methods:** Youth baseball players (9-12 years old) were examined and then followed for approximately 6 months via coach/parent surveys. Athletic exposure (AE) and presence of UE injury was tracked per player. All athletes who reported injuries were re-examined by a licensed physical therapist. Athletes were classified as specialized or multi-sport using 2 methods: self-classification and research-based classification, however research-based results were used for data analysis. Chi square analyses were used to compare the frequency of UE injuries based on position group (pitchers vs. position players only), sports specialization status and participation in additional specialty training. Odds ratios,

absolute, and absolute risk differences with 95% confidence intervals (CI) were calculated between groups.

Results: Uninjured male baseball players (n=159) were prospectively followed for an average of  $6.7 \pm 1.5$  months in this study. The UE injury incidence rate was 16.3/1000 AEs (95% CI=9.3, 23.3). The majority of athletes (83.0%) were classified as specialized in this cohort. Specialized athletes demonstrated a 15.9% increase in absolute risk for developing an UE injury when compared to multi-sport counterparts ( $P=0.03$ ).

Conclusion: Sport specialization impacts an athlete's UE injury risk during youth baseball. USA Baseball's pitch count limitations were designed to decrease overuse injuries at the shoulder and elbow by requiring more athletes to pitch. This may have inadvertently had the opposite effect by increasing the rate of specialty training outside of competition.

Clinical Relevance: Participation in specialty training as a pitcher may influence the development of UE injuries in youth populations. Youth pitchers who took pitching lessons demonstrated a significant increase in absolute injury risk, which may be related to increased athletic exposure.

Keywords: youth baseball; risk factors; early sports specialization; early position specialization

## 4.2 Introduction

Baseball is a popular sport to play in the United States, with approximately 13-17 million athletes under the age of 18 participating at the club and high school levels.<sup>4,31,51,143</sup> The incidence of baseball-related overuse injuries is comparatively low in adolescent players (13-18 years old) with 1.3 – 4.0 injuries per 1,000 athletic exposures recorded.<sup>31,137</sup> It is unknown in youth baseball players (9-12 years old) as there is a significant lack of epidemiologic data in this population. The majority of baseball-related overuse injuries are reported in the upper extremity (UE), specifically at the shoulder and elbow, however little is known about the etiology and development of these injuries at the youth level.<sup>31,137</sup>

One potential explanation for the lack of epidemiologic data may be related to difficulties in injury surveillance, particularly in the younger age groups.<sup>43,75</sup> Unlike the collegiate and professional ranks, who employ athletic trainers to record and treat their injuries, youth and adolescent injuries are inconsistently reported, and often treated, by the athletes' parents and coaches.<sup>43,113</sup> The burden of identifying and recording injuries is much greater at this level as the majority of youth players participate on multiple teams, and in some cases multiple sports, throughout the year.<sup>75</sup>

Despite evidence suggesting that sport specialization may be related to the development of overuse injuries in youth and adolescent athletes, its prevalence continues to increase in the U.S.<sup>72,107</sup> Prior research has defined sport specialization based on a battery of criteria including year-round training in a single sport (>8 months/year), identification of a primary sport over additional sports and the cessation of additional sports to focus on a primary sport.<sup>8,71,72,84</sup> Despite established definitions in the literature,

less is understood about the public perception of sports specialization with parents and coaches.<sup>71,72,84,107,108</sup> The effects of sport specialization on UE injury risk has not been previously established in the baseball literature. Other risk factors, such as excessive pitch counts, varied pitch types and faulty throwing mechanics, have been linked to the development of shoulder and elbow pain in youth throwers using self-report and survey data.<sup>12,71,72,90,91,116,162</sup> The USA Baseball Medical & Safety Advisory Committee used this research to establish age-appropriate guidelines for pitch counts, pitch types, throwing mechanics and most recently a long term athlete development model released in 2017.<sup>12,19,90,91,109,136,140,148,154,155</sup> The effectiveness of these recommendations on the reduction of baseball-related overuse injuries is unknown at this time.

The paucity of data describing UE injury incidence in youth baseball players poses a significant knowledge gap in the literature as the extent of the problem has not been accurately established in this population. Additional research is also needed to better understand the impact of sport specialization on shoulder and elbow injury risk in youth baseball players. The purposes of this study were to 1) establish UE injury incidence in a cohort of 9-12 year old male baseball players, and 2) examine the effects of sport specialization on the development of UE injuries in a cohort of 9-12 year old male baseball players.

### **4.3 Methods**

#### **4.3.a Study Population**

This is a prospective cohort study of competitive male youth baseball players (9-12 years old) recruited in the spring of 2016. Two hundred and sixty one players were recruited from local baseball clubs, baseball tournaments and little leagues in South

Carolina and invited to participate in this study. One hundred and fifty nine athletes consented to participate. Asymptomatic competitive youth baseball players were followed for a 6-month period after baseline examination by a research team of licensed physical therapists. All players were male, between the ages of 9-12 years and participating in all baseball activities without restriction at the time of baseline examination. Pitchers and position players were recruited for this study, however youth teams are predominantly comprised of pitchers (Table 4.1). Position players were identified as any athlete who did not report pitching for an organized baseball team.

One hundred and two players were excluded from this study because they (1) reported any injuries that currently restricted their ability to participate in baseball activities, (2) reported a shoulder or elbow injury that required medical attention during the 3 months prior to initial examination or (3) did not respond the required number of times during the follow up period (3 times throughout the study with each response being <2 months apart). The University of South Carolina's Institutional Review Board (IRB) approved this study. Parental consent and athlete assent were obtained for each participant enrolled.

#### 4.3.b Data Collection

At the time of enrollment (Spring 2016), each participant completed a study questionnaire using the RedCap online survey system prior to initial examination and throughout the 6-month follow up period. Participants were then contacted twice a month using online survey methods followed by phone and in-person interviews for improved response rates.

All participants completed a baseline online study questionnaire with the help of a parent or team coach. The initial questionnaire surveyed baseline characteristics, current sports participation, baseball-specific playing history and training history. Participants were asked to self-classify as either specialized or multi-sport athletes at the time of study enrollment (Figure 4.1). The research team then re-classified each participant as specialized or multi-sport, using previously stated research criteria (See Appendix A).<sup>8,71,72</sup> Follow up questionnaires surveyed baseball-related athletic exposure (i.e. team practice and game counts) first by team and then confirmed through individual report over the course of 6 months. Any shoulder or elbow impairments including pain, injury, tightness or weakness reported during that time were also recorded. Any player that reported a baseball-related shoulder or elbow impairment via survey was contacted and then physically examined by the lead researcher, a licensed physical therapist, to confirm the presence of injury. Acute trauma such as acute fractures, lacerations and abrasions were not included in this study.

Injuries were defined as any shoulder or elbow impairment that resulted in either: 1) an athlete missing  $\geq 1$  practice(s) or game(s), or 2) an athlete experiencing a reduction in their performance (i.e. decreased ball control or velocity) or change in position (i.e. moving from pitcher to 1<sup>st</sup> base) related to the UE complaint.<sup>43,75,113,137</sup> Following the physical examination, athletes who required additional medical care were referred to a board certified, fellowship-trained, sports medicine physician for continued evaluation.

#### 4.3.c Statistical Analysis

Incidence rates were calculated per 1,000 athletic exposures for all UE injuries. Rate ratios (RR) and 95% confidence intervals were then calculated to determine injury

rates in pitchers and position players only. Frequency counts were also calculated for each categorical variable. Chi square analyses were used to compare the frequency of all shoulder and elbow injuries between the following groups: sports specialization status, position played and participation in specialty training outside of team-sanctioned practices or games. Odds ratios (OR), absolute risk (AR) and absolute risk difference (ARD) with 95% confidence intervals (CI) were then respectively calculated for each of these groups. Absolute risk difference was determined by subtracting the AR of the exposed group (athletes who demonstrated the risk factor) from the AR of the unexposed group (athletes who did not demonstrate the risk factor). Statistical significance was set a priori at  $\alpha=0.05$ . All statistical analyses were performed using SPSS Statistics 22.0 (SPSS Inc., Chicago, IL, USA) software.

#### **4.4 Results**

##### **4.4.a Injury Incidence**

The UE injury incidence rate in a cohort of healthy 9-12 year old male baseball players was 16.3/1000 AEs (95% CI = 9.3, 23.3). Twenty-one injuries were reported during the 6-month follow up period, 14 in pitchers (13.2%) and 7 in position players (13.2%). Pitchers represented 66.7% (n = 106) of the cohort and demonstrated an UE incidence rate of 16.6/1000 AEs (95% CI 7.9, 25.3) while position players represented 33.3% (n = 53) of the cohort and demonstrated 15.8/1000 AEs (95% CI 4.1, 27.5). The difference in incidence rates was not significant (RR = 1.1, 95% CI = 0.4, 2.6;  $P = 0.91$ ). The proportion of UE injuries was highest at the shoulder (61.9%, n = 13), followed by the elbow (38.1%, n = 8) in youth baseball players. This pattern was consistent in pitchers (shoulder: 66.7%, n = 10; elbow: 33.3%, n = 5) while position players

demonstrated an equal injury distribution across both sites (shoulder: 50.0%, n = 3; elbow: 50.0%, n = 3). Despite these observations, location of UE injury was not significantly different between pitchers and position players ( $P = 0.48$ ). Of the 21 athletes who sustained injuries in the study, only 1 reported a history of overuse UE injury prior to the onset of symptoms.

#### 4.4.b Early Sport Specialization

At the start of the study, participants were asked to self-classify as either specialized or multi-sport athletes based on their or their parents' perceptions of sport specialization in youth baseball. Thirty-one percent of youth athletes (n = 49) self-classified as specialized in baseball while the remaining 69.0% (n = 110) identified as multi-sport athletes (Figure 4.1). The research team then re-classified each participant as specialized or multi-sport, using research-based criteria found in the literature.<sup>8,71,72</sup> Based on these criteria, 83.0% of the cohort (n = 132) qualified as specialized athletes while only 17.0% (n = 27) were classified as true multi-sport athletes (Figure 4.1). The study results showed that a significant number of youth baseball players were misclassified as multi-sport athletes yet participated and competed as specialized athletes (57.9%, n = 92;  $P = 0.001$ ). The research-based methods were used for athlete classification the remainder of the statistical analyses. Youth baseball players that competed as specialized athletes were at a significantly greater risk for developing a shoulder or elbow injury when compared to their multi-sport counterparts (OR = 1.2, 95% CI 1.1, 1.3;  $P = 0.03$ ) (Table 4.1).



#### 4.4.c Early Position Specialization

The majority of youth baseball players in this cohort participated in some type of additional sport-related training (84.9%,  $n = 135$ ). One hundred and eleven players took formal hitting lessons (69.8%), 72 took formal pitching lessons (67.9%) and 36 participated in generalized strength and conditioning programs (22.6%) (Table 4.1). Youth baseball players who participated in formal hitting lessons demonstrated no differences in injury frequency compared to players who did not ( $P = 0.86$ ) (Table 4.1). Players who participated in formal pitching lessons did show an increased frequency of shoulder and elbow injuries compared to those who did not ( $P = 0.04$ ) (Table 4.1). Youth athletes who demonstrated this early position specialization as a pitcher were found to be 2.8 times as likely (95% CI 1.1, 7.3;  $P = 0.04$ ) to experience a shoulder or elbow injury when compared to athletes who did not specialize early as a pitcher (Table 4.1). Participation in generalized strength and conditioning programs did not impact injury frequencies in youth baseball players when compared to the athletes who participated solely in baseball-specific training ( $P = 0.60$ ) (Table 4.1).

#### 4.5 Discussion

The most important result of our study shows that youth baseball players who specialize in baseball displayed greater shoulder and elbow injury risk compared to those who did not specialize. Interestingly, youth players who also participated in additional specialty training, particularly as a pitcher, were at the greatest risk for sustaining a shoulder or elbow injury in this cohort. This is concerning given that two-thirds of youth baseball players identified as pitchers and reported participating as position players when

not pitching. These findings suggest that not only is specialization an issue but that physical overtraining may also contribute to the observed disparities in injury rates.

#### 4.5.a Injury Incidence

During the 6-month study period, youth baseball players demonstrated an UE injury rate of 16.3/1000 AEs, markedly higher than the injury rates previously reported in high school (4.0/1000 AEs) and collegiate players (5.83/1000 AEs).<sup>36,137</sup> The higher injury rate may be, in part, due to differences in the injury definitions between studies. Previous research examining injury profiles in youth baseball players relied on self-report measures from survey data to establish risk factors associated with UE pain in this population.<sup>90,91,116,162</sup> This research indicated that nearly 50% of youth athletes reported experiencing shoulder or elbow pain during the course of a baseball season, however a licensed medical professional did not physically confirm these reports as was done in our study.<sup>90,91,116,162</sup> Prospectively examining UE injury profiles has allowed us to build on the knowledge gained from previous studies and generate a more complete picture of UE injury development in youth baseball.

In contrast to previous studies on high school, collegiate and professional pitchers, the majority of youth baseball players pitch in some capacity and play additional field positions when not pitching.<sup>31,36,137</sup> Based on discussions with Little League and competitive travel team coaches, the majority of youth baseball teams that participated in this study included a mean of 12 players per roster with 8 or more competing as pitchers throughout the season. This is likely a result of the USA Baseball pitch count limitations, which require teams to distribute the physical demands of pitching across multiple players in an attempt to limit overuse.<sup>32</sup> Based on data collected from team coaches,

teams typically play 2-4 tournaments per month with 4-5 games per tournament during competitive seasons. Each youth baseball game lasts 4-6 innings with pitch counts that may exceed 100 pitches per game.<sup>86</sup> Depending on the number of batters faced and the number of innings played, a minimum of 6 pitchers is required to participate in 1 youth baseball tournament. The pitch count recommendations stem from previous research that linked excessive pitch counts and improper throwing mechanics to shoulder and elbow pain in a cohort of youth baseball pitchers.<sup>53,90,91</sup>

This study was the first to examine UE injuries in a cohort of youth baseball players, irrespective of position. While pitchers sustained the majority of the injuries reported, injury proportions were equal between positions, as pitchers also comprised the majority of the sample. We found no differences in absolute risk for developing a shoulder and elbow injury in youth pitchers when compared to their position player counterparts (Table 4.1). These findings were in contrast to previous research performed in the high school, collegiate and professional ranks which indicated that pitchers were at a significantly higher risk for sustaining an UE injury when compared to position players.<sup>36,123,137</sup>

#### 4.5.b Early Sport Specialization

Specialization in a single sport, prior to the onset of adolescence, has been repeatedly identified as a risk factor for injury across multiple sports.<sup>39,71,72,84,99,107,108</sup> Current research also suggests early sport specialization does not correlate with an athlete's long-term success in sport and that early diversification may be more beneficial to their physical development.<sup>84</sup> USA Baseball recently released a Long Term Athlete Development Model which advises athletes to avoid specializing in a single sport prior to

14 years of age.<sup>109</sup> Despite these widespread recommendations, youth athletes continue to specialize in a single sport at alarming rates. Based on well-established research criteria, 83% of the youth baseball players in this study were classified as specialized athletes (Table 4.1).<sup>71,72</sup> Our results showed that sport specialization in baseball significantly increased an athlete's absolute risk for sustaining a shoulder or elbow injury when compared to sport diversification (Table 4.1). These findings are consistent with previous research and support the recommendations put forth by USA Baseball in their Long Term Athlete Development Model.<sup>72,99,109</sup>

One potential reason for the continuation of sport specialization in youth athletes, despite the acknowledged risks associated with it, is a lack of understanding of its definition. Sport specialization has been defined in the literature as 'intense training for >8 months per year in a single sport to the exclusion of other sports.'<sup>71,84</sup> At the start of our study, athletes, typically with the help of parents and coaches, were asked to self-classify as either specialized or multi-sport athletes. Approximately 70% of the players classified themselves as multi-sport athletes based on their own perceptions (Figure 4.1). Based on previous research, 83% of this youth cohort were classified and competed as specialized athletes suggesting a significant discrepancy in athlete (and parental) perception of what constituted sport specialization (Figure 4.1). When asked to expand upon their views on sport specialization, parents and coaches stated that they encouraged their athletes to participate in multiple sports in addition to competing in year-round baseball activities. These attitudes seem to contribute to overscheduling with little time for rest and recovery and may prove detrimental to an athlete's health and long-term success in sport.<sup>84</sup>

#### 4.5.c Early Position Specialization

The effects of early specialty training, or position specialization, were examined in this study. Athletes who took formal hitting lessons or any variation of baseball-specific training, prior to the onset of adolescence, did not demonstrate an increased risk for injury when compared to those who did not (Table 4.1). Athletes who participated in formal pitching lessons however, did demonstrate a greater absolute risk for sustaining a shoulder or elbow injury in baseball when compared to those who did not (Table 4.1). With the advent of age-based pitch count restrictions, youth teams are required to carry more pitchers than their high school counterparts. In this cohort, the majority of youth pitchers sought out formal pitching lessons (67.9%) in an effort to improve athletic performance. Specialty training as a pitcher, in addition to pitching in practices and games, may result in increased physical loads being placed across an athlete's shoulder and elbow joints. Participating in formal pitching lessons, prior to the onset of adolescence, may derail the original purpose of age-based pitch count restrictions, which was to reduce physical loads across a growing athlete's body. Also, per USA Baseball and their Long Term Athlete Development Model, specialty training may not be as beneficial as generalized physical training for youth athletes at this stage of physical development.<sup>71,84,107,108</sup> Despite age-specific recommendations that emphasize the importance of physical literacy and functional skill acquisition, remarkably few youth baseball players took generalized strength and conditioning lessons in this cohort (22.6%). Future authors should examine the impact of participation in generalized physical training programs over specialty training programs on injury risk and performance in youth athletes.

#### 4.5.d Strengths

To our knowledge, this is the first study to report UE injury incidence in a cohort of youth baseball players. This research is novel with respect to both the study population and its prospective design. Less than 10% of all injury risk and prevention data has been collected in athletes <18 years old, with even less data describing youth athletes (9-12 years old).<sup>98</sup> Despite the fact that youth and adolescent athletes comprise the majority of the population competing in sports, they continue to be significantly understudied.<sup>85</sup> The prospective design allowed UE injury incidence in a cohort of youth baseball players to be established. In addition, the study was strengthened because there was confirmation of all injuries reported in this study via physical examination as opposed to self-report measures. This study provides insight into the unique injury profiles and actual participation levels of youth baseball players compared to well-studied collegiate and professional baseball players.

#### 4.5.e Limitations

Certain limitations should be noted while interpreting the results of this research. Previous epidemiological studies have typically included injury rates for all injuries across an athlete's body while this study reported injury rates solely at the shoulder and elbow.<sup>31,43,124,137</sup> This decision was based on previous data that shows the shoulder and elbow to be two of the most commonly injured body parts in competitive baseball players.<sup>80,124,137</sup> Additional limitations in this study include the obvious disparities in the number of youth pitchers versus position players reported as well as the number of specialized athletes versus multi-sport athletes reported in this study. While the groups were uneven for statistical analysis, the proportions for each position and participation

level were representative of the population being studied. Lastly, to capture an adequately sized sample of youth baseball players, baseline evaluations required a 10-week period to complete throughout the competitive season. Athletes were then tracked for 6 months following individual baseline evaluations. This resulted in varied baseball participation rates among youth athletes that were likely impacted by competition level (i.e. little league baseball vs. tournament team baseball) and participation in additional sport seasons (i.e. football season, basketball season). The variability in baseball participation over a 6-month calendar period may have influenced the consistency of our athletic exposure data however was representative of youth athlete participation rates in baseball.

#### **4.6 Conclusion**

Youth baseball players demonstrated higher UE injury incidence rates than previously reported in adolescent baseball populations. Sport and position specialization, prior to the onset of adolescence, increases an athlete's absolute risk for developing an UE injury during youth baseball. USA Baseball's age-based pitch count limitations were designed to decrease overuse injuries at the shoulder and elbow by requiring more athletes to pitch. This may have inadvertently had the opposite effect by increasing the rate of specialty training outside of competition.




Figure 4.1 Sport Specialization – Perception vs. Reality



**Table 4.1. Analysis of Risk Factors for Upper Extremity Injuries**

<b>Variable</b>	<b><i>n</i> (% of total cohort)</b>	<b><i>Odds Ratio</i> (95% CI)</b>	<b><i>Absolute Risk</i> (%)</b>	<b><i>Absolute Risk Difference<sup>b</sup></i> (%) (95% CI)</b>	<b><i>P</i> value</b>
<b>Position</b>					
- Pitchers	106 (66.7)	1.0 (0.4, 2.7)	13.2	0.0	1.00
- Position Players Only	53 (33.3)	Referent	13.2	(-11.1, 11.1)	
<b>Sport Specialization</b>					
- Specialized Athlete	132 (83.0)	1.2 (1.1, 1.3)	15.9	15.9	<b>0.03*</b>
- Multi-Sport Athlete	27 (17.0)	Referent	0.0	(9.7, 22.1)	
<b>Position-Specific Training</b>					
- Hitting Lessons	111 (69.8)	1.1 (0.4, 3.0)	13.5	1.0	0.86
- No Hitting Lessons	48 (30.2)	Referent	12.5	(-10.3, 12.2)	
- Pitching Lessons	72 (45.3)	2.8 (1.1, 7.3)	19.4	11.3	<b>0.04*</b>
- No Pitching Lessons	87 (54.7)	Referent	8.1	(6.2, 22.2)	
- Baseball-Specific Training <sup>a</sup>	118 (74.2)	2.3 (0.6, 8.2)	15.2	7.9	0.20
- No Baseball-Specific Training <sup>a</sup>	41 (25.8)	Referent	7.3	(-2.3, 18.2)	
<b>General Physical Training</b>					
- Strength & Conditioning Lessons	36 (22.6)	1.4 (0.5, 4.0)	16.7	4.5	0.49
- No Strength & Conditioning Lessons	123 (77.4)	Referent	13.9	(-9.0, 17.9)	

\*Indicates a statistically significant difference ( $P < 0.05$ ).

<sup>a</sup> Baseball-Specific Training was comprised of players who took only hitting lessons ( $n = 46$ ), only pitching lessons ( $n = 7$ ) or a combination of both ( $n = 65$ ).

<sup>b</sup> Absolute Risk Difference = Absolute Risk of Non-Exposed Group – Absolute Risk of Exposed Group.

## CHAPTER 5

### NORMALIZATION METHODS IN ISOMETRIC SHOULDER STRENGTH IN YOUTH BASEBALL PLAYERS: A COMPARISON ACROSS 5 METHODS

#### 5.1 Abstract

**Introduction:** The measurement of shoulder muscle strength is an important component in the physical assessment of overhead athletes. Although several measures have been described, isometric testing using hand held dynamometry (HHD) has proven to be a reliable, low cost and portable method in this population. The use of this procedure in youth athletes (ages 9-12 years) is challenging because of the wide variations observed in strength testing performance. These variations may result from substantive differences in anthropometric characteristics such as height and weight. Considering this, ‘normalized’ strength measures that account for an individual’s current body size may be of great use in understanding the relationship between shoulder strength and athletic performance in youth baseball players.

**Objective:** The purposes of this study were to 1) compare the measurement properties of 5 potential methods for normalizing isometric shoulder muscle strength and 2) examine the relationship between normalized isometric shoulder muscle strength and ball velocity in a cohort of 9-12 year old male baseball players.

**Methods:** One hundred and fifty nine male youth baseball players (mean age  $11.1 \pm 1.1$  years) volunteered for this study. Baseline and follow up height, weight and ulnar length

measurements were assessed followed by isometric strength in both the dominant and non-dominant shoulders. Ball velocity was assessed as a measure of throwing performance. Intraclass correlation coefficients (ICC), standard errors of measurement (SEM) and minimal detectable change (MDC) were calculated for all baseline and follow-up strength measures. Repeated measures analyses of variance (ANOVA) were conducted to compare changes in isometric shoulder strength at 2 time points after normalizing to 5 separate measures of body size. Linear regression models were used to examine the relationships between normalized isometric shoulder torque measures and ball velocity. Statistical significance was set a priori at  $\alpha=0.05$ .

Results: Torque, defined as the measure of shoulder strength divided by the corresponding ulnar length, was the only method that demonstrated excellent reliability (ICC<sub>2,1</sub> 0.98-0.99) and detected significant changes between shoulder strength in each of the 4 measures tested (SEM 0.39-0.69 Nm). Modest but significant correlations were observed between scaption torque and ball velocity ( $r^2 = 0.27, P < 0.001$ ) and external rotation (ER) torque at 0° and ball velocity ( $r^2 = 0.23, P < 0.001$ ).

Conclusion: The normalization method that demonstrated the most consistent measurement properties for the assessment of isometric shoulder strength in a youth baseball player was torque. Ulnar length is the most stable and reliable anthropometric measure evaluated in this study. Once normalized, isometric shoulder scaption strength was the most significant predictor of ball velocity, followed by ER strength at 0° in 9-12 year old baseball players.

Keywords: torque; youth; baseball; normalization; ball velocity

## 5.2 Introduction

Upper extremity muscle strength is an important component in the assessment of athletic performance and injury prevention in baseball players.<sup>17,28,44,50,52,61,97,105,148,156,160</sup>

Strength is defined as the amount of force a muscle can maximally produce during a single repetition.<sup>67,68,70</sup> Clinicians and researchers routinely use a battery of strength measures in performance assessments, injury diagnostics and return to sport decisions following injury.<sup>17,33,61,148</sup> While upper extremity strength measures have been widely reported at the collegiate and professional levels, little to no evidence is available describing these measures at the youth and adolescent levels.<sup>17,28,33,44,50,61,97,105,148,156</sup>

Establishing an objective and reliable method for evaluating strength is imperative in understanding shoulder function and injury risk in youth baseball players.<sup>33,148</sup>

A variety of methods, including isokinetic, isometric and functional testing, have been used to measure shoulder strength in athletic populations.<sup>17,28,44,61,97,148,160</sup> While isokinetic testing is considered the gold standard in strength assessment, the high equipment costs and lack of portability make it impractical for use outside of laboratory settings.<sup>33,52,142</sup> Isometric testing using hand held dynamometry (HHD) has proven to be a reliable alternative to isokinetics in assessing strength at the shoulder.<sup>17,33,142,148</sup> Hand held dynamometry is low cost, portable and easy to use however it does have acknowledged limitations including investigator strength, lack of stabilization and inconsistencies in testing procedures.<sup>33,52</sup>

The majority of isometric strength testing has been conducted in collegiate and professional baseball players with few studies examining shoulder strength in younger players.<sup>17,44,61,97,105,148</sup> Despite acknowledging that anthropometric measurements, such as

height and weight, influence the body's ability to produce force and thereby muscle strength, normalization methods accounting for body size are inconsistently reported in the literature.<sup>44,67,68,97,105</sup> The evaluation of isometric strength in youth and adolescent athletes is inherently different from that of collegiate and professional athletes.<sup>67,68,92</sup> Height, weight and neuromuscular control can fluctuate frequently in physically developing populations with the potential to rapidly change over short periods of time. Performance assessments that rely solely upon absolute measures, without normalization, may lack the ability to discern changes in muscle strength from changes in body size in youth populations.<sup>67,68,70</sup> Accounting for these alterations in growth and development through normalization is critical to accurately assessing muscle function and injury risk in young athletes.<sup>67,68,92</sup>

There is a notable gap in the literature surrounding the evaluation and normalization of shoulder strength in youth baseball players. Research suggests normalization methods, which include body mass, body mass index (BMI), height, torque and percent of non-dominant shoulder strength described by Trakis<sup>148</sup> may be potential methods for assessing muscle strength and changes in muscle strength over time in this population.<sup>63,67,68</sup> The purposes of this study were to 1) compare the measurement properties of 5 potential methods for normalizing isometric shoulder strength in a cohort of 9-12 year old male baseball players and 2) examine the relationship between normalized isometric shoulder strength and ball velocity in a cohort of 9-12 year old male baseball players.

## 5.3 Methods

### 5.3.a Study Population

One hundred and fifty nine competitive male youth baseball players with a mean age of  $11.1 \pm 1.1$  years volunteered to participate in this study (Table 5.1). All players in this study were recruited from local baseball clubs, baseball tournaments and little leagues in the Upstate Region of South Carolina. All players were male, between the ages of 9-12 years and uninjured at the time of initial examination. Players were excluded from the study if they (1) reported any injuries that currently restricted their ability to participate in baseball activities or (2) reported a shoulder or elbow injury that required medical attention during the 3 months prior to initial examination. The University of South Carolina's Institutional Review Board (IRB) approved this study. Parental consent and athlete assent were obtained for each participant enrolled.

### 5.3.b Instruments

Height, weight and ulnar length were measured with using a portable stadiometer, digital weight scale and body tape measure, respectively. Athletes were asked to remove their footwear for anthropometric measurements. Height and ulnar length were recorded to the nearest 0.5 centimeter (cm) while weight was recorded to the nearest 0.1 kilogram (kg). Isometric shoulder strength was measured using a Lafayette Manual Muscle Tester hand-held dynamometer (Lafayette Instrument Company, Lafayette, IN, USA). All isometric strength measurements were performed by the lead researcher who demonstrated excellent intra-rater reliability prior to initial data collection ( $ICC_{2,1} = 0.94-0.99$ ). Ball velocity was assessed using a Stalker Sport Radar Gun (Stalker Radar, Richardson, TX, USA).

### 5.3.c Procedures

At the time of study enrollment, baseline height, weight and ulnar length measurements were assessed for each participant followed by isometric shoulder strength in both the dominant and non-dominant arms (Table 5.2). Two values of each strength measure were recorded per arm and averaged for statistical analysis. Isometric shoulder strength was assessed bilaterally using a Lafayette Manual Muscle Tester hand-held dynamometer and methods previously reported in the literature.<sup>17</sup> Isometric shoulder strength measures included abduction in the scapular plane (scaption) at 90°, external rotation (ER) at 0°, ER at 90° and internal rotation (IR) at 90° for the dominant and non-dominant arms (Table 5.2). Make tests were used for each isometric strength measure based on higher reliability when compared to break tests in hand-held dynamometers.<sup>146</sup> Scaption and ER at 0° forces were measured in the seated position (Figure 5.1). Scaption was measured with the dynamometer placed 5 cm distal to the cubital fossa while ER at 0° was measured with the dynamometer placed on the dorsal aspect of the forearm, 2 cm proximal to ulnar styloid process (Figure 5.2). External rotation at 90° and IR at 90° was measured with the shoulder in 90° of abduction, 90° of ER and 90° of elbow flexion (Figure 5.2). The dynamometer was placed on the dorsal aspect of the forearm, 2 cm proximal to ulnar styloid process. Internal rotation at 90° was assessed in a similar fashion to ER at 90°, however the shoulder was in a state of neutral rotation and the dynamometer was placed on the volar aspect of the forearm (Figure 5.2). Each participant was asked to provide maximal effort throughout each trial during examination.<sup>17,46</sup> Isometric shoulder strength was then normalized prior to statistical analysis using 5

separate methods: body mass, body mass index (BMI), height, torque and the Trakis Method (Table 5.3).<sup>148</sup>

Height, weight, ulnar length and isometric shoulder strength measurements were re-assessed in a subset of participants (n = 58) to examine changes in body size and strength over the 6-month period (Table 5.2). Isometric shoulder strength was again normalized for statistical analysis using the 5 previously stated methods: body mass, BMI, height, torque and the Trakis Method (Table 5.3).<sup>148</sup>

Throwing performance was assessed using ball velocity during an overhead throw. This measure was assessed in a subset of participants (n = 80). Following a warm up period during team practice, participants were asked to throw 3 balls from a distance of 46 feet on flat ground to a specified target. A Stalker Sport Radar Gun (Stalker Radar, Richardson, TX, USA) was used to record the velocity of each throw in miles per hour (mph). The 3 throws were recorded and averaged for statistical analysis.

#### 5.3.d Statistical Analysis

Means and standard deviations (SD) were calculated across all participants for the dependent variables: height, weight, BMI and normalized shoulder strength measures. Reliability was assessed for all baseline and follow-up strength measures using intraclass correlation coefficients (ICC) with corresponding 95% confidence intervals (CI).<sup>122</sup> Standard errors of measurement (SEM) were also calculated to determine the absolute reliability of each strength measure using the largest SD in the formula  $SD \times \sqrt{1 - ICC}$ .<sup>122</sup> Individual SEMs were then used to calculate corresponding minimal detectable change (MDC) values for each of the normalized strength measures using the formula  $SEM \times 1.96 \times \sqrt{2}$ .<sup>122</sup> Repeated measures analyses of variance (ANOVA) were conducted



to compare changes in isometric shoulder strength at 2 time points (baseline and follow-up) after co-varying for physical growth and body size. Effect sizes were calculated to identify the magnitude of change detected between the 2 time points for each of the normalized strength measures. Linear regression models were used to examine the relationships between the normalized isometric shoulder strength measures and ball velocity in youth baseball players. The method with the most consistent measurement properties for normalizing isometric shoulder strength in youth baseball players was determined based on each measure's test-retest reliability, ability to detect changes over time and strength of association with ball velocity. Statistical significance was set a priori at  $\alpha=0.05$ . All statistical analyses were performed using SPSS Statistics 21.0 (SPSS Inc., Chicago, IL, USA) software.

## 5.4 Results

### 5.4.a Normalization Methods Reliability

Baseline anthropometric characteristics of youth baseball players are reported in Table 5.1. Reliability data for the torque normalization method are reported in Table 5.4. The intra-rater reliability for the remaining 4 normalization methods were: body mass  $ICC_{2,1}$  0.97-0.98, BMI  $ICC_{2,1}$  0.95-0.98, height  $ICC_{2,1}$  0.94-0.98 and Trakis  $ICC_{2,1}$  0.80-0.98. Their respective SEM values were: body mass 0.46-0.63%, BMI 0.95-1.16 kg/m<sup>2</sup>, height 0.12-0.28 kg/m and Trakis 4.15-15.00% (Figure 5.3). High inter-participant variability was apparent in each of the strength normalization methods, with the exception of torque, which suggests a lack of stability in these measures in test-retest situations (Figure 5.3). Torque was the only method to demonstrate good-to-excellent

reliability and detect significant changes in shoulder strength over time in each of the 4 measures tested based on corresponding  $MDC_{95}$  values (Table 5.4).

#### 5.4.b Shoulder Torque and Ball Velocity

The relationship between normalized isometric shoulder torque and ball velocity was examined using stepwise linear regression models with forward selection to determine the impact of each measure. Four normalized shoulder strength measures were entered into the model. Scaption torque demonstrated a high correlation with ball velocity and was entered into the model first followed by ER at 0° torque, ER at 90° torque and IR at 90° torque. A significant relationship was observed between scaption torque and ball velocity ( $r^2 = 0.27$ ,  $P < 0.001$ ) (Figure 5.4). The remaining measures demonstrated non-significant relationships with ball velocity when scaption was entered first into the model (ER at 0° torque  $r^2 = 0.27$ ,  $P = 0.59$ ; ER at 90° torque  $r^2 = 0.28$ ,  $P = 0.69$ ). Internal rotation torque was completely removed from the model. A second model was run with ER at 0° torque entered first followed by the 3 remaining measures. Significant relationships were observed between ER at 0° torque and ball velocity ( $r^2 = 0.23$ ,  $P < 0.001$ ) and scaption torque and ball velocity ( $r^2 = 0.23$ ,  $P = 0.04$ ) in this model (Figure 5.4). The remaining measures demonstrated non-significant relationships with ball velocity when ER at 0° torque was entered first into the model (IR at 90° torque  $r^2 = 0.28$ ,  $P = 0.30$ ). External rotation torque at 90° was completely removed from the model. Tests for collinearity indicated that a high level of collinearity was present between scaption torque and ER torque at 0° measures (*Eigenvalue* = 1.96) in the regression models. Based on these findings, scaption torque alone predicts 27% of ball velocity in

youth baseball players while ER torque at 0° predicts 23% of ball velocity when measured in these athletes.

## 5.5 Discussion

This study investigated five distinct methods for normalizing isometric shoulder strength to determine which had the best measurement properties for youth baseball players. The torque method demonstrated excellent intra-rater reliability, with the lowest reported SEM and MDC values of any method examined. Once normalized, dominant shoulder scaption torque was the most predictive measure with respect to ball velocity, followed by dominant shoulder ER torque at 0°. Dominant shoulder scaption torque alone predicted 27% of the variation in ball velocity in a cohort of youth baseball players. Results also indicated that dominant shoulder ER torque at 0° was strong measure and could be used independently of dominant shoulder scaption torque to predict 23% of the variation in ball velocity in this cohort.

### 5.5.a Normalization Method Types

Few original research studies have employed normalization methods when examining muscle strength measures and none have compared findings to determine the most appropriate method based on a specific population.<sup>132,148</sup> Previous literature impresses the importance of normalizing strength measures for accurate comparison across multiple time points, particularly in longitudinal and repeated measures study designs.<sup>67,68</sup> In the absence of normalization, any observed changes in muscle strength may be misinterpreted as simply functions of growth and physical development as opposed to definitive changes in the measures themselves.<sup>67,68</sup> A study by Trakis<sup>148</sup> using isometric testing and HHD referenced dominant shoulder strength measures to non-

dominant shoulder strength measures as a means of normalizing muscle strength in adolescent baseball players. The theory was based on the concept of using non-dominant shoulder strength values as internal reference points for each athlete.<sup>148</sup> When the Trakis method<sup>148</sup> was applied in younger players, ages 9-12 years old, excessive inter-participant variability was noted both in single session measures and repeated measures over time. Neuromuscular control patterns in youth athletes are not as well developed as their adolescent and adult counterparts, which may impact their ability to reproduce consistent results with isometric muscle strength measures.<sup>54,62,63,84</sup>

Biomechanical studies have suggested the use of body mass and derivations of body mass, including BMI, as potential normalizing factors though few studies have formally tested those theories on youth and adolescent athlete populations.<sup>70,132</sup> Frequent fluctuations in body mass measures imply that, while the weight of a youth athlete certainly contributes to their ability to produce muscle force, is may not possess the stability required to accurately detect changes in muscle strength measures over time.<sup>68,70</sup> Height and torque, a derivation of height, appeared to be more stable choices for normalization factors as the measures only increase over time in youth athlete populations. When the normalization methods containing height and torque were compared, the limb-specific torque method outperformed the more generalized height method in test-retest reliability as well as internal consistency as evidenced by lower SEM and MDC values. These findings indicate that torque was the most consistent normalization method for assessing isometric muscle strength at the shoulder in youth baseball players.

### 5.5.b Shoulder Torque and Ball Velocity

Dominant shoulder scaption strength and ER strength at 0° have been previously linked to throwing performance and upper extremity injury risk in baseball players.<sup>17,44,61,97,105,148</sup> The majority of studies examining upper extremity strength as a factor for performance and injury risk were performed in collegiate and professional athletes.<sup>44,50,97,105</sup> Studies have found little conclusive evidence supporting the theory of a direct relationship between baseline shoulder muscle strength and ball velocity in throwers, however shoulder muscle weakness has been repeatedly linked to injury throughout the baseball literature.<sup>17,28,97,105,133</sup> Magnusson<sup>97</sup>, Mullaney<sup>105</sup> and Byram<sup>17</sup> have all shown that pitchers demonstrated deficits in dominant shoulder scaption and ER muscle strength. While we did not examine the effects of normalized dominant shoulder scaption torque on injury risk in this study, our results do support the significance of measuring dominant shoulder scaption torque with regards to performance measures in competitive baseball players.

To our knowledge, this is the first study to show a positive association between normalized isometric shoulder torque and ball velocity, an acknowledged performance measure in youth baseball players.<sup>89,90,116</sup> Minimal data exists examining upper extremity muscle strength and injury risk in youth athletes. One study exists by Harada<sup>61</sup> compared a battery of shoulder muscle strength measures and injury risk in a cohort of Japanese youth baseball players. They found that injured athletes demonstrated greater dominant shoulder strength compared to uninjured athletes.<sup>61</sup> These findings suggest that the connection between shoulder strength and upper extremity injury risk may be different in

youth athletes when compared to their collegiate and professional counterparts, however further studies are needed to better understand these relationships.<sup>17,61,97,105</sup>

#### 5.5.c Limitations

Isometric muscle strength testing using hand held dynamometry has several clinical advantages such as low cost, portability and ease of use however it also has acknowledged limitations.<sup>33,52</sup> While extreme effort was expended to standardize all measurements and testing procedures including using a single investigator with excellent intra-rater reliability, no external stabilization methods were applied to the athletes during the assessments. This decision was based on feasibility and applicability in clinical settings however may have influenced our results.

Another potential limitation of this study was the collinearity between the isometric shoulder scaption torque and ER torque at 0° measures in this population. Our results indicate that either measure is predictive of ball velocity however further research is needed to determine what additional variables should be included in the model to best explain this performance measure in youth athletes. Overhead throwing is a complex motor skill that requires coordination and the proper sequencing of a series of linked movements that start in the lower extremities and ultimately culminate in ball release.<sup>157</sup> This statement supports our findings that isometric shoulder strength explains only a portion of the variability observed in ball velocity in young throwers.

Lastly, the high levels of inter-participant variability observed in isometric shoulder ER and IR torque at 90° may have negatively influenced the predictability of these measures in youth athletes. The variability may be attributable to age-appropriate deficits in neuromuscular control in the prone overhead position however further study is

needed to better understand the role neuromuscular control plays in youth baseball throwing mechanics. Future studies should consider the use of isometric shoulder torque measures, not only in performance assessments of youth athletes, but in injury prevention programs as well. The relationship between upper extremity injury risk and isometric shoulder strength is not well understood in youth populations suggesting that further study is warranted.

## 5.6 Conclusion

The normalization method with the best measurement properties for assessing isometric shoulder strength in youth baseball players was torque. Ulnar length is the most stable and reliable anthropometric measure evaluated in this study. Once normalized, isometric shoulder scaption strength was the most significant predictor of ball velocity, followed by ER strength at 0° in 9-12 year old baseball players. Muscle strength assessments performed in 90° of shoulder abduction demonstrated high inter-subject variability and provided minimal information concerning the shoulder function and athletic performance of youth baseball players.

**Table 5.1. Baseline Characteristics of Youth Baseball Players**

	<b>N</b>	<b>Mean</b>	<b>SD<sup>a</sup></b>
<b>Age, years</b>	159	11.1	1.1
<b>Height, cm</b>	159	146.8	8.3
<b>Weight, kg</b>	159	41.6	10.1
<b>BMI<sup>b</sup>, kg/m<sup>2</sup></b>	159	19.1	3.4
<b>Arm Dominance Right, %</b>	137	86.2%	-

<sup>a</sup>SD, standard deviation.

<sup>b</sup>BMI, body mass index.

**Table 5.2. Overview of Anthropometric & Isometric Shoulder Strength Measures**

Type of Measure	Testing Equipment	Body Position	Arm Position	Examiner Position	
<b>Anthropometric</b>					
1. Height	Stadiometer	Standing	-	Standing	
2. Weight	Digital Scale		-		
3. Ulnar Length	Tape Measure	Supine	-		
<b>Isometric Strength</b>					
4. Scaption	Hand Held Dynamometer	Sitting	Shoulder Abducted 90°	Standing Anterior to Testing Arm	
5. External Rotation at 0°			Horizontally Adducted 45°		Rotation Neutral
6. External Rotation at 90°	Hand Held Dynamometer	Prone	Shoulder Abducted 0°	Standing Perpendicular to Dorsal Forearm	
7. Internal Rotation at 90°			Elbow Flexed 90°		Forearm Rotation Neutral
			Shoulder Abducted 90°		Externally Rotated 90°
			Elbow Flexed 90°	Kneeling Inferior to Testing Arm	
			Shoulder Abducted 90°		
			Rotation Neutral		
			Elbow Flexed 90°		



**Table 5.3. Normalization Methods for Isometric Shoulder Strength**

Method	Calculation
<b>Body Mass (kg)</b>	= $\frac{\text{Shoulder Strength Measure}}{\text{Body Mass}}$
<b>BMI (kg/m<sup>2</sup>)</b>	= $\frac{\text{Shoulder Strength Measure}}{\text{BMI}}$
<b>Height (m)</b>	= $\frac{\text{Shoulder Strength Measure}}{\text{Height (m)}}$
<b>Muscle torque (Nm)</b>	= $\frac{\text{Shoulder Strength Measure (N)}}{\text{Ulnar Length (m)}}$
<b>Percent Non-Dominant Strength Method (%)<sup>148</sup></b>	= $\frac{(\text{Dominant Shoulder Strength} - \text{Non-Dominant Shoulder Strength})}{\text{Non-Dominant Shoulder Strength}} \times 100$

**Table 5.4. Reliability of Normalized Dominant Isometric Shoulder Strength using Torque Method**

	Mean	SD <sup>a</sup>	ICC <sub>2,1</sub> (95% CI) <sup>b</sup>	SEM <sup>c</sup>	MDC <sub>95</sub> <sup>d</sup>	Effect Size <sup>e</sup>
<b>Baseline Strength (n=159)</b>						
- Scaption	17.9	5.2	0.99 (0.94, 0.99)	0.57	1.6	-
- External Rotation at 0°	15.2	4.1	0.99 (0.99, 1.00)	0.52	1.4	-
- External Rotation at 90°	11.9	3.0	0.98 (0.93, 0.99)	0.39	1.1	-
- Internal Rotation at 90°	16.3	4.0	0.99 (0.95, 0.99)	0.45	1.3	-
<b>Follow-Up Strength (n=58)</b>						
- Scaption	20.7	6.2	0.99 (0.98, 0.99)	0.69	1.9	-
- External Rotation at 0°	19.0	5.5	0.99 (0.97, 0.99)	0.54	1.5	-
- External Rotation at 90°	14.1	3.3	0.98 (0.98, 0.99)	0.46	1.3	-
- Internal Rotation at 90°	19.0	4.5	0.99 (0.96, 0.99)	0.55	1.5	-
<b>Strength Change over Time (n=58)</b>						
- Scaption	2.8	3.4	-	-	-	0.41
- External Rotation at 0°	3.8	3.6	-	-	-	0.54
- External Rotation at 90°	2.2	2.2	-	-	-	0.51
- Internal Rotation at 90°	2.7	2.7	-	-	-	0.51

<sup>a</sup>SD, standard deviation.

<sup>b</sup>ICC<sub>2,1</sub> (95% CI), intraclass correlation coefficient with 95% confidence interval.

<sup>c</sup>SEM, standard error of the mean.

<sup>d</sup>MDC<sub>95</sub>, minimal detectable change.

<sup>e</sup>Effect Size, calculated using Partial Eta squared statistics from ANOVA analysis.



Figure 5.1 Examination of Isometric (A) Shoulder Scaption and (B) Shoulder External Rotation at 0° Strength in Seated Position



Figure 5.2 Examination of Isometric (A) Shoulder External Rotation at 90° and (B) Shoulder Internal Rotation at 90° Strength in Prone Position




Figure 5.3 Changes in Normalized Dominant Shoulder Strength by Method over 6 Month Period




Figure 5.4 Relationship Between Normalized Dominant Shoulder Scaption Torque, External Rotation at  $0^\circ$  Torque and Ball Velocity

## CHAPTER 6

### DISCUSSION

Youth and adolescent baseball players comprise the majority of the 13-17 million athletes that participate in this sport in the U.S. each year.<sup>4,31,51,143</sup> The benefits of sports participation are well documented throughout the literature, however risks are also associated with these activities.<sup>18,21,26,38,71,72</sup> Despite increased awareness surrounding the nature of these risks, youth and adolescent athletes are reporting baseball-related overuse injuries at alarming rates.<sup>12,19,90,91,136,140,148,154,155</sup> The most common overuse injuries reported in baseball players are at the shoulder and elbow however there is a significant lack of epidemiologic data establishing the magnitude of upper extremity overuse injuries, particularly at the youth level.<sup>31,137</sup>

Determining the extent of the problem is principal in unraveling UE injury risk in youth baseball players.<sup>151</sup> The next step is to understand the etiology surrounding UE injury development in youth athletes by identifying population-specific risk factors, such as player position, sport specialization and participation in specialty training. Research on athlete-dependent risk factors is also needed in youth athletes, as <10% of studies include participants <18 years old.<sup>98</sup>

Evaluating physical measures in youth and adolescent athletes is inherently different when compared to collegiate and professional athletes.<sup>67,68,92</sup> This is especially true with regards to strength and performance assessments.<sup>67,68,92</sup> Anthropometric measures have been repeatedly linked to the body's ability to produce force and therefore

muscle strength.<sup>67-69</sup> As height and weight fluctuate more frequently, and much more rapidly, in youth and adolescence versus adulthood, these measures become integral to the assessment of strength and performance in physically developing athletes.<sup>70</sup> Independent, or absolute, measures that do not take into account the current anthropometric measures of an athlete may be appropriate for use in the collegiate and professional ranks who typically demonstrate minimal changes in these measures over time.<sup>68,70</sup> Relying on assessment data gleaned solely from absolute measures in youth and adolescent athletes is less appropriate as they lack the ability to discern changes in physical measures from changes in body size which commonly occur in physically developing populations.<sup>67,68,70</sup> Accounting for these alterations in growth and development through anthropometric normalization is critical to accurately assessing athletic performance and injury risk in young athletes.<sup>67,68,92</sup> This led to the current study, which established UE injury incidence, examined the effects of population-specific risk factors and identified the most consistent method for normalizing isometric shoulder strength in a cohort of youth baseball players.

### **6.1 Upper Extremity Injuries in Youth Baseball Players**

Youth baseball players demonstrated an UE injury rate of 16.3/1000 AEs, markedly higher than the injury rates previously reported in high school (4.0/1000 AEs) and collegiate players (5.83/1000 AEs).<sup>36,137</sup> The higher injury rate may be, in part, due to differences in the types of injuries included in each of the studies. Previous research focused on examining time-loss injuries however, based on the target population of this study, both time-loss and non-time-loss injuries were included.<sup>90,91,116,162</sup> Another factor that may have contributed to a higher UE injury rate in youth baseball players was the

decreased number of athletic exposures recorded at this level of competition. Based on similar studies performed at the high school and collegiate levels, injury incidence was calculated using team activities (i.e. practices and games).<sup>36,137</sup> This study found that youth baseball players, while in season, typically attended team practices twice a week in addition to the varied number of games and tournaments played by each team. A novel finding within this study population was that the majority of athletes also participated in individual specialty training outside of team events. These activities were not included in the incidence calculations, however in retrospect, excluding individual activities may not have adequately captured the true number of athletic exposures for each youth athlete. When comparing the UE injury frequencies of this study with prior youth baseball studies, injury frequencies were lower than the previously reported values despite demonstrating a high UE injury rate (Table 6.1).<sup>90,91,116,162</sup> Based on the youth baseball participation patterns observed in this study, future projects should include individual training sessions in addition to team activities when capturing athletic exposure.

The most important result of our study shows that youth baseball players who specialize in baseball demonstrated greater UE injury risk compared to those who did not specialize. These findings support recent work conducted in high school athletes, that displayed increased LE injury risk in individuals who specialized in a single sport compared to those who did not.<sup>8,99</sup> Despite evidence showing the widespread risks associated with sport specialization in youth and adolescent athletes, its prevalence continues to increase in the U.S.<sup>72,107</sup> The evolution of youth sports has developed into a multi-billion dollar business as more parents and athletes aspire to achieve elite levels of play and competition.<sup>58,71</sup> A commonly held misconception in youth sports is that



focusing on a single sport early in athlete's career will improve their future performances in that sport.<sup>71</sup> Current research does not support that theory and has shown that early sport specialization is not associated with an athlete's long-term success in sport and that early diversification may be more beneficial to their physical development.<sup>84,109</sup>

Another interesting finding of this study was that youth players who participated in additional specialty training, particularly as a pitcher, were at the greatest risk for sustaining an upper extremity injury in this cohort. This is concerning given that two-thirds of youth baseball players identified as pitchers and reported playing other positions when not pitching. Participating in formal pitching lessons, in addition to team practices and games, may derail the original purpose of age-based pitch count restrictions, by increasing the physical loads placed across a growing athlete's body. This data suggests that not only is specialization an issue but that overtraining may also contribute to the observed disparities in injury rates. Prospectively examining UE injury profiles allowed us to build on the knowledge gained from previous studies and generate a more complete picture of UE injury development in youth baseball.

## **6.2 Normalization Methods for Isometric Shoulder Strength in Youth Baseball Players**

This study investigated five distinct methods for normalizing isometric shoulder strength to determine which had the best measurement properties for youth baseball players. The torque method demonstrated excellent intra-rater reliability, with the lowest reported SEM and MDC values of any method examined. Once normalized, dominant shoulder scaption torque was the most predictive measure with respect to ball velocity, followed by dominant shoulder ER torque at 0°. When examined individually, dominant

shoulder scaption torque predicted 27% of the variation in ball velocity while dominant shoulder ER torque at 0° predicted 23% in a cohort of youth baseball players.

Dominant shoulder scaption strength and ER strength at 0° have been previously linked to upper extremity injury risk and throwing performance in baseball players.<sup>17,44,61,97,105,148</sup> The majority of studies examining upper extremity strength as a factor for performance and injury risk were performed in collegiate and professional athletes.<sup>44,50,97,105</sup> Studies have found little conclusive evidence supporting the theory of a direct relationship between baseline shoulder muscle strength and ball velocity in throwers, however shoulder muscle weakness has been repeatedly linked to injury throughout the baseball literature.<sup>17,28,97,105,133</sup> Magnusson<sup>97</sup>, Mullaney<sup>105</sup> and Byram<sup>17</sup> have all shown that pitchers demonstrated deficits in dominant shoulder scaption and ER strength. While minimal data exists examining upper extremity muscle strength and injury risk in youth athletes, one study by Harada<sup>61</sup> compared a battery of shoulder muscle strength measures to injury risk in a cohort of Japanese youth baseball players. They found that injured athletes demonstrated greater dominant shoulder strength compared to uninjured athletes.<sup>61</sup>

The contradictory nature of the findings above suggests that the connection between shoulder strength and upper extremity injury risk may be different in youth athletes when compared to their collegiate and professional counterparts.<sup>17,61,97,105</sup> The results of the current study also indicate that the relationship between shoulder strength and throwing performance may be different in youth baseball players when compared to the collegiate and professional ranks. Youth baseball players demonstrated a positive relationship between normalized isometric shoulder torque and ball velocity while

previously studies have shown no such relationship in collegiate and professional players.<sup>17,28,97,105</sup>

### **6.3 Limitations**

Limitations in study design should be noted when interpreting the results of this research. Previous epidemiological studies have focused on the etiology and factors surrounding time-loss injuries and typically include all injuries across an athlete's body in their data analyses.<sup>31,43,124,137</sup> The current study included both time-loss and non-time-loss injuries at the shoulder and elbow when calculating injury rates. This decision was based on the target population and previous data that showed the shoulder and elbow to be two of the most commonly injured body parts in competitive baseball players.<sup>80,124,137</sup> The purpose of the study was to examine UE overuse injuries in youth throwers so the decision was made to exclude all non-throwing-related injuries from the data analyses. The exclusion of non-throwing injuries is an acknowledged limitation of the study as any injury has the potential to impact participation levels in baseball activities.

Sampling limitations were also present, specifically the obvious disparities in the number of youth pitchers versus position players as well as the number of specialized athletes versus multi-sport athletes reported in this study. While the groups were uneven for statistical analysis, the proportions for each position and participation level were representative of the target population being studied. To capture an adequately sized sample of youth baseball players, a 10-week rolling enrollment period was required to complete baseline evaluations. These evaluations were performed throughout the competitive baseball season. Athletes were then tracked for 6 months following individual baseline evaluations. This resulted in varied baseball participation rates among

youth athletes that were likely impacted by competition level (i.e. little league baseball vs. tournament team baseball) and participation in additional sport seasons (i.e. football season, basketball season). The variability in the number of athletic exposures recorded per athlete over the 6-month calendar period, while representative of youth baseball participation rates, may have influenced our athletic exposure data collection. Gaps in baseball participation, that are unrelated to injury, can be multi-factorial in youth populations resulting in a decreased number of total exposures and an increased UE injury rate.

Hand held dynamometry has several clinical advantages such as low cost, portability and ease of use however it also has acknowledged instrumental limitations.<sup>33,52</sup> While extreme effort was expended to standardize all measurements and testing procedures including using a single investigator with excellent intra-rater reliability, no external stabilization was applied to the athletes during the assessments. This decision was based on feasibility and the ability to generalize our results to clinical settings, however it may have influenced our results.

One potential statistical limitation of this study was the collinearity observed between the isometric shoulder scaption torque and ER torque at 0° measures in this population. Our results indicate that either measure is predictive of ball velocity however further research is needed to determine what additional variables should be included in the model to best explain this performance measure in youth athletes. Overhead throwing is a complex motor skill that requires coordination and the proper sequencing of a series of linked movements that start in the lower extremities and ultimately culminate in ball release.<sup>157</sup> This statement supports our findings that isometric shoulder strength explains

only a portion of the variability observed in ball velocity in young throwers. Lastly, the high levels of inter-participant variability observed in isometric shoulder ER and IR torque at 90° may have negatively influenced the predictability of these measures in youth athletes. The variability may be attributable to age-appropriate deficits in neuromuscular control in the prone overhead position however further study is needed to better understand the role neuromuscular control plays in youth baseball throwing mechanics.

#### **6.4 Clinical Implications**

The current study established UE injury incidence and identified population-specific risk factors, such as sport specialization and participation in specialty training, which increased UE injury risk in a cohort of youth baseball players. The torque normalization method for isometric shoulder strength exhibited the most consistent measurement properties in young throwers. Future longitudinal and repeated measures studies can utilize this method in physical developing populations as torque was proven to be a reliable means of assessing strength while accounting for changes in body size over time. Dominant shoulder scaption torque and shoulder ER at 0° torque were the most useful measures in predicting throwing performance in youth baseball players. The results of the current study suggest that participation patterns, risk factors and performance measures in youth baseball players are inherently different from those reported in collegiate and professional players. Additional population-specific research studies are needed to better understand injury risk, prevention and player performance in youth athletes.

## 6.5 Future Studies

Future epidemiologic studies should seek to capture specific sports participation and physical activity levels in youth athletes to better identify the factors and workloads associated with overtraining in sports. Identifying individual as well as team-based activities is imperative when assessing athletic exposure in these populations. Studies should also examine the effects of athlete-specific risk factors, such as height, weight, flexibility and strength, on UE injury risk and athletic performance in youth athletes, as these relationships are not well established in the literature. This data is imperative for the development and successful implementation of physical training and injury prevention programs in active youth populations.

**Table 6.1. Injury Frequency Comparisons Across Multiple Youth Baseball Epidemiologic Studies**

<b>Author</b>	<b>Sample Size</b>	<b>Age Range</b>	<b>Follow Up Period</b>	<b>Injury Description</b>	<b>Injury Location</b>	<b>Injury Frequency</b>
<b>Lyman (2001)</b>	298	9-12	2 seasons	Self-Reported Pain	Shoulder, Elbow	Shoulder 32% Elbow 26%
<b>Lyman (2002)</b>	476	9-14	1 season	Self-Reported Pain	Shoulder, Elbow	Shoulder 35% Elbow 28%
<b>Olsen (2006)</b>	95 – Surgical Group 45 – Control Group	14-20	12 months	Self-Reported Pitching w/Pain	Shoulder, Elbow	Surgical 67% Control 42%
<b>Yang (2014)</b>	754	9-18	12 months	Self-Reported Injury, Self-Reported Pain	Shoulder, Elbow	Injury 31.3% Pain 37.9%
<b>Current Study (2017)</b>	159	9-12	6 months	PT* -Confirmed Injury	Shoulder, Elbow	Shoulder 8% Elbow 5%

\* PT, physical therapist.

∞

## REFERENCES

1. Adirim TA, Cheng TL. Overview of injuries in the young athlete. *Sports Med.* 2003;33(1):75-81.
2. Akgul S, Dilicikik U, Kanbur NO, Kaya D, Donmez G, Doral MN. Proximal humeral physeal widening: little leaguer's shoulder or a variation of normal development? *Turk J Pediatr.* 2011;53(6):711-714.
3. Albanese SA, Palmer AK, Kerr DR, Carpenter CW, Lisi D, Levinsohn EM. Wrist pain and distal growth plate closure of the radius in gymnasts. *J Pediatr Orthop.* 1989;9(1):23-28.
4. American Academy of Pediatrics CoSMaF. Policy statement: Baseball and Softball. *Pediatrics.* 2012;129:842-856.
5. Anton C, Podberesky DJ. Little League shoulder: a growth plate injury. *Pediatr Radiol.* 2010;40 Suppl 1:S54.
6. Arnold A, Thigpen CA, Beattie PF, Kissenberth MJ, Shanley E. Overuse Physeal Injuries in Youth Athletes. *Sports Health.* 2017;9(2):139-147.
7. Bailey LB, Shanley E, Hawkins R, et al. Mechanisms of Shoulder Range of Motion Deficits in Asymptomatic Baseball Players. *Am J Sports Med.* 2015;43(11):2783-2793.
8. Bell DR PE, Trigsted SM, Hetzel S, McGuine TA and Brooks MA. Prevalence of Sport Specialization in High School Athletics: A 1-Year Observational Study. *Am J Sports Med.* 2016;44(6):1469-1474.
9. Beovich RF, P.A. Osgood-Schlatter's disease: A review of the literature and an Australian series. *Australian Journal of Science, Medicine & Sport.* 1988;20:11-13.



10. Binder H, Schurz M, Aldrian S, Fialka C, Vecsei V. Physeal injuries of the proximal humerus: long-term results in seventy two patients. *Int Orthop*. 2011;35(10):1497-1502.
11. Bloom OJ, Mackler L, Barbee J. Clinical inquiries. What is the best treatment for Osgood-Schlatter disease? *J Fam Pract*. 2004;53(2):153-156.
12. Bohne C, George SZ, Zeppieri G, Jr. Knowledge of injury prevention and prevalence of risk factors for throwing injuries in a sample of youth baseball players. *Int J Sports Phys Ther*. 2015;10(4):464-475.
13. Boyd KT, Batt ME. Stress fracture of the proximal humeral epiphysis in an elite junior badminton player. *Br J Sports Med*. 1997;31(3):252-253.
14. Brooks MA, Schiff MA, Rivara FP. Identifying previous sports injury among high school athletes. *Clin Pediatr (Phila)*. 2009;48(5):548-550.
15. Bruns W, Maffulli N. Lower limb injuries in children in sports. *Clin Sports Med*. 2000;19(4):637-662.
16. Butler TA, Yingling VR. The effects of delayed puberty on the growth plate. *J Pediatr Orthop*. 2013;33(1):99-105.
17. Byram IR, Bushnell BD, Dugger K, Charron K, Harrell FE, Jr., Noonan TJ. Preseason shoulder strength measurements in professional baseball pitchers: identifying players at risk for injury. *Am J Sports Med*. 2010;38(7):1375-1382.
18. Caine D, Caine C, Maffulli N. Incidence and distribution of pediatric sport-related injuries. *Clin J Sport Med*. 2006;16(6):500-513.
19. Caine D, DiFiori J, Maffulli N. Physeal injuries in children's and youth sports: reasons for concern? *Br J Sports Med*. 2006;40(9):749-760.
20. Caine D, Lewis R, O'Connor P, Howe W, Bass S. Does gymnastics training inhibit growth of females? *Clin J Sport Med*. 2001;11(4):260-270.

21. Caine D, Maffulli N, Caine C. Epidemiology of injury in child and adolescent sports: injury rates, risk factors, and prevention. *Clin Sports Med.* 2008;27(1):19-50, vii.
22. Caine D, Roy S, Singer KM, Broekhoff J. Stress changes of the distal radial growth plate. A radiographic survey and review of the literature. *Am J Sports Med.* 1992;20(3):290-298.
23. Caine DB, S.; Daly, R. Does elite competition inhibit growth and delay maturation in some gymnasts? Quite possibly. *Pediatric exercise science.* 2003;15(4):360-372.
24. Caine DH, W.; Ross, W.; Bergman, R. Does repetitive physical loading inhibit radial growth in female gymnasts. *Clin J Sports Med.* 1997;7(4):302-308.
25. Caine DJ, Golightly YM. Osteoarthritis as an outcome of paediatric sport: an epidemiological perspective. *Br J Sports Med.* 2011;45(4):298-303.
26. Caine DJ, Maffulli N. Epidemiology of children's individual sports injuries. An important area of medicine and sport science research. *Med Sport Sci.* 2005;48:1-7.
27. Carson WG, Jr., Gasser SI. Little Leaguer's shoulder. A report of 23 cases. *Am J Sports Med.* 1998;26(4):575-580.
28. Carter AB, Kaminski TW, Douex AT, Jr., Knight CA, Richards JG. Effects of high volume upper extremity plyometric training on throwing velocity and functional strength ratios of the shoulder rotators in collegiate baseball players. *J Strength Cond Res.* 2007;21(1):208-215.
29. Carter SR, Aldridge MJ, Fitzgerald R, Davies AM. Stress changes of the wrist in adolescent gymnasts. *Br J Radiol.* 1988;61(722):109-112.
30. Cohen E, Sala DA. Rehabilitation of pediatric musculoskeletal sport-related injuries: a review of the literature. *Eur J Phys Rehabil Med.* 2010;46(2):133-145.
31. Collins CL, Comstock RD. Epidemiological features of high school baseball injuries in the United States, 2005-2007. *Pediatrics.* 2008;121(6):1181-1187.

32. USA Baseball Medical & Safety Advisory Committee. Youth Baseball Pitching Injuries. 2008;  
[http://web.usabaseball.com/news/article.jsp?ymd=20090813&content\\_id=640950](http://web.usabaseball.com/news/article.jsp?ymd=20090813&content_id=640950)  
8. Accessed April 2, 2016.
33. Cools AM, De Wilde L, Van Tongel A, Ceyskens C, Ryckewaert R, Cambier DC. Measuring shoulder external and internal rotation strength and range of motion: comprehensive intra-rater and inter-rater reliability study of several testing protocols. *J Shoulder Elbow Surg.* 2014;23(10):1454-1461.
34. de Lucena GL, dos Santos Gomes C, Guerra RO. Prevalence and associated factors of Osgood-Schlatter syndrome in a population-based sample of Brazilian adolescents. *Am J Sports Med.* 2011;39(2):415-420.
35. De Smet L, Claessens A, Lefevre J, Beunen G. Gymnast wrist: an epidemiologic survey of ulnar variance and stress changes of the radial physis in elite female gymnasts. *Am J Sports Med.* 1994;22(6):846-850.
36. Dick R, Sauers EL, Agel J, et al. Descriptive epidemiology of collegiate men's baseball injuries: National Collegiate Athletic Association Injury Surveillance System, 1988-1989 through 2003-2004. *J Athl Train.* 2007;42(2):183-193.
37. DiFiori JP. Overuse injury and the young athlete: the case of chronic wrist pain in gymnasts. *Curr Sports Med Rep.* 2006;5(4):165-167.
38. DiFiori JP, Benjamin HJ, Brenner J, et al. Overuse injuries and burnout in youth sports: a position statement from the American Medical Society for Sports Medicine. *Clin J Sport Med.* 2014;24(1):3-20.
39. DiFiori JP, Benjamin HJ, Brenner JS, et al. Overuse injuries and burnout in youth sports: a position statement from the American Medical Society for Sports Medicine. *Br J Sports Med.* 2014;48(4):287-288.
40. DiFiori JP, Caine DJ, Malina RM. Wrist pain, distal radial physeal injury, and ulnar variance in the young gymnast. *Am J Sports Med.* 2006;34(5):840-849.
41. DiFiori JP, Puffer JC, Aish B, Dorey F. Wrist pain, distal radial physeal injury, and ulnar variance in young gymnasts: does a relationship exist? *Am J Sports Med.* 2002;30(6):879-885.

42. DiFiori JP, Puffer JC, Mandelbaum BR, Dorey F. Distal radial growth plate injury and positive ulnar variance in nonelite gymnasts. *Am J Sports Med.* 1997;25(6):763-768.
43. Dompier TP, Powell JW, Barron MJ, Moore MT. Time-loss and non-time-loss injuries in youth football players. *J Athl Train.* 2007;42(3):395-402.
44. Donatelli R, Ellenbecker TS, Ekedahl SR, Wilkes JS, Kocher K, Adam J. Assessment of shoulder strength in professional baseball pitchers. *J Orthop Sports Phys Ther.* 2000;30(9):544-551.
45. Doral MN, Aydog ST, Tetik O, Atay OA, Turhan E, Demirel HA. Multiple osteochondroses and avulsion fracture of anterior superior iliac spine in a soccer player. *Br J Sports Med.* 2005;39(3):e16.
46. Dover GC, Kaminski TW, Meister K, Powers ME, Horodyski M. Assessment of shoulder proprioception in the female softball athlete. *Am J Sports Med.* 2003;31(3):431-437.
47. Drescher WR, Falliner A, Zantop T, Oehlert K, Petersen W, Hassenpflug J. Little league shoulder syndrome in an adolescent cricket player. *Br J Sports Med.* 2004;38(4):E14.
48. Eastwood D, Bijlsma, P. Physeal injuries in children. *Surgery.* 2014;32(1):1-8.
49. Ebell MH, Siwek J, Weiss BD, et al. Strength of recommendation taxonomy (SORT): a patient-centered approach to grading evidence in the medical literature. *J Am Board Fam Pract.* 2004;17(1):59-67.
50. Ellenbecker TS, Mattalino AJ. Concentric isokinetic shoulder internal and external rotation strength in professional baseball pitchers. *J Orthop Sports Phys Ther.* 1997;25(5):323-328.
51. ESPN. Hidden Demographics of Youth Sports. 2013; [http://www.espn.com/espn/story/\\_/id/9469252/hidden-demographics-youth-sports-espn-magazine](http://www.espn.com/espn/story/_/id/9469252/hidden-demographics-youth-sports-espn-magazine). Accessed September 1, 2017.
52. Fieseler G, Molitor T, Irlenbusch L, et al. Intrarater reliability of goniometry and hand-held dynamometry for shoulder and elbow examinations in female team

- handball athletes and asymptomatic volunteers. *Arch Orthop Trauma Surg.* 2015;135(12):1719-1726.
53. Fleisig GS, Andrews JR, Cutter GR, et al. Risk of serious injury for young baseball pitchers: a 10-year prospective study. *Am J Sports Med.* 2011;39(2):253-257.
54. Ford P, De Ste Croix M, Lloyd R, et al. The long-term athlete development model: physiological evidence and application. *J Sports Sci.* 2011;29(4):389-402.
55. Frush TJ, Lindenfeld TN. Peri-epiphyseal and Overuse Injuries in Adolescent Athletes. *Sports Health.* 2009;1(3):201-211.
56. Gottschalk AW, Andrish JT. Epidemiology of sports injury in pediatric athletes. *Sports Med Arthrosc.* 2011;19(1):2-6.
57. Greenberg EM, Fernandez-Fernandez A, Lawrence JT, McClure P. The Development of Humeral Retrotorsion and Its Relationship to Throwing Sports. *Sports Health.* 2015;7(6):489-496.
58. Gregory S. How Kids' Sports Became a \$15 Billion Industry. *TIME Magazine Sports*; 2017.
59. Hajdu S, Kaltenecker G, Schwendenwein E, Vecsei V. Apophyseal injuries of the proximal tibial tubercle. *Int Orthop.* 2000;24(5):279-281.
60. Hang DW, Chao CM, Hang YS. A clinical and roentgenographic study of Little League elbow. *Am J Sports Med.* 2004;32(1):79-84.
61. Harada M, Takahara M, Mura N, Sasaki J, Ito T, Ogino T. Risk factors for elbow injuries among young baseball players. *J Shoulder Elbow Surg.* 2010;19(4):502-507.
62. Hebert LJ, Maltais DB, Lepage C, Saulnier J, Crete M. Hand-Held Dynamometry Isometric Torque Reference Values for Children and Adolescents. *Pediatr Phys Ther.* 2015;27(4):414-423.

63. Hebert LJ, Maltais DB, Lepage C, Saulnier J, Crete M, Perron M. Isometric muscle strength in youth assessed by hand-held dynamometry: a feasibility, reliability, and validity study. *Pediatr Phys Ther.* 2011;23(3):289-299.
64. Hibberd EE, Oyama S, Myers JB. Increase in humeral retrotorsion accounts for age-related increase in glenohumeral internal rotation deficit in youth and adolescent baseball players. *Am J Sports Med.* 2014;42(4):851-858.
65. Hurd WJ, Kaplan KM, Elattrache NS, Jobe FW, Morrey BF, Kaufman KR. A profile of glenohumeral internal and external rotation motion in the uninjured high school baseball pitcher, part I: motion. *J Athl Train.* 2011;46(3):282-288.
66. Hussain AH, G.A. Osgood-Schlatter disease. *Sports Exercise Injury.* 1996;2:202-206.
67. Jaric S. Muscle strength testing: use of normalisation for body size. *Sports Med.* 2002;32(10):615-631.
68. Jaric S. Role of body size in the relation between muscle strength and movement performance. *Exerc Sport Sci Rev.* 2003;31(1):8-12.
69. Jaric S, Radosavljevic-Jaric S, Johansson H. Muscle force and muscle torque in humans require different methods when adjusting for differences in body size. *Eur J Appl Physiol.* 2002;87(3):304-307.
70. Jaric S, Ugarkovic D, Kukolj M. Evaluation of methods for normalizing muscle strength in elite and young athletes. *J Sports Med Phys Fitness.* 2002;42(2):141-151.
71. Jayanthi N PC, Dugas L, Patrick B and LaBella C. Sports Specialization in Young Athletes: Evidence-Based Recommendations. *Sports Health.* 2013;5(3):251-257.
72. Jayanthi NA, LaBella CR, Fischer D, Pasulka J, Dugas LR. Sports-specialized intensive training and the risk of injury in young athletes: a clinical case-control study. *Am J Sports Med.* 2015;43(4):794-801.
73. Kaplan KM, Elattrache NS, Jobe FW, Morrey BF, Kaufman KR, Hurd WJ. Comparison of shoulder range of motion, strength, and playing time in uninjured

- high school baseball pitchers who reside in warm- and cold-weather climates. *Am J Sports Med.* 2011;39(2):320-328.
74. Kennedy JG, Knowles B, Dolan M, Bohne W. Foot and ankle injuries in the adolescent runner. *Curr Opin Pediatr.* 2005;17(1):34-42.
75. Kerr ZY, Lynall RC, Roos KG, Dalton SL, Djoko A, Dompier TP. Descriptive Epidemiology of Non-Time-Loss Injuries in Collegiate and High School Student-Athletes. *J Athl Train.* 2017;52(5):446-456.
76. Kerssemakers SP, Fotiadou AN, de Jonge MC, Karantanas AH, Maas M. Sport injuries in the paediatric and adolescent patient: a growing problem. *Pediatr Radiol.* 2009;39(5):471-484.
77. Klingele KE, Kocher MS. Little league elbow: valgus overload injury in the paediatric athlete. *Sports Med.* 2002;32(15):1005-1015.
78. Kocher MS, Waters PM, Micheli LJ. Upper extremity injuries in the paediatric athlete. *Sports Med.* 2000;30(2):117-135.
79. Kolt GS, Kirkby RJ. Epidemiology of injury in elite and subelite female gymnasts: a comparison of retrospective and prospective findings. *Br J Sports Med.* 1999;33(5):312-318.
80. Krajnik S, Fogarty KJ, Yard EE, Comstock RD. Shoulder injuries in US high school baseball and softball athletes, 2005-2008. *Pediatrics.* 2010;125(3):497-501.
81. Krueger-Franke M, Siebert CH, Pfoerringer W. Sports-related epiphyseal injuries of the lower extremity. An epidemiologic study. *J Sports Med Phys Fitness.* 1992;32(1):106-111.
82. Kujala UMK, M.; Heinonen, O. Osgood-Schlatter's disease in adolescent athletes: Retrospective study of incidence and duration. *Am J Sports Med.* 1985;13:236-241.
83. Laor T, Wall EJ, Vu LP. Physeal widening in the knee due to stress injury in child athletes. *AJR Am J Roentgenol.* 2006;186(5):1260-1264.

84. LaPrade RF, Agel J, Baker J, et al. AOSSM Early Sport Specialization Consensus Statement. *Orthop J Sports Med.* 2016;4(4):2325967116644241.
85. Lawson BR, Comstock RD, Smith GA. Baseball-related injuries to children treated in hospital emergency departments in the United States, 1994-2006. *Pediatrics.* 2009;123(6):e1028-1034.
86. Little League Baseball. Little League Regulations, Rules and Policies. 2017.
87. LeVeau BF, Bernhardt DB. Developmental biomechanics. Effect of forces on the growth, development, and maintenance of the human body. *Phys Ther.* 1984;64(12):1874-1882.
88. Liebling MS, Berdon WE, Ruzal-Shapiro C, Levin TL, Roye D, Jr., Wilkinson R. Gymnast's wrist (pseudorickets growth plate abnormality) in adolescent athletes: findings on plain films and MR imaging. *AJR Am J Roentgenol.* 1995;164(1):157-159.
89. Lyman S, Fleisig GS. Baseball injuries. *Med Sport Sci.* 2005;49:9-30.
90. Lyman S, Fleisig GS, Andrews JR, Osinski ED. Effect of pitch type, pitch count, and pitching mechanics on risk of elbow and shoulder pain in youth baseball pitchers. *Am J Sports Med.* 2002;30(4):463-468.
91. Lyman S, Fleisig GS, Waterbor JW, et al. Longitudinal study of elbow and shoulder pain in youth baseball pitchers. *Med Sci Sports Exerc.* 2001;33(11):1803-1810.
92. De Ste Croix, M. Advances in paediatric strength assessment: changing our perspective on strength development. *Journal of Sports Science and Medicine.* 2007;6:292-304.
93. Maffulli N, Bruns W. Injuries in young athletes. *Eur J Pediatr.* 2000;159(1-2):59-63.
94. Maffulli N, Longo UG, Gougoulas N, Caine D, Denaro V. Sport injuries: a review of outcomes. *Br Med Bull.* 2011;97:47-80.



95. Maffulli N, Longo UG, Gougoulas N, Loppini M, Denaro V. Long-term health outcomes of youth sports injuries. *Br J Sports Med.* 2010;44(1):21-25.
96. Maffulli N, Longo UG, Spiezia F, Denaro V. Sports injuries in young athletes: long-term outcome and prevention strategies. *Phys Sportsmed.* 2010;38(2):29-34.
97. Magnusson SP, Gleim GW, Nicholas JA. Shoulder weakness in professional baseball pitchers. *Med Sci Sports Exerc.* 1994;26(1):5-9.
98. McBain K, Shrier I, Shultz R, et al. Prevention of sports injury I: a systematic review of applied biomechanics and physiology outcomes research. *Br J Sports Med.* 2012;46(3):169-173.
99. McGuine TA, Post EG, Hetzel SJ, Brooks MA, Trigsted S, Bell DR. A Prospective Study on the Effect of Sport Specialization on Lower Extremity Injury Rates in High School Athletes. *Am J Sports Med.* 2017:363546517710213.
100. McHugh MP, Tyler TF, Tetro DT, Mullaney MJ, Nicholas SJ. Risk factors for noncontact ankle sprains in high school athletes: the role of hip strength and balance ability. *Am J Sports Med.* 2006;34(3):464-470.
101. Meister K, Day T, Horodyski M, Kaminski TW, Wasik MP, Tillman S. Rotational motion changes in the glenohumeral joint of the adolescent/Little League baseball player. *Am J Sports Med.* 2005;33(5):693-698.
102. Mirtz TA, Chandler JP, Eysers CM. The effects of physical activity on the epiphyseal growth plates: a review of the literature on normal physiology and clinical implications. *J Clin Med Res.* 2011;3(1):1-7.
103. Mital MA, Matza RA, Cohen J. The so-called unresolved Osgood-Schlatter lesion: a concept based on fifteen surgically treated lesions. *J Bone Joint Surg Am.* 1980;62(5):732-739.
104. Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Int J Surg.* 2010;8(5):336-341.

105. Mullaney MJ, McHugh MP, Donofrio TM, Nicholas SJ. Upper and lower extremity muscle fatigue after a baseball pitching performance. *Am J Sports Med.* 2005;33(1):108-113.
106. Murray, J.; Morscher, E.; Rahn, B.A.; Kaslin, M. Bone growth and remodeling after fracture. *J Bone Joint Surg Br.* 1996;78(1):42-50.
107. Myer GD, Jayanthi N, Difiori JP, et al. Sport Specialization, Part I: Does Early Sports Specialization Increase Negative Outcomes and Reduce the Opportunity for Success in Young Athletes? *Sports Health.* 2015;7(5):437-442.
108. Myer GD, Jayanthi N, DiFiori JP, et al. Sports Specialization, Part II: Alternative Solutions to Early Sport Specialization in Youth Athletes. *Sports Health.* 2016;8(1):65-73.
109. Myers JB, DeFreese JD, DiStefano L, et al. USA Baseball Long Term Athlete Development Model. 2017; <http://usabltd.com>. Accessed September 25, 2017.
110. Myers JB, Oyama S, Clarke JP. Ultrasonographic assessment of humeral retrotorsion in baseball players: a validation study. *Am J Sports Med.* 2012;40(5):1155-1160.
111. Myers JB, Oyama S, Goerger BM, Rucinski TJ, Blackburn JT, Creighton RA. Influence of humeral torsion on interpretation of posterior shoulder tightness measures in overhead athletes. *Clin J Sport Med.* 2009;19(5):366-371.
112. Myers JB, Oyama S, Rucinski TJ, Creighton RA. Humeral retrotorsion in collegiate baseball pitchers with throwing-related upper extremity injury history. *Sports Health.* 2011;3(4):383-389.
113. Nabhan D, Walden T, Street J, Linden H, Moreau B. Sports injury and illness epidemiology during the 2014 Youth Olympic Games: United States Olympic Team Surveillance. *Br J Sports Med.* 2016;50(11):688-693.
114. Nanni M, Butt S, Mansour R, Muthukumar T, Cassar-Pullicino VN, Roberts A. Stress-induced Salter-Harris I growth plate injury of the proximal tibia: first report. *Skeletal Radiol.* 2005;34(7):405-410.

115. Obembe OO, Gaskin CM, Taffoni MJ, Anderson MW. Little Leaguer's shoulder (proximal humeral epiphysiolysis): MRI findings in four boys. *Pediatr Radiol.* 2007;37(9):885-889.
116. Olsen SJ, 2nd, Fleisig GS, Dun S, Loftice J, Andrews JR. Risk factors for shoulder and elbow injuries in adolescent baseball pitchers. *Am J Sports Med.* 2006;34(6):905-912.
117. Orava S, Virtanen K. Osteochondroses in athletes. *Br J Sports Med.* 1982;16(3):161-168.
118. Oyama S, Hibberd EE, Myers JB. Changes in humeral torsion and shoulder rotation range of motion in high school baseball players over a 1-year period. *Clin Biomech (Bristol, Avon).* 2013;28(3):268-272.
119. Patel DR, Nelson TL. Sports injuries in adolescents. *Med Clin North Am.* 2000;84(4):983-1007, viii.
120. Paterno MV, Taylor-Haas JA, Myer GD, Hewett TE. Prevention of overuse sports injuries in the young athlete. *Orthop Clin North Am.* 2013;44(4):553-564.
121. Petty DH, Andrews JR, Fleisig GS, Cain EL. Ulnar collateral ligament reconstruction in high school baseball players: clinical results and injury risk factors. *Am J Sports Med.* 2004;32(5):1158-1164.
122. Portney LW, MP. *Foundations of Clinical Research: Applications to Practice.* 3rd ed. Upper Saddle River, NJ: Pearson Education, Inc.; 2009.
123. Posner M, Cameron KL, Wolf JM, Belmont PJ, Jr., Owens BD. Epidemiology of Major League Baseball injuries. *Am J Sports Med.* 2011;39(8):1676-1680.
124. Powell JW, Dompier TP. Analysis of Injury Rates and Treatment Patterns for Time-Loss and Non-Time-Loss Injuries Among Collegiate Student-Athletes. *J Athl Train.* 2004;39(1):56-70.
125. Rauck RC, LaMont LE, Doyle SM. Pediatric upper extremity stress injuries. *Curr Opin Pediatr.* 2013;25(1):40-45.

126. Rauh MJ, Koepsell TD, Rivara FP, Margherita AJ, Rice SG. Epidemiology of musculoskeletal injuries among high school cross-country runners. *Am J Epidemiol.* 2006;163(2):151-159.
127. Ray TR. Youth baseball injuries: recognition, treatment, and prevention. *Curr Sports Med Rep.* 2010;9(5):294-298.
128. Reinold MM, Gill TJ. Current concepts in the evaluation and treatment of the shoulder in overhead-throwing athletes, part 1: physical characteristics and clinical examination. *Sports Health.* 2010;2(1):39-50.
129. Reinold MW, K.; Hooks, T.; Andrews, J. Comparison of bilateral shoulder external and internal rotation strength at 0 degrees and 90 degrees of abduction in professional baseball pitchers. *Journal of Orthopaedic and Sports Physical Therapy.* 2003;33(2):A51.
130. Rossi F, Dragoni S. Acute avulsion fractures of the pelvis in adolescent competitive athletes: prevalence, location and sports distribution of 203 cases collected. *Skeletal Radiol.* 2001;30(3):127-131.
131. Roy S, Caine D, Singer KM. Stress changes of the distal radial epiphysis in young gymnasts. A report of twenty-one cases and a review of the literature. *Am J Sports Med.* 1985;13(5):301-308.
132. Sabick MB, Kim YK, Torry MR, Keirns MA, Hawkins RJ. Biomechanics of the shoulder in youth baseball pitchers: implications for the development of proximal humeral epiphysiolysis and humeral retrotorsion. *Am J Sports Med.* 2005;33(11):1716-1722.
133. Sabick MB, Torry MR, Lawton RL, Hawkins RJ. Valgus torque in youth baseball pitchers: A biomechanical study. *J Shoulder Elbow Surg.* 2004;13(3):349-355.
134. Scharfbillig RW, Jones S, Scutter SD. Sever's disease: what does the literature really tell us? *J Am Podiatr Med Assoc.* 2008;98(3):212-223.
135. Severini GC, A.; Campana, V.; Milano, G. Prevention strategies of shoulder injuries. *Sports Injuries: Prevention, Diagnosis, Treatment and Rehabilitation.* 2015:279-290.

136. Shanley E, Kissenberth MJ, Thigpen CA, et al. Preseason shoulder range of motion screening as a predictor of injury among youth and adolescent baseball pitchers. *J Shoulder Elbow Surg.* 2015;24(7):1005-1013.
137. Shanley E, Rauh MJ, Michener LA, Ellenbecker TS. Incidence of injuries in high school softball and baseball players. *J Athl Train.* 2011;46(6):648-654.
138. Shanley EB, L.B.; Rauh, M.J.; et al. Influence of a prevention program on arm injury risk: An RCT in adolescent pitchers. *Orthop J Sports Med.* 2014;2(2).
139. Shanley ER, M.J.; Michener, L.A.; Ellenbecker, T.S.; Garrison, J.C.; Thigpen, C.A. Shoulder range of motion measures as risk factors for shoulder and elbow injuries in high school softball and baseball players. *Am J Sports Med.* 2011;39(9):1997-2006.
140. Shanley ET, C.A. Throwing injuries in the adolescent athlete. *Int J Sports Phys Ther.* 2013;8(5):630-640.
141. Sharma PL, K.L.; Maffulli, N. Sports injuries in children. *Trauma.* 2003;5(4):245-259.
142. Stark T, Walker B, Phillips JK, Fejer R, Beck R. Hand-held dynamometry correlation with the gold standard isokinetic dynamometry: a systematic review. *PM R.* 2011;3(5):472-479.
143. Statista. Number of Baseball & Softball Players in the U.S. 2017; <https://www.statista.com/statistics/227429/number-of-softball-players-and-baseball-players-usa/>. Accessed September 1, 2017.
144. Stracciolini A, Casciano R, Levey Friedman H, Meehan WP, 3rd, Micheli LJ. Pediatric sports injuries: an age comparison of children versus adolescents. *Am J Sports Med.* 2013;41(8):1922-1929.
145. Stracciolini A, Casciano R, Levey Friedman H, Stein CJ, Meehan WP, 3rd, Micheli LJ. Pediatric sports injuries: a comparison of males versus females. *Am J Sports Med.* 2014;42(4):965-972.

146. Stratford PW, Balsor BE. A comparison of make and break tests using a hand-held dynamometer and the Kin-Com. *J Orthop Sports Phys Ther.* 1994;19(1):28-32.
147. Torg JS, Pollack H, Sweterlitsch P. The effect of competitive pitching on the shoulders and elbows of preadolescent baseball players. *Pediatrics.* 1972;49(2):267-272.
148. Trakis JE, McHugh MP, Caracciolo PA, Busciacco L, Mullaney M, Nicholas SJ. Muscle strength and range of motion in adolescent pitchers with throwing-related pain: implications for injury prevention. *Am J Sports Med.* 2008;36(11):2173-2178.
149. Tyler TF, McHugh MP, Mirabella MR, Mullaney MJ, Nicholas SJ. Risk factors for noncontact ankle sprains in high school football players: the role of previous ankle sprains and body mass index. *Am J Sports Med.* 2006;34(3):471-475.
150. Valentino M, Quiligotti C, Ruggirello M. Sinding-Larsen-Johansson syndrome: A case report. *J Ultrasound.* 2012;15(2):127-129.
151. van Mechelen WH, H; Kemper H. Incidence, Severity, Aetiology and Prevention of Sports Injuries: A Review of Concepts. *Sports Med.* 1992;14(2):82-99.
152. Vandervliet EJ, Vanhoenacker FM, Snoeckx A, Gielen JL, Van Dyck P, Parizel PM. Sports-related acute and chronic avulsion injuries in children and adolescents with special emphasis on tennis. *Br J Sports Med.* 2007;41(11):827-831.
153. Whiteley R, Ginn K, Nicholson L, Adams R. Indirect ultrasound measurement of humeral torsion in adolescent baseball players and non-athletic adults: reliability and significance. *J Sci Med Sport.* 2006;9(4):310-318.
154. Whiteley RJ, Adams RD, Nicholson LL, Ginn KA. Reduced humeral torsion predicts throwing-related injury in adolescent baseballers. *J Sci Med Sport.* 2010;13(4):392-396.
155. Whiteley RJ, Ginn KA, Nicholson LL, Adams RD. Sports participation and humeral torsion. *J Orthop Sports Phys Ther.* 2009;39(4):256-263.

156. Wilk KE, Andrews JR, Arrigo CA, Keirns MA, Erber DJ. The strength characteristics of internal and external rotator muscles in professional baseball pitchers. *Am J Sports Med.* 1993;21(1):61-66.
157. Wilk KE, Meister K, Andrews JR. Current concepts in the rehabilitation of the overhead throwing athlete. *Am J Sports Med.* 2002;30(1):136-151.
158. Wilk KE, Reinold MM, Macrina LC, et al. Glenohumeral internal rotation measurements differ depending on stabilization techniques. *Sports Health.* 2009;1(2):131-136.
159. Wong D.L. H-EM, Wilson D., Winkelstein M.L., Schwartz P. *Wong's Essentials of Pediatric Nursing, 6th ed.* St. Louis 2001.
160. Wooden MJ, Greenfield B, Johanson M, Litzelman L, Mundrane M, Donatelli RA. Effects of strength training on throwing velocity and shoulder muscle performance in teenage baseball players. *J Orthop Sports Phys Ther.* 1992;15(5):223-228.
161. Yamamoto N, Itoi E, Minagawa H, et al. Why is the humeral retroversion of throwing athletes greater in dominant shoulders than in nondominant shoulders? *J Shoulder Elbow Surg.* 2006;15(5):571-575.
162. Yang J, Mann BJ, Guettler JH, et al. Risk-Prone Pitching Activities and Injuries in Youth Baseball: Findings From a National Sample. *Am J Sports Med.* 2014;42(6):1456-1463.
163. Zaremski JL, Krabak BJ. Shoulder injuries in the skeletally immature baseball pitcher and recommendations for the prevention of injury. *PM R.* 2012;4(7):509-516.

# APPENDIX A – SAGE PUBLISHING PERMISSION TO REPRINT POLICY



DISCIPLINES PRODUCTS RESOURCES ABOUT

Search: keyword, title, author, ISBN



0

## Authors Re-Using Their Own Work

SAGE Journal authors are able to re-use their Contribution in certain circumstances without requiring permission from SAGE. It is important to check your signed Contributor Agreement and/or the submission guidelines for the journal to which you submitted your Contribution to review the journal's policy regarding author re-use.

### Gold Open Access and SAGE Choice

If you wish to re-use an Open Access Contribution published under a Creative Commons License, please see [Re-use of Open Access Content](#) for more information.

### Green Open Access: SAGE's archiving policy

Most SAGE journals are published under SAGE's Green Open Access policy, which allows you, as author, to re-use your Contribution as indicated below. For a list of titles that are exceptions to this policy, please scroll down to the bottom of the page.

#### Green Open Access policy:

- You may share the version of the Contribution you submitted to the journal (**version 1**) anywhere at any time.
- Once the Contribution has been accepted for publication, you may post the accepted version (**version 2**) of the Contribution on your own personal website, your department's website or the repository of your institution without any restrictions.
- You may not post the accepted version (**version 2**) of the Contribution in any repository other than those listed above (i.e. you may not deposit in the repository of another institution or a subject repository) until 12 months after first publication of the Contribution in the journal.
- You may use the published Contribution (**version 3**) for your own teaching needs or to supply on an individual basis to research colleagues, provided that such supply is not for commercial purposes.
- You may use the published Contribution (**version 3**) in a book authored or edited by you at any time after publication in the journal. This does not apply to books where you are contributing a chapter to a book authored or edited by someone else.
- You may not post the published Contribution (**version 3**) on a website or in a repository without permission from SAGE.

When posting or reusing your Contribution under this policy, appropriate credit must be given to the SAGE journal where the Contribution has been published, as the original source of the content, as follows: **Author(s), Article Title, Journal Title (Journal Volume Number and Issue Number) pp. xx-xx. Copyright © [year] (Copyright Holder). Reprinted by permission of SAGE Publications.** Additionally, please provide a link to the appropriate DOI for the published version of the Contribution on the SAGE Journals website (<http://journals.sagepub.com>).

The chart below includes common requests and an explanation of which 'version' of the article can be used in each circumstance. If the table below indicates that permission is required, please see the instructions for requesting permission at [Journals Permissions](#).

I WANT TO:	CLEARED PERMISSION	REQUIRES PERMISSION
include up to one full article in my unpublished dissertation or thesis	Version 3	
supply my article to my students or use the article for teaching purposes	Version 3	



## APPENDIX B – REDCAP ONLINE SURVEY SAMPLE

### Youth Baseball Questionnaire

Please answer the following questions as completely as possible. If parents are completing this questionnaire in place of their child, please respond based your child's information. If you have any questions, we would be happy to answer them at any time.

Name: \_\_\_\_\_ Today's Date: \_\_\_\_\_

What is your age? \_\_\_\_\_

Date of Birth: \_\_\_\_\_ Height: \_\_\_\_\_ Weight: \_\_\_\_\_ Mothers Height: \_\_\_\_\_ Father's Height: \_\_\_\_\_

What grade are you in?  3<sup>rd</sup>  4<sup>th</sup>  5<sup>th</sup>  6<sup>th</sup>  7<sup>th</sup>  8<sup>th</sup>  9<sup>th</sup>  10<sup>th</sup>  11<sup>th</sup>  12<sup>th</sup>

What organized sports do you play?  Baseball  Basketball  Any other Sports

(List all): \_\_\_\_\_

How long have you played these sports? (in years)  Baseball \_\_\_\_\_  Basketball \_\_\_\_\_

Any other Sports \_\_\_\_\_

How many months of the year do you compete at your other sport(s)?  1  2  3  4  5  6  7  8  9  10  11

12

How many months of the year do you compete in more than one sport?  1  2  3  4  5  6  7  8  9  10  11

12

How many teams do you play your baseball with during the year?  1  2  3  4  5

Do you play on an elite level (invited team) or travel team?  no  yes Name of Team(s)? \_\_\_\_\_

How many months of the year do you practice/play baseball?  1  2  3  4  5  6  7  8  9  10  11  12

How many practices per week do you average for baseball?  1  2  3  4  5  6  7  8  9  10  11  12

How many games per week do you average for baseball?  1  2  3  4  5  6  7  8  9  10  11  12

What is the maximum # of games per week you have played over the last year for baseball?  1  2  3  4  5  6  7  8

9  10  11  12  >12

What is the minimum recovery time between baseball tournaments? In Days  1  2  3  4  5  6  7  >7

How many months of the year do you completely rest from baseball?  1  2  3  4  5  6  7  8  9  10  11  12

What position do you play most often? (Choose one)

**Baseball:**  catcher  infield  outfield  pitcher

What other positions do you play? (Choose all)

**Baseball:**  catcher  infield  outfield  pitcher

Other than sport specific training, do you participate in an **in season** conditioning program?  no  yes

If so, is the training sport specific (e.g. agility, hitting/pitching lessons, sport specific weight lifting or conditioning)

no  yes Please specify, type of training: \_\_\_\_\_

Location of training: \_\_\_\_\_

Other than sport specific training, do you participate in an **off-season** conditioning program?  no  yes

If so, is the training sport specific (e.g. agility, hitting/pitching lessons, sport specific weight lifting or conditioning)

no  yes Please specify, type of training: \_\_\_\_\_

Location of training: \_\_\_\_\_

Do you have pain in any joint, muscle or body part while playing baseball now or within the last 6 months?

Now?  no  yes ... Last 6 months?  no  yes ... . If yes, which body part? \_\_\_\_\_

Did you see a doctor for the injury?  no  yes

Ever had an injury that caused you to miss a practice or game?  no  yes Change position or alter practice?  no  yes

## APPENDIX C – DATA COLLECTION SHEET

Name: \_\_\_\_\_ Date: \_\_\_\_\_ Birth Date: \_\_\_\_\_

Weight: \_\_\_\_\_ Height: \_\_\_\_\_

Mother's Height: \_\_\_\_\_ Father's Height: \_\_\_\_\_ Dominant Arm:    **R**    **L**

		LEFT		RIGHT	
		Trial 1	Trial 2	Trial 1	Trial 2
ROM	HRT				
	P ER@90				
	P IR@90				
	Elbow Ext				
	P HA				
STRENGTH		Trial 1	Trial 2	Trial 1	Trial 2
	Scaption				
	ER@0				
	ER@90				
	IR@90				
	Ulnar length				

**Current Pain Level:**



Parent's Names: \_\_\_\_\_ Phone: \_\_\_\_\_

Email: \_\_\_\_\_

Current Level/# of teams \_\_\_\_\_ Position (1<sup>st</sup>, 2<sup>nd</sup>): \_\_\_\_\_

Additional Sport(s): \_\_\_\_\_

**Injury History:**

Side of Injury	Body Part (Ex: elbow, shoulder)	Type of Injury	Sport/Position Played (when injured)	Month/Year of Injury