



Phosphorylation-guarded light-harvesting complex II contributes to broad-spectrum blast resistance in rice

Muxing Liu^{a,b}, Suobing Zhang^c, Jiexiong Hu^{a,b}, Wenxian Sun^d, Jonas Padilla^c, Yanglan He^{a,b}, Ying Li^{a,b}, Ziyi Yin^{a,b}, Xinyu Liu^{a,b}, Wenhao Wang^{a,b}, Danyu Shen^{a,b}, Dayong Li^d, Haifeng Zhang^{a,b}, Xiaobo Zheng^{a,b}, Zhongli Cui^e, Guo-Liang Wang^f, Ping Wang^{g,h}, Bo Zhou^{c,i}, and Zhengguang Zhang^{a,b,1}

^aDepartment of Plant Pathology, College of Plant Protection, Nanjing Agricultural University, 210095 Nanjing, China; ^bKey Laboratory of Integrated Management of Crop Diseases and Pests, Ministry of Education, College of Plant Protection, Nanjing Agricultural University, 210095 Nanjing, China; ^cGenetics and Biotechnology Division, International Rice Research Institute, Los Baños, 4031 Laguna, Philippines; ^dCollege of Plant Protection, Jilin Agricultural University, 130118 Changchun, China; ^eKey Laboratory of Agricultural Environmental Microbiology, Ministry of Agriculture, College of Life Sciences of Nanjing Agricultural University, 210095 Nanjing, China; ^fDepartment of Plant Pathology, The Ohio State University, Columbus, OH 43210; ^gDepartment of Pediatrics, Louisiana State University Health Sciences Center, New Orleans, LA 70112; ^hDepartment of Microbiology, Immunology, and Parasitology, Louisiana State University Health Sciences Center, New Orleans, LA 70112; and ⁱInternational Rice Research Institute–Jiangsu Academy of Agricultural Sciences Joint Laboratory, Jiangsu Academy of Agricultural Sciences, 210095 Nanjing, China

Edited by Sheng Yang He, Department of Energy, Plant Research Laboratory, Michigan State University, East Lansing, MI, and approved July 22, 2019 (received for review March 25, 2019)

Environmental conditions are key factors in the progression of plant disease epidemics. Light affects the outbreak of plant diseases, but the underlying molecular mechanisms are not well understood. Here, we report that the light-harvesting complex II protein, LHCB5, from rice is subject to light-induced phosphorylation during infection by the rice blast fungus *Magnaporthe oryzae*. We demonstrate that single-nucleotide polymorphisms (SNPs) in the *LHCB5* promoter control the expression of *LHCB5*, which in turn correlates with the phosphorylation of LHCB5. LHCB5 phosphorylation enhances broad-spectrum resistance of rice to *M. oryzae* through the accumulation of reactive oxidative species (ROS) in the chloroplast. We also show that LHCB5 phosphorylation-induced resistance is inheritable. Our results uncover an immunity mechanism mediated by phosphorylation of light-harvesting complex II.

light-harvesting complex II | phosphorylation | *Magnaporthe oryzae* | broad-spectrum resistance

Environmental conditions strongly influence pathogen–host plant interactions, and changes in a range of environmental factors can directly determine the outcomes of such interactions. In the 1960s, George McNew proposed a “disease triangle” concept, comprising the host, pathogen, and environment as the 3 fundamental components that affect crop diseases. Many studies have subsequently demonstrated that environmental conditions, such as light, temperature, and humidity, can affect disease development and plant innate immunity. For example, the wheat (*Triticum aestivum*) *Yr36* gene confers broad-spectrum resistance to stripe rust at high temperatures, but not at low temperatures (1), while moderate temperatures have been shown to enhance pathogen-associated molecular patterns (PAMPs)-triggered immunity (PTI) in *Arabidopsis thaliana* (2). Other studies have shown that high humidity promotes bacterial infection by also modulating plant immunity (3). Rice (*Oryza sativa*) blast caused by *Magnaporthe oryzae* is often triggered by the overcast weather and lack of sunlight conditions. A study documenting rice blast of over 55 y in Jiangsu, China, showed that fewer hours of sunlight directly correlated with the seriousness of the blast disease (4, 5). However, the underlying molecular mechanisms linking light to infection are not well understood.

Notably, chloroplasts are not only the organelles where light sensing and photosynthesis take place, but also the site of biosynthesis of defense-related molecules, such as salicylic acid (SA) and jasmonic acid (JA), as well as secondary messengers, including calcium and reactive oxygen species (ROS) (6–9). Photosynthetic electron transduction in the chloroplast can result in the reduction of O₂ and the formation of ROS, such as singlet oxygen (¹O₂), which is generated from the chlorophyll and photosystems II (PSII) antenna complex (10). PSII, which is central to

the conversion of light into chemical energy, is surrounded by the light-harvesting complex II (LHCII) and the monomeric light-harvesting proteins LHCB4 (CP29), LHCB5 (CP26), and LHCB6 (CP24) (11, 12). It has been established that LHCB4, LHCB5, and LHCB6 are all involved in energy dissipation in *A. thaliana* (13, 14) and that LHCB5 functions as a trimeric protein complex rescuing a defect in LHCII function due to the absence of LHCB1 and LHCB2 (15). However, in contrast to its known function in energy transduction, the role of LHCII in plant immunity has not been well elucidated.

Results

LHCB5 Is Involved in Light Regulation of Resistance to *M. oryzae* in Different Rice Varieties. To investigate whether rice resistance to *M. oryzae* is light-dependent, 25 rice varieties, collected worldwide, were screened for blast resistance under 2 different light intensities. These varieties showed various degrees of susceptibilities to the wild-type *M. oryzae* Guy11 under the low light intensity of 50 μmol photons m⁻²s⁻¹ (Fig. 1), based on both the

Significance

Environmental conditions strongly influence pathogen–host plant interactions. Rice blast erupts in overcast and rainy conditions, due not only to favorable environmental conditions but also to insufficient light, which reduces host resistance. Due to the importance of breeding of blast-resistant rice varieties, elucidation of light-regulated rice immunity is an important research goal. We revealed that light induces protein phosphorylation of a harvesting complex II protein, LHCB5, upon infection by the rice blast fungus *Magnaporthe oryzae*. Resistance governed by LHCB5 phosphorylation cosegregates with the progenies harboring the desirable haplotype promoter. This establishes the genetic basis of LHCB5-regulated resistance mediated by phosphorylation. Our study highlights a mechanism for light-dependent rice blast resistance that promises future breeding of blast-resistant rice varieties.

Author contributions: M.L., W.S., X.Z., Z.C., G.-L.W., P.W., B.Z., and Z.Z. designed research; M.L., S.Z., B.Z., and Z.Z. performed research; M.L. and Z.Z. contributed new reagents/analytic tools; M.L., S.Z., J.H., J.P., Y.H., Y.L., Z.Y., X.L., W.W., D.S., D.L., H.Z., B.Z., and Z.Z. analyzed data; and M.L. and Z.Z. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Published under the PNAS license.

¹To whom correspondence may be addressed. Email: zhgzhang@njau.edu.cn.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1905123116/-DCSupplemental.

Published online August 12, 2019.




Fig. 1. Identification of blast-resistant rice varieties in a light-dependent manner. (A) The wild-type TP309 and 25 rice varieties collected worldwide were used for blast resistance screen. The rice varieties inoculated with Guy11 were cultivated in a light incubator with different light intensity (50 and 200 $\mu\text{mol photons m}^{-2}\text{s}^{-1}$). Red boxes indicate rice varieties with resistant variations under different light intensity. (B) Disease lesion area (DLA) was assessed by Image J. Lesions were photographed and measured or scored at 7 d postinoculation (dpi). The experiments were repeated twice with similar results. (C) Fungal growth and severity of blast were evaluated by quantifying *M. oryzae* genomic 28S rDNA relative to rice genomic Rubq1 DNA. The mean values of 3 determinations with SDs are shown. The asterisks indicate a significant difference according to Student's *t* test ($P < 0.01$).

diseased leaf area (DLA) and the pathogen biomass assay. However, 8 varieties showed significantly increased resistance when the assessment was conducted under the higher light intensity of 200 $\mu\text{mol photons m}^{-2}\text{s}^{-1}$ (Fig. 1). In contrast, the remaining 17 varieties did not show any significant differences in light-associated blast resistance (Fig. 1).

This light intensity-associated resistance prompted us to investigate the possible roles of the LHCII proteins, which are required for absorbing and transferring the light energy (16, 17). We first tested the transcript levels of 6 *LHCB* genes in the nonresponsive TP309 variety rice and found that all of them were significantly up-regulated under the lighted conditions, compared to the dark, suggesting that all *LHCB* genes are light-inducible (SI Appendix, Fig. S1). We also observed that the *LHCB5* transcript level was uniformly up-regulated in all 8 light-dependent resistant varieties, but not in TP309 (SI Appendix, Fig. S2), as was the accumulation of *LHCB5* protein (SI Appendix, Fig. S3). *LHCB5* transcript levels gradually increased at different time points until 96 h postinoculation (hpi) (SI Appendix, Fig. S4), and this accumulation also corresponded with the levels of the *LHCB5* protein. These results indicated that *LHCB5* is light-

and blast pathogen-responsive and may play a role in light-responsive blast resistance.

***LHCB5*-Mediated Resistance Is Highly Associated with Promoter Sequence Divergence Leading to Different Expression Levels among Diverse Rice Germplasm.** The various levels of the *LHCB5* transcript and protein in different rice varieties prompted us to investigate the variance of its promoter and the coding sequences from 3,000 sequenced rice genomes (18). No nonsynonymous sequence mutations were found in *LHCB5* coding sequences, suggesting high conservation (Dataset S1). In contrast, 11 single-nucleotide polymorphisms (SNPs) were identified that were separated into 18 groups (SNP ≥ 1) using *O. sativa* ssp *japonica* cv. Nipponbare (NPB) as the control (SNP = 0) (Dataset S2). The SNPs were mainly discovered in the *indica* rice varieties, and most of these varieties showed variations in 7 of the SNPs identified (SI Appendix, Fig. S5A and B). The difference in SNP variation suggested a causal relationship with a lower expression in *indica* than in *japonica* (SI Appendix, Fig. S6A). To test this hypothesis, 238 rice varieties were selected for analyzing the relationship between SNPs and transcription levels (Dataset S3). The expression of *LHCB5* in SNP variation rice lines (SNP ≥ 1) was significantly lower than that in nonvariation varieties (SNP = 0) (SI Appendix, Fig. S5C), and this was, in particular, true among 7 of the 11 SNP positions (1–4, 9–11) (SI Appendix, Fig. S5D). To validate the correlation between SNPs and the *LHCB5* expression levels, we expressed GFP (green fluorescent protein) reporter gene driven by different promoters (p35S:GFP, pLHCB5^{SNP=0}:GFP and pLHCB5^{SNP=7}:GFP) in rice protoplast. The result showed that constructs p35S:GFP and pLHCB5^{SNP=0}:GFP had a much higher level of green fluorescent; the Western blot also showed a higher GFP protein level under the control of 35S and SNP = 0 promoter than SNP = 7 promoter, indicating that the promoter SNPs are indeed associated with higher *LHCB5* expression (SI Appendix, Fig. S7). When 238 rice varieties were inoculated with 2 rice blast isolates, Mo15-125 and Mo15-9, a correlation between higher *LHCB5* expression and smaller disease lesion area was found (SI Appendix, Fig. S8). These data indicated that SNP variations in *indica* rice varieties result in reduced *LHCB5* expression and lower resistance to blast.

To further investigate the genetic association between the SNP pattern and expression of *LHCB5*, we used an F2 population of 44 progenies derived from a cross between YG456 (Yangeng456, SNP = 0) and LTH (Lijiangxintuanheigu, SNP ≥ 1 , containing all 7 candidate SNPs mentioned above). As indicated in Fig. 2A and B, the parental line YG456 and 32 progenies containing the YG456-type promoter showed significantly higher expression levels than LTH and the 12 progenies containing the LTH-type promoter. This revealed a cosegregation between the expression level and the promoter type in the F2 population.

YG456 was also found to be resistant to the virulent rice blast isolate Guy11, which had a lesion length 40% smaller than the lesion of the susceptible rice variety LTH (Fig. 2C and F). Moreover, all of the progenies with a higher *LHCB5* expression level showed resistance to Guy11, whereas those with a lower *LHCB5* expression level were susceptible (Fig. 2E and F). Taken together, these results revealed that the elevated level of *LHCB5* governed by the promoter (SNP = 0) contributes to enhanced resistance to blast, and the trait is genetically transferable.

Expression of *LHCB5* in Transgenic Rice Lines Correlates with Resistance to Infection. To further investigate the relationship between *LHCB5* expression and resistance, we characterized transgenic rice lines with expression differences. We first generated transgenic *LHCB5* RNAi-silenced (*lhcb5*-RNAi) and over-expression (*LHCB5*-OX) rice lines in the TP309 background. The *lhcb5*-RNAi lines #1 and #2 showed significantly reduced *LHCB5* transcript levels in the leaves, whereas *LHCB5*-OX lines #1 and #2 showed significantly increased levels, compared to TP309 (SI Appendix, Fig. S9A). Western blot analysis showed that the amount of




Fig. 2. Genetic association between the SNP pattern and expression of *LHCBS* and the inheritance of *LHCBS* phosphorylation. (A) The transcript analysis of *LHCBS* in YG456 and LTH. qRT-PCR on *LHCBS* in YG456 in comparison with LTH. (B) The transcript analysis of *LHCBS* in an F2 population derived from a cross between YG456 and LTH. The expression level of 44 progeny was analyzed by qRT-PCR. (C) Blast resistance of LTH and YG456 plants. The leaves of 4-wk-old plants were inoculated using method of punch. Photos were taken at 6 dpi. (D) Detection of *LHCBS* phosphorylation in LTH and YG456. LTH and YG456 plants were inoculated with (infection) or without (noninfection) Guy11 after 48 hpi. The protein extracts were subjected to Phos-tag SDS/PAGE and normal SDS/PAGE followed by immunoblotting with the anti-*LHCBS* polyclonal antibody. (E) Rice blast resistance is correlated with phosphorylation assess in the F2 generation of LTH and YG456. Blast resistance of 44 F2 generations using punch inoculation. Phosphorylation assay was performed as above. (F) Lesion length was measured 6 dpi. Values are the means of 3 replications, and error bars represent the SD ($n = 3$). The asterisks indicate a significant difference according to Student's *t* test ($P < 0.01$).

the *LHCBS* protein was 3- to 4-fold higher in the *LHCBS*-OX lines than in TP309, while the *lhcbs*-RNAi lines had 20 to 25% of the levels of TP309 (SI Appendix, Fig. S9B).

In infection, the *LHCBS*-OX rice lines showed strong resistance to blast, with punctate and significantly reduced lesion areas, whereas the *lhcbs*-RNAi lines were more susceptible (Fig. 3 A and B). In addition, lesions on *LHCBS*-OX lines failed to produce any conidia (Fig. 3C). A rice sheath infection assay for invasive hyphal growth (19) revealed that >90% of penetration was at the level I in the *LHCBS*-OX rice lines, in contrast to 11% and 10% in TP309 and *lhcbs*-RNAi lines at 24 hpi, respectively. The infectious hyphae on the *LHCBS*-OX rice lines failed to expand, even at 48 hpi, and showed level II penetration (Fig. 3 D and E). Furthermore, we used the CRISPR/Cas9 technology to knock out the endogenous *LHCBS* gene (*LHCBS*-KO) in TP309, and the infection phenotype of *LHCBS*-KO lines was as susceptible as *lhcbs*-RNAi lines (SI Appendix, Fig. S10). Together, these data indicated that *LHCBS* overexpression enhances basal resistance.

Induced Immune Response in *LHCBS*-OX Lines Is Broad Spectrum to *M. oryzae*. To explore the mechanism by which *LHCBS* mediates resistance, we inoculated TP309, *LHCBS*-OX, and *lhcbs*-RNAi

leaves with Guy11 and the incompatible strain 51# and stained the leaf cells with 3, 3'-diamino-benzidine (DAB) and Trypan Blue (TB). We observed that *LHCBS*-OX accumulated higher levels of H_2O_2 , accompanied by cell death (SI Appendix, Fig. S11A). We also used a luminol-based chemiluminescence assay to monitor the generation of ROS induced by purified mycelia (PM) as the elicitor (20) and obtained a similar result (SI Appendix, Fig. S11B). Finally, we performed a rice sheath infection assay to examine ROS accumulation and cell death and found that the DAB and TB stain signals were 4 to 5 times stronger in *LHCBS*-OX sheaths than in those of TP309 and *lhcbs*-RNAi. Pretreating *LHCBS*-OX rice sheaths with the catalase of *Aspergillus niger* (CAG) (21) and reduced glutathione (GSH) (22) partially suppressed the production of ROS and rescued the invasive growth (SI Appendix, Fig. S11 C-E). Moreover, qRT-PCR revealed that the expression levels of PR genes, including *PR1*, *PBZ1*, *AOS2*, and *LOX1*, as well as 2 NADPH oxidases (*RBOHA* and *RBOHB*), were significantly up-regulated in the *LHCBS*-OX lines (SI Appendix, Fig. S11 F and G).

We next examined TP309, *lhcbs*-RNAi, and *LHCBS*-OX resistant to 21 *M. oryzae* isolates containing various *AVR* genes (SI Appendix, Fig. S12) (23) and found that the *LHCBS*-OX lines showed greater resistance to all 21 isolates (SI Appendix, Fig. S13). To investigate whether these lines have also been more resistant to other pathogens, we inoculated them with the pathogenic fungus *Bipolaris oryzae* and the pathogenic bacterium *Xanthomonas oryzae* pv *oryzae* PXO99. We observed that the *LHCBS*-OX lines were not resistant to either of these pathogens (SI Appendix, Fig. S14A), suggesting that the basal resistance in




Fig. 3. The function of *LHCBS* in blast resistance. (A) Blast resistance of *lhcbs*-RNAi and *LHCBS*-OX plants using spraying inoculation. Numbers "1" and "2" mean 2 independent transformants. (B) The DLA of the leaves infected by Guy11 was assessed by Image J. Lesions were photographed and measured or scored at 7 dpi, and experiments were repeated twice with similar results. Values are the means of 3 replications, and error bars represent the SD ($n = 3$). The asterisks indicate a significant difference according to Student's *t* test ($P < 0.01$). (C) Assays for fungal growth on surface-sterilized rice leaves inoculated by spray with Guy11. (D) Typical infection sites of TP309, *lhcbs*-RNAi, and *LHCBS*-OX leaf sheath inoculated with Guy11 strain, showing greater fungal proliferation and tissue invasion in TP309 and *lhcbs*-RNAi by the wild-type strain and restricted in *LHCBS*-OX leaf sheath. Infectious growth was observed at 30 hpi. (Scale bar, 10 μ m). (E) Statistics of invasive hyphal growth at 100 appressorium penetration sites by rating the hyphal growth from level I to IV (I, no penetration; II, with primary invasive hypha; III, secondary invasive hypha does not extend to the neighboring plant cells; IV, invasive hypha extended into neighboring plant cells).

the *LHCBS*-OX lines, associated with ROS production, was *M. oryzae*-specific.

LHCBS Phosphorylation Regulates Basal Immunity Independent of *R* Genes *Pia* and *Pizt*. Previous studies have suggested that LHCII is phosphorylated to balance light excitation energy between photosystem I and photosystem II (24). In the alga *Chlamydomonas reinhardtii*, a marked increase in phosphorylation of the chlorophyll protein CP26 (*LHCBS* homolog) was observed in state II (25). We analyzed the phosphorylation of *LHCBS* in TP309 and *LHCBS*-OX lines that were uninfected, or that had been infected by Guy11, using Mn²⁺-Phos-tag gel electrophoresis (26). We found that *LHCBS* phosphorylation was induced by Guy11 in the *LHCBS*-OX lines, but not in TP309 (Fig. 4A), and that this phosphorylation was *M. oryzae*-specific (SI Appendix, Fig. S14B). To further test whether the *LHCBS* phosphorylation was light-dependent, we examined *LHCBS* phosphorylation at various light intensities and found that it was induced at 100 and 200 μmol photons m⁻²s⁻¹, but not below 50 μmol photons m⁻²s⁻¹ (Fig. 4B).

We next carried out a blast-resistance assay with 14 randomly selected lines out of 3,000 rice accessions (18) and 45 lines of the Chinese origin to determine whether phosphorylation was associated with resistance. We observed that only *LHCBS*-phosphorylated lines exhibited resistance (Fig. 4C), and a total of 12 lines showed no observable lesions, indicative of the typical *R* resistance gene-mediated immune response (corresponding to the AVR protein in Guy11) (Fig. 4C). Further, we observed that this correlates with *LHCBS* not being phosphorylated during interactions between *Avr-Pia/Pia* and *AvrPiz-t/Piz-t* (Fig. 4D). These findings suggested that *LHCBS* phosphorylation regulates basal immunity and that the resistance is independent of the *R* genes *Pia* and *Pizt*.

Finally, we examined the expression of *LHCBS* in the above-mentioned 47 rice lines (excluding the 12 lines showing no lesions) and found higher *LHCBS* expression levels correlated with smaller lesion sizes (0.622 Pearson Correlation Coefficient) (Fig. 4E), which is consistent with the results shown in SI Appendix, Fig. S7. Since *LHCBS* expression was significantly reduced in *indica* compared to *japonica* (SI Appendix, Fig. S6A), we next determined whether *LHCBS* phosphorylation similarly differed,




Fig. 4. Phosphorylation of *LHCBS* regulates blast resistance. (A) *LHCBS* was phosphorylated in *LHCBS*-OX plants inoculated with Guy11. Proteins from TP309 and the *LHCBS*-OX cell extracts were subjected to Phos-tag SDS/PAGE and normal SDS/PAGE followed by immunoblotting with *LHCBS* polyclonal antibody. (B) *LHCBS* phosphorylation was light-dependent. The *LHCBS*-OX plants inoculated with or without Guy11 were cultivated in a light incubator with different light intensity (0, 50, 100, and 200 μmol photons m⁻²s⁻¹). (C) The phosphorylation is correlated with blast resistance. Blast resistance of 59 rice varieties using spraying inoculation. The phosphorylation of *LHCBS* was detected at 48 hpi. Lesions were photographed and measured or scored at 7 dpi. (D) Phosphorylation assay of *LHCBS* during interaction between *Avr-Pia/Pia* and *AvrPiz-t/Piz-t*. *LTH-Pia* and *LTH-Pizt* plants were inoculated with Guy11/*Avr-Pia* and Guy11/*AvrPiz-t*, respectively. (E) The expression level is associated with blast resistance. The DLA was measured by Image J, and the expression (Exp) level was detected by qRT-PCR. The Pearson correlation coefficient was -0.622. Values are the means of 3 replications, and error bars represent the SD ($n = 3$).

which we tested using 9 high and low *LHCb5* expression varieties from each subgroup. After *M. oryzae* infection, LHCb5 phosphorylation was found to mainly occur in the highly expressing *japonica* varieties (SI Appendix, Fig. S6B).

Phosphorylation of LHCb5 Correlates with Enhanced Resistance of Progeny Containing the YG456-Type Promoter. To confirm the relationship between the phosphorylation of LHCb5 and resistance to *M. oryzae*, we investigated LHCb5 phosphorylation in the same F2 population as used for the genetic association experiments. First, we analyzed the phosphorylation of LHCb5 in the LTH and YG456 lines, with or without infection, and found that LHCb5 was phosphorylated in infected YG456 (Fig. 2D). Of all 42 progenies, LHCb5 phosphorylation was only detected in resistant progenies (Fig. 2E), indicating that these traits cosegregated. Taken together with the observed close association between the YG456-type promoter and resistance to rice blast, we speculate that the resistance governed by the phosphorylation of LHCb5 is genetically controlled by the *LHCb5* promoter in YG456.

Phosphorylation of LHCb5 24th Threonine (T24) Activates an ROS Burst in Chloroplasts. To study the phosphorylation mechanism, we first compared the LHCb5 sequence to those of other plant species and predicted 3 possible phosphorylation sites at the less conserved N-terminal domain, using DISPHOS 1.3 and NetPhos 3.1 server (SI Appendix, Fig. S15A). We showed that the constitutively activated LHCb5 T24 (LHCb5^{T24D}) allele induced ROS production and cell death in *Nicotiana benthamiana* leaves, in contrast to the nonphosphorylated LHCb5 (LHCb5^{T24A}) (SI Appendix, Fig. S15 B and C), or LHCb5 with constitutive phosphorylation at 2 other sites (SI Appendix, Fig. S16). The phosphorylation of LHCb5 was observed in the cytoplasm, but not in the chloroplast (SI Appendix, Fig. S17). We further expressed the rice LHCb5 and LHCb5^{T24A} protein with a FLAG peptide tag at the C terminus (OsLHCb5:Flag and OsLHCb5^{T24A}:Flag) in *lhc5*-RNAi protoplasts induced by PM and detected mobility shifts representing phosphorylated LHCb5 in OsLHCb5, but not OsLHCb5^{T24A} (SI Appendix, Fig. S15D). Using a luminol-based chemiluminescence assay, we observed higher ROS levels in the presence of constitutively activated LHCb5 (LHCb5^{T24D}) than with inactivated LHCb5 (LHCb5^{T24A}) in protoplasts (SI Appendix, Fig. S15E). These results further suggested that LHCb5 24T phosphorylation is important for LHCb5 function.

To confirm the elevated production of ROS in chloroplasts, we expressed the empty plasmid pBIN:GFP (as a marker for subcellular localization), as well as OsLHCb5^{T24D}:GFP, and OsLHCb5^{T24A}:GFP in rice protoplasts, and found that both OsLHCb5^{T24D}:GFP and OsLHCb5^{T24A}:GFP were localized to the chloroplast. Nitroblue tetrazolium (NBT) staining revealed an accumulation of superoxide (O₂^{•-}) in OsLHCb5^{T24D}:GFP expressing protoplasts, but not in pBIN:GFP or OsLHCb5^{T24A}:GFP expressing protoplasts (SI Appendix, Fig. S15F), consistent with LHCb5 phosphorylation that results in O₂^{•-} accumulation in the chloroplast and activation of the immune response.

Phosphorylation Facilitates LHCb5 Accumulation and Trimerization in Chloroplasts. We noticed that the phosphorylation site was located within the region of the LHCb5 polypeptide responsible for chloroplast transit, which is important for the import of precursor proteins (27, 28). To investigate whether phosphorylation influences the accumulation of LHCb5 in the chloroplast, we separated chloroplast and cytoplasmic protein fractions extracted from TP309, *lhc5*-RNAi, and *LHCb5*-OX lines, with or without Guy11 inoculation. Western blot analysis indicated the increased accumulation of LHCb5 in the chloroplasts of *LHCb5*-OX lines inoculated with Guy11 (SI Appendix, Fig. S18). Furthermore, we expressed pBIN-LHCb5-Flag, pBIN-LHCb5^{T24A}-Flag, and pBIN-LHCb5^{T24D}-Flag in the *LHCb5*-KO rice protoplast treated with or without PM. As shown in SI Appendix, Fig. S19 A and B, inactivated LHCb5 (LHCb5^{T24A}) cannot stimulate

host immunity. Moreover, we found that pBIN-LHCb5-Flag and pBIN-LHCb5^{T24D}-Flag showed higher protein accumulation in chloroplast than pBIN-LHCb5^{T24A}-Flag when treated with PM (SI Appendix, Fig. S19C). Together, the results indicated that phosphorylation of 24th threonine facilitates LHCb5 accumulation in chloroplast to stimulate the immune response.

Since a previous study found that *A. thaliana* LHCb5 is organized into trimeric complexes in the absence of LHCb1 and LHCb2 (15), we next tested whether increased LHCb5 accumulation also led to the accumulation of trimers. Using native-PAGE analysis, we found that LHCb5 is mostly present in the form of monomers and dimers in TP309 and that trimeric forms were only founded in infected *LHCb5*-OX lines (SI Appendix, Fig. S20A). When a His-tagged form of LHCb5, LHCb5-His, was expressed in *Escherichia coli* BL21 and then purified (SI Appendix, Fig. S20B), fractionated, and analyzed by native-PAGE analysis, we also observed LHCb5-His in a trimeric complex (SI Appendix, Fig. S20C). Together, these data indicated that phosphorylation facilitates the accumulation of LHCb5 into chloroplasts and that LHCb5 exists in multimeric conformations, including a trimer.

Trimeric LHCb5 Conformation Affects PsbS Binding and the Electron Transport Rate in Chloroplasts. It was reported that rice plants deficient in the PSII protein PsbS had higher levels of chloroplastic superoxide and hydrogen peroxide and were more resistant to *M. oryzae* infection (29, 30). In *A. thaliana*, PsbS binds to LHCb5 to control nonphotochemical quenching (NPQ) (31). We therefore tested the interaction between LHCb5 and PsbS in the *LHCb5*-OX lines and found that this is indeed also the case in rice. However, the Co-IP analysis showed that the LHCb5-PsbS interaction does not occur upon infection by Guy11 (SI Appendix, Fig. S21). We hypothesized that multimeric forms of LHCb5 interact less with PsbS than does the LHCb5 monomer. To test this, we performed an in vitro pull-down assay with varying amounts of LHCb5. Consistent with our hypothesis, PsbS binding decreased as LHCb5 concentration increased (SI Appendix, Fig. S22).

To validate O₂^{•-} accumulation in the chloroplast, we measured photosynthetic parameters in the TP309, G-TP309 (Guy11-infected), *lhc5*-RNAi, G-*lhc5*-RNAi (Guy11-infected), *LHCb5*-OX, and G-*LHCb5*-OX (Guy11-infected) lines using a Chlorophyll Fluorescence Imager (Ecotek, Beijing). The electron transport rate (ETR) was significantly reduced in G-*LHCb5*-OX lines (SI Appendix, Fig. S23A). Moreover, a transmission microscopic analysis of chloroplast morphology revealed that the structure was less compact in G-TP309 and G-*lhc5*-RNAi lines when compared to noninfected lines, while no difference was detected in the G-*LHCb5*-OX lines (SI Appendix, Fig. S23B). These data suggested that LHCb5 may form trimeric complexes during infection to maintain normal morphology and may affect PsbS binding, thereby influencing O₂^{•-} accumulation.

Discussion

Rice blast threatens rice production worldwide, and serious outbreaks can destroy whole crop harvests. Rice blast erupts in overcast and rainy conditions, due not only to favorable environmental conditions but also to insufficient light, which reduces host resistance (32). The mechanistic basis of the relationship between resistance and light levels has not been elucidated until now. Our study establishes the relationships among light conditions, light-harvesting complex protein LHCb5, and disease resistance.

It is known that transcription is often associated with SNP variations in the promoter and that nucleotide polymorphism in either the promoter or gene can affect resistance (33, 34). For example, it was reported that an A to G transition in the promoter region of the *BSR-D1* gene results in lower expression, resulting in elevated resistance (35). A survey of SNPs among *LHCb5* promoter sequences of 3,000 sequenced rice genomes suggested that SNP variation is associated with the differential

expression of *LHCb5* and, accordingly, resistance (Fig. 4E and *SI Appendix*, Fig. S8). It has been demonstrated that *japonica* rice varieties from the Chinese Yuanyang terraces have higher basal immunity, but a lower content of major resistance (*Pi*) genes than do *indica* varieties (36). Our results suggested that *LHCb5* expression was higher in *japonica* (*SI Appendix*, Fig. S6A). Moreover, a higher *LHCb5* expression leads to its phosphorylation upon challenge with the blast fungus, particularly in *japonica* (*SI Appendix*, Fig. S6B). We propose that SNP variation among *japonica* and *indica* varieties causes differences in *LHCb5* expression patterns, resulting in the differential phosphorylation of *LHCb5*. This is consistent with the elevated basal immunity observed in *japonica* rice. Whereas the evidence for a direct connection remains to be established, our studies suggest a strong correlation between promoter SNP and *LHCb5* phosphorylation.

Biotic stress typically stimulates the production of ROS, such as $O_2^{\cdot-}$, in the chloroplast, (37) and ROS generation is associated with resistance to rice blast (35). Here, we showed that the ROS burst in the chloroplast is correlated with the phosphorylation of *LHCb5*, and that mediated broad-spectrum blast resistance (*SI Appendix*, Fig. S15). A widely used strategy to protect rice against pathogens is through the breeding of resistant varieties (38, 39), despite that the rapid variation of rice blast can lessen the effectiveness of this approach (40). To evaluate *LHCb5* phosphorylation-induced resistance, we assessed the inheritance of this trait and found that it cosegregated with resistance (Fig. 2). Our studies indicated no differences between *LHCb5* transgenic and wild-type rice lines in the plant height, grain weight, and seed-setting rate (*SI Appendix*, Fig. S24), which

promises for future breeding of high yield and blast-resistant rice varieties.

We propose a model where during the interaction between rice and *M. oryzae*, the host monitors the progression of infection through the phosphorylation of *LHCb5* in a light-dependent manner. *M. oryzae* challenge results in phosphorylated *LHCb5* accumulation in chloroplasts, which helps maintain chloroplast function by reducing the binding of *LHCb5* to PsbS, which in turn leads to reduced ETR for ROS production (*SI Appendix*, Fig. S25).

Materials and Methods

The plant strains and blast isolates are listed in *SI Appendix, Materials and Methods*. Resistance test and infection assessment were performed as described (19). The ROS and cell death observation was performed as described (19). The phosphorylation assay was performed as described (41). Purified recombinant proteins from *Escherichia coli* were used for in vitro pull-down assays (41). All other materials and details of experimental methods can be found in *SI Appendix, Materials and Methods*.

ACKNOWLEDGMENTS. This research was supported by the Key program of Natural Science Foundation of China (NSFC) (Grant No: 31530063, to Z.Z.), Innovation Team Program for NSFC (Grant No: 31721004), Innovation Team Program for Jiangsu Universities (2017), and Jiangsu Agricultural Science and Technology Innovation Fund (CX[18]1003). B.Z. acknowledges the support of the Swiss Programme for Research on Global Issues for Development (IZ07Z0-160877). We thank Dr. Jianmin Wan of Nanjing Agricultural University and Dr. Zhongqiang Qi of Jiangsu Academy of Agricultural Sciences for providing rice seeds.

- D. Fu *et al.*, A kinase-START gene confers temperature-dependent resistance to wheat stripe rust. *Science* **323**, 1357–1360 (2009).
- C. Cheng *et al.*, Plant immune response to pathogens differs with changing temperatures. *Nat. Commun.* **4**, 2530 (2013).
- X. F. Xin *et al.*, Bacteria establish an aqueous living space in plants crucial for virulence. *Nature* **539**, 524–529 (2016).
- Y. Ren, P. Gao, F. Zhu, Meteorological grade prediction of occurrence degree of rice blast in Jiangsu. *Jiangsu Agr. Sci.* **44**, 151 (2016).
- H. Zhang, X. Zheng, Z. Zhang, The *Magnaporthe grisea* species complex and plant pathogenesis. *Mol. Plant Pathol.* **17**, 796–804 (2016).
- D. A. Dempsey, A. C. Vlot, M. C. Wildermuth, D. F. Klessig, Salicylic Acid biosynthesis and metabolism. *Arabidopsis Book* **9**, e0156 (2011).
- S. Fonseca *et al.*, (+)-7-iso-Jasmonoyl-L-isoleucine is the endogenous bioactive jasmonate. *Nat. Chem. Biol.* **5**, 344–350 (2009).
- M. A. Strawn *et al.*, *Arabidopsis* isochlorismate synthase functional in pathogen-induced salicylate biosynthesis exhibits properties consistent with a role in diverse stress responses. *J. Biol. Chem.* **282**, 5919–5933 (2007).
- A. Shapiguzov, J. P. Vainonen, M. Wrzaczek, J. Kangasjärvi, ROS-talk—How the apoplast, the chloroplast, and the nucleus get the message through. *Front. Plant Sci.* **3**, 292 (2012).
- P. Pospíšil, Molecular mechanisms of production and scavenging of reactive oxygen species by photosystem II. *Biochim. Biophys. Acta* **1817**, 218–231 (2012).
- B. Hankamer *et al.*, Isolation and biochemical characterisation of monomeric and dimeric photosystem II complexes from spinach and their relevance to the organisation of photosystem II in vivo. *Eur. J. Biochem.* **243**, 422–429 (1997).
- E. J. Boekema, J. F. van Breemen, H. van Roon, J. P. Dekker, Arrangement of photosystem II supercomplexes in crystalline macrodomains within the thylakoid membrane of green plant chloroplasts. *J. Mol. Biol.* **301**, 1123–1133 (2000).
- J. Andersson, R. G. Walters, P. Horton, S. Jansson, Antisense inhibition of the photosynthetic antenna proteins CP29 and CP26: Implications for the mechanism of protective energy dissipation. *Plant Cell* **13**, 1193–1204 (2001).
- S. de Bianchi, L. Dall'Osto, G. Tognon, T. Morosinotto, R. Bassi, Minor antenna proteins CP24 and CP26 affect the interactions between photosystem II subunits and the electron transport rate in grana membranes of *Arabidopsis*. *Plant Cell* **20**, 1012–1028 (2008).
- A. V. Ruban *et al.*, Plants lacking the main light-harvesting complex retain photosystem II macro-organization. *Nature* **421**, 648–652 (2003).
- S. Jansson, The light-harvesting chlorophyll a/b-binding proteins. *Biochim. Biophys. Acta* **1184**, 1–19 (1994).
- P. Horton, A. V. Ruban, R. G. Walters, Regulation of light harvesting in green plants. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **47**, 655–684 (1996).
- J. Y. Li, J. Wang, R. S. Zeigler, The 3,000 rice genomes project: New opportunities and challenges for future rice research. *Gigascience* **3**, 8 (2014).
- Z. Qi *et al.*, The syntaxin protein (MoSyn8) mediates intracellular trafficking to regulate conidiogenesis and pathogenicity of rice blast fungus. *New Phytol.* **209**, 1655–1667 (2016).
- A. R. Ayers, J. Ebel, B. Valent, P. Albersheim, Host-pathogen interactions: X. Fractionation and biological activity of an elicitor isolated from the mycelial walls of *phytophthora megasperma* var. *sojae*. *Plant Physiol.* **57**, 760–765 (1976).
- S. Tanabe, Y. Nishizawa, E. Minami, Effects of catalase on the accumulation of H_2O_2 in rice cells inoculated with rice blast fungus, *Magnaporthe oryzae*. *Physiol. Plant.* **137**, 148–154 (2009).
- J. Li *et al.*, Hydrogen peroxide regulates elicitor PB90-induced cell death and defense in non-heading Chinese cabbage. *Physiol. Mol. Plant Pathol.* **67**, 220–230 (2005).
- Q. H. Chen, Y. C. Wang, X. B. Zheng, Genetic diversity of *Magnaporthe grisea* in China as revealed by DNA fingerprint haplotypes and pathotypes. *J. Phytopathol.* **154**, 361–369 (2006).
- S. Lemeille *et al.*, Analysis of the chloroplast protein kinase Stt7 during state transitions. *PLoS Biol.* **7**, e45 (2009).
- N. Depège, S. Bellafiore, J. D. Rochaix, Role of chloroplast protein kinase Stt7 in LHClI phosphorylation and state transition in *Chlamydomonas*. *Science* **299**, 1572–1575 (2003).
- E. Kinoshita, E. Kinoshita-Kikuta, K. Takiyama, T. Koike, Phosphate-binding tag, a new tool to visualize phosphorylated proteins. *Mol. Cell. Proteomics* **5**, 749–757 (2006).
- G. Lamberti, C. Drurey, J. Soll, S. Schwenkert, The phosphorylation state of chloroplast transit peptides regulates preprotein import. *Plant Signal. Behav.* **6**, 1918–1920 (2011).
- T. May, J. Soll, 14-3-3 proteins form a guidance complex with chloroplast precursor proteins in plants. *Plant Cell* **12**, 53–64 (2000).
- I. S. Zulfugarov *et al.*, Production of superoxide from Photosystem II in a rice (*Oryza sativa* L.) mutant lacking PsbS. *BMC Plant Biol.* **14**, 242 (2014).
- I. S. Zulfugarov *et al.*, Enhanced resistance of PsbS-deficient rice (*Oryza sativa* L.) to fungal and bacterial pathogens. *J. Plant Biol.* **59**, 616–626 (2016).
- J. Sacharz, V. Giovagnetti, P. Ungerer, G. Mastroianni, A. V. Ruban, The xanthophyll cycle affects reversible interactions between PsbS and light-harvesting complex II to control non-photochemical quenching. *Nat. Plants* **3**, 16225 (2017).
- Y. Zhu *et al.*, Genetic diversity and disease control in rice. *Nature* **406**, 718–722 (2000).
- W. Bimolata *et al.*, Nucleotide diversity analysis of three major bacterial blight resistance genes in rice. *PLoS One* **10**, e0120186 (2015).
- M. A. Rashid, Y. Zhao, H. Zhang, J. Li, Z. Li, Nucleotide diversity, natural variation, and evolution of flexible culm-1 and Strong culm-2 lodging resistance genes in rice. *Genome* **59**, 473–483 (2016).
- W. Li *et al.*, A natural allele of a transcription factor in rice confers broad-spectrum blast resistance. *Cell* **170**, 114–126.e15 (2017).
- J. Liao *et al.*, Pathogen effectors and plant immunity determine specialization of the blast fungus to rice subspecies. *eLife* **5**, e19377 (2016).
- J. Su *et al.*, Active photosynthetic inhibition mediated by MPK3/MPK6 is critical to effector-triggered immunity. *PLoS Biol.* **16**, e2004122 (2018).
- S. Fukuoka *et al.*, Loss of function of a proline-containing protein confers durable disease resistance in rice. *Science* **325**, 998–1001 (2009).
- B. Zhou *et al.*, The eight amino-acid differences within three leucine-rich repeats between Pi2 and Pi2-t resistance proteins determine the resistance specificity to *Magnaporthe grisea*. *Mol. Plant Microbe Interact.* **19**, 1216–1228 (2006).
- J. Huang, W. Si, Q. Deng, P. Li, S. Yang, Rapid evolution of avirulence genes in rice blast fungus *Magnaporthe oryzae*. *BMC Genet.* **15**, 45 (2014).
- L. Li *et al.*, MoCAP proteins regulated by MoArk1-mediated phosphorylation coordinate endocytosis and actin dynamics to govern development and virulence of *Magnaporthe oryzae*. *PLoS Genet.* **13**, e1006814 (2017).