


 PROFILE

Profile of Julian I. Schroeder

Sandeep Ravindran, *Science Writer*

Julian Schroeder's research career might have unfolded differently if he had not run into Linus Pauling in the coffee room. Schroeder was a graduate student at the Max Planck Institute for Biophysical Chemistry in Göttingen, Germany, and was torn between attending a seminar on plant science and a speech by Linus Pauling on nuclear disarmament. As he pondered this dilemma, Schroeder took a break to grab a cup of tea. "I opened the door to our lab's coffee room, and who's sitting there? Linus Pauling!" says Schroeder. Pauling was by himself, and Schroeder got to discuss everything from nuclear disarmament and physics to chemistry.

Thanks to his chance encounter with Pauling, Schroeder decided to forgo Pauling's speech and attended the plant science seminar, which would set the ball rolling for a long and fruitful career studying plants. Schroeder has spent most of his career characterizing plant ion channels and elucidating the mechanisms by which their regulation helps plants resist environmental stresses. He has described the signaling pathways that regulate stomatal guard cell movements and has improved our understanding of how plants protect themselves from drought, high salt, and heavy metals, with practical implications for improving crop plants. Now a Novartis Distinguished Professor in Plant Sciences at the University of California, San Diego, Schroeder was elected to the National Academy of Sciences in 2015. In his Inaugural Article (IA), Schroeder describes a mechanism by which guard cells sense carbon dioxide, which is relevant for understanding how plants cope with increasing levels of atmospheric carbon dioxide (1).

Circuitous Path to Plant Research

Schroeder says his mother instilled a sense of self-sufficiency in him. She became separated from her Bulgarian/Czech family in Bulgaria during World War II at the age of 15 years and ended up on her own in the West at the end of the war. She eventually immigrated to the United States by herself and has retained a positive and accepting outlook.

Schroeder says he has long been interested in plants, beginning with the time he and a friend cultivated a vegetable garden as undergraduates at the



Julian I. Schroeder. Image courtesy of Erik Jepsen (University of California, San Diego, La Jolla, CA). Copyright Regents of the University of California.

University of Göttingen. "I bought an old truck and converted it into a tiny house," he says. "I lived for a year-and-a-half in this abandoned overgrown apple and plum-tree orchard, and that's where we had our vegetable garden and where I first started working with plants."

It would take a few years before plants became the focus of Schroeder's research. Schroeder studied physics as an undergraduate, and after completing the second-year examination, decided to take a semester off and get a job, an experience that would guide him toward a new interest. "I worked with physically and mentally handicapped persons, and I became fascinated by the question of how the brain works."

Upon resuming his studies, Schroeder started doing internships in biophysics and neurophysiology laboratories. "I realized I could focus on the area of biophysics that looks at the inner workings of neurons," he says. Schroeder began work on a master's degree at the Max Planck Institute with biophysicist Erwin Neher, who had recently developed the patch-clamp technique for studying electric currents in individual cells. Neher would later share the Nobel Prize in Physiology or Medicine with colleague Bert Sakmann for developing this technology. "I was

Published under the [PNAS license](#).

This is a Profile of a recently elected member of the National Academy of Sciences to accompany the member's Inaugural Article on page 11129.

Published online October 15, 2018.

www.pnas.org/cgi/doi/10.1073/pnas.1815757115

PNAS | October 30, 2018 | vol. 115 | no. 44 | 11109–11111

www.manaraa.com

The researchers used a combination of computer-aided molecular dynamics modeling and experimental assays providing evidence for the model that SLAC1 plays an important role in carbon dioxide/bicarbonate sensing in guard cells. They found that mutating some predicted putative bicarbonate-interacting sites in the SLAC1 channel impaired the carbon dioxide/bicarbonate response in *Xenopus* oocytes. Using patch-clamp and gas exchange analyses in transgenic plants, the researchers identified a specific SLAC1 residue required for carbon dioxide/bicarbonate enhancement of slow anion channel activity and carbon dioxide-induced, but not ABA-induced, stomatal closing.

These findings point to SLAC1 being one of the carbon dioxide/bicarbonate sensors that mediates carbon dioxide signaling during control of stomatal movements, and provide important insights into how plants respond to rising carbon dioxide. Understanding the pathways involved may aid the development of plants that can adapt to climate change.

Basic Research with Applications

Studying carbon dioxide in guard cells is not the only avenue of Schroeder's research with practical applications. "We are often motivated by questions that are relevant for present-day problems, and we focus on the fundamental underlying mechanisms," he says.

Schroeder's research on the HKT sodium transporter family has helped plant scientists and breeders boost the salt tolerance of crop plants, such as rice and wheat. "It turns out that this HKT1 transporter mechanism is of major importance in crops for breeding enhanced salt tolerance, as others have now shown," he says.

Schroeder and collaborators also discovered plant transporters that could play a role in how plants accumulate and protect themselves from heavy metals. "Some of the genes and mechanisms could potentially be used to avoid accumulation of toxic metals and arsenic in the edible portions of plants, or could have an impact on removing heavy metals or arsenic from soils," says Schroeder.

Additionally, Schroeder and his colleagues have long been working on how plants resist drought. "The research community has discovered various mechanisms that you can manipulate to obtain plants that are more drought-resistant," he says. Applying these mechanisms to crop plants could have a huge impact, but Schroeder notes that translational research is in its early stages.

"There are many avenues to affect the way a plant responds to drought emanating from this research, but it could take 5 to 10 years or so to see which ones may be working best in combination in which crops in the field," Schroeder says. "Clearly, there will be many useful tools that build on basic discovery and knowledge."

- 1 Zhang J, et al. (2018) Identification of SLAC1 anion channel residues required for CO₂/bicarbonate sensing and regulation of stomatal movements. *Proc Natl Acad Sci USA* 115:11129–11137.
- 2 Schroeder JI, Hedrich R, Fernandez JM (1984) Potassium-selective single channels in guard cell protoplasts of *Vicia faba*. *Nature* 312:361–362.
- 3 Schroeder JI, Raschke K, Neher E (1987) Voltage dependence of K channels in guard-cell protoplasts. *Proc Natl Acad Sci USA* 84:4108–4112.
- 4 Pei ZM, Kuchitsu K, Ward JM, Schwarz M, Schroeder JI (1997) Differential abscisic acid regulation of guard cell slow anion channels in *Arabidopsis* wild-type and *abi1* and *abi2* mutants. *Plant Cell* 9:409–423.
- 5 Schachtman DP, Schroeder JI (1994) Structure and transport mechanism of a high-affinity potassium uptake transporter from higher plants. *Nature* 370:655–658.
- 6 Rubio F, Gassmann W, Schroeder JI (1995) Sodium-driven potassium uptake by the plant potassium transporter HKT1 and mutations conferring salt tolerance. *Science* 270:1660–1663.
- 7 Mäser P, et al. (2002) Altered shoot/root Na⁺ distribution and bifurcating salt sensitivity in *Arabidopsis* by genetic disruption of the Na⁺ transporter *AtHKT1*. *FEBS Lett* 531:157–161.
- 8 Schroeder JI, Hagiwara S (1989) Cytosolic calcium regulates ion channels in the plasma membrane of *Vicia faba* guard cells. *Nature* 338:427–430.
- 9 Negi J, et al. (2008) CO₂ regulator SLAC1 and its homologues are essential for anion homeostasis in plant cells. *Nature* 452:483–486.
- 10 Vahisalu T, et al. (2008) SLAC1 is required for plant guard cell S-type anion channel function in stomatal signalling. *Nature* 452:487–491.