

SOME METABOLIC CHANGES IN NEWLY BORN PUPPIES OF DIABETIC BITCHES

¹Hussein A. A. Abdel-Maksoud, ¹Yakout A. Y. El-Senousy and ¹Mohamed M.
Ghanem

¹Dept. Biochemistry and ¹Dept Animal Med. Fac. Vet. Med. (Moshtohor)
Zagazig University / Benha Branch.

The present study is designed to explain some of the major metabolic complications in puppies of diabetic bitches. For this aim, twelve newly born puppies (n = ١٢) of six pregnant diabetic bitches and six non-diabetic ones were collected in El-Bayda pet animals clinic at Omer El-Mokhtar University - Faculty of Veterinary Medicine – Republic of Libya. Venous blood samples were collected from each puppy during the ٢٤ hours after birth. The present data revealed hypoglycemia, hyperinsulinemia, hyperlactacidemia, hypocalcemia, hypomagnesemia, hyperbilirubinemia. Also, a significant increase was observed in hepatic function enzymes [alanine aminotransferase (ALT), aspartate aminotransferase (AST) and gamma glutamyl transferase (GGT)], alkaline phosphatase (ALP) and hematocrit values (PCV) in puppies of diabetic dams on comparison with the mean values of the puppies of non-diabetic ones. On the other hand, manganese (Mn), zinc (Zn) and copper (Cu) levels were non-significantly changed. Moreover, acid phosphatase (ACP), urea, creatinine and plasma immunoglobulins (IgG, IgA and IgM) concentrations were non-significantly changed. This study highlights the great risk of maternal diabetes on the newly born puppies.

INTRODUCTION

Diabetes mellitus (DM) is a state of chronic hyperglycemia. Many factors have been involved in the pathogenesis of DM including environmental, immunological and genetic. These factors may cause hyperglycemia by reducing endogenous insulin or by opposing its action. The lack of insulin leads to abnormalities of carbohydrate, protein and lipid metabolism.

The increased incidence of DM that is complicating pregnancy is of concern since it is associated with an increase in mortality and morbidity of the fetus and neonate (Whu, ٢٠٠٣). Despite the current improvement of diabetes care in pregnancy, neonatal complications are still more frequent than in the general population (Oberhoffer *et al*, ١٩٩٧). Many of these complications are related to the severity of the maternal hyperglycemia during pregnancy (Weintrob *et al*, ٢٠٠٢) and to the metabolic status of the diabetic mother (Rosem *et al*, ٢٠٠٣). Diabetes in pregnant females can be detrimental to her fetus for many reasons. First, she has an increased spontaneous abortion rate (Mills *et al*, ١٩٩٨). Second, ٦ % to ٨ % prevalence of major congenital anomalies has been found in the newborns of diabetic dams mainly cardiovascular, central nervous and musculoskeletal systems. (Kitzmilller *et al*, ٢٠٠٢).

The major negative consequences of neonates of diabetic mothers are macrosomia, neonatal morbidities and respiratory distress syndrome (Hod *et al*, ١٩٩١), hypoglycemia (Otaga, ٢٠٠١), hyperbilirubinemia, disturbed hepatic function and hypocalcemia (Whu, ٢٠٠٣) and cardiomyopathy which

is secondary to the anabolic effect of fetal hyperinsulinemia (Cowell and Schwartz, ٢٠٠٢).

Accordingly, the aim of the present investigation is to study some of the major metabolic changes that may occur in the newly born puppies of diabetic bitches after birth.

MATERIALS AND METHODS

Animals

Twelve newly born puppies of six pregnant diabetic bitches and six non-diabetic ones were admitted at El-Bayda pet animals clinic at Omer El-Mokhtar University - Faculty of Veterinary Medicine – Republic of Libya. Diabetes mellitus in bitches was diagnosed on the basis of the clinical signs and determination of fasting blood glucose level and insulin level. The major clinical signs observed were polyuria, polydipsia, obesity and cataract. All diabetic bitches had hyperglycemia, glucosuria and hypoinsulinemia.

Samples

Venous blood samples were collected from each puppy during the first ٢٤ hours after birth. Blood samples were withdrawn by a vein puncture from the cephalic veins according to **Kirk and Bistner (١٩٨٩)**. The blood samples were divided into ٢ portions, the first on EDTA for whole blood analysis, the other one was allowed to coagulate at room temperature and then the clear sera were separated by aspiration after centrifugation at ٣٠٠٠ rpm for ١٠ minutes.

Analysis of Sera and Whole Blood Samples

The collected sera were freshly used for quantitative determination of glucose (Barham and Trinder, ١٩٧٢), lactate (Tietz ١٩٧٦), Insulin (Mullner, *et*

al, ١٩٩١), calcium (Grindler and King, ١٩٧٢), magnesium (Bohuon, ١٩٦٢), total bilirubin (Jendrassik and Grof, ١٩٨٣), urea (Patton and Crouch, ١٩٧٧), creatinine (Henry, ١٩٧٤), immunoglobulin concentration (Macini, *et al*, ١٩٦٥), gamma glutamyl transferase (GGT) {EC ٢,٣,٢,٢} (Persijn and Van, ١٩٧٦), alanine amino transferase (ALT) {EC ٢,٦,١,٢}, aspartate amino transferase (AST) {EC ٢,٦,١,١}, alkaline phosphatase (ALP){EC ٣,١,٣,١} (Belifield and Goldberg, ١٩٧١), acid phosphatase (ACP) {EC ٣,١,٣,٢} (Zoss, ١٩٨٤) and trace elements (Zn, Cu, and Mn) by using ^oPg atomic absorption spectrophotometer (Pye Unicam) model ٣٣٠٠ USA according to Fernandy and Kahen (١٩٧١). The whole blood EDTA samples used for determination of Packed cell volume (PCV) (Dacie and Lewis, ١٩٧٥)

Statistical Analysis

The obtained data were statistically analyzed and the significant difference between groups was evaluated by t-test as explained by Snedecor and Cochran (١٩٨٢). All differences were considered significant at $p < ٠,٠٥$.

RESULTS

The recorded data in the provided tables revealed significant increases ($P < ٠,٠٥$) in the mean values of serum insulin, PCV % (table ١), total bilirubin, AST, ALT and GGT (table ٣) in the experimental group (newly born puppies of diabetic dams) compared to the control group (newly born puppies of non-diabetic dams). Moreover, there was a highly significant increase ($P < ٠,٠١$) in serum lactate (table ١) and ALP (table ٣) in the experimental group compared to control group.

On the other hand, there were significant decreases ($P < 0.05$) in the mean values of serum glucose (table ١), calcium and magnesium (table ٢) of the experimental group (newly born puppies of diabetic dams) on comparison with the values recorded in the control (newly born puppies of non-diabetic dams).

Meanwhile the reported data showed non-significant increases in serum copper (table ٣), ACP (table ٤) and IgA (table ٥); and non-significant decreases in serum zinc, manganese (table ٦), IgG and IgM (table ٧) of the experimental group (newly born puppies of diabetic dams) on comparison with the values recorded in the control (newly born puppies of non-diabetic dams).

Table (١):

The mean values \pm S.E of Glucose, Lactate, Insulin and PCV% in newly born puppies of non- diabetic dams and of diabetic dams.

Parameter Groups	Glucose (mg / dl)	Lactate (mg / dl)	Insulin (μ IU /ml)	PCV %
Control (puppies of non – diabetic dams)	٧٩,٨٠ \pm ١,٢	٤١,٩٠ \pm ١,٠٩	٩,٣٣ \pm ٠,٢١	٣١,٩١ \pm ٢,٦٠
Experimental (puppies of diabetic dams)	٥٠,٠٣ \pm ١,٠٩*	٦٩,٣٠ \pm ٢,١١**	١٨,٣٩ \pm ١,٠٧*	٥٥,١٨ \pm ٣,١١*

* Significantly different at $P < 0.05$

** Highly significantly different at $P < 0.01$

Table (٢):-

The mean values \pm S.E of Calcium, Magnesium, Zinc, Copper and Manganese in newly born puppies of non- diabetic dams and of diabetic dams.

Parameters Groups	Calcium (mg / dl)	Magnesium (mg / dl)	Zinc (μ g/dl)	Copper (μ g/dl)	Manganese (μ g/dl)
Control (puppies of non – diabetic dams)	٩,٣٠ \pm ٠,٨٧	٢,٦٦ \pm ٠,١٦	١٩,٢٠ \pm ٠,٥٠	٢٦,٣٣ \pm ١,١٠	٨,١٠ \pm ٠,١٨

Experimental (puppies of diabetic dams)	٦,٢١ ± ٠,٨٣*	١,٠٨ ± ٠,١٣**	١٨,٦ ± ٠,٦١	٢٧,١١ ± ١,٤٠	٧,٧٥ ± ٠,٢٠
---	--------------	---------------	-------------	--------------	-------------

* Significantly different at $P < ٠,٠٥$

Table (٣):-

The mean Values \pm S.E of Total bilirubin, AST, ALT, GGT, ALP and ACP in newly born puppies of non- diabetic dams and of diabetic dams.

Parameters Groups	Total bilirubin (mg / dl)	AST (U/L)	ALT (U/L)	GGT (U/L)	ALP (U/L)	ACP (U/L)
Control (puppies of non – diabetic dams)	٠,٩٦ ± ٠,٠٣	١٨,٤٠ ± ٠,٩٣	٢٢,١١ ± ١,١٦	٧,٣٠ ± ٠,٦٩	١,١٧ ± ٠,٠٨	٠,٦٩ ± ٠,٠٢
Experimental (puppies of diabetic dams)	٢,٦٣ ± ٠,٢٦*	٣٨,٦١ ± ٢,١٠*	٤٧,١٦ ± ٢,٢٢*	١٧,١١ ± ١,١١*	٧,١٥ ± ١,٠٢**	١,٠٥ ± ٠,١١

* Significantly different at $P < ٠,٠٥$

** Highly significantly different at $P < ٠,٠١$

Table (٤):-

The mean values \pm S.E of Immunoglobulins (IgA, IgG and IgM) in newly born puppies of non- diabetic dams and of diabetic dams.

Parameters Groups	IgA (mg / dl)	IgG (mg/dl)	IgM (mg/dl)
Control (puppies of non – diabetic dams)	٦٣,٢٥ ± ٢,١١	٢٦١,٤٠ ± ٦,١١	٤٥,٣٧ ± ٢,١٣
Experimental (puppies of diabetic dams)	٧١,٦٠ ± ٢,٩٠	٢٥٠,١٩ ± ٤,١٢	٣٩,١١ ± ٢,٧١

DISCUSSION

The recorded data revealed that there was hypoglycemia in the experimental group which coincided with the results obtained by Amina *et al*,

(٢٠٠٠) and Cordero *et al*, (٢٠٠٣) who stated that hypoglycemia has been observed in newly born infants of diabetic mothers. The detected hypoglycemia could be attributed to the hypertrophy and hyperplasia of the Langerhans islets, which might be connected with insulin resistance in peripheral tissues as a significant risk associated with diabetic mother in pregnancy (Dooley and Sugamori, ٢٠٠١). This result was also confirmed by the opinion of Rakhab and Chernev (٢٠٠٠) who proved that the hypoglycemia of the infants of diabetic mothers is probably due to the high insulin levels caused by the hyperplasia of the pancreatic beta-cells that was normally found in these infants.

The observed hyperinsulinemia was similar to the recorded data in infants of diabetic mothers by Oberhoffer *et al*, (١٩٩٧) who showed a proportional increase in serum insulin with maternal glycated hemoglobin in them, it was also stated by Simmons, (١٩٩٩) that the hyperinsulinemia depended on the degree of maternal hyperglycemia which might be related to the primary source of glucose in the early postnatal hours is mobilized from the hepatic glycogen with appropriated catecholamins and glucagon response however this response is blunted in diabetic mothers. Furthermore the conversion of triacylglycerol to fatty acids and glycerol which is ultimately converted to glucose is prevented because of the high insulin and low catecholamine and glucagon levels in the circulation (Whu, ٢٠٠٣).

Regarding the highly significant elevation of the serum lactate level, the presented data agreed with the results of Pribylova and Dovrakova (٢٠٠٣) who found a positive correlation between maternal glycated

hemoglobin and plasma lactate in all diabetic females and their infants. The recorded hyperlactacidemia might be attributed to that the maternal hyperglycemia resulted in fetal hyperinsulinemia thereby, increasing metabolic rate and oxygen demand which exceeds oxygen availability leading to fetal hypoxia and increase lactic acid (Amina *et al*, ٢٠٠٠).

The recorded hypocalcemia, hypomagnesemia and increased ALP activity were similar to the results observed by Rakhab and Chernev (٢٠٠٠) who found a negative correlations between maternal hyperglycemia and serum calcium, magnesium and the activity of serum ALP. These changes could be probably secondary to the transient functional hypoparathyroidism that might be also secondary to the maternal and fetal hypomagnesemia

The recorded significant increase in serum total bilirubin and the activities of the hepatic function enzymes AST, ALT and GGT indicated the direct evidence for fetal hypoxia and polythychemia which are hepatic stressors as stated by Ylevin *et al*, (٢٠٠١) who observed that the polythychemia resulted from increased hemoglobin which then increase bilirubin due to the elevated turnover of heme and decrease clearance of bilirubin

These changes could also be related to the increased hemolysis, elevated red cells masses, ineffective erythroblisis, prematurity, increased bruising and trauma and retardation of enzymes system maturation (Weintrob *et al*, ٢٠٠٢).

The regarded significant increase in PCV % was similar to the data recorded by Cordero *et al*, (٢٠٠٣) who stated that there were a significant increase in Hb, PCV, and RBCs in infants of diabetic mothers which might be attributed to the fetal hypoxia and polycythemia.

٨

The detected non-significant changes in serum IgA, copper, ACP, IgG, IgM, zinc and manganese suggest that these changes could be independent on the maternal diabetes, which means that maternal diabetes of bitches is not associated with changes of these parameters in newly born puppies.

In conclusion, this study provided an insight on the great risk of maternal diabetes of bitches on their newly born puppies. Therefore, it is recommended that the thorough control of diabetes mellitus in pregnant bitches is crucial order to avoid the short-term complications in the newly born puppies.

REFERENCES

1. Amina, M., Azza, A., Hisham, A. and Khaled, F. (٢٠٠٠). Some metabolic complications in infants of diabetic mothers. The Egy. J. Bio. ١٨: ١١١-١٢٧.
2. Barham, D. and Trinder, P. (١٩٧٢). An improved colour reagent for the determination of blood glucose by the oxidase system. Analyst ٧٩: ١٤٢-٤٩.
3. Belifield, D. and Goldberg, S. (١٩٧١). In Enzymes. ١٢th Ed. pp. ٥٦١. (Cited in Biochemical Kits).
4. Bohuon, C. (١٩٦٢). Anal. Chim. Acta. ٧: ٨١١-٨١٧.
5. Cordero, L., Treuer, S., Landon, M and Gabbe, S. (٢٠٠٣). Management of infants of diabetic mothers. Arch. Pediatr. Adolesc. Med. ١٥٢: ٢٤٩-٥٤.
6. Cowett, R., and Schwartz, R. (٢٠٠٢). The infants of diabetic mother. Pediatr. Clin. North. Am. ٤٩: ١٢١٣-٢٤.
7. Dacie, J. V. and Lewis, S. M. (١٩٧٥). In 'Practical Haematology ٥th Ed., Longman group limited, Churchill living stone, chapter ٢, pp. ٢٣٤

٨. Dooley, J. and Sugamori, M. (٢٠٠١). Pregnancy outcome in aboriginal women with NIDDM in the Sioux Lookout Zone. *Int. J. Circumpolar. Health* ٥٧: ٣٥٩-٦٤.
٩. Fernandy, F. and Kahen, H. (١٩٧١). Clinical methods for atomic absorption spectrophotometry. *Clin. Chem. News* ٣: ٢٤.
١٠. Grindler, E. M. and King, J. D. (١٩٧٢). Rapid colorimetric determination of calcium in biologic fluids with methythymol blue. *Am. J. Clin. Pathol.* ٥٨: ٣٧٦.
١١. Henry, R. (١٩٧٤). In' *Clinical chemistry principles and techniques* ' ٢nd Ed. Harper and Row :pp.٥٤٣.
١٢. Hod, M., Merlob, P. and Ovadia, J. (١٩٩١). Gestational diabetes mellitus : A survey of perinatal complications in the ١٩٨٠s. *Diabetes* ٤٠: ٧٤-٩٠.
١٣. Kirk, R. and Bistner, S. (١٩٨٩). In “ *Handbook of veterinary procedures and emergency treatment* .٥th Ed.W.B. Saunders Co., Philadelphia, London, Toronto, Mexico . pp. ٦٥.
١٤. Jendrassik, L. and Grof, P. (١٩٨٣). *Biochem.J.* ٢٩٧: ٨٢-٨٩.
١٥. Kitzmiller, H., Buchanan, T., Kjos, S. and Ratner, R. (٢٠٠٢). Preconception care of diabetes, congenital malformatios and spontaneous abortion .*Diabetes Care* ١٩: ٥١٤- ٢٠.
١٦. Macini, G., Carbonara, A. O. and Heremans, J. E. (١٩٦٥). Immunochemical quantitation of antigens by single radial immunodiffusion. *Immunochem.* ٢: ٢٣٥-٢٥٤.
١٧. Mills, J., Simpson, L., Driscoll, D. and Knopp, R. (١٩٩٨). Incidence of spontaneous abortion among normal women and insulin-dependent diabetic women whose pregnancies were identified within ٢١ days of conception. *N. Engl. J. Med.* ٣١٩: ١٦١-٧٠.
١٨. Mullner, S., Naubauer, H. and Koning, W. (١٩٩١). A radioimmunoassay for the determination of insulin in several animal species .*Insulin derivatives. J. Immunol. Meth.* ١٤٠: ٢١١-١٩٨

١٩. Oberhoffer, R., Tsng, R. and Miodovnik, M. (١٩٩٧). Cardiac and extracardiac complications in infants of diabetic mothers and their relation to parameters of carbohydrate metabolism. *Eur. J. Pediatr.* ١٥٦: ٧٩٨-٨٠٣.
٢٠. Otaga, S. E. (٢٠٠١). The infants of diabetic mothers: Pregnancy as a tissue culture experience. *Isr. G. Med. Sci.* ٣٧: ٥٢٤-٣٣.
٢١. Patton, C. F. and Crouch, S. R. (١٩٧٧). *Anal. Chem.* ٤٩: ٤٦٤-٦٩.
٢٢. Persijn J. P. and van der Slik, W. (١٩٧٦). A new method for the determination of gamma glutamyl transferase in serum. *J. Clin. Chem. Clin. Biochem.* ١٤: ٤٢١ - ٧.
٢٣. Pribylova, H. and Dovrakova, L. (٢٠٠٣). Long-term prognosis of infants of diabetic mothers: Relationship between metabolic disorders in newborns and adult offspring. *Acta Diabetologica* ٣٣: ٣٠-٣٩.
٢٤. Rakhab, M. and Chernev, T. (٢٠٠٠). Pregnancy outcome (perinatal mortality and morbidity) in women with diabetes. *Akush. Ginekol. Sofia* ٣٧: ١٧-٢٣
٢٥. Rosenn, B., Miodovnik, M. and Tsand, R. (٢٠٠٣). Common clinical manifestations of maternal diabetes in newborn infants: implications for the practicing pediatrician. *Pediatr. Ann.* ٢٥: ٢١٥-٢١
٢٦. Simmons, D. (١٩٩٩). Association between neonatal blood pressure and umbilical cord insulin concentration. *Diabet. Med.* ١٤: ١٩٦-٢٠٣.
٢٧. Snedecor, G. W. and Cochran, W. G. (١٩٨٢). "Statistical methods" ٦th Ed. The Iowa State Univ. Press, Ames, Iowa, USA.
٢٨. Tietz N. (١٩٨٧): Determination of Lactate. *Fundamentals of clin. chem.*. Pp. ٥٠٨, W.B. Sanders Comp. Londone, Philadelphia.
٢٩. Weintrob, N., Karp, M. and Hod, M. (٢٠٠٢). Short and long range complications in offspring of diabetic mothers. *J. Diabetes Complications* ١٠: ٢٩٤-٣٠١

٣٠. Whu, P. Y.(٢٠٠٣). Infant of diabetic mother : A continuing challenge for perinatal-neonatal medecine. Acta Pediatr.Sin. ٣٧: ٣١٢-١٩.
٣١. Ylevin, L., Rigg, A. and Marshall, R. (٢٠٠١). Pregnancy and diabetes. Arch. Intern. Med. ١٤٦: ٧٦٥-٧
٣٢. Zoss, D. W. (١٩٨٤). Methods of enzymatic analysis. Am. J. Clin. Pathol. ٥٤: ٩٢-١٠٦.

الملخص العربي

بعض التغيرات الايضية في الكلاب المولودة حديثا لأمهات مصابة بداء السكري

حسين عبد المقصود على* , ياقوت عبد الفتاح السنوسي* , محمد محمدى غانم**

قسم الكيمياء الحيوية و. ط. طب الحيوان ، بكلية الطب البيطري بمشهور جامعة الرقازيق - فرع بنها

صممت هذه الدراسة لمعرفة تأثير الإصابة بمرض السكري لاناث الكلاب الحوامل على بعض التغيرات الايضية في الكلاب المولودة حديثا لذلك تم اختيار اثني عشر من عشائر اناث الكلاب بمستشفى البيضا للحيوانات الاليفة بكلية الطب البيطري بجامعة عمر المختار بليبيا - قسمت الى مجموعتين كلا منهما تحتوى على ستة كلابات الاولى امهات سليمة و الثانية مصابة بداء السكري - جمعت عينات الدم من الكلاب المولودة خلال ال ٢٤ ساعة الاولى بعد الولادة. و قد اظهرت النتائج زيادة معنوية فى الانسولين و حمض الاكتيك و البيلبيروبين و نشاط انزيمات (الجاما جلوتاميل ترانسفيريز و الالانين امينو ترانسفيريز و الاسبارات امينو ترانسفيريز و الالكالين فوسفاتيز) و الهيماتوكريت و نقصا معنويا فى مستويات الكالسيوم و الماغنسيوم . بالاضافة لذلك سجلت تغيرات غير معنوية فى مستويات المنجنيز و الزنك و النحاس و الاسيد فوسفاتيز و كذلك البولينيا و الكرياتينين و الاجسام المناعية IgG, IgA, IgM . من هذه الدراسة اتضح الخطورة الكبيرة للامهات المصابة بالسكري ان لم يتم التحكم فيه على الصغار المولودين حديثا .