



Seeking natural capital projects: Forest fires, haze, and early-life exposure in Indonesia

Jie-Sheng Tan-Soo^a and Subhrendu K. Pattanayak^{b,c,1}

^aLee Kuan Yew School of Public Policy, National University of Singapore, Singapore 259772; ^bSanford School of Public Policy, Economics Department, Duke University, Durham, NC 27708; and ^cGlobal Health Institute, Duke University, Durham, NC 27708

Edited by S. Polasky, University of Minnesota, St. Paul, MN, and approved January 15, 2019 (received for review March 26, 2018)

Natural capital will be depleted rapidly and excessively if the long-term, offsite impacts of depletion are ignored. By examining the case of tropical forest burning, we illustrate such myopia: Pursuit of short-term economic gains results in air pollution that causes long-term, irreversible health impacts. We integrate longitudinal data on prenatal exposure to the 1997 Indonesian forest fires with child nutritional outcomes and find that mean exposure to air pollution during the prenatal stage is associated with a half-SD decrease in height-for-age z score at age 17, which is robust to several statistical checks. Because adult height is associated with income, this implies a loss of 4% of average monthly wages for approximately one million Indonesian workers born during this period. To put these human capital losses in the context of policy making, we conduct social cost-benefit analyses of oil palm plantations under different scenarios for clearing land and controlling fires. We find that clearing for oil palm plantations using mechanical methods generates higher social net benefits compared with clearing using fires. Oil palm producers, however, would be unwilling to bear the higher private costs of mechanical clearing. Therefore, we need more effective fire bans, fire suppression, and moratoriums on oil palm in Indonesia to protect natural and human capital, and increase social welfare.

sustainable development | environmental health | oil palm | cost-benefit analysis | health irreversibility

Economists will argue that natural capital has been depleted rapidly and excessively because the offsite lagged impacts of depletion are either ignored or remain unmeasured (1–4). We use the case of Indonesia to illustrate the extent of such oversight. Despite its vast tropical forests, forest loss is rapid because forests are burned to clear land cheaply and plant lucrative job-friendly export crops such as oil palm (5, 6). Unfortunately, such economic development is unsustainable because we ignore externalities of forest fires—air pollution and biodiversity loss, chief among others (4, 7). For example, forest fires in Indonesia, started to establish estate crops, burned out of control due to the El Niño-induced abnormally dry weather in 1997. These fires destroyed habitat (around 11 million hectares of forests), compromised hydrological services, and generated health-damaging air pollution (around 25% of global carbon emissions from fossil fuels came from this single event) (8–11). Despite the severity of this event, we do not know the full extent of health damages, especially the irreversible, offsite lagged human capital impacts. In addition, we do not know whether the costs (in terms of jobs and profits) of policies to avoid the haze will outweigh the social benefits (12).

In the post-2015 sustainable development goals (SDGs) era, debates rage in the global community about how best to protect natural capital, promote health, mitigate climate change, and reduce poverty (13, 14). The main question posed in our paper—do the economic benefits of avoiding health damages of haze from forest fires outweigh the economic costs of alternative policies—is relevant to these debates. Specifically, our research illustrates why SDGs should focus on reducing negative externalities, promoting intergenerational equity, and improving capabilities (15). We focus on the 1997 forest fires in Indonesia and the associated haze. The 1997 fires were one of the largest in recent history, but

unfortunately, such forest fires have become even more frequent lately, including a round of devastating forest fires in 2015. Despite their magnitude and frequency, we know surprisingly little about the full social costs of these fires. While there is evidence on short-run health damages of air pollution, little is known about the long-term and intergenerational costs of early-life exposure to air pollution.

Most studies of early-life exposure to air pollution are conducted in high- or middle-income countries and focus on immediate birth outcomes (16, 17). From past analyses, there is strong evidence showing that early-life exposure to air pollutants is associated with low birthweight and preterm birth (16, 17). The suspected pathways from air pollutants to birth outcomes are inflammation and direct toxic effects to the placenta and fetus, oxygen supply to the fetus, and DNA expression (17). With respect to longer-term outcomes, the literature on the “fetal origins” hypothesis suggests that intrauterine health insults can cause lasting and irreversible damage to cardiovascular and respiratory health and that low birthweight is associated with shorter height in adulthood (18–21). Still, there are very few studies that specifically make the connection between environmental exposure to air pollutants at early-life and long-term outcomes (22).

Demonstrating that a short-term episode of extremely high air pollution has long-term health impacts is salient to many low- and middle-income countries, especially as many parts of Asia face frequent “airpocalypses” in recent years. In our study setting, the 24-h total suspended particulate (TSP) concentration reached as high as 4,000 $\mu\text{g}/\text{m}^3$ during multiple days in October 1997 in parts of Sumatra—the World Health Organization (WHO) recommended 24-h maximum for TSPs is 120 $\mu\text{g}/\text{m}^3$ (23). More recently, New Delhi experienced 24-h $\text{PM}_{2.5}$ levels of 700–1,200 $\mu\text{g}/\text{m}^3$ in November 2016—the WHO daily guideline is 25 $\mu\text{g}/\text{m}^3$. Also in late 2016, multiple cities in northern China attained their highest level of air pollution warning. Because earlier studies were mainly conducted in rich countries that experience very different exposure profiles, economic development, and environmental institutions, there is little evidence to guide policy makers and practitioners in low- and middle-income contexts.

This paper results from the Arthur M. Sackler Colloquium of the National Academy of Sciences, “Economics, Environment, and Sustainable Development,” held January 17–18, 2018, at the Arnold and Mabel Beckman Center of the National Academies of Sciences and Engineering in Irvine, CA. The complete program and video recordings of most presentations are available on the NAS website at www.nasonline.org/economics-environment-and.

Author contributions: J.-S.T.-S. and S.K.P. designed research; J.-S.T.-S. and S.K.P. performed research; J.-S.T.-S. analyzed data; and J.-S.T.-S. and S.K.P. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Published under the PNAS license.

¹To whom correspondence should be addressed. Email: subhrendu.pattanayak@duke.edu.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1802876116/-DCSupplemental.

Published online February 19, 2019.

It is not enough to stop at estimating environmental health damages, however, without considering how they stack up against the costs (and other benefits) of policies to control fires. The political economy of policies to control forest fires in Indonesia is complex and covers the entire gamut from government regulation to market incentives to voluntary control by firms and landowners (24–27). Currently, key policies include blanket bans on fires for clearing land and moratoriums on expanding oil palm (28, 29). Because oil palm is extremely lucrative in terms of foreign exchange revenue (e.g., it accounted for US\$18.6 billion, or about 10% of Indonesia's total exports in 2015) and jobs (e.g., sector employs around 4.5 million workers), such moratoriums may be politically impractical and unsustainable. Moreover, firms will not comply with bans and moratoriums if enforcement is not strict (30, 31). Hence, we must consider costs and benefits of policies to detect and suppress forest fires before they spread, and to enforce bans and moratoriums (32). Such cost–benefit analyses (CBAs) can identify the parameters driving the maximizations of social welfare, given the health benefits of reduced fires and haze (7). All of these options are complicated by the fact that fire severity depends on the location, and the costs and benefits vary by stakeholders (33).

Study Rationale and Design

First, it is difficult to know the extent of irreversible and offsite lagged human capital impacts because high-quality longitudinal data that faithfully depict long-term impacts have been difficult

to find in environmentally vulnerable and economically poor locations (7, 34). Thus, we carefully integrate longitudinal data on prenatal exposure to the pollution from the 1997 Indonesian forest fires with data on nutritional outcomes, genetic inheritance, climatic factors, household inputs, and various sociodemographic factors. Second, methodologically flexible approaches are needed because of the interdisciplinary nature of issues in sustainable development and planetary health (4). We show how rigorous statistical analyses and numerous robustness checks can be applied to a multisectoral panel dataset to control for potential sources of confounding the correlation of prenatal haze exposure with adult height. While previous research has drawn attention to the deaths caused by the forest fires, we show that survivors also suffer large and irreversible losses of human capital because of haze exposure (35, 36). Third, scientists are often unwilling (because of norms or incentives) to provide approximate, if imperfect, practical advice that policy makers seek, preferring to provide precise answers to questions, even if these questions are irrelevant or untimely for policy (37). To avoid such “type III” errors and to put such human capital losses in context, we conduct a CBA of various policies to avoid fires in oil palm plantations (38–40). Specifically, we combine our haze-height results with other costs and benefits, and also account for the heterogeneity and uncertainty inherent in the model parameters.




Fig. 1. Spatial and temporal distribution of air quality across Indonesia.




Fig. 2. Impact of early-life air pollution on HAZ for different regression specifications.

Results

Our data show that the average 1997 air pollution exposure [aerosol index (AI)] is 0.1 (where 0.01 represents crystal-clear sky with maximum visibility and 0.4 represents trouble seeing the midday sun). As shown in Fig. 1, however, AI varies significantly in space and time, with exposure ranging from 0 (18% of sample, from Sulawesi and Nusa Tenggara) to 0.3 (18% of sample, from Kalimantan and Sumatra). Average height-for-age z score (HAZ) is negative, suggesting that Indonesian children are shorter than the reference group from the United Kingdom. The other environmental, household, and parental variables are summarized in *SI Appendix, Table S1*.

Fig. 2 shows effects of AI, including the 95% confidence interval, on child's HAZ 17 y after exposure [i.e., fourth round of the Indonesian Family and Life Survey (IFLS)]. The first, second, and third coefficient in each set of results represent the overall, prenatal, and postnatal impact, respectively, of early-life exposure (*Materials and Methods*). Results from the main specification

show that the mean level of exposure (i.e., AI = 0.1) translates to a 0.41 decrease in HAZ (or about 3.4 cm, equivalently) by 2014. The full results are also presented in *SI Appendix, Table S2*. These results show that the decrease in HAZ is statistically consistent across earlier waves of the IFLS. In other words, children in our analysis experienced a decrease in height from 3 y of age, and this impact persisted through the age of 17.

We further examined whether this early-life effect was attributable to prenatal or postnatal exposure. When exposed during the prenatal stage, the impact on HAZ is essentially unchanged—the average effect is 0.43. In contrast, while the coefficient for postnatal exposure is negative, it is statistically insignificant. Thus, the relationship we detect appears to be completely driven by prenatal exposure.

To ensure our results are not driven by confounders and spurious correlations, we undertake a series of robustness checks (*Materials and Methods*). We confirm that our findings are not driven by (i) high levels of pollution in later years, (ii) an indirect effect of severe air pollution on a family's ability to work and to earn income, (iii) something about the location, rather than the exposure per se, and (iv) overall reductions in food consumption during the forest fire months.

To put these estimates of irreversible health impacts in context, we conduct a CBA of oil palm plantations by including social externalities (*Materials and Methods*). Briefly, the analysis of haze-reducing policies considers benefits to health (both avoided losses in income and in mortality), tourism, and transportation, and costs to firms (land preparation) and to agencies for program implementation. To acknowledge and model the heterogeneity and uncertainty inherent in the parameters used in the CBA, we conduct Monte Carlo simulations by allowing each parameter to take on a range of values. This process yields 10,000 net present values (NPVs) for each scenario, summarized in a cumulative distribution function (CDF) for each scenario.

Starting with the baseline scenario of fire-based clearing, we see that 30% of the CDF is negative (Fig. 3). This finding is particularly sensitive to emissions attributed to oil palm fires, exposure to air pollution, income growth, and mortality and human capital losses (*SI Appendix, Fig. S1*). First, we consider the main alternative of clearing mechanically (and avoiding fires) to establish plantations and find that the CDF of social NPV is




Fig. 3. Comparing social welfare of land clearing for oil palm using fire, mechanical options, and public policies (e.g., ban and suppression).

Statistical Approach. The impact of AI on height, β_1 , is obtained by regressing child's health, y_{ijt} , on AI, AI_{ijt} , and a host of controls using least-squares regression method:

$$y_{ijt} = \beta_0 + \beta_1 AI_{ijt} + \beta_2 AI_{ijt} \cdot I(Y_{2000}) + \beta_3 AI_{ijt} \cdot I(Y_{2007}) + I(Y_{2000}) + I(Y_{2007}) + X_i \gamma + \delta_j + \alpha_t + \varepsilon_{ijkt} \quad [1]$$

where i , j , and t represent child i from location j and born in period t . The subscript k denotes the particular survey in which height was measured (i.e., IFLS 2000, IFLS 2007, or IFLS 2014). X is a vector of parent and household characteristics that could impact child's HAZ, for example, parents' education, parents' heights (a genetic contribution to the child outcomes), and household inputs at birth such as sanitation and clean cooking fuels. $I(Y_{2000})$ is an indicator variable equal to 1 if the height was obtained from IFLS 2000. δ and α are district—the same scale at which AI is measured—and birth month fixed effects, respectively. The birth month-by-birth year fixed effects are included to control for unobserved factors that are constant across all individuals born in the same year and month, such as macroeconomic conditions (e.g., currency devaluation related to the Asian financial crisis) or seasonal weather patterns, which might otherwise confound the relationship between AI and HAZ. Similarly, district fixed effects are included to control for any unobserved factors constant across individuals born in the same location, such as access to local nutrition programs. Thus, following an established empirical method in applied statistics, the relationship between air quality and height is identified by removing any confounding differences attributable to location and season. Last, ε is an idiosyncratic error term. To guard against potential biases from the "Moulton" effect due to coarseness of the AI data, we cluster SEs at both the district and birth month-by-birth year levels (62). Three rounds of data from the same individuals are pooled. Therefore, when AI is interacted with the year in which the survey was administered, β_2 and β_3 represent changes in impact of early-life AI on height in the 2000 and 2007 surveys. In other words, the impact of early-life exposure of AI on height at 17 y is β_1 , whereas the impact of early-life exposure of AI on height at 3 y is $\beta_1 + \beta_2$, and $\beta_1 + \beta_2 + \beta_3$ at 10 y. By way of summary, note that the AI coefficients are estimated using variation in air quality at the spatial-temporal level. This means that any potential confounders (e.g., the Asian financial crisis) would need to covary at the same temporal (by month) and spatial (by districts) scale with air pollution to bias the estimates of the AI coefficients.

Eq. 1 can be modified to separate the effects of prenatal versus postnatal exposure AI on height:

$$y_{ijt} = \beta_0 + \beta_1 preAI_{ijt} + \beta_2 postAI_{ijt} + \beta_3 preAI_{ijt} \cdot I(Y_{2000}) + \beta_4 preAI_{ijt} \cdot I(Y_{2007}) + \beta_5 postAI_{ijt} \cdot I(Y_{2000}) + \beta_6 postAI_{ijt} \cdot I(Y_{2007}) + I(Y_{2000}) + I(Y_{2007}) + X_i \gamma + \delta_j + \alpha_t + \varepsilon_{ijkt} \quad [2]$$

The main difference between Eqs. 1 and 2 is that β_1 and β_2 in Eq. 2 report the effects of prenatal and postnatal exposure to AI, respectively, when the respondents are at 17 y, whereas β_1 in Eq. 1 reports the overall effect of any exposure to AI. Similarly, the coefficients for the other interaction terms in Eq. 2 report any remaining effects of AI from prenatal and postnatal AI exposure. Statistical analyses are conducted using Stata 14.

Robustness Checks for Statistical Analysis. To confirm that our results are not driven by spurious correlations and confounding factors, we conduct four robustness checks.

First, it is possible that the height impacts are driven by high levels of pollution in later years. *SI Appendix, Table S1* demonstrates that the 1997 pollution was unprecedented, and AI levels were much higher in 1997 compared with later years. Moreover, the district fixed effects control for later years' exposure for all individuals within the same district. The district fixed-effects strategy would not work, however, if those exposed at birth were systematically more likely to migrate to heavily polluted, dirtier locations. Thus, we gathered further information on households' migration history and computed the AI exposure for 1998 and 1999 for their updated locations. In regression analysis that includes 1998 and 1999 AI as additional explanatory variables, only the 1997 exposure is statistically significant (Fig. 2; full results in *SI Appendix, Table S2*). We are not claiming that exposure to air pollution later in life does not impact height, rather that the initial (1997) early-life exposure is the dominant channel.

Second, we consider other nonpollution mechanisms by which height could be affected by the fires. For example, all else being equal, severe air pollution may have reduced a family's ability to work, and, in turn, this would decrease household income and, consequently, caloric intake. This is especially true

for those engaged in outdoor work. This loss-of-income mechanism can be tested by interacting the AI variable with proportion of adult household members engaged in outdoor work. Indeed, the impact on HAZ is stronger in this subsample of households with outdoor workers; that is, while the AI coefficient mostly retains similar magnitude and statistical significance from the main results, there seem to be additional impacts on HAZ for households with higher proportion of outdoor workers (Fig. 2 and *SI Appendix, Table S2*). This means that this channel of outdoor work partially explains HAZ differences. However, the key AI coefficients remain significant for all three waves, signaling that the loss-of-income mechanism still leaves room for a large, direct impact of exposure on growth.

Third, to test whether there is something about the location, rather than the exposure itself, that is driving the association, we used the true exposure group but assigned a placebo AI exposure (1996 AI, true exposed cohort). That is, we assigned the 1996 AI exposure to our existing sample of exposed children. These children could not have been exposed to the pollution in 1996, as this is the year before their conception. The placebo test demonstrates this effect: The total impact of 1996 AI on HAZ is insignificant for each of the three rounds (Fig. 2 and *SI Appendix, Table S2*).

Last, we consider another source of potential confounding: if the forest fires influenced reduced food consumption. We can test this channel because of the timing of the surveys, as some household were surveyed during and others after the forest fires. Regression analysis of food consumption shows no statistical difference in consumption during and after the forest fires (*SI Appendix, Table S3*).

CBA. CBAs essentially compare the discounted stream of costs and benefits arising from a project or policy. In this setting, the CBA puts our estimates of health irreversibilities in context but is also essential for understanding the implementation of conservation policies in general (12). First, we conduct a social CBA of oil palm plantations by including social externalities. Second, because the private optimum will diverge from what is best for society, we also conduct the CBA from firms' perspectives to learn, for example, if credible enforcement can incentivize firms to behave in a way so that higher social welfare is achieved. Briefly, the analysis of haze-reducing policies considers benefits to health (both reductions in mortality and morbidity), tourism, and transportation, and costs to firms (land preparation) and to agencies for program implementation. While we included the ecological costs of mechanical clearing, we are not able to include ecosystem-related costs of fire clearing related to carbon and habitat because we could not find conclusive quantitative data on these benefits. Including these other costs would make the net social benefits of oil palm cleared using fire even more negative.

Third, to acknowledge and model the heterogeneity and uncertainty inherent in the parameters used in the CBA, we conduct Monte Carlo simulations (63). That is, we allow each parameter to take on a range of values (obtained from the literature) and specify a statistical distribution for these values (either uniform or normal distribution). For example, the AI-HAZ relationship is estimated in this study with a SD. To fully utilize the range behind each parameter, we ran 10,000 trials in the Monte Carlo simulation whereby, in each trial, a value for each parameter is randomly from the specified statistical distribution. Eventually, this process yields 10,000 different NPVs, which constitute a CDF.

The 26 parameters that underlie these computations and the eight primary equations that combine these parameters to estimate benefits and costs are described in *SI Appendix, Tables S3 and S4*. Without describing each and every parameter and equation (*SI Appendix*), we briefly summarize some of the key computations here.

Haze attributed to oil palm. Neither are forest fires the only source of air pollution in Indonesia, nor are all forest fires caused by establishing oil palm plantations. While there is a large literature examining the role of the causes of the 1997 forest fires and its effects, there is no specific study that directly quantifies the pollution attributable to oil palm or any concessions (38, 39). However, we can draw on this literature and estimate the attributable fraction as follows. First, we compare the AI in the August to October 1997 period to both August to October 1996 and to January to June 1997 periods, when there were fewer forest fires. Thus, we estimate that 57–77% of air pollution is from forest fires during the August to October 1997 period. Second, we rely on a recent study to approximate the forest fire emissions attributable to oil palm plantations, which suggests the range 10–60% of all pollution emissions is because of oil palm plantations (48). Third, we multiply these two fractions to compute the proportion of air pollution attributed to oil palm-related forest fires. Critically, recognizing that this attributable fraction can vary (as with other causes) from the low to high range of this product, we build this variability into the 10,000 Monte Carlo simulations, introduced previously.

Avoided mortality damages. Epidemiological models estimate the total mortality burden for the 1997 fires in Indonesia to be around 15,000 deaths (64). In monetary terms, we assign a value-of-statistical life (VSL) of US\$108,900 (in 2008 US dollars) to these deaths. This VSL estimate is obtained by starting first with OECD baseline VSL of US\$3.3 million (in 2008 US dollars) and then applying “benefits transfer” logic.

Avoided income loss. For loss of income, we start with a well-established literature that shows that relative height is correlated with adult mortality, morbidity, neurodevelopmental, and economic capability (56). Specifically, height has been shown to be correlated with earnings; for example, 1 SD in HAZ at adulthood is associated with 8% increase in adult income (57). Thus, our estimate of a 0.41 SD HAZ decrease (i.e., from the 2014 wave when the cohort is about 17 or 18 y old) translates into ~3.3% decrease in income through human capital channels. Consider the fact that 1.13 million individuals were in their prenatal stage during August through October 1997 in the impacted provinces of Sumatra or Kalimantan, where the air pollution and fires were most intense (that is, what we report next is a conservative estimate because others in other provinces were partially exposed but we do not count them). Assuming, (i) their working age spans 21–58 (the official retirement age), (ii) the average annual wage for blue-collar work at US\$860 (from Indonesian Statistical Department), (iii) a social discount rate of 8%, and (iv) an annual real wage increment of 2% for the first 15 working years (from Indonesian Statistical Department), we estimate that the lifetime productivity loss for this exposed population of 1.13 million is about US\$392 for each individual.

Profits of oil palm plantations. The operating costs and revenue of oil palm plantations are obtained from Butler et al. (65). The plantations are evaluated on a 25-y basis and the range of operating costs and revenue is based off the high- and low-yield scenarios in the original analysis. The size of the oil palm plantation is assumed to be 100,000 ha, based off new plantation area in 1998 (66). Clearing by fire costs between \$82/ha to US\$320/ha (47,

67). In contrast, clearing using mechanical means is much higher and ranges from low of \$200/ha to \$990/ha (47, 67).

Fire detection and suppression. Our analysis of the early detection and suppression policy rests primarily on the related parameters of program costs and effectiveness, and program life. The effectiveness of this policy ranges from 0 to 0.6. That is, emissions from oil palm plantations is reduced by 60% if fully effective. Additionally, to allow for spatial targeting of this policy, we model positive correlation between program effectiveness and both the levels of emissions from oil palm and the income growth. The Indonesian government approximates that the costs of early-detection and suppression program would range between US\$450 million to US\$2.3 billion. Last, we assume this program will last between 2 and 10 y.

Enforcing the fire ban. We model an enforcement policy for fire bans and the firm's reaction on three parameters—probability of detecting which plantations are using fires, the resulting penalties, and firms' risk aversion. We conservatively estimate the probability of identifying fires to range from 0 to 20%. The penalties are derived from recent court judgements on plantation owners that were found guilty of starting fires and are estimated to range from US\$300/ha to US\$40,000/ha. A risk-averse firm would reduce fires more than proportional to the ratio of expected penalty vs. additional costs of mechanical clearing. We consider the risk aversion to range between 0.1 (risk averse) and 3 (risk taking).

ACKNOWLEDGMENTS. For excellent comments, we are grateful to the seminar participants at Duke University, Indian Statistical Institute Delhi (Centre for Research on the Economics of Climate, Food, Energy and Environment's inaugural meeting), National University of Singapore, Zhejiang University, and the National Academy of Sciences (Sackler Colloquium), and to Marc Jeuland, Emily Pakhtigian, Steve Polasky, Alberto Salvo, Chen Shuai, E. Somanathan, Anant Sudarshan, Faraz Usmani, and two anonymous reviewers. Ruoming Fang provided excellent research assistance.

- Dasgupta P (2010) Nature's role in sustaining economic development. *Philos Trans R Soc Lond B Biol Sci* 365:5–11.
- Daily GC, Kareiva PM, Polasky S, Ricketts TH, Tallis H (2011) Mainstreaming natural capital into decisions. *Natural Capital: Theory and Practice of Mapping Ecosystem Services* (Oxford Univ Press, Oxford), pp 3–14.
- Bauch SC, Birkenbach AM, Pattanayak SK, Sills EO (2015) Public health impacts of ecosystem change in the Brazilian Amazon. *Proc Natl Acad Sci USA* 112: 7414–7419.
- Whitmee S, et al. (2015) Safeguarding human health in the Anthropocene Epoch: Report of the Rockefeller Foundation-Lancet Commission on Planetary Health. *Lancet* 386:1973–2028.
- Koh LP (2009) Calling Indonesia's US\$13 billion bluff. *Conserv Biol* 23:789.
- Cochrane MA, et al. (1999) Positive feedbacks in the fire dynamic of closed canopy tropical forests. *Science* 284:1832–1835.
- Pattanayak SK, Haines A (2017) Implementation of policies to protect planetary health. *Lancet Planet Health* 1:e255–e256.
- Aiken SR (2004) Runaway fires, smoke-haze pollution, and unnatural disasters in Indonesia. *Geogr Rev* 94:55–79.
- Frankenberg E, McKee D, Thomas D (2005) Health consequences of forest fires in Indonesia. *Demography* 42:109–129.
- Tacconi L (2003) *Fires in Indonesia: Causes, Costs and Policy Implications* (CIFOR, Bogor, Indonesia).
- Page SE, et al. (2002) The amount of carbon released from peat and forest fires in Indonesia during 1997. *Nature* 420:61–65.
- Pattanayak SK, Kramer RA, Vincent JR (2017) Ecosystem change and human health: Implementation economics and policy. *Philos Trans R Soc Lond B Biol Sci* 372: 20160130.
- Davis A, Matthews Z, Szabo S, Fogstad H (2015) Measuring the SDGs: A two-track solution. *Lancet* 386:221–222.
- Dora C, et al. (2015) Indicators linking health and sustainability in the post-2015 development agenda. *Lancet* 385:380–391.
- Horton R (2014) Offline: Why the sustainable development goals will fail. *Lancet* 383: 2196.
- Maisonet M, Correa A, Misra D, Jaakkola JJ (2004) A review of the literature on the effects of ambient air pollution on fetal growth. *Environ Res* 95:106–115.
- Shah PS, Balkhair T; Knowledge Synthesis Group on Determinants of Preterm/LBW births (2011) Air pollution and birth outcomes: A systematic review. *Environ Int* 37: 498–516.
- Barker DJ (1990) The fetal and infant origins of adult disease. *BMJ* 301:1111.
- Sørensen HT, et al. (1999) Birth weight and length as predictors for adult height. *Am J Epidemiol* 149:726–729.
- Barker DJ (2007) Obesity and early life. *Obes Rev* 8:45–49.
- Chen Y, Zhou L-A (2007) The long-term health and economic consequences of the 1959–1961 famine in China. *J Health Econ* 26:659–681.
- Bharadwaj P, Gibson M, Zivin JG, Neilson C (2017) Gray matters: Fetal pollution exposure and human capital formation. *J Assoc Environ Resour Econ* 4:505–542.
- Heil A, Goldammer J (2001) Smoke-haze pollution: A review of the 1997 episode in Southeast Asia. *Reg Environ Change* 2:24–37.
- Glover D, Jessup T (2006) *Indonesia's Fires and Haze: The Cost of Catastrophe* (International Development Research Centre, Ottawa).
- Murdiyarto D, et al. (2004) Policy responses to complex environmental problems: Insights from a science-policy activity on transboundary haze from vegetation fires in Southeast Asia. *Agric Ecosyst Environ* 104:47–56.
- Quah E (2002) Transboundary pollution in Southeast Asia: The Indonesian fires. *World Dev* 30:429–441.
- Chander P (2015) A political economy analysis of the Southeast Asian haze and some solutions. *RSIS Working Paper* 313 (S. Rajaratnam School of International Studies, Singapore).
- Busch J, et al. (2015) Reductions in emissions from deforestation from Indonesia's moratorium on new oil palm, timber, and logging concessions. *Proc Natl Acad Sci USA* 112:1328–1333.
- Sloan S (2014) Indonesia's moratorium on new forest licenses: An update. *Land Use Policy* 38:37–40.
- Dauvergne P (1998) The political economy of Indonesia's 1997 forest fires. *Aust J Int Aff* 52:13–17.
- Jepson P, Jarvie JK, MacKinnon K, Monk KA (2001) Essays on science and society. The end for Indonesia's lowland forests? *Science* 292:859–861.
- Tacconi L, Jotzo F, Grafton RQ (2008) Local causes, regional co-operation and global financing for environmental problems: The case of Southeast Asian haze pollution. *Int Environ Agreement Polit Law Econ* 8:1–16.
- Purnomo H, et al. (2017) Fire economy and actor network of forest and land fires in Indonesia. *For Policy Econ* 78:21–31.
- Young HS, et al. (2017) Conservation, biodiversity and infectious disease: Scientific evidence and policy implications. *Philos Trans R Soc Lond B Biol Sci* 372:20160124.
- Jayachandran S (2009) Air quality and early-life mortality evidence from Indonesia's wildfires. *J Hum Resour* 44:916–954.
- Marlier ME, et al. (2013) El Niño and health risks from landscape fire emissions in Southeast Asia. *Nat Clim Chang* 3:131–136.
- Schwartz S, Carpenter KM (1999) The right answer for the wrong question: Consequences of type III error for public health research. *Am J Public Health* 89:1175–1180.
- Barber CV, Schweithelm J (2000) *Trial by Fire: Forest Fires and Forestry Policy in Indonesia's Era of Crisis and Reform* (World Resources Institute, Forest Frontiers Initiative, Washington, DC).
- Glastra R, Wakker E, Richert W (2002) Oil palm plantations and deforestation in Indonesia. What role do Europe and Germany play? Report for WWF (AIDEnvironment, Amsterdam).
- Harrison ME, Page SE, Limin SH (2009) The global impact of Indonesian forest fires. *Biologist* 56:156–163.
- Koplitz SN, et al. (2016) Public health impacts of the severe haze in equatorial Asia in september–October 2015: Demonstration of a new framework for informing fire management strategies to reduce downwind smoke exposure. *Environ Res Lett* 11: 094023.
- Sen A (2001) *Development as Freedom* (Oxford Univ Press, Oxford).
- Dasgupta P (2009) *Poverty Traps: Exploring the Complexity of Causation. The Poorest and Hungry: Assessments, Analyses, and Actions* (International Food Policy Research Institute, Washington, DC), pp 129–146.

44. Ruyschaert D, Salles D (2014) Towards global voluntary standards: Questioning the effectiveness in attaining conservation goals: The case of the roundtable on sustainable palm oil (RSPO). *Ecol Econ* 107:438–446.
45. Bateman IJ, et al. (2015) Conserving tropical biodiversity via market forces and spatial targeting. *Proc Natl Acad Sci USA* 112:7408–7413.
46. Naidoo R, et al. (2006) Integrating economic costs into conservation planning. *Trends Ecol Evol* 21:681–687.
47. Tacconi L, Moore PF, Kaimowitz D (2007) Fires in tropical forests—what is really the problem? Lessons from Indonesia. *Mitig Adapt Strategies Glob Change* 12:55–66.
48. Marlier ME, et al. (2015) Fire emissions and regional air quality impacts from fires in oil palm, timber, and logging concessions in Indonesia. *Environ Res Lett* 10:085005.
49. Ismail S (2017) Peatland restoration work on track despite slow progress: Indonesia. *Channel NewsAsia*. Available at <https://www.channelnewsasia.com/news/asia/peatland-restoration-work-on-track-despite-slow-progress-8856536>. Accessed January 24, 2019.
50. Keeler BL, et al. (2017) Society is ready for a new kind of science—is academia? *Bioscience* 67:591–592.
51. Polasky S, Carpenter SR, Folke C, Keeler B (2011) Decision-making under great uncertainty: Environmental management in an era of global change. *Trends Ecol Evol* 26:398–404.
52. Kinzig A, et al. (2003) Coping with uncertainty: A call for a new science-policy forum. *Ambio* 32:330–335.
53. Guerry AD, et al. (2015) Natural capital and ecosystem services informing decisions: From promise to practice. *Proc Natl Acad Sci USA* 112:7348–7355.
54. Qiu J, et al. (2018) Evidence-based causal chains for linking health, development, and conservation actions. *Bioscience* 68:182–193.
55. Strauss J, Witoelar F, Sikoki B, Wattie AM (2009) The fourth wave of the Indonesian family life survey (IFLS4): Overview and field report, Technical Report WR-675/1-NIA/NICHD (RAND Corporation, Santa Monica, CA).
56. Prendergast AJ, Humphrey JH (2014) The stunting syndrome in developing countries. *Paediatr Int Child Health* 34:250–265.
57. Victora CG, et al.; Maternal and Child Undernutrition Study Group (2008) Maternal and child undernutrition: Consequences for adult health and human capital. *Lancet* 371:340–357.
58. Hsu NC, et al. (1999) Comparisons of the TOMS aerosol index with sun-photometer aerosol optical thickness: Results and applications. *J Geophys Res D Atmos* 104: 6269–6279.
59. Maccini S, Yang D (2009) Under the weather: Health, schooling, and economic consequences of early-life rainfall. *Am Econ Rev* 99:1006–1026.
60. Thompson AM, et al. (2001) Tropical tropospheric ozone and biomass burning. *Science* 291:2128–2132.
61. Guan H, et al. (2010) A multi-decadal history of biomass burning plume heights identified using aerosol index measurements. *Atmos Chem Phys* 10:6461–6469.
62. Moulton BR (1990) An illustration of a pitfall in estimating the effects of aggregate variables on micro units. *Rev Econ Stat* 72:334–338.
63. Jeuland MA, Pattanayak SK (2012) Benefits and costs of improved cookstoves: Assessing the implications of variability in health, forest and climate impacts. *PLoS One* 7:e30338.
64. Tacconi L (2016) Preventing fires and haze in Southeast Asia. *Nat Clim Chang* 6: 640–643.
65. Butler RA, Koh LP, Ghazoul J (2009) REDD in the red: Palm oil could undermine carbon payment schemes. *Conserv Lett* 2:67–73.
66. Wicke B, Sikkema R, Dornburg V, Faaij A (2011) Exploring land use changes and the role of palm oil production in Indonesia and Malaysia. *Land Use Policy* 28:193–206.
67. Simorangkir D (2007) Fire use: Is it really the cheaper land preparation method for large-scale plantations? *Mitig Adapt Strategies Glob Change* 12:147–164.