


1-2017

Synthesis of some glucose-fatty acid esters by lipase from *Candida antarctica* and their emulsion functions

Kangzi Ren
Iowa State University

Buddhi P. Lamsal
Iowa State University, lamsal@iastate.edu

Follow this and additional works at: https://lib.dr.iastate.edu/fshn_ag_pubs

 Part of the [Comparative Nutrition Commons](#), [Food Chemistry Commons](#), [Food Processing Commons](#), [Human and Clinical Nutrition Commons](#), and the [Molecular, Genetic, and Biochemical Nutrition Commons](#)

The complete bibliographic information for this item can be found at https://lib.dr.iastate.edu/fshn_ag_pubs/205. For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

This Article is brought to you for free and open access by the Food Science and Human Nutrition at Iowa State University Digital Repository. It has been accepted for inclusion in Food Science and Human Nutrition Publications by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

Synthesis of some glucose-fatty acid esters by lipase from *Candida antarctica* and their emulsion functions

Abstract

The synthesis of glucose esters with palmitic acid, lauric acid and hexanoic acid using lipase enzyme was studied and their emulsion functionality in oil-in-water system were compared. Reactions at 3:1 M ratio of fatty acids-to-glucose had the highest conversion percentages (over 90% for each of the fatty acid). Initial conversion rate increased as substrate solubility increased. Ester bond formation was confirmed by nuclear magnetic resonance technique that the chemical shifts of glucose H-6 and α -carbon protons of fatty acids in the ester molecules shifted to the higher fields. Contact angle of water on esters' pelleted surface increased as the hydrophobicity increased. Glucose esters' and commercial sucrose esters' functionality as emulsifiers were compared. Glucose esters delayed, but did not prevent coalescence, because the oil droplets diameter doubled during 7 days. Sucrose esters prevented coalescence during 7 days since the droplets diameter did not have significant change.

Keywords

Glucose ester, Synthesis, Lipase, Contact angle, Emulsifier, Sucrose ester

Disciplines

Comparative Nutrition | Food Chemistry | Food Processing | Food Science | Human and Clinical Nutrition | Molecular, Genetic, and Biochemical Nutrition

Comments

This accepted article is published as Ren, K. and Lamsal, B.P; 2017. Synthesis of some glucose-fatty acid esters by lipase from *Candida Antarctica* and their emulsion functions. *Food Chemistry*, 214 (1); 556-563. Doi: [10.1016/j.foodchem.2016.07.031](https://doi.org/10.1016/j.foodchem.2016.07.031).

**SYNTHESIS OF SOME GLUCOSE-FATTY ACID ESTERS BY LIPASE FROM
CANDIDA ANTARCTICA, AND THEIR EMULSION FUNCTIONS**

Kangzi Ren, Buddhi P Lamsal*

Department of Food Science and Human Nutrition, Iowa State University.
2312 Food Science Building, Iowa State University, Ames, IA 50010, USA.

* Corresponding author:

Department of Food Science and Human Nutrition, Iowa State University
2312 Science Building, Iowa State University, Ames, IA 20010, USA

lamsal@iastate.edu; +1 (515) 294-8681

Published in the *Food Chemistry*, 214 (2017): 556-563

3.1 Abstract

The synthesis of glucose esters with palmitic acid, lauric acid and hexanoic acid using lipase enzyme were studied and their emulsion functionality in oil-in-water system were compared. Reactions at 3:1 molar ratio of fatty acids-to-glucose had the highest conversion percentages (over 90% for each of the fatty acid). Initial conversion rate increased as substrate solubility increased. Ester bond formation was confirmed by nuclear magnetic resonance technique that the chemical shifts of glucose H-6 and α -carbon proton of fatty acid in the esters shifted to the higher fields. Contact angle of water on esters' pelleted surface increased as the hydrophobicity increased. Glucose esters' and commercial sucrose esters' functionality as emulsifiers were compared. Glucose esters delayed, but did not prevent coalescence, because the oil droplets diameter doubled during 7 days. Sucrose esters prevented coalescence during 7 days since the droplets diameter did not have significant change.

Key words: glucose ester; synthesis; lipase; contact angel; emulsifier; sucrose ester.

3.2. Introduction

Surfactants are amphiphilic molecules with both hydrophilic and hydrophobic moieties that can adsorb at the interface between different polarity phases and reduce interfacial tension. Thus, they have functionality in detergency, emulsifying, dispersion, foaming industry (Greek, 1991, 1990). Traditional surfactants are mainly derived from petroleum industry products, which requires unfavorable reaction conditions such as high temperature, high acidity, alkalinity , organic solvent, and have low biodegradability and high aquatic toxicity (Deleu & Paquot, 2004). However, surfactants can also be produced via enzymatic reactions or microbial fermentation utilizing biological feedstocks. These environmental compatible surfactants, also called biobased surfactants, are biodegradable and environmental friendly. Some of the

microorganisms that produce surfactants during fermentation are *Pseudomonas*, *Rhodococcus*, *Mycobacterium*, *Toruplopsis*, *Bacillus*, *Thiobacillus*, etc (Desai & Banat, 1997). The surfactants derived from microorganisms are glycolipid, lipopeptides, lipoproteins, fatty acids and phospholipids, etc (Desai & Bannat 1997). In enzyme-catalyzed production of biobased surfactants, the common enzymes utilized are lipase, proteinase and glucosidase (Hayes, 2011). Lipase catalyzes ester bond formation between fatty acyl groups and hydroxyl group of alcohols or polyols; amino acid also can act as acyl donor and form ester or amide bond if proteinase are used; glycosidases catalyze the acetal bond formation between saccharides and fatty alcohols (Van Rantwijk, Woudenberg-van Oosterom, & Sheldon, 1999). Lipase-catalyzed reactions were studied in terms of reaction solvent, substrate ratio, reaction time etc. Commonly used acyl acceptors are carbohydrate, sugar alcohol. Acyl donors are various fatty acids or fatty acid esters. Enzymatic synthesis of esters is one of the major methods due to the higher selectivity, relatively lower temperatures (lower than 70°C), lower solvent toxicity, and easier separation of products. Enzyme that have been used for synthesis of esters are subtilisin from *Bacillus amyloliquefaciens* (Rich, Bedell, & Dordick, 1995), lipase from *Candida antarctica* (Pedersen, Wimmer, Emmersen, Degn, & Pedersen, 2002), *Candida rugosa* (Zaidan, Abdul Rahman, Othman, Basri, Abdulmalek, Abdul Rahman, et al., 2012) *Mucor miehei* (Degn, Pedersen, & Zimmermann, 1999), *Humicola lanuginosa* (Ferrer, Cruces, Bernabe, Ballesteros, & Plou, 1999), *Thermomyces lanuginosus* (Tsavas, Polydorou, Fafli, Voutsas, Tassios, Flores, et al., 2002), and alkaline protease from *Streptomyces* spp (Kitagawa, Tokiwa, Fan, Raku, & Tokiwa, 2000). Sugar- fatty acid esters are non-ionic surfactants with a wide range of hydrophilic-lipophilic balance (HLB) values. Since they are biodegradable, non-toxic (Ferrer, Cruces, Bernabe, Ballesteros, & Plou, 1999), non-irritant to skin (Plat & Linhardt, 2001) and odorless, they are widely used in food,

pharmaceutical, cosmetic and detergent industries. Sucrose esters have been approved by Food and Drug Administration and are widely used in food industry, such as wheat products, confectioneries, and dairy products, etc. The functions are various, such as increasing dough resistance to kneading, increasing cake volume, prevent stickiness to the machine, make stable emulsion, improve mouthfeel, prevent staling, etc. (Mitsubishi-Kagaku Food Corporation, <http://www.mfc.co.jp/english/infor.htm>).

The challenge to synthesize sugar-fatty acid ester enzymatically is to find good solvent(s) to solubilize the substrates that have different polarities, at the meantime, not deactivating enzymes. It has been extensively studied in different medium, for example, single phase organic solvent systems (Degn, Pedersen, & Zimmermann, 1999; Ljunger, Adlercreutz, & Mattiasson, 1994;), two organic solvent systems (Kitagawa, Tokiwa, Fan, Raku, & Tokiwa, 2000; Reyes-Duarte, López-Cortés, Ferrer, Plou, & Ballesteros, 2005), non-solvent systems (Martin-Arjol, Isbell, & Manresa, 2015), ionic liquid systems, supercritical carbon dioxide (Habulin, Šabeder, & Knez, 2008) and deep eutectic systems (Pohnlein, Ulrich, Kirschhofer, Nusser, Muhle-Goll, Kannengiesser, et al., 2015). Mixed organic solvents were preferred than single solvent since by varying the ratio of each solvents, solubility of acyl acceptor and enzyme activity can be controlled. To avoid the use of organic solvent and address the solubility issues, solvent-free system were used (Fregapane, Sarney, & Vulfson, 1991; Ward, Fang, & Li, 1997). However, the reaction system had high viscosity and low miscibility (Wei, Yu, Song, & Su, 2003). Ionic solvents were used because of their advantage of low vapor pressures and tunable chemical structure that can solubilize different substrates (Park & Kazlauskas, 2001) while they have disadvantage of extra steps to synthesize and purify of ionic solvents, and some ionic liquid were reported to deactivate the enzyme (Schöfer, Kaftzik, Wasserscheid, & Kragl, 2001). Dimethyl

sulfoxide (DMSO) and 2-methyl-2-butanol (2M2B) were chosen as reaction media to synthesize glucose esters due to relatively high solubility of sugars and being benign to enzymes.

Although the synthesis of sugar fatty acid esters has been studied extensively, their functionality as emulsifiers in the basic oil-in-water systems has not been looked in-depth. Glucose is a cheap carbohydrate with only one primary hydroxyl group, therefore, the high selectivity is expected. Also, not many research (Arcos, Bernabe, & Otero, 1998; Degn, Pedersen, & Zimmermann, 1999; Ljunger, Adlercreutz, & Mattiasson, 1994) studied the synthesis condition and their functionalities. The objectives of this study are to: 1) optimize the synthesis of glucose esters with respect to substrate ratio and fatty acid types, and 2) evaluate the functionality of glucose esters as emulsifiers and compare with commercial sucrose esters with different HLB values.

3.3. Materials and Methods

3.3.1 Reagents

DMSO, 2M2B, D-glucose, HPLC-grade methanol, and molecular sieves (3Å) were purchased from Fisher Scientific (Fair Lawn, NJ). Palmitic acid (98%), lauric acid (>98%), hexanoic acid (>99.5%), and immobilized lipase from *Candida Antarctica* were purchased from Sigma-Aldrich (St. Louis, MO). Pure canola oil was purchased from a local grocery store. Commercial sucrose esters SP30 (sucrose distearate, HLB 6, monoester content 30%), SP50 (sucrose stearate, HLB 11, monoester content 50%), PS750 (sucrose palmitate, HLB 16, monoester content 75%) were donated.

3.3.2 Synthesis of glucose esters

The lipase-catalyzed synthesis of glucose esters was carried out in 50-mL Erlenmeyer flasks following published method with some modifications (Ferrer, Cruces, Bernabe,

Ballesteros, & Plou, 1999). Palmitic acid, lauric acid, and hexanoic acid were used as acyl donor and glucose was used as acyl acceptor for esterification reactions. Molar ratios of fatty acid and sugar were 0.3 mM: 0.1 mM, 0.2 mM: 0.1 mM, 0.1 mM: 0.1 mM, and 0.1 mM: 0.3 mM. One g molecular sieves and 0.25 g immobilized lipases were added in 10 mL solvent mix (80% DMSO and 20% 2M2B). The flasks were incubated at 55°C in a water bath with shaking at 96 rpm for 48 h.

After 48 h of reaction, reactants were centrifuged to obtain the supernatant. The supernatants were placed under a fume hood overnight to evaporate 2M2B. Water (approximately 10:1 v/v of solvent) was added to the medium to precipitate the fatty acid residue and esters. The viscous white slurry were filtered to obtain white solid. The solids were washed with 10 volumes of methanol for 3-4 times to dissolve free fatty acid residue and obtain highly pure esters. The purity of glucose esters was determined with NMR 1D proton test to obtain the area ratio of proton of α carbon ($-\text{CH}_2\text{-COOH}$) of bonded fatty acid and free fatty acid.

3.3.3 Quantitation of fatty acid conversion

The *quantitation* of fatty acids by HPLC method followed a previous study (Reyes-Duarte, López-Cortés, Ferrer, Plou, & Ballesteros, 2005) with slight modification. At 12th, 24th, 36th and 48th h, aliquots of reactant mix were withdrawn and measured for residual free fatty acid by high performance liquid chromatography (Thermo Scientific, ACCELA 1250 HPLC) using a C18 column (Hypersil Gold, 50×2.1 mm, 1.9 μm), a PDA detector (at 200 nm), and EZChrom Elite software (Agilent, Version3.2.1). For palmitate acid detection, methanol: water 70/30 (v/v) with 0.1% v/v acetic acid was used as mobile phase A for the first 2 min, then a gradient from this eluent to pure methanol (B) was continued for 5 min, after which the gradient was changed back to the original mobile phase for 5 min. The flow rate was 0.5 mL/min and the

temperature was 45°C. The method to detect lauric acid was the same as it was for palmitic acid. For hexanoic acid detection, the mobile phase A was methanol: water 80/20 (v/v) with 0.1% acetic acid for 4 min, then a gradient to pure methanol was last for 5min, then it changed back to A. The flow rate of mobile phase was 0.5 mL/min for all fatty acids. The conversion of fatty acid was calculated as:

$$\text{Conversion of fatty acid} = (X_0 - X_1) / X_0 \times 100\%$$

X_0 - fatty acid concentration at the start of reaction

X_1 -residual fatty acid concentration in reaction mix at different time points.

X_0 - theoretical concentration of fatty acid that can be fully converted to ester.

From stoichiometry of reaction systems, X_0 is the 1/3 of initial concentration at 3:1 acid:sugar ratio, and 1/2 of initial concentration for 2:1 ratio.

3.3.4 Identification of esters

Agilent Quadrupole Time-of-Flight (QTOF) 6540 liquid chromatography mass spectrometry (LC/MS) was used to identify reaction products. A XDB C18 column (4.6×50 mm, 1.8 μm) and an electrospray ionization detector were used. The products were scanned in the negative mode from 100-1000 Daltons. A gradient from 95% mobile phase A (water, 100%) and 5% B (methanol, 100%) to 95% B and 5% A was applied for 20 min and kept for another 5 min. The flow rate of mobile phase was 0.8 mL/min. Nuclear magnetic resonance (NMR) (Bruker Avance III 600, Billerica, MA and Karlsruhe, Germany) was used to confirm the formation of ester bonds and chemical shifts of important carbon and hydrogen atoms. The products were dissolved in deuterated DMSO to achieve a concentration range of 50-800 mg/mL. Heteronuclear single quantum coherence (HSQC) and heteronuclear multiple bond correlation

(HMBC) spectroscopy for ^1H and ^{13}C were used to determine the ester bond formation. The data were analyzed with the TopSpin software (Bruker, Billerica, MA).

Chemical shifts, splitting patterns, J-coupling and positions of hydrogen and carbon for reactants and products are as follows:

Glucose, the spectrum showed it was a mixture of α - and β -anomers. ^1H NMR (600 MHz, $\text{DMSO-}d_6$) δ 4.91 (t, $J = 4.2$ Hz, 1H, H-1), 3.42 (dt, $J = 9.1, 4.3$ Hz, 1H, H-2), 3.11 (ddd, $J = 10.0, 6.7, 3.6$ Hz, 1H, H-3), 3.04 (td, $J = 9.3, 5.4$ Hz, 1H, H-4), 3.58 – 3.52 (m, 1H, H-5), 3.48 – 3.44 (m, 1H, H-6a), 3.60 (dd, $J = 11.5, 5.9$ Hz, 1H, H-6b). ^{13}C NMR (151 MHz, $\text{DMSO-}d_6$) δ 92.67 (C-1), 72.809 (C-2), 73.541 (C-3), 71.03 (C-4), 72.403 (C-5), 61.676 (C-6).

Palmitic acid. ^1H NMR (600 MHz, $\text{DMSO-}d_6$) δ 11.93 (s, 1H, -COOH), 2.166 (t, $J = 7.5$ Hz, 2H, -CH₂CO-), 1.486 (t, $J = 7.2$ Hz, 2H, -CH₂-CH₂-CO-), 1.205-1.294 (s, 24H, chain), 0.853 (m, 3H, -CH₃). ^{13}C NMR (151 MHz, $\text{DMSO-}d_6$) δ 174.75 (C=O), 34.119 (-CH₂-CO-), 31.87 (-CH₂-CH₂-CO-), 29.661-29.661, 29.31, 29.16, 24.997, 22.617 (-CH₂- palmitic acid backbone), 14.263 (-CH₃).

Lauric acid. ^1H NMR (600 MHz, $\text{DMSO-}d_6$) δ 11.95 (s, 1H, -COOH), 2.176 (t, $J = 7.5$ Hz, 2H, -CH₂CO-), 1.495 (t, $J = 7.4$ Hz, 2H, -CH₂-CH₂-CO-), 1.244-1.297 (s, 16H, chain), 0.86 (d, $J = 7.5$ Hz, 3H, -CH₃). ^{13}C NMR (151 MHz, $\text{DMSO-}d_6$) δ 174.85 (C=O), 34.12 (-CH₂-CO-), 31.816, 29.541, 29.528, 29.449, 29.285, 29.243, 29.079, 24.979, 22.597 (-CH₂- lauric acid backbone), 14.344 (-CH₃).

Hexanoic acid. ^1H NMR (600 MHz, $\text{DMSO-}d_6$) δ 11.889 (s, 1H, -COOH), 2.172 (t, $J = 7.6$ Hz, 2H, -CH₂CO-), 1.502-1.264 (6H, chain), 0.859 (t, $J = 7.1$ Hz, 3H, -CH₃). ^{13}C NMR (151 MHz, $\text{DMSO-}d_6$) δ 174.86 (C=O), 34.05 (-CH₂-CO-), 31.25, 24.64, 22.31 (-CH₂- hexanoic acid backbone), 14.16 (-CH₃).

6-O-Palmitoylglucopyranose. ^1H NMR (600 MHz, $\text{DMSO-}d_6$) δ 4.91 (t, $J = 4.4$ Hz, 1H, H-1), 3.137 (m, 1H, H-2), 3.44 (td, $J = 9.1, 4.5$ Hz, 1H, H-3), 3.042 (td, $J = 9.3, 5.5$ Hz, 1H, H-4), 3.78 (dd, $J = 9.5, 6.5$ Hz, 1H, H-5), 4.28 (d, $J = 11.5$ Hz, 1H, H-6a), 4.00 (dd, $J = 11.7, 6.2$ Hz, 1H, H-6b), 2.28 (t, $J = 7.4$ Hz, 2H, -CH₂-CO-), 1.51 (q, $J = 7.2$ Hz, 2H, -CH₂-CH₂-CO-), 1.25 (s, 24H, chain), 0.87 (t, $J = 6.9$ Hz, 3H, -CH₃). ^{13}C NMR (151 MHz, $\text{DMSO-}d_6$) δ 92.755 (C-1), 72.66 (C-2), 73.336 (C-3), 71.035 (C-4), 69.601 (C-5), 64.355 (C-6), 173.363 (C=O), 33.906 (-CH₂-CO-), 31.778 (-CH₂-CH₂-CO-), 29.534 – 28.931, 24.94, 22.569 (-CH₂- palmitoyl backbone), 14.341 (-CH₃).

6-O-Lauroylglucopyranose. ^1H NMR (600 MHz, $\text{DMSO-}d_6$) δ 4.909 (d, $J = 4.7$ Hz, 1H, H-1), 3.137 (m, 1H, H-2), 3.439 (td, $J = 9.2, 4.3$ Hz, 1H, H-3), 3.043 (m, 1H, H-4), 3.774 (dd, $J = 9.7, 6.8$ Hz, 1H, H-5), 4.277 (d, $J = 11.9$ Hz, 1H, H-6a), 4.006 (dd, $J = 11.7, 6.2$ Hz, 1H, H-6b), 2.28 (t, $J = 7.3$ Hz, 2H, -CH₂-CO-), 1.52 (t, $J = 7.1$ Hz, 2H, -CH₂-CH₂-CO-), 1.25 (s, 16H, chain), 0.87 (t, $J = 6.8$ Hz, 3H, -CH₃). ^{13}C NMR (151 MHz, $\text{DMSO-}d_6$) δ 92.754 (C-1), 72.659 (C-2), 73.334 (C-3), 71.027 (C-4), 69.602 (C-5), 64.362 (C-6), 173.272 (C=O), 33.909 (-CH₂-CO-), 31.814, 29.541, 29.526, 29.434, 29.268, 29.24, 28.964, 24.94, 22.592 (-CH₂- lauroyl backbone), 14.42 (-CH₃).

6-O-Hexanoylglucopyranose

^1H NMR (600 MHz, $\text{DMSO-}d_6$) δ 4.911 (d, $J = 3.9$ Hz, 1H, H-1), 3.132 (m, 1H, H-2), 3.43 (m, 1H, H-3), 3.055 (dd, $J = 17.2, 7.4$ Hz, 1H, H-4), 3.771 (s, 1H, H-5), 4.29 (dd, $J = 19.7, 11.6$ Hz, 1H, H-6a), 3.997 (m, 1H, H-6b), 2.293, 1.51 (dt, $J = 14.7, 7.9$ Hz, 2H, $-\text{CH}_2\text{-CO-}$), 1.576-1.191 (m, 6H, caproyl backbone), 0.873 (d, $J = 8.3$ Hz, 3H, $-\text{CH}_3$). ^{13}C NMR (151 MHz, $\text{DMSO-}d_6$) δ 173.444 (C=O), 34.177 ($-\text{CH}_2\text{-CO-}$), 31.097, 24.643, 22.287 ($-\text{CH}_2\text{-}$ hexanoyl backbone).

3.3.5 Initial sugar solubility

Initial sugar solubility of ester was tested using an HPLC. After mixing reactants except lipase in the solvents as described in synthesis method above, flasks were incubated in the water bath at 55°C at 3.8 rpm for 30 min. After cooling down, aliquots from each flask were run through HPLC. A carbohydrate column (HyperRez XP Carbohydrate H+, 300×7.7 mm, $8 \mu\text{m}$), a guard column (HyperRez XP Carbohydrate H+, 50×7.7 mm, $8 \mu\text{m}$) and a RI detector were used. The temperature for guard column and carbohydrate column were 65°C and 70°C , the flow rate was 0.4 mL/min.

3.3.6 Emulsion stability Index

0.01% w/w, 0.1% w/w and 0.5 % w/w 100g ester solutions were prepared and mixed with 10g canola oil, control treatment did not contain any esters. The mixtures were sonicated for 10 min to improve ester dispersion. The mixture was homogenized for 2 min at 15,000 rpm using a blender (Bamix Type M 150) and 10 μL - 20 μL aliquots of emulsion were diluted with 2 mL deionized water in the spectrophotometer cuvette. Absorbance of emulsion at 500 nm was measured at 0, and 20 min. The emulsion stability index (ESI) was calculated as:

$ESI=A_0*20/[A_0-A_{20}]$. A_0 and A_{20} were the absorbance obtained at 0 min and 20 min (Pearce & Kinsella, 1978). The higher ESI values indicated higher emulsion stability.

3.3.7 Emulsion droplet size distribution

The emulsion droplet size and distribution was measured by Malvern Particle Size analyzer (Mastersizer Hydro 2000). Emulsions with 0.5% w/w level of esters were prepared as above and were introduced into the instrument until a laser obscuration of 10-20% was achieved. Measurements were taken at time 0 h, 6th h, 24th h, 3rd d and 7th d.

3.3.8 Contact angle measurement

Contact angle measurement followed a previous research (Crowley, Desautel, Gazi, Kelly, Huppertz, & O'Mahony, 2015). Pellets of each of the esters were prepared with 0.08g powdered ester, placing them on a 13 mm pellet die and pressing under a force of 5000 kg for 2 min in Carver Press (model 3619, Carver Inc, Wabash, IN). Contact angle measurement was conducted using a goniometer (Rame-Hart Model 250 Standard Goniometer). Approximately 4 μ L water droplet was dispensed on the pellet's surface placed on a stage. Side view pictures were taken immediately after the water droplet left the syringe tip using a high-resolution camera.

3.3.9 Statistical analysis

Statistic test was conducted using SAS 9.4 software (SAS Institute Inc., Cary, NC). Proc GLIMMIX test was used to determine significant difference between treatments ($P<0.05$). At least three observations for each treatment were measured for conversion, sugar solubility, contact angle, emulsion stability and droplet size analysis.

3.4. Results and Discussion

3.4.1 Fatty acid conversion

Figure 3.1 (a), (b) and (c) show the conversion percentages of palmitic acid, lauric acid and hexanoic acid over 48 h, respectively. The fatty acid/glucose molar ratios were compared for optimum ratio for the highest conversion. For each glucose-fatty acid ester studied, the highest conversion of fatty acid was achieved when the ratio was 3:1 (97.17%, 76.57%, 113.11% for glucose palmitate, laurate and hexanoate, respectively). The second highest conversions were at the ratio of 2:1 (75.96%, 62.82%, 73.66%, respectively). The 1:1 and 1:3 ratios did not have any significantly differences; the conversion was around 50%, 40%, 30%, respectively for palmitic acid, lauric acid and hexanoic acid. Higher concentrations of FA favored formation of the products as explained by the equilibrium constant:

$$k = \frac{[\text{ester}][\text{water}]}{[\text{fatty acid}][\text{glucose}]} \quad (\text{Equation 1});$$

$$\text{or, } [\text{ester}] = k \times \frac{[\text{fatty acid}][\text{glucose}]}{[\text{water}]} \quad (\text{Equation 2}).$$

At a given temperature (55°C in this study), k is a constant. At higher molar ratios of fatty acid to glucose (2:1 or 3:1), the limiting reactant is glucose, thus, only 1/3rd of available fatty acid is converted to ester and water. Therefore, the increase in fatty acid concentration in reaction mix would be larger than the increase in water amount. Meanwhile, the glucose concentration remained at a similar level compared to the reaction when the ratio was 1:1. Therefore, the increase of fatty acids amount increased the ester amount according to Equation 2, as reflected by the conversion percentage. For the reactants ratio of 1:1 and 1:3 of fatty acid to glucose, glucose solubility was limited in the medium as glucose crystals could be seen throughout the reaction, and fatty acid were completely dissolved. For these conditions, terms in Equation 2 did not have significant changes, indicating similar conversions for these two ratios. Among various fatty acids, the hexanoic acid had the highest conversion percentage at 3:1 ratio (113%). This could be due to the formation of diesters, along with monoesters, but at low levels

since they were not detected by LCMS and NMR. Two reasons could contribute to the higher conversion of hexanoic acid: shorter carbon chains making the solvent more polar resulting in higher glucose solubility (data will be shown in later section) than other two fatty acids, and possible steric hindrance- the smaller molecule would have easier access to enzyme active sites and result in more esterification.

3.4.2 Initial substrate concentration and initial conversion rates

Influence of reactant concentrations on the initial conversion rates (linear range for first three hours) is presented in Table 3.1. For the molar ratios of 3:1, 2:1, 1:1 (fatty acid/glucose) in esterification of palmitic acid and lauric acid, fatty acids were completely soluble in solvent, so, the fatty acids concentration increased as molar ratio increased leading to higher initial fatty acid conversion rates. For molar ratio increase from 1:1 to 3:1 (fatty acid concentration increase from 74.3 mM to 243.7 mM), the initial conversion rates also increased threefold from 3.3 $\mu\text{mol}/(\text{min} \cdot \text{g})$ to 9.6 $\mu\text{mol}/(\text{min} \cdot \text{g})$. Glucose did not completely solubilize for these molar ratios of 1:1 and 1:3 (fatty acid/glucose), because sugar crystals were seen in the medium, thus we examined the soluble glucose concentration in reactant. For fatty acid: glucose molar ratios of 1:1 and 1:3 with increase in glucose concentration from 21 mM to 28.2 mM, the initial conversion rates increased from 3.3 $\mu\text{mol}/(\text{min} \cdot \text{g})$ to 5.2 $\mu\text{mol}/(\text{min} \cdot \text{g})$. From these two comparisons, it is seen that the concentration of both fatty acid and glucose had direct and proportional relationship with the initial conversion rate, indicating the reaction to be a first-order reaction in terms of either fatty acid or glucose (Degn & Zimmermann, 2001). For hexanoic acid, the increase of fatty acid concentration from 146.5 mM to 235.2 mM almost quadrupled the rate (Table 3.1), and the glucose solubility did not change over the three ratios and did not affect the initial rate. The highest conversion rate obtained was 11.83 $\mu\text{mol}/(\text{min} \cdot \text{g})$, which is similar to reported 15.2

$\mu\text{mol}/(\text{min}\cdot\text{g})$ for synthesis of glucose myristate (Degn & Zimmermann, 2001) that was conducted in t-butanol: pyridine system. Although the glucose solubility was higher in our system, the deactivation of lipase in DMSO is stronger than pyridine. They reported that below molar ratio of 10:1 up to 20 mg/mL soluble glucose concentration, the initial reaction rate increased as the ratio increased, which was consistent with our study when we kept the sugar level constant. The effect of lauric acid and glucose concentration on initial conversion rates in the single solvent 2M2B was reported to increase when either of the two substrate concentrations increased, however, lauric acid was saturated at 140 mM, glucose was not saturated up to 50 mM (Flores, Naraghi, Engasser, & Halling, 2002). In our study, we did not find saturation levels for both substrates even at much higher fatty acid concentration of 259.6 mM. The reason for the difference in solubility could be that our bi-solvent system dissolved more fatty acids boosting the initial rates.

Initial sugar solubility can also be associated with medium hydrophobicity (Pedersen, Wimmer, Emmersen, Degn, & Pedersen, 2002; Reyes-Duarte, López-Cortés, Ferrer, Plou, & Ballesteros, 2005). Glucose solubility increased as the chain length of the fatty acid decreased at each substrate ratio. The solubility is also associated with the amount of sugar that was put into the medium, as it increased when the ratio of sugar increased, since they themselves created a more polar environment. Though not measured in our study, previous studies indicated as the esters were being produced, glucose solubility would increase through hydrophobic interaction (Degn & Zimmermann, 2001; Tsavas, et al., 2002).

The initial conversion rates may or may not relate to the length of the acyl donor, as variously reported. Some studies reported the reaction rate with was faster for shorter chain length of fatty acid (C4-C12) that were esterified with disaccharide (Pedersen, Wimmer,

Emmersen, Degn, & Pedersen, 2002). Adelhorst, Bjokling, Godtfredsen and Kirk (1990) reported the enzyme showed faster reaction with longer fatty acids (C12-C18) than shorter acids (C8-C10) in solvent free condition. However, Degn, Pedersen, & Zimmermann, (1999) found that the initial reaction rate was independent within chain length C2-C20 of acyl donors for glucose. Our study also indicates that the initial rates and chain length were independent. The difference could be due to the difference in substrate and reaction conditions.

3.4.3 Product identification and complete ¹H and ¹³C assignment for reactants and products.

The formation of glucose palmitate, glucose laurate and glucose hexanoate were confirmed by LCMS (data not shown) and NMR techniques. In the HMBC graph (Figure 3.2) of reaction mixture of glucose and palmitic acid, the sixth protons (4.01 and 4.27 ppm) of glucose were seen to have interacted with carbonyl carbon indicating the formation of ester bonds. The chemical shifts of H-6 of glucose and α -carbon proton of fatty acid in the esters to the higher fields indicated the chemical environment change due to esterification that caused de-shielding effect (Kitagawa, Tokiwa, Fan, Raku, & Tokiwa, 2000; Pedersen, Wimmer, Emmersen, Degn, & Pedersen, 2002; Walsh, Bombyk, Wagh, Bingham, & Berreau, 2009). We had two chemical shifts for the sixth proton in glucose indicating the alpha and beta conformation of the D-glucose (Roslund, Tähtinen, Niemitz, & Sjöholm, 2008). Similar NMR graph for glucose laurate and glucose hexanoate esters were obtained.

We successfully obtained the purity of 95.50%, 98.97% of glucose palmitate and glucose laurate, respectively. Due to relatively high solubility of glucose hexanoate in both hydrophobic and hydrophilic environment, it was not possible to purify the ester by solvent extraction. NMR data for glucose hexanoate was obtained from reaction mixture rather than pure product.

3.4.4 Contact angle on product surface

Due to the low solubility of glucose palmitate in water, it was hard to measure the critical micelle concentration and surface tension; therefore, contact angle was measured for the esters to compare the relative hydrophobicity. The hydrophobicity is generally positively related to contact angle (Daffonchio, Thaveesri, & Verstraete, 1995) and reversely related to HLB value (Griffin, 1949). The contact angles and HLB values for glucose palmitate, glucose laurate, SP30, SP50 and PS750 were 98.6 (HLB 8.6), 93.6 (HLB 9.9), 76.9 (HLB 6), 44.8 (HLB 11) and 28.1 (HLB 16) respectively. The contact angles of our products followed the trend: the longer alkyl chain of palmitic acid made glucose palmitate more hydrophobic. The HLB values for these two products were calculated respectively according to the method of Griffin (1955). The HLB value indicates that both of the esters can perform as emulsifiers and wetting agents for oil-in-water system (value from 7-18, according to Griffin (1946)). Commercial SP50 and PS750 esters may have more function in detergent and solubilizing application because they had higher HLB value (Griffin, 1949). Overall, the larger the contact angle the ester had, the lower the HLB value they were, except that SP30 had lower HLB value than glucose palmitate but it had lower contact angle than the other.

3.4.5 Emulsion stability of esters

The emulsion stability index (ESI) for esters are shown in Figure 3.3. Glucose esters were not water-soluble and they stayed on top of aqueous phase; whereas, sucrose esters were dispersible but not soluble. Glucose esters demonstrated stabilizing effect compared to control treatment. The ESI increased as the concentration increased for all esters, which indicated lower concentrations of esters did not completely cover the oil surface to prevent creaming, the addition of more esters covered more surface area so the creaming process were retarded. At low

concentrations of 0.01% for glucose palmitate and glucose laurate, the ESI values (171.4, 178.0 respectively) were slightly higher than control treatment (138.8). At medium concentration of 0.1%, glucose palmitate ESI (315.3) was twice as much as control, whereas glucose laurate ESI (163.4) did not increase compared to 0.01%. At 0.5%, glucose palmitate again showed stronger stabilizing effect (ESI 664.7) than glucose laurate (460.8). These data show that glucose palmitate had better stabilizing effect compared to glucose laurate. One reason for this could be the stronger hydrophobic interactions by longer alkyl chains in the molecule with each other and with oil droplets, which can form a more compact structure than glucose laurate (Ferrer, Comelles, Plou, Cruces, Fuentes, Parra, et al., 2002).

For commercial sucrose esters, similarly, the ESI increased as the concentration increased. Compared with glucose esters, the sucrose esters had better stabilizing effect since the ESI were higher than those of in-house glucose esters at every concentration. Particularly, at 0.5%, the ESI were significantly higher (1351.8, 1212.5, 1492.3 for SP30, SP50, PS750 respectively) than glucose esters.

3.4.6 Emulsion droplet size distributions

Droplet distribution and diameter parameters for control, glucose palmitate, glucose laurate, and sucrose esters SP30 are presented in Figure 3.4 and Table 3.2 respectively. It was obvious for control and glucose esters that the distribution had undergone from single-modal to bio-modal or tri-modal change during 7-day storage, indicating the size were diverging into bigger or smaller. The appearance of the peaks to the right side of the distribution indicated the presence of droplets that have not been completely covered by the surfactants have experienced coalescence (McClement, 2004a). The sucrose esters emulsified systems were relatively stable as

the distributions were bio-modal or tri-modal throughout. Only one for the sucrose esters is presented because the distribution of the patterns were similar.

Mean and standard deviation of the droplet diameters for each treatment for 7-day time points are reported in Table 3.2; for multiple comparison, data were transformed to a natural log scale to fit the normal distribution. For all the diameters, both fixed effects (type of esters and time) had significant effects on change in droplet diameters ($P < 0.05$). The interaction of treatment and time was also significant ($P < 0.05$), meaning the diameters changed differently among all the treatments at different times. From 0 h to 7th day, $D[0.1]$ decreased gradually for control (5.2 μm to 2.0 μm), glucose palmitate (1.7 μm to 0.3 μm) and glucose laurate-stabilized emulsions (2.0 μm to 0.3 μm), respectively; multiple comparison of log-transformed data indicated that the change was significant ($P < 0.05$). However, droplet size almost did not change for sucrose esters. This indicated that smaller droplets were decreasing in control and glucose esters and were undergoing coalescence. This can be confirmed by $D[0.9]$ and volume mean diameter $D[4,3]$ that the size of droplets for these three esters were increasing. However, they increased differently – for the volume mean diameter, control group experienced the greatest change (from 37.2-74.7 μm), glucose palmitate stabilized droplet changed from 18.2 μm to 32.7 μm , glucose laurate stabilized droplet changed from 16.6 μm to 47.4 μm . This indicated that glucose palmitate and glucose laurate generated smaller droplets overall and they had stabilizing effect on droplet size, but could not completely prevent coalescence. Compared with glucose laurate, glucose palmitate is relatively more effective in stabilizing the oil-in-water emulsions. The commercial sucrose esters can effectively prevent coalescence, as indicated by no significant changes for diameter parameters over time. Three mechanisms can explain the phenomenon we observed: hydrophobic interaction (Ferrer, et al., 2002), steric stabilization (Nilsson &


Bergenståhl, 2007) and adsorption on the interfacial surface (McClement, 2004b). Between glucose palmitate and glucose laurate, the former has longer alkyl chain, likely with the stronger hydrophobic interaction with oil droplets resulting in larger area coverage on the droplet to prevent coalescence. Also, the bigger molecule of glucose palmitate has stronger steric hindrance that prevent droplet from aggregating. For sucrose esters, both steric hindrance and adsorption contribute to the better stabilization effect. Sucrose esters has bigger size because the presence of a fructose moiety in addition to glucose, meanwhile the presence of more hydroxyl group made it easier to solubilize in the continuous phase and easier to adsorb at the interface, whereas glucose esters had much lower solubility in the continuous phase, their adsorption at the interface were much slower.

3.5 Conclusion


Glucose-fatty acid monoesters were successfully synthesized in tert-amyl butanol and DMSO mixture solvent system in lipase catalyzed reactions with high level of conversion. Products were purified with solvent extraction. Synthesized and commercial esters were compared for emulsion capabilities. Glucose esters stabilized oil droplets to some extent, but could not completely prevent coalescence compared to commercial sucrose esters. The relatively smaller sizes of glucose esters and low aqueous solubility can explain their difference of emulsifying property.

Acknowledgement


Iowa State University Agriculture Experiment Station and USDA/NIFA multistate project S1041 partially contributed to the project. We express sincere appreciation to Iowa State University Chemical Instrumentation Facility staff member Sarah Cady for training and assistance to interpret the AVIII-600 results included in this publication.



a



b



c

Fig 3.1 Conversion percentage of palmitic acid (a), lauric acid (b) and hexanoic acid (c) during 48 h. Lines with diamond, square, triangle and cross represents molar ratio of 3:1, 2:1, 1:1 and 1:3 of fatty acid/glucose respectively. Standard deviations are shown.




Fig 3.2 Heteronuclear multiple bond correlation (HMBC) of reaction mixture of palmitic acid (PA) and glucose. The horizontal and vertical axis indicate ^1H proton and ^{13}C chemical shift (ppm) respectively. The interaction of proton of C6 of glucose ring with the carbonyl carbon demonstrated ester bond has been formed. The ester bond also caused the α -carbon (the one next to the carbonyl carbon) chemical shift to the higher field.







Fig 3.3 Emulsion stability index of glucose esters and sucrose esters. Mean value and standard deviation are shown.




a



b



c



d

Fig 3.4 Droplet size distribution of emulsions of control treatment (a), glucose palmitate (b), glucose laurate (c), sucrose ester SP30 (d) at 0 h, 6th h, 24th h, 3rd day, 7th day.

Table 3.1 Initial reaction rate ($\mu\text{mol}/\text{min}\cdot\text{g}$) and substrate solubility (mM)

Ratio of fatty acid/glucose	PA initial conversion rate ($\mu\text{mol}/(\text{min} \cdot \text{g})$)	Initial sugar solubility (mM)	PA concentration (mM)	LA initial conversion rate ($\mu\text{mol}/(\text{min} \cdot \text{g})$)	Initial sugar solubility (mM)	LA concentration (mM)	HA initial conversion rate ($\mu\text{mol}/(\text{min} \cdot \text{g})$)	Initial sugar solubility(mM)	HA concentration (mM)
3:1	9.6 ^a	16.9 ^c	243.7 ^a	7.2 ^a	21.1 ^b	259.6 ^a	11.8 ^a	26.2 ^a	235.2 ^a
2:1	4.8 ^b	18.3 ^{bc}	158.0 ^b	5.3 ^a	22.5 ^b	178.3 ^b	3.8 ^b	22.8 ^a	146.5 ^b
1:1	3.3 ^b	21.0 ^b	74.3 ^c	2.2 ^b	21.6 ^b	103.8 ^c	2.3 ^b	28.1 ^a	75.5 ^c
1:3	5.2 ^{ab}	28.2 ^a	72.9 ^c	2.6 ^b	33.6 ^a	103.6 ^c	2.7 ^b	30.5 ^a	74.4 ^c

PA- palmitic acid, LA- lauric acid, HA- hexanoic acid. Different letters indicate significantly difference in a row ($P < 0.05$).

Table 3.2 Diameter summary of droplet size distribution

	D[0.1]					D[0.5]				
	0h	6h	24h	3d	7d	0h	6h	24h	3d	7d
Control	5.2±1.2	1.2±0.3	1.2±0.1	2.3±1.2	2.0±2.0	26.8±4.6	10.4±6.5	58.0±28.3	65.5±41.9	17.1±2.2
GP	1.7±0.1	0.6±0.1	0.6±0.0	0.4±0.1	0.3±0.0	6.6±0.5	2.4±0.1	2.5±0.1	2.8±0.3	2.6±0.8
GL	2.0±0.3	0.4±0.1	0.4±0.1	0.4±0.0	0.3±0.0	9.8±0.8	2.4±0.3	2.6±1.0	5.9±4.2	3.1±1.2
SP30	0.4±0.1	0.3±0.1	0.4±0.0	0.3±0.1	0.3±0.0	6.4±1.8	4.7±2.4	8.3±2.7	6.9±2.3	7.1±2.0
SP50	0.3±0.1	0.3±0.0	0.3±0.0	0.2±0.0	0.3±0.1	5.4±0.8	2.8±1.4	4.3±2.1	2.1±1.4	4.5±2.8
PS750	0.5±0.2	0.3±0.0	0.3±0.1	0.3±0.0	0.2±0.0	4.7±2.1	1.3±0.4	4.5±1.4	8.1±3.5	4.1±1.0

	D[0.9]					D[4.3]				
	0h	6h	24h	3d	7d	0h	6h	24h	3d	7d
Control	83.2±15.3	201.0±3.6	232.0±24.1	314.1±91.4	210.7±55.3	37.2±6.8	58.5±6.7	94.6±17.3	124.1±34.2	74.7±17.5
GP	34.1±4.1	20.6±22.1	148.5±93.5	91.3±23.6	99.5±96.7	18.2±4.8	17.8±6.4	42.3±29.2	23.7±5.8	32.7±30.4
GL	36.8±2.8	35.5±41.5	117.6±30.0	172.5±17.0	160.5±34.2	16.6±1.5	21.7±10.1	32.5±9.2	57.2±10.6	47.4±12.3
SP30	23.2±4.0	26.1±5.6	30.9±4.8	24.0±6.6	25.9±3.1	8.1±1.6	5.2±3.3	6.9±2.0	5.5±5.0	6.4±5.9
SP50	19.3±5.1	12.9±6.0	17.4±0.6	15.3±1.6	15.2±1.6	10.3±1.7	11.1±2.0	17.7±0.8	12.0±0.8	17.0±1.6
PS750	24.4±7.6	22.5±6.8	29.8±3.1	64.1±57.2	34.5±10.1	10.1±4.8	6.6±2.4	10.7±1.0	21.9±17.9	12.9±3.9

Mean value and standard deviation of diameters are shown in the table. For data analysis, data were transformed to a log scale to fit a normal distribution.

3.6 Reference

- Adelhorst, K., Bjokling, F., Godtfredsen, S. E., & Kirk, O. (1990). Enzyme catalysed preparation of 6-O-acylglucopyranosides. *Synthesis*(2), 112-115.
- Arcos, J., Bernabe, M., & Otero, C. (1998). Quantitative enzymatic production of 6 - O - acylglucose esters. *Biotechnology and Bioengineering*, 57(5), 505-509.
- Crowley, S. V., Desautel, B., Gazi, I., Kelly, A. L., Huppertz, T., & O'Mahony, J. A. (2015). Rehydration characteristics of milk protein concentrate powders. *Journal of Food Engineering*, 149, 105-113.
- Daffonchio, D., Thaveesri, J., & Verstraete, W. (1995). Contact angle measurement and cell hydrophobicity of granular sludge from upflow anaerobic sludge bed reactors. *Applied and environmental microbiology*, 61(10), 3676-3680.
- Degn, P., Pedersen, L. H., & Zimmermann, W. (1999). Lipase-catalysed synthesis of glucose fatty acid esters in tert-butanol. *Biotechnology letters*, 21(4), 275-280.
- Degn, P., & Zimmermann, W. (2001). Optimization of carbohydrate fatty acid ester synthesis in organic media by a lipase from *Candida antarctica*. *Biotechnology and bioengineering*, 74(6), 483-491.
- Deleu, M., & Paquot, M. (2004). From renewable vegetables resources to microorganisms: new trends in surfactants. *Comptes Rendus Chimie*, 7(6), 641-646.
- Desai, J. D., & Banat, I. M. (1997). Microbial production of surfactants and their commercial potential. *Microbiology and Molecular biology reviews*, 61(1), 47-64.
- Ferrer, M., Comelles, F., Plou, F. J., Cruces, M. A., Fuentes, G., Parra, J. L., & Ballesteros, A. (2002). Comparative surface activities of di-and trisaccharide fatty acid esters. *Langmuir*, 18(3), 667-673.
- Ferrer, M., Cruces, M. A., Bernabe, M., Ballesteros, A., & Plou, F. J. (1999). Lipase-catalyzed regioselective acylation of sucrose in two-solvent mixtures. *Biotechnology and bioengineering*, 65(1), 10-16.
- Flores, M. V., Naraghi, K., Engasser, J. M., & Halling, P. J. (2002). Influence of glucose solubility and dissolution rate on the kinetics of lipase catalyzed synthesis of glucose laurate in 2 - methyl 2 - butanol. *Biotechnology and bioengineering*, 78(7), 815-821.
- Fregapane, G., Sarney, D. B., & Vulfson, E. N. (1991). Enzymic solvent-free synthesis of sugar acetal fatty acid esters. *Enzyme and microbial technology*, 13(10), 796-800.
- Greek, B. (1991). Sales of detergents growing despite recession. *Chemical & Engineering News*, 69(4), 25-&.
- Greek, B. F. (1990). Detergent industry ponders products for new decade. *Chemical & Engineering News*, 68(5), 37-&.
- Griffin, W. C. (1946). Classification of surface-active agents by " HLB". *J Soc Cosmetic Chemists*, 1, 311-326.
- Habulin, M., Šabeder, S., & Knez, Ž. (2008). Enzymatic synthesis of sugar fatty acid esters in organic solvent and in supercritical carbon dioxide and their antimicrobial activity. *The Journal of Supercritical Fluids*, 45(3), 338-345.
- Hayes, D. G. (2011). LIPIDS-Bioprocessing Methods to Prepare Biobased Surfactants for Pharmaceutical Products. *American Pharmaceutical Review*, 14(3), 8.
- Kitagawa, M., Tokiwa, T., Fan, H., Raku, T., & Tokiwa, Y. (2000). Transesterification of divinyladipate with glucose at various temperatures by an alkaline protease of *Streptomyces* sp. *Biotechnology letters*, 22(10), 879-882.

- Lee, S. H., Ha, S. H., Hiep, N. M., Chang, W.-J., & Koo, Y.-M. (2008). Lipase-catalyzed synthesis of glucose fatty acid ester using ionic liquids mixtures. *Journal of biotechnology*, 133(4), 486-489.
- Ljunger, G., Adlercreutz, P., & Mattiasson, B. (1994). Lipase catalyzed acylation of glucose. *Biotechnology letters*, 16(11), 1167-1172.
- McClement, D.J. (2004a). Food Emulsion. (2 nd.). Boca Raton: CRC Press, (Chapter 1).
- McClements, D. J. (2004b). Food emulsions: principles, practice and techniques (2nd ed.) Boca Raton: CRC Press. (Chapter 3).
- Mitsubishi-Kagaku Food Corporation. (2016). Available from <http://www.mfc.co.jp/english/infor.htm>. Accessed 2.21.16.
- Martin-Arjol, I., Isbell, T. A., & Manresa, A. (2015). Mono-Estolide Synthesis from trans-8-Hydroxy-Fatty Acids by Lipases in Solvent-Free Media and Their Physical Properties. *Journal of the American Oil Chemists' Society*, 92(8), 1125-1141.
- Nilsson, L., & Bergenståhl, B. (2007). Emulsification and adsorption properties of hydrophobically modified potato and barley starch. *Journal of agricultural and food chemistry*, 55(4), 1469-1474.
- Park, S., & Kazlauskas, R. J. (2001). Improved preparation and use of room-temperature ionic liquids in lipase-catalyzed enantio- and regioselective acylations. *The Journal of organic chemistry*, 66(25), 8395-8401.
- Pearce, K. N., & Kinsella, J. E. (1978). Emulsifying properties of proteins: evaluation of a turbidimetric technique. *Journal of Agricultural and Food Chemistry*, 26(3), 716-723.
- Pedersen, N. R., Wimmer, R., Emmersen, J., Degn, P., & Pedersen, L. H. (2002). Effect of fatty acid chain length on initial reaction rates and regioselectivity of lipase-catalysed esterification of disaccharides. *Carbohydrate Research*, 337(13), 1179-1184.
- Plat, T., & Linhardt, R. J. (2001). Syntheses and applications of sucrose-based esters. *Journal of Surfactants and Detergents*, 4(4), 415-421.
- Pohnlein, M., Ulrich, J., Kirschhofer, F., Nusser, M., Muhle-Goll, C., Kannengiesser, B., Brenner-Weiss, G., Luy, B., Liese, A., Syldatk, C., & Hausmann, R. (2015). Lipase-catalyzed synthesis of glucose-6-O-hexanoate in deep eutectic solvents. *European Journal of Lipid Science and Technology*, 117(2), 161-166.
- Reyes-Duarte, D., López-Cortés, N., Ferrer, M., Plou, F. J., & Ballesteros, A. (2005). Parameters affecting productivity in the lipase-catalysed synthesis of sucrose palmitate. *Biocatalysis and Biotransformation*, 23(1), 19-27.
- Rich, J. O., Bedell, B. A., & Dordick, J. S. (1995). Controlling enzyme-catalyzed regioselectivity in sugar ester synthesis. *Biotechnology and Bioengineering*, 45(5), 426-434.
- Roslund, M. U., Tähtinen, P., Niemitz, M., & Sjöholm, R. (2008). Complete assignments of the ¹H and ¹³C chemical shifts and J H, H coupling constants in NMR spectra of d-glucopyranose and all d-glucopyranosyl-d-glucopyranosides. *Carbohydrate research*, 343(1), 101-112.
- Schöfer, S. H., Kaftzik, N., Wasserscheid, P., & Kragl, U. (2001). Enzyme catalysis in ionic liquids: lipase catalysed kinetic resolution of 1-phenylethanol with improved enantioselectivity. *Chemical Communications*(5), 425-426.
- Tsavas, P., Polydorou, S., Fafila, I., Voutsas, E. C., Tassios, D., Flores, M. V., Naraghi, K., Halling, P. J., Chamouleau, F., & Ghoul, M. (2002). Solubility of glucose in mixtures containing 2-methyl-2-butanol, dimethyl sulfoxide, acids, esters, and water. *Journal of Chemical & Engineering Data*, 47(4), 807-810.

- Van Rantwijk, F., Woudenberg-van Oosterom, M., & Sheldon, R. (1999). Glycosidase-catalysed synthesis of alkyl glycosides. *Journal of Molecular Catalysis B: Enzymatic*, 6(6), 511-532.
- Walsh, M. K., Bombyk, R. A., Wagh, A., Bingham, A., & Berreau, L. M. (2009). Synthesis of lactose monolaurate as influenced by various lipases and solvents. *Journal of Molecular Catalysis B: Enzymatic*, 60(3), 171-177.
- Ward, O. P., Fang, J., & Li, Z. (1997). Lipase-catalyzed synthesis of a sugar ester containing arachidonic acid. *Enzyme and microbial technology*, 20(1), 52-56.
- Wei, D., Yu, Y., Song, Q., & Su, W. (2003). Enzymatic synthesis of ethyl-glucoside monooleate with lipase in solvent-free medium. *Biocatalysis and Biotransformation*, 21(3), 135-139.
- Zaidan, U. H., Abdul Rahman, M. B., Othman, S. S., Basri, M., Abdulmalek, E., Abdul Rahman, R. N. Z. R., & Salleh, A. B. (2012). Biocatalytic production of lactose ester catalysed by mica-based immobilised lipase. *Food Chemistry*, 131(1), 199-205.