

# TELE-HEALTHCARE AND THE USE OF VIRTUAL COMMUNICATION TECHNOLOGIES IN MEDICAL RESEARCH AND APPLICATION: THE FUTURE OF TELEMEDICINE IS NOW!

WILLIAM E. HILLS<sup>1 A-F</sup>

• ORCID: 0000-0001-7319-9629

KAREN T. HILLS<sup>2 E,F</sup>

<sup>1</sup> Coastal Carolina University, Conway, SC, USA

<sup>2</sup> Beaufort Jasper Hampton Comprehensive Health Services, Ridgeland, SC, USA

A – study design, B – data collection, C – statistical analysis, D – interpretation of data, E – manuscript preparation, F – literature review, G – sourcing of funding

## ABSTRACT

Contemporary world events are demonstrating the need to embrace and further develop tele-health options for assessment and delivery of biopsychosocial healthcare services. This is now possible, given advances in communication technologies allowing virtual connections of medical personnel with constituents, as well as necessary, in light of recent challenges posed by infectious conditions and growing needs for travel restrictions, social distancing, and isolation of large portions of populations. Moreover, the opportunity to virtually connect with persons through ubiquitous computer-based and handheld communication devices allows comprehensive care provision to include underserved areas, where restricted, walk-in access to brick-and-mortar establishments has long been recognized as a limiting factor in healthcare. This review examines evolving approaches of tele-healthcare, with a specific focus on telemedicine as a bridge between traditional, in-person approaches to diagnose and treat medical conditions and new healthcare opportunities developing to meet changing societal needs. The three purposes of the review are: discuss background information, with a brief look at policy and procedure guiding applications of tele-techniques in healthcare practices; identify relevant scientific studies to show the breadth of new evidence-based research for telemedicine practices; and, discuss challenges for the further development of telemedicine as healthcare systems in the United States evolve to meet current and projected healthcare needs.

**KEYWORDS:** telemedicine, tele-healthcare, tele-techniques, virtual communication technologies, integrated primary care-behavioral healthcare, mental health

## BACKGROUND

The need for healthcare to be responsive to environmental concerns has never been greater. As this article is being written, challenges for healthcare providers are growing exponentially, with calls for wide-ranging solutions to meet unprecedented needs posed by the novel coronavirus – COVID-19 [1]. Among these concerns are: more efficient testing procedures to pinpoint virus spread, community containment and contact tracing strategies to isolate hot spots of contagion [1,2], and nuanced implementation of strategies to minimize iatrogenic effects of widespread economic disruption. In addition, the need to better protect first-responders and front-line medical personnel has been pushing the use of virtual communication for remote diagnoses, advice,

and triage systems [3]. In the United States, persons experiencing symptoms are told to avoid, if possible, going to the hospital emergency room and to first call the doctor or clinic before showing up in person; following a consultation and diagnosis, persons with less than severe symptoms are advised to isolate at home. Telemedical services are increasingly being utilized and developed to handle these initial communications with patients as well as a wide variety of needed follow-ups. While many of the adjustments made to healthcare are being forced by the logistics of COVID-19 treatment and care, the further development of tele-procedures is expected to continue to reshape the biopsychosocial delivery of services beyond the current crisis and into the future [4–6].

## TELEMEDICINE AND GROWTH OF TELE-HEALTHCARE

Telemedicine includes the use of information and communication technology devices to deliver professional services across geographical distance and time [7,8]. While the concept of telemedicine has been discussed as potentially useful in practice for decades [9,10], the continuing lack of empirical validation for tele- techniques, overall, has led many physicians to move cautiously toward modification of traditional practice guidelines. In short, not all physicians have been convinced that patients would be well served (not to mention better served) by significant alterations of traditional face-to-face interactions occurring inside brick-and-mortar establishments [11]. The circumstances prompting the use of tele-techniques, however, has now changed with the exigencies of the COVID-19 crisis. Against the backdrop of ever-rising healthcare costs [12], projected shortages of professionals to meet growing healthcare challenges [13], and increasing needs to manage chronic health-related conditions [13], physicians are being encouraged to consider whether more rapid incorporation of telemedicine into medical practices is warranted [1].

Consumer-demand issues are also driving the market toward the adoption of tele- techniques in healthcare [14]. Use of virtual technologies outside the realm of healthcare has become common and familiar; phones are now computers, and comparison shopping has become an everyday activity for many people. Research shows that consumers now expect choices and often look beyond price to range of services offered [15]. The healthcare industry has shown sensitivity to this trend, with hospitals and physicians developing and using websites to advertise and gauge demand. In turn, just as retail consumers have shown a preference for comprehensive business systems – consider Walmart and Amazon, the ultimate one stop shops – consumers in the healthcare marketplace are now similarly beginning to look for comprehensive care provision. As an example, the development of integrated primary care-behavioral healthcare (PCBH) models over the last decade in the United States has broadened many medical service delivery practices to include psychological and social forms of care previously ignored or accessed only through off-site referrals by medical practitioners [16,17]. The more comprehensive PCBH systems have provided choice for persons with mental health issues, who still rely on physicians as a first line of defense [18]. Rather than not knowing where else to go or having no choices, though, data now show that persons in need of mental health interventions can and do search online for inclusive service delivery, where medical, psychological and social care are accessible in a single practice [19,17]. The PCBH systems have been early in the move to adopt virtual care delivery, sometimes using tele-health techniques to facilitate service delivery onsite but, more typically, using tele-techniques such as vid-

eoconferencing when too few professionals are available onsite and services must be accessed and shared across virtual space [16]. The sharing of resources has been particularly relevant for rural areas, where medical and mental health professionals are in great demand, although there are shortages of psychiatrists and psychologists in many urban areas, as well [20]. As mental illness is now recognized as a leading cause of disability [21], and the conjoint presence of mental and physical health problems affects one in four persons in the United States [22], a persuasive argument can be made that use of tele-techniques offers potential to alleviate suffering and aid in provision of treatments for a significant number of people.

These shifts in societal needs and consumer expectations toward a virtual consumer market for medical services have recently been supported by reformulations of policy by professional organizations [23,12], resulting in a broader range of practitioner credentials approved for tele-practice modalities [17]. In March 2020, the United States Federal Government, under the direction of the president, declared that COVID-19 represented a national emergency and invoked the National Emergencies Act (NEA) to temporarily loosen restrictions on use of telemedicine and telehealth interventions, broadly defined [1]. Among those professionals moving quickly to incorporate and/or further expand use of tele-techniques were physicians, psychiatrists, psychologists, nurses, counselors, and social workers, with additional professionals (e.g., occupational therapists) operating under the aegis of integrated practices [10]. This rush to market for virtual service delivery has not, however, been ignored or quietly accepted by all healthcare professionals. Editorials and commentary in medical circles have been issued and note the continuing need to hold the empirical line for science to guide the direction and future development of telemedicine [24,25]. This response is consistent with the NEA, which holds as a first principle that national emergency responses must strike a balance between public health and individual rights and insure that interventions are evidence-based through scientific studies and not representative of political concerns [1].

## TELEMEDICINE PRACTICE AREAS

As tele-service provision is relatively new, overall, literature reviews to gauge its use and effectiveness typically cover studies across various uses of the term tele-. For example, a search for the term “telemedicine” in medical databases results in research-based articles for telemedicine [26], but also articles for tele-health [27,10,28], tele-rehabilitation [29,30], e-consults [31], eHealth [8], mHealth [32], and tele-practice [33]. Similarly, a search for “telemedicine” across science and health literature databases, in general, results in studies that employ such comprehensive terms as technology-driven interventions [34], information and commu-

nication technology devices [35], innovative assistive technologies [36], and digital health interventions [37]. This abbreviated list does not include research for mental health issues, typically covered under such terms as tele-psychology [38] and tele-mental health [39,40], nor does it address studies increasingly identifying roles in virtual delivery of mental health services by practitioners (e.g., social workers) operating as behavioral health providers [16,17]. For purposes of clarity, this overview will use the term telemedicine and primarily focus on studies clearly indicative of use of tele-techniques for medically-directed interventions.

Research literature made available since the declaration of the pandemic by the World Health Organization on March 11, 2020 [41] highlights evidence for when telemedicine service delivery might be useful as stand-alone or adjunctive therapies and, further, defines areas of medical practice for which telemedicine may not be appropriate [23,42]. An overview of these recent, evidence-based studies shows that tele-interventions are useful in: allergy/immunology practices [8]; cancer and radiotherapy treatments [43]; chronic respiratory disease treatment [44]; palliative care for Parkinson's and neurocognitive disorders [45]; and, treatment, monitoring, and/or rehabilitation of persons with cardiovascular conditions [46], type 2 diabetes [47,43]), stroke conditions [48], asthma [42], and chronic obstructive pulmonary disease [49]. Further, recent studies have supported the use of telemedicine techniques in ophthalmology [25], treatments of opioid use disorder [50], chronic pain care [51], anxiety and depression [52], and in post-trauma treatments within a primary care-behavioral health framework of integrated care [17]. In addition, tele-techniques have been shown to be efficacious: for children [53], adolescents [54,55], and adults of all ages [43]; in urban [43] as well as rural areas [56]; for medical [47], psychological [52], and social problems [57]; with acceptance by clinicians [58], nurses [59], and clients and their caregivers [58,44]; when used in primary care [23] and home settings [60]; and, for acute [17], postacute [61], and long term care [56].

In addition to treatment outcome studies, the telemedicine literature also covers a wide variety of practice issues as providers share knowledge for what needs attention. Assistive-technology needs of caregivers are discussed to support aging-in-place and independent living in light of the burgeoning and coming need for long-term care based on population aging [36,13]. Studies are available that call for "reimagining" medical education [5, p. 1127] and how best to prepare nurses for the "uncharted waters" of the COVID-produced "transformed workplace" [59, p. 288]. Issues such as licensing [8], ethics of practice [44], and reimbursement for specialty consultations outside of the "traditional inpatient consult structure" are discussed [31, p. 399] and provide insight for providers on how to best move forward. An examination of rural-urban disparities in care gives insight into the potential of tele-techniques

to address perceived inequities in mental health care practice [62]. Some authors have provided in-depth examinations for what tele-techniques work in their practices – yes: smartphone monitoring in cardiology emergencies [46]; no: "breaking bad news" in an oncology setting [23, p. e879]. In an insightful commentary entitled, "Telemedicine: The unsung corona warrior", issues in the legal, technological, financial, ethical, and scientific domains were discussed as barriers that must be addressed before telehealth can reach its full potential [25]. It seems apparent that this burgeoning literature, albeit perhaps as a result of societal restrictions somewhat limiting access to traditional care, attests to a growing interest in tele-healthcare options. This cannot be determined at the present, however, and will most likely not be determined until the pandemic ends and full empirical assessments of the new treatment modalities are conducted.

### CHALLENGES FOR TELEMEDICINE

There are numerous issues to address before use of tele-techniques can become the "new normal". There is an expressed need for training and education, necessary to manage complex aspects of technology of telehealthcare, although data exist showing that technology issues diminish for healthcare personnel with a higher frequency of usage [58]. Nonetheless, depending on the type of technology involved, virtual service delivery may require the assistance of an IT (information technology) person to set up and manage equipment used [16]. This is an issue with synchronous (i.e., real time) connections in a supervised setting, such as when videoconferencing allows face-to-face connections within a provider system; even with commercially-available programs designed for virtual connections, training for personnel responsible for scheduling and executing sessions [58], as well as a knowledgeable person to upload program updates and troubleshoot disconnections (e.g., during power surges and outages), may be useful. Similar technology concerns surround asynchronous communications, defined as automated and pre-programmed content of computer-based applications, when store-and-forward health provider information and remote-patient monitoring [8] are administered within practices by persons without backgrounds in information technology areas. Both synchronous and asynchronous issues are compounded when service delivery involves direct-to-consumer care and clients located offsite use their own equipment [8]. Although most people today have access to and know how to use smart phones, the types of technology involved in healthcare delivery, with stringent standards for such practice issues as informed consent and confidentiality, are often complex and beyond the capabilities of many consumers. This is particularly true for older adults, who are making gains in the virtual world but still lag behind younger persons in utilization of internet-based technologies [63,64].

A similar issue involves the non-uniform access in the United States to broadband, a problem highlighted through the pandemic by unmet needs of adults working remotely and children attending internet-based classes. An often cited advantage of tele-communications for healthcare is provision of services in rural areas, where access to brick-and-mortar institutions is restricted. Although phone service and internet-based computer connections have improved greatly in recent years and communication devices meeting healthcare standards (i.e., computers, laptops, tablets) are increasingly affordable, access is still limited or completely unavailable in many remote areas [62].

There are also practice issues associated with telehealth that may require further resolution, such as when boundaries overlap and professionals working in interdisciplinary settings adhere to different sets of practice guidelines [16]. While physicians are most often the acknowledged leaders of telemedicine teams, there are other providers working in close association with physicians, sometimes as employees but also through consultations, who may have their own guidelines for practice and ability to seek reimbursement for services rendered. One would think that the practice guidelines for different disciplines should be consistent in instances of team treatments of clients – for example, in primary care-behavioral health integrated practices – but preemptive conversations for clarification of possible differences can be useful. Licensure issues appear to be reaching consensus for telehealth providers in the United States, with professional licensure most typi-

cally required in the state of the client, regardless of practice and/or provider location [8]. Similarly, reimbursement for tele-services, at least during the temporary conditions granted by the NEA, are consistent with previous rates.

## CONCLUSION

It is clear that the use of tele-techniques in practice settings predate the pandemic [10], but it is equally true that there has been an acceleration of telehealth service delivery driven by need and the rapid production and expansion of policies necessary to guide and protect providers. There is some concern, perhaps well-founded, that technological advances may outpace ethical guideline development [65]. It is true that the accelerated pace of change for technologies has become the expectation, particularly among young consumers, and it may be difficult to return from this forced immersion in the tele-health world to care approaches that are perceived as less convenient. In any evaluation, the pandemic has lasted a sufficient length of time for researches to begin establishing scientific validation for what aspects of telemedicine do and do not work in practice settings. It is doubtful that the insights gained through this process will be lost as COVID-19 is finally brought to heel. What is more certain is that the key to the continuation of telehealth lessons learned during the pandemic is acceptance by physicians [58], and acceptance by physicians depends upon establishing the empirical bases for tele-healthcare techniques [1].

## REFERENCES

1. Gostin LO, Hodge JG, Wiley LF. Presidential powers and response to COVID-19. *JAMA* 2020 Apr 28; 323(16): 1547–1548.
2. Parodi SM, Liu VX. From containment to mitigation of COVID-19 in the US. *JAMA* 2020 Apr 21; 323(15): 1441–1442.
3. Adams JG, Walls RM. Supporting the health care workforce during the COVID-19 global epidemic. *JAMA* 2020 Apr 21; 323(15): 1439–1440.
4. Ducharme J. The coronavirus outbreak could finally make telemedicine mainstream in the U.S. *Time* [online] 3.03.2020 [cit. 4.03.2020]. Available from URL: <https://time.com/5793535/coronavirus-telemedicine-telehealth/>.
5. Emanuel EJ. The inevitable reimagining of medical education. *JAMA* 2020; 323(12): 1127–1128.
6. Green M, Wayne DB, Neilson EG. Medical education 2020 – charting a path forward. *JAMA* 2019; 322(10): 934–935.
7. Timpel P, Oswald S, Schwarz PEH, Harst L. Mapping the evidence on the effectiveness of telemedicine interventions in diabetes, dyslipidemia, and hypertension: an umbrella review of systematic reviews and meta-analyses. *J Med Internet Res* 2020; 22(3): e16791.
8. Waller MR, Taylor LM, Portnoy JM. Benefits and risks of incorporating virtual visits into an allergy/immunology practice. *Allergy Asthma Proc* 2020; 41(2): 76–81.
9. Nickelson DW. Behavioral telehealth: emerging practice, research, and policy opportunities. *Behav Sci Law* 1996; 14: 443–457.
10. McClellan MJ, Florell D, Palmer J, Kidder C. Clinician telehealth attitudes in a rural community mental health center setting. *J Rural Ment Health* 2020; 44(1): 62–73.
11. Barnett ML, Ray KN, Souza J, Mehrotra A. Trends in telemedicine use in a large commercially insured population, 2005–2017. *JAMA* 2018; 320(20): 2147–2149.
12. Shekelle PG, Goldman DP. Development of a group judgment process for forecasts of health care innovations. *JAMA Netw Open* 2018; 1(7): e185108.
13. Hills WE. Behavioral health and new models of service delivery for an aging world: public/private partnerships to develop best practices of care for older adults. *Med Sci Pulse* 2019; 13(1): 29–33.
14. Øyen KR, Sunde OS, Solheim M, Moricz M, Ytrehus S. Understanding attitudes toward information and communication technology in home-care: information and communication technology as a market good within Norwegian welfare services. *Informatics for Health and Social Care* 2018; 43(3): 300–309.
15. Bamberger KA, Egelman S, Han C, Bar On AE, Reyes I. Can you pay for privacy? Consumer expectations and the behavior of free and paid apps. *Berkeley Technology Law Journal* 2020; 35: 327–365.
16. Hills WE, Hills KT. Virtual treatments in an integrated primary care-behavioral health practice: an overview of synchronous

- telehealth services to address rural-urban disparities in mental health care. *Med Sci Pulse* 2019; 13(3): 54–59.
17. Kanzler KE, Ogbeide S. Addressing trauma and stress in the COVID-19 pandemic: challenges and the promise of integrated primary care. *Psychol Trauma* 2020; 12(S1): S177–S179.
  18. Olfson M. Building the mental health workforce capacity needed to treat adults with serious mental illnesses. *Health Affairs* 2016; 35(6): 983–990.
  19. Bierman AS. Preventing and managing multimorbidity by integrating behavioral health and primary care. *Health Psychol* 2019; 38(9): 851–854.
  20. Hoeft TJ, Fortney JC, Patel V, Unutzer J. Task-sharing approaches to improve mental healthcare in rural and other low-resource settings: a systematic review. *J Rural Ment Health* 2018; 34: 48–62.
  21. Hughes MC, Gorman JM, Ren Y, Khalid S, Clayton C. Increasing access to rural mental health care using hybrid care that includes telepsychiatry. *J Rural Ment Health* 2019; 43(1): 30–37.
  22. Suls J, Green PA, Boyd CM. Multimorbidity: implications and directions for health psychology and behavioral medicine. *Health Psychology* 2019; 38(9): 772–782.
  23. Wolf I, Waissengrin B, Pelles S. Breaking bad news via telemedicine: a new challenge at times of an epidemic. *The Oncologist* 2020; 25: e879–e880.
  24. Dinesen B, Nonnecke B, Lindeman D, Toft E, Kidholm K, Jethwani K, et al. Personalized telehealth in the future: a global research agenda. *J Med Internet Res* 2016; 18(3): e53.
  25. Biswas P, Batra S. Telemedicine: the unsung corona warrior. *Indian J Ophthalmol* 2020; 68(6): 112–113.
  26. Costanzo MC, Arcidiacono C, Rodolico A, Panebianco M, Aguglia E, Signorelli MS. Diagnostic and interventional implications of telemedicine in Alzheimer's disease and mild cognitive impairment: a literature review. *Int J Geriatr Psychiatry* 2020 Jan; 35(1): 12–28.
  27. Frey SM, Milne Wenderlich A, Halterman JS. New opportunities with school-based telehealth: convenient connections to care. *JAMA Pediatr* 2019; 173(11): 1017–1018.
  28. Wu YP, Boucher K, Hu N, Hay J, Kohlmann W, Aspinwall LG, et al. A pilot study of a telehealth family-focused melanoma preventive intervention for children with a family history of melanoma. *Psycho-Oncology* 2020; 29: 148–155.
  29. Piotrowicz E, Piotrowski W, Piotrowicz R. Positive effects of the reversion of depression on the sympathovagal balance after telerehabilitation in heart failure patients. *Ann Noninvasive Electrocardiol* 2016; 21(4): 358–368.
  30. Lewis JA, Boian RF, Burdea G, Deutsch JE. Remote console for virtual telerehabilitation. *Stud Health Technol Inform* 2005; 111: 294–300.
  31. Wachter RM, Judson TJ, Mourad M. Reimagining specialty consultation in the digital age: the potential role of targeted automatic electronic consultations. *JAMA* 2019; 322(5): 399–400.
  32. Bateman DR, Srinivas B, Emmett TW, Schleyer TK, Holden RJ, Hendrie HC, et al. Categorizing health outcomes and efficacy of mHealth apps for persons with cognitive impairment: a systematic review. *J Med Internet Res* 2017; 19(8): e301.
  33. Stewart SE, Neal C. Consultants' use of telepractice: Practitioner survey, issues, and resources. *Consult Psychol J Pract Res* 2015; 67(2): 85–99.
  34. Godwin KM, Mills WL, Anderson JA, Kunik ME. Technology-driven interventions for caregivers of persons with dementia: a systematic review. *Am J Alzheimers Dis Other Demen* 2013; 28(3): 216–222.
  35. Kim D, Choi J, Han K. Risk management-based security evaluation model for telemedicine systems. *BMC Med Inform Decis Mak* 2020; 20: 106.
  36. Thordardottir B, Fänge AM, Lethin C, Gatta DR, Chiatti C. Acceptance and use of innovative assistive technologies among people with cognitive impairment and their caregivers: a systematic review. *Biomed Res Int* 2019 Mar 6; 2019: 9196729.
  37. McVay MA, Bennett GG, Steinberg D, Voils CI. Dose–response research in digital health interventions: concepts, considerations, and challenges. *Health Psychol* 2019; 38(12): 1168–1174.
  38. Varker T, Brand RM, Ward J, Terhaag S, Phelps A. Efficacy of synchronous telepsychology interventions for people with anxiety, depression, posttraumatic stress disorder, and adjustment disorder: a rapid evidence assessment. *Psychol Serv* 2019 Nov; 16(4): 621–635.
  39. Gonzalez MLS, McCord CE, Dopp AR, Tarlow KR, Dickey NJ, McMaughan DK, et al. Telemental health training and delivery in primary care: a case report of interdisciplinary treatment. *J Clin Psychol* 2019 Feb; 75(2): 260–270.
  40. Douglas S, Jensen-Doss A, Ordorica C, Comer JS. Strategies to enhance communication with telemental health measurement-based care (tmBC). *Pract Innov* 2020; 5(2): 143–149.
  41. Jordan V. Coronavirus (COVID-19): remote care through telehealth. *J Prim Health Care* 2020; 12(2): 184–185.
  42. Kew KM, Cates CJ. Home telemonitoring and remote feedback between clinic visits for asthma. *Cochrane Database Syst Rev* 2016 Aug 3; 2016(8): CD011714.
  43. Nguyen NP, Vinh-Hung V, Baumert BG, Zamagni A, Arenas M, Motta M, et al. Older cancer patients during the COVID-19 epidemic: practice proposal of the international geriatric radiotherapy group. *Cancers* 2020; 12: 1287.
  44. McGee JS, Meraz R, Myers DR, Davie MR. Telehealth services for persons with chronic lower respiratory disease and their informal caregivers in the context of the COVID-19 pandemic. *Pract Innov* 2020; 5(2): 165–177.
  45. Kluger BM, Miyasaki J, Katz M, Galifianakis N, Hall K, Pantilat S, et al. Comparison of integrated outpatient palliative care with standard care in patients with Parkinson disease and related disorders: a randomized clinical trial. *JAMA Neurol* 2020 May 1; 77(5): 551–560.
  46. Koulaouzidis G, Charisopoulou D, Wojakowski W, Koulaouzidis A, Marlicz W, Jadczyk T. Telemedicine in cardiology in the time of coronavirus disease 2019: a friend that everybody needs. *Pol Arch Intern Med* 2020 Jun 25; 130(6): 559–561.
  47. Klösch M, Klösch C, Kundt FS, van der Zee-Neuen A, Dieplinger AN. eHealth systems for the optimised care of patients with type 2 diabetes. *Br J Nurs* 2020; 29(5): 274–278.
  48. Garcia-Esperon C, Dinkelspiel FS, Miteff F, Gangadharan S, Wellings T, O'Brien B, et al. Implementation of multimodal computed tomography in a telestroke network: five-year experience. *CNS Neurosci Ther* 2020; 26: 367–373.
  49. McCabe C, McCann M, Brady AM. Computer and mobile technology interventions for self-management in chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 2017 May 23; 5(5): CD011425.

50. Rubin R. Using telemedicine to treat opioid use disorder in rural areas. *JAMA* 2019; 322(11): 1029–1031.
51. Glynn LH, Chen JA, Dawson TC, Gelman H, Zeliadt SB. Bringing chronic-pain care to rural veterans: a telehealth pilot program description. *Psychol Serv* 2020 Jan 16; doi: 10.1037/ser0000408 [ahead of print].
52. Hull TD, Malgaroli M, Connolly PS, Feuerstein S, Simon NM. Two-way messaging therapy for depression and anxiety: longitudinal response trajectories. *BMC Psychiatry* (2020); 20: 297.
53. Gurwitsch RH, Salem H, Nelson MM, Comer JS. Leveraging parent–child interaction therapy and telehealth capacities to address the unique needs of young children during the COVID-19 public health crisis. *Psychol Trauma* 2020 Aug; 12(S1): S82–S84.
54. Michael KD. Youth mental health in North Carolina: creative innovations in challenging times. *North Carolina Med J* 2020; 81(2): 101–105.
55. North S. Addressing students' mental health needs via telehealth. *North Carolina Med J* 2020; 81(2): 112–113.
56. Riboni FV, Comazzi B, Bercovitz K, Castelnuovo G, Molinari E, Pagnini F. Technologically-enhanced psychological interventions for older adults: a scoping review. *BMC Geriatrics* 2020; 20: 191.
57. Carballea D, Rivera RM. Coronavirus and interpersonal violence: a need for digital mental health resources. *Psychol Trauma* 2020 Aug; 12(S1): S208–S209.
58. Vidal-Alaball J, Seguí FL, Domingo JLG, Mateo GF, Valmaña GS, Ruiz-Comellas A, et al. Primary care professionals' acceptance of medical record-based, store and forward provider-to-provider telemedicine in Catalonia: results of a web-based survey. *Int J Environ Res Public Health* 2020; 17: 4092.
59. Boland MG. Navigating uncharted waters: preparing COVID-19 capable nurses to work in a transformed workplace. *Hawaii J Health Soc Welf* 2020 Sep 1; 79(9): 288–289.
60. Almathami HKY, Win KT, Vlahu-Gjorgievska E. Barriers and facilitators that influence telemedicine-based, real-time, online consultation at patients' homes: systematic literature review. *J Med Internet Res* 2020 Feb 20; 22(2): e16407.
61. Grabowski DC, Joynt Maddox KE. Postacute care preparedness for COVID-19: thinking ahead. *JAMA* 2020 May 26; 323(20): 2007–2008.
62. Summers-Gabr NM. Rural–urban mental health disparities in the United States during COVID-19. *Psychol Trauma* 2020 Aug; 12(S1): S222–S224.
63. Lepkowsky CM, Arndt S. The internet: barrier to health care for older adults? *Pract Innov* 2019; 4(2): 124–132.
64. Walker J, Crotty BH, O'Brien J, Dierks MM, Lipsitz L, Safran C. Addressing the challenges of aging: how elders and their care partners seek information. *Gerontologist* 2017; 57(5): 955–962.
65. Edwards-Stewart A, Alexander C, Armstrong CM, Hoyt T, O'Donohue W. Mobile applications for client use: ethical and legal considerations. *Psychol Serv* 2019; 16(2): 281–285.

Word count: 2405

• Tables: –

• Figures: –

• References: 65

**Sources of funding:**

The research was funded by the authors.

**Conflicts of interests:**

The authors report that there were no conflicts of interest.

**Cite this article as:**

Hills WE, Hills KT.

Tele-healthcare and the use of virtual communication technologies in medical research and application: the future of telemedicine is now! *Med Sci Pulse* 2020; 14, 3: 78–83.**Correspondence address:**William E. Hills, PhD, MSW  
Professor of Psychology  
Coastal Carolina University  
Conway, South Carolina, USA  
Phone: (+1) 843-349-2276  
E-mail: hillsw@coastal.edu

Received: 25.09.2020

Reviewed: 29.09.2020

Accepted: 30.09.2020

Copyright of Medical Science Pulse is the property of Public Higher Medical Professional School in Opole and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.