

DOCUMENT RESUME

ED 401 508

CG 027 125

AUTHOR            Beaty, Lee A.  
 TITLE            Is Relaxation Training Effective in the Treatment of  
                   Clinical Depression?  
 PUB DATE         [96]  
 NOTE             10p.  
 PUB TYPE         Information Analyses (070)

EDRS PRICE       MF01/PC01 Plus Postage.  
 DESCRIPTORS      Cognitive Restructuring; \*Depression (Psychology);  
                   \*Emotional Disturbances; Medical Care Evaluation;  
                   Psychopathology; \*Psychophysiology; Psychotherapy;  
                   \*Relaxation Training; \*Stress Management

ABSTRACT

The process of relaxation is a complex triarchic phenomenon that incorporates behavioral, cognitive, and physiological components. Existing literature is surveyed in order to determine the efficacy of treating various forms of depression with cognitive-behavioral relaxation strategies. Relaxation training has been shown to be effective in treating a variety of psychosomatic problems, including muscle contraction headaches, asthma, hypertension, and panic attacks. Successful treatment of these problems, which have a physical and psychological overlap, indicates that relaxation training may be a beneficial intervention strategy. It also may produce acceptable success rates in treating depressive disorders when used in combination with other therapies. However, empirical findings have been mixed in regard to relaxation training's effectiveness in treating various types of depression, and more research is clearly needed before therapists can be confident that it is really effective in treating clinical depression. Contains 15 references. (TS)

\*\*\*\*\*  
 \* Reproductions supplied by EDRS are the best that can be made \*  
 \* from the original document. \*  
 \*\*\*\*\*

U.S. DEPARTMENT OF EDUCATION  
Office of Educational Research and Improvement  
EDUCATIONAL RESOURCES INFORMATION  
CENTER (ERIC)

- This document has been reproduced as received from the person or organization originating it.
- Minor changes have been made to improve reproduction quality.
- Points of view or opinions stated in this document do not necessarily represent official OERI position or policy.

"PERMISSION TO REPRODUCE THIS MATERIAL HAS BEEN GRANTED BY

L. Beaty

TO THE EDUCATIONAL RESOURCES INFORMATION CENTER (ERIC)."

IS RELAXATION TRAINING EFFECTIVE  
IN THE TREATMENT OF CLINICAL DEPRESSION?

by

Lee A. Beaty, Ph.D.

Counselor Education Program, Northeastern Illinois University

Running Head: Relaxation Training

Abstract: During the past two decades, relaxation training has been advocated in the treatment of various psychological disorders. This paper explores the extant literature on the efficacy of relaxation in the treatment of depression.

BEST COPY AVAILABLE

## Relaxation Training/1

The process of relaxation is a complex triarchic phenomenon that incorporates behavioral, cognitive, and physiological components. Until the 1980's, relaxation training generated relatively little research, although anecdotal evidence suggested the potential benefits of incorporating relaxation training into treatment plans for various types of psychological disorders. Despite the fact that therapeutic gains have been associated with the reduction of physiological stress, many treatment programs have concentrated on behavioral/cognitive interventions, thereby ignoring an important feature of successful relaxation: reduction of stress itself. However, stress reduction may be but one of the therapeutic variables at work in the relaxation response. More recent approaches (e.g., Smith, 1990) have proposed a cognitive-behavioral relaxation training model. This model challenges the traditional arousal reduction model by incorporating stress reduction into a more holistic framework that views relaxation as a complex cognitive-behavioral process. Thus, this newer view of relaxation may make it more compatible with cognitive approaches to the treatment of depression. The purpose of this paper is to survey existing literature in order to determine the efficacy of treating various forms of depression with cognitive-behavioral relaxation strategies.

Relaxation training has been shown to be effective in treating a variety of psychosomatic problems, including muscle contraction headaches, asthma, hypertension, and panic attacks. Successful

## Relaxation Training/2

treatment of these problems which have a physical and psychological overlap indicates that relaxation training may be a beneficial intervention strategy.

Relaxation training has been shown to relieve high levels of anxiety and to reduce associated physiological and muscular tension (Goldfried & Trier, 1974). Most traditional treatments for depression have been concerned with cognitive/behavioral interventions that sought to influence targeted cognitions or behaviors. However, studies of treatment outcomes indicate that interventions were more likely to affect non-targeted outcomes (Reynolds & Coats, 1986). The clear implication of this finding is that the type of treatment does not ensure that the specifically targeted cognitions or behaviors will be changed. Rather, one could hypothesize that a nonspecific change occurs in the general level of depression.

The outcomes of relaxation therapy indicate that numerous nonspecific factors influence the efficacy of treatments for depression (McLean & Hakstian, 1979). Furthermore, Smith (1990) states that "A relaxer may be quite unaware of the implications of his actions, that is, the chain of potential changes he is capable of setting into motion. Similarly, each supporting structure may imply additional structures outside the awareness of the relaxer (p. 44)." Smith believes that a complete assessment of relaxation must take into account the different levels of depth revealed through the cognitive processes of focusing, passivity, and receptivity.

### Relaxation Training/3

Interestingly, there is some empirical evidence for the effectiveness of relaxation training in treating various types of depression. Relaxation training was used to reduce postpartum depression in first-time mothers (Halonen & Passman, 1985). Relaxation training paired with biofeedback significantly reduced levels of depression in chronic pain patients (Peniston, Hughes, & Kulkosky, 1986). Relaxation training was found to be as effective as cognitive-behavioral therapy in a group counseling setting designed to eliminate moderate levels of depression for adolescents (Reynolds & Coats, 1986). Meditation, a specific form of relaxation training, was used to significantly reduce depression levels in a sample of geriatric patients (Deberry, Davis, & Reinhard, 1989). Relaxation training was found to neutralize stress factors and thereby reduce levels of depression when compared to a control condition (Sroota & Dhir, 1990).

Short-term relaxation training, cognitive-behavioral treatment, and self-modeling therapy were equally effective in significantly reducing levels of depression and increasing levels of self-esteem (Kahn, Kehle, Jenson, & Clark, 1990). In a study measuring clients' perceptions of multiple treatment regimens, relaxation training was rated as being among the most effective treatments and was determined to be more effective than psychotherapy or drug therapy (Rokke, Carter, Rehm, and Veltum, 1990).

However, other researchers have not found relaxation training to be so effective. For example, behavior therapy was found to be

#### Relaxation Training/4

significantly more effective in treating moderate levels of depression than either relaxation training or drug therapy (Rehm, Kaslow, & Rabin, 1987). A longitudinal assessment of four types of therapy (psychotherapy, behavior therapy, drug therapy, and relaxation training) determined that only behavior therapy was effective in producing long-term reductions in depression (McLean & Hakstian, 1990).

In order for therapies to be effective, whether specific or nonspecific, it has been argued that improving the client's sense of self-efficacy is of vital importance (Zeiss, Lewinsohn, & Munoz, 1979). According to Sowers (1990), a consistent therapeutic framework offers clients a reference point from which to understand the changes occurring in their depressive symptoms, which may in turn enhance their feelings of self-efficacy. The state of physiological and psychological relaxation that is the goal of relaxation training would presumably be consistent with an increased level of personal effectiveness, at least in terms of the targeted symptoms.

Thus, the answer to the question which the title of this article poses is difficult to ascertain. It may be that relaxation training in combination with other therapies (particularly cognitive/behavioral ones) will produce acceptable success rates in treating depressive disorders, which are, according to Beck (1970), one of the most prevalent and insidious forms of psychological distress. Sroota and Dhir's (1990) research suggests that relaxation may have a mediating effect in reducing stress

## Relaxation Training/5

which in turn produces reductions in depression. This possibility lends support to the notion that there may be some overlap in the symptomatology of depressive disorders and anxiety disorders. Another interesting point to ponder is whether relaxation training is differentially effective in treating milder or more severe forms of depression. Most of the studies reviewed here focus on clients with depressions in the milder range. It may be that relaxation techniques are relatively ineffective for those clients with more severe forms of depression. Clearly, more empirical research is needed before therapists can be confident that relaxation training is really effective in treating clinical depression.

## Relaxation Training/6

### References

Beck, A.T. (1970). Depression: Causes and treatment. Philadelphia, PA: University of Pennsylvania Press.

Deberry, S., Davis, S., & Reinhard, K.E. (1989). A comparison of meditation-relaxation and cognitive-behavioral techniques of reducing anxiety and depression in a geriatric population. Journal of Geriatric Psychiatry, 22(2), 231-247

Goldfried, M.R. & Trier, C. (1974). Effectiveness of relaxation as an active coping skill. Journal of Abnormal Psychology, 83(4), 348-355.

Halonen, J.S. & Passman, R.H. (1985). Relaxation training and expectation in treatment of postpartum distress. Journal of Consulting & Clinical Psychology, 53(6), 839-845.

Kahn, J.S., Kehle, T.J., Jenson, W.R., & Clark, E. (1990). Comparison of cognitive-behavioral, relaxation, and self-modeling interventions for depression among middle-school students. School Psychology Review, 19(2), 196-211.

McLean, P.D. & Hakstian, A.R. (1979). Clinical depression: Comparative efficacy of outpatient treatments. Journal of Consulting & Clinical Psychology, 47(5), 818-836.



## Relaxation Training/7

- McLean, P.D. & Hakstian, A.R. (1990). Relative endurance of unipolar depression treatment effects: Longitudinal follow-up. Journal of Consulting & Clinical Psychology, 58(4), 482-488.
- Peniston, E.G., Hughes, R.s., & Kulkosky, P.J. (1986). EMG biofeedback-assisted relaxation training in the treatment of reactive depression in chronic pain patients. The Psychological Record, 36, 471-482.
- Rehm, L.P., Kaslow, N.J. & Rabin, A.S. (1987). Cognitive and behavioral targets in a self-control therapy program for depression. Journal of Consulting & Clinical Psychology, 55, 60-67.
- Reynolds, W.M. & Coats, K.I. (1986). A comparison of cognitive-behavioral therapy and relaxation training for the treatment of depression in adolescents. Journal of Consulting & Clinical Psychology, 54(5), 653-660.
- Rokke, P.D., Carter, A.S., & Rehm, L.P. (1990). Comparative credibility of current treatments for depression. psychotherapy, 27(2), 235-242.
- Smith, J.C. (1990). Cognitive-behavioral relaxation training: A new system for treatment and assessment. New York, NY: Springer.

Relaxation Training/8

sowers, W.A. (1990). Treatment of depressed in-patients: Cognitive therapy plus medication, relaxation plus medication, and medication alone. Swedish Journal of Psychiatry, 156, 73-78.

Sroota, A. & Dhir, R. (1990). Efficacy of two relaxation techniques in depression. Journal of Personality & Clinical Studies, 6(1), 83-90.

Zeiss, A.M., Lewinsohn, P.M., & Munoz, R.F. (1979). Nonspecific improvement effects in depression using interpersonal skills training, pleasant activity schedules, or cognitive training. Journal of Consulting & Clinical Psychology, 47(3), 427-439.



U.S. Department of Education  
Office of Educational Research and Improvement (OERI)  
Educational Resources Information Center (ERIC)



# REPRODUCTION RELEASE

(Specific Document)

## I. DOCUMENT IDENTIFICATION:

Title: "Is Relaxation Therapy Effective in the Treatment of Clinical Depression?"	
Author(s): Lee A. Beaty	
Corporate Source:	Publication Date: 1996

## II. REPRODUCTION RELEASE:

In order to disseminate as widely as possible timely and significant materials of interest to the educational community, documents announced in the monthly abstract journal of the ERIC system, *Resources in Education* (RIE), are usually made available to users in microfiche, reproduced paper copy, and electronic/optical media, and sold through the ERIC Document Reproduction Service (EDRS) or other ERIC vendors. Credit is given to the source of each document, and, if reproduction release is granted, one of the following notices is affixed to the document.

If permission is granted to reproduce and disseminate the identified document, please CHECK ONE of the following two options and sign at the bottom of the page.



Check here

### For Level 1 Release:

Permitting reproduction in microfiche (4" x 6" film) or other ERIC archival media (e.g., electronic or optical) and paper copy.

The sample sticker shown below will be affixed to all Level 1 documents

PERMISSION TO REPRODUCE AND DISSEMINATE THIS MATERIAL HAS BEEN GRANTED BY

\_\_\_\_\_  
Sample  
\_\_\_\_\_

TO THE EDUCATIONAL RESOURCES INFORMATION CENTER (ERIC)

Level 1

The sample sticker shown below will be affixed to all Level 2 documents

PERMISSION TO REPRODUCE AND DISSEMINATE THIS MATERIAL IN OTHER THAN PAPER COPY HAS BEEN GRANTED BY

\_\_\_\_\_  
Sample  
\_\_\_\_\_

TO THE EDUCATIONAL RESOURCES INFORMATION CENTER (ERIC)

Level 2



Check here

### For Level 2 Release:

Permitting reproduction in microfiche (4" x 6" film) or other ERIC archival media (e.g., electronic or optical), but *not* in paper copy.

Documents will be processed as indicated provided reproduction quality permits. If permission to reproduce is granted, but neither box is checked, documents will be processed at Level 1.

"I hereby grant to the Educational Resources Information Center (ERIC) nonexclusive permission to reproduce and disseminate this document as indicated above. Reproduction from the ERIC microfiche or electronic/optical media by persons other than ERIC employees and its system contractors requires permission from the copyright holder. Exception is made for non-profit reproduction by libraries and other service agencies to satisfy information needs of educators in response to discrete inquiries."

Sign here → please

Signature: Lee A Beaty	Printed Name/Position/Title: LEE A BEATY ASST PROF
Organization/Address: Northeastern Ill. University 5500 N. St. Louis Chicago IL 60625	Telephone: 312-794-2880 FAX: E-Mail Address: Date: 7/9/96



### III. DOCUMENT AVAILABILITY INFORMATION (FROM NON-ERIC SOURCE):

If permission to reproduce is not granted to ERIC, or, if you wish ERIC to cite the availability of the document from another source, please provide the following information regarding the availability of the document. (ERIC will not announce a document unless it is publicly available, and a dependable source can be specified. Contributors should also be aware that ERIC selection criteria are significantly more stringent for documents that cannot be made available through EDRS.)

Publisher/Distributor:
Address:
Price:

### IV. REFERRAL OF ERIC TO COPYRIGHT/REPRODUCTION RIGHTS HOLDER:

If the right to grant reproduction release is held by someone other than the addressee, please provide the appropriate name and address:

Name:
Address:

### V. WHERE TO SEND THIS FORM:

Send this form to the following ERIC Clearinghouse:	<b>ERIC/CASS School of Education Park 101, UNCG Greensboro NC 27412</b>
---	---

However, if solicited by the ERIC Facility, or if making an unsolicited contribution to ERIC, return this form (and the document being contributed) to:

**ERIC Processing and Reference Facility**  
1100 West Street, 2d Floor  
Laurel, Maryland 20707-3598

Telephone: 301-497-4080

Toll Free: 800-799-3742

FAX: 301-953-0263

e-mail: [ericfac@inet.ed.gov](mailto:ericfac@inet.ed.gov)

WWW: <http://ericfac.piccard.csc.com>