

2015

The Effects Of Curcumin On The Expression Of P53 And Bcl-2 Oncoproteins On The Human Breast Carcinoma Cell Line

Angelin S. Shajan

Follow this and additional works at: https://csuepress.columbusstate.edu/theses_dissertations



Part of the [Biology Commons](#)

Recommended Citation

Shajan, Angelin S., "The Effects Of Curcumin On The Expression Of P53 And Bcl-2 Oncoproteins On The Human Breast Carcinoma Cell Line" (2015). *Theses and Dissertations*. 180.
https://csuepress.columbusstate.edu/theses_dissertations/180

This Thesis is brought to you for free and open access by the Student Publications at CSU ePress. It has been accepted for inclusion in Theses and Dissertations by an authorized administrator of CSU ePress.

**THE EFFECTS OF CURCUMIN ON THE EXPRESSION OF P53 AND BCL-2
ONCOPROTEINS ON THE HUMAN BREAST CARCINOMA CELL LINE
MDA-MB 468**

Angelin S. Shajan

COLUMBUS STATE UNIVERSITY
THE EFFECTS OF CURCUMIN ON THE EXPRESSION OF P53 AND BCL-2
ONCOPROTEINS ON THE HUMAN BREAST CARCINOMA CELL LINE
MDA-MB 468

A THESIS SUBMITTED TO
THE HONORS COLLEGE AT COLUMBUS STATE UNIVERSITY IN
PARTIAL FULFILLMENT OF THE REQUIREMENTS FOR THE DEGREE
WITH HONORS OF

BACHELOR OF SCIENCE
DEPARTMENT OF BIOLOGY

BY
ANGELIN S. SHAJAN
COLUMBUS, GEORGIA
2015

The copyright in this work is the property of the author. This document is the
property of the author and is not to be distributed or reproduced without the
author's permission.

Copyright © 2015 Angelin S. Shajan
All Rights Reserved




A Thesis Submitted in Partial Fulfillment of
Requirements for the M.A. Degree in
Education in the degree of
Master of Arts
in
Education
College of Letters & Science
California State University

Author: Angelin S. Shajan Date: 12/15/15
Chair: Dr. [Signature]
Committee: [Signature] Date: 12/15/15
Date of Thesis: 12/15/15
Dr. [Signature]

The Effects of Curcumin on the Expression of p53 and Bcl-2 Oncoproteins on the Human Breast Carcinoma Cell Line MDA-MB 468

by
Angelin S. Shajan

A Thesis Submitted in Partial Fulfillment of
Requirements of the CSU Honors College
for Honors in the degree of
Bachelor of Science
in
Biology
College of Letters & Sciences
Columbus State University

Thesis Advisor  Date 4/29/15
Dr. Monica Frazier
Committee Member  Date 4/29/15
Dr. Kathleen Hughes
Dean of Honors College  Date 5/4/15
Dr. Cindy Ticknor

INDEX WORDS: Apoptosis, bcl-2 protein, curcumin, p53 protein.

ABSTRACT

Advisors Dr. Monica Frazier and Dr. Kathleen Hughes

We studied the effects of curcumin on the proliferation and viability of MDA-MB-468 breast cancer cells and its effects on the expression of Bcl-2 and p53 proteins in these cells. Based on previous studies, we predicted a dose-dependent increase in p53 (tumor suppressor) expression, and a decrease in Bcl-2 (anti-apoptotic oncoprotein) expression. Briefly, MDA-MB-468 cells were seeded into 12-well plates at 2.0×10^4 cells/well and allowed to grow for 72 hrs. Cells were treated with curcumin at 0, 25, 50 and 100 μ M in triplicate at time zero and cultured for another 24 hours. All data were obtained using a Muse Cell Analyzer. Proliferation and viability data were determined using an Annexin V and Dead Cell Kit, p53 expression was determined using a Multi-Colored DNA Damage Kit, and Bcl-2 expression was determined using a Bcl-2 Activation Dual Detection Kit. Results show a significant dose-dependent increase in both cell death and proliferation. However, there was no significant difference in expressions of both p53 and Bcl-2 oncoproteins across the four treatments. Future studies will investigate additional mechanisms by which curcumin induces these apoptotic effects.

INDEX WORDS: Apoptosis, bcl-2 protein, curcumin, p53 protein.

TO MY BIGGEST SUPPORTERS

Dear mom and dad: Words cannot begin to describe my appreciation and gratitude for all you have done for me. Your love and support has helped me not only get through college, but also through life. While I honestly do not think I will ever comprehend the true depth of your sacrifice, I can promise that I will continuously try. I will always strive to make you proud because you truly deserve the best. I love you both so much. Love, Angelin.

ACKNOWLEDGMENTS

I would like to thank:

Dr. Monica Frazier, for being a brilliant, extraordinary, and inspiring professor, research mentor, and advisor. You are an amazing and wonderful person and was always available to help me when needed. I feel very happy and blessed for the opportunities to take classes with you and work with you for my senior research. Thank you for all the learning experiences and invaluable guidance over the past four years. I appreciate you very much for all you have done to support me. You rock!

My advisor, Dr. Kathleen Hughes, for always there for me when I needed advice. I owe a lot of my success to you. I am grateful for your guidance, constructive criticism, generosity, and emotional support. I will forever appreciate that fact that you always find ways to help students grasp the material and succeed. You are very much appreciated!

Dean Cindy Ticknor, for all the guidance and knowledge that has deeply enriched my academic experience as an Honors College student. Thank you so much for everything you have done to help me with my transition to college and everything you continue to do for honor students like myself. You are a valuable asset to our institution!

Finally, to the Department of Biology and Chemistry for funding my research through the Student Research And Creative Endeavors (SRACE) Grant, the Flora Clark Research Grant, the Beta Beta Beta Research Grant. To all of my biology and chemistry professors who have taught me everything I have needed to be a successful graduate of the College of Letters and Sciences, and who have been positive supporters of my goals and pursuits.

TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS.....	v
LIST OF FIGURES.....	vii
Chapter I. Introduction.....	1
1.1.1. Breast Cancer.....	2
1.1.2. Curcumin.....	2, 3
1.1.3. p53 protein and cancer.....	3, 4
1.1.4. Bcl-2 protein and cancer.....	4, 5
1.1.5. Aim of our study.....	6
Chapter II. Materials and Methods.....	7
2.1.1. Cell Treatment.....	8
2.1.2. Annexin V and Dead Cell Kit.....	9
2.1.3 Multi Color DNA Damage Kit, Bcl-2 Activation Dual Detection Kit.....	9, 10
2.1.4 Statistical Analyses.....	10
Chapter III. Results and Discussion.....	11
3.1.1. Future Directions.....	18
3.1.2 References.....	19

LIST OF FIGURES

Figure 1. Mechanism of p53 activation.....	4
Figure 2. Mechanism of Bcl-2 activation.....	5
Figure 3. Proposed pathways for p53 and Bcl-2 activation.....	6
Figure 4. Cell treatment in 12-well plate.....	9
Figure 5. Multi-Color DNA Damage assay sample screenshot.....	12
Figure 6. Bcl-2 Activation Dual Detection assay sample screenshot.....	12
Figure 7. Cell death graphical representation.....	13
Figure 8. Cell proliferation graphical representation.....	14
Figure 9. Percentage of DSBs graphical representation.....	15
Figure 10. Percentage of inactive Bcl-2 graphical representation.....	16

1.1.1. *Project Overview*

Project Overview: This project aims to develop a comprehensive system for the management of organizational resources. The system will be designed to support the strategic and operational needs of the organization, providing a central platform for data collection, analysis, and reporting. The project is expected to be completed by the end of 2023, with a budget of approximately \$1.5 million.

Project Objectives: The primary objectives of this project are to: 1) Improve the efficiency of resource allocation; 2) Enhance the accuracy of financial reporting; 3) Provide real-time insights into organizational performance; and 4) Ensure the system is scalable and secure.

Chapter I

Introduction

The purpose of this chapter is to provide an overview of the project and its significance. It will discuss the background of the organization, the current challenges it faces, and the proposed solution. The chapter will also outline the project's scope, objectives, and the expected benefits. The project is a strategic initiative for the organization, and its successful implementation will have a significant impact on its long-term success.

1.1.2. *Context*

Context: The current landscape of business is characterized by rapid technological change and increasing competition. Organizations are required to be more agile and data-driven to remain competitive. This project is a response to these challenges, aiming to provide a robust and scalable solution for resource management. The project is supported by senior management and has a strong track record of successful implementation. The system will be developed using the latest technologies and will be designed to be user-friendly and easy to integrate with existing systems.

1.1.1 Breast Cancer

Recent statistics show that breast cancer is the most common and the second leading cause of malignancy-related deaths among women worldwide (Crawford and Richard, 2013; Jemal et al., 2011). In the United States alone, more than 200,000 new cases are diagnosed annually and approximately 40,000 deaths occur as a result of this malignancy (Crawford and Richard, 2013; Jemal et al., 2011).

Cancer occurs when cells begin to proliferate uncontrollably due to genetic mutations and/or abnormal proliferation signals (Watts, 2013). During puberty, the hormone estrogen signals the development of breast tissue and the hormone progesterone stimulates the growth of lobular tissue (BreastCancer.org, Watts, 2013). When abnormal cell growth arises in ductal breast tissue, it is termed ductal carcinoma in situ, DCIS and when abnormalities occur in lobular tissue, it is termed lobular carcinoma in situ, LCIS (BreastCancer.org, Watts, 2013). Breast cancer onset can take a few years, beginning with a cluster of cells that are unable to follow the “rules” of the cell cycle progression, to a later accumulation of increased mutations that make these cells capable of unrestricted proliferation (BreastCancer.org).

1.1.2 Curcumin

Curcumin, the active ingredient of turmeric (curry spice), is believed to be associated with reduced incidence rates of breast cancer among women in Southeast Asian countries (Nagaraju et al. 2012). Curcumin (diferuloylmethane) is a yellow-colored polyphenol that is extracted from the *Curcuma longa* plant, commonly known as turmeric (Liu and Chen, 2013; Welkin et al., 2011). It has been consumed as a dietary spice for centuries and has played a key role in Ayurvedic medicine. Curcumin is known to possess a variety of beneficial and

therapeutic properties such as anti-oxidant, anti-inflammatory, anti-metastatic and anti-proliferative activities (Liu and Chen, 2013; Wilken et al., 2011). Recent studies have also proven curcumin's anti-cancer properties via its suppression of gene products such as c-myc, Bcl-2, and interleukins, its target of biological pathways, inhibition of growth factors, transcription factors and cell adhesion molecules involved in mutagenesis, cell cycle regulation, gene expression, apoptosis, metastasis, and tumorigenesis (Wilken et al., 2011, Liu and Chen 2013; Cine et al., 2013).

1.1.3 p53 protein and cancer

The tumor suppressor gene p53, often referred to as the “guardian of the genome”, is required for checkpoint control during cell cycle progression (Meikrantz and Schlegel, 1995). During the cell cycle, DNA damage is detected at two checkpoints: the G₁-S phase and the G₂-M phase (Merritt et al., 1997). Depending on the type and extent of DNA damage, p53 can initiate one of three pathways: cell cycle arrest at a checkpoint, repair of the damaged cell, or apoptosis of the cell (Elmore, 2007; Figure 1). However, in cancer cells, these pathways are defective and therefore the cells lack the ability to detect damage and thereby initiate an appropriate response as needed. This can eventually lead to genetic instability and uncontrolled cell growth resulting in tumors (Elmore, 2007). When DNA damage occurs, the kinase ATM is the first protein to be phosphorylated. Once phosphorylated, ATM can activate various downstream gene products, including p53, SMC1, NBS1, and Histone H2A.X (Burma et al., 2001; Figure 1). One of these factors, the H2A.X, is a key indicator of the levels of DNA damage, specifically double-stranded DNA breaks. As the level of DNA damage rises, the level of phosphorylated H2A.X also rises and begins to aggregate at locations where damage is detected (Burma et al., 2001). H2A.X can

then activate various downstream factors such as p53 to initiate repair of the damaged cell. The major role of the checkpoint controlling entry into S phase of the cell cycle is to prevent damaged cells from replicating damaged DNA. This is controlled via the p53 protein, which induces either cell cycle arrest in G₁ or cause apoptosis (Merritt et al., 1997; Figure 1). Although the p53-mediated G₁ arrest has been commonly studied, p53 regulation at the G₂ phase has also been found to occur (Guillouf et al., 1995).




Figure 1. A proposed pathway by which ATM and H2A.X activate p53 to induce DNA repair, cell cycle arrest, and apoptosis¹⁷.

1.1.4 Bcl-2 protein and cancer

Bcl-2 is a member of the much larger Bcl-2 family that controls apoptosis by regulating the proteins responsible for mitochondrial membrane permeability (Jin and El-Deiry, 2005). The Bcl-2 family consists of two protein sub-groups, the pro-apoptotic proteins and the anti-apoptotic

proteins. Bcl-2 is an anti-apoptotic family member that acts by forming a complex with pro-apoptotic proteins such as Bax, thereby sequestering them and inhibiting their insertion into the mitochondrial membrane (Adams and Cory, 2007). When an apoptotic signal is received, Bcl-2 is supposed to release Bax, allowing Bax to insert into the mitochondria and release cytochrome C into the cytoplasm, promoting caspase activation and cell death (Figure 2). However, in cancer cells, apoptosis is often prevented due to an up-regulation of anti-apoptotic Bcl-2 proteins or dysfunction of pro-apoptotic proteins (Adams and Cory, 2007). The ratio of anti-apoptotic Bcl-2 proteins is often very high in various hematologic malignancies such as non-Hodgkin's lymphoma (Jin and El-Deiry, 2005). Preclinical studies have shown that Bcl-2 serves a major role in regulating the intrinsic apoptotic signaling pathway by binding pro-apoptotic molecules and thereby preventing them from carrying out their specific functions within the mitochondria (Jin and El-Deiry, 2005). Therefore, Bcl-2 is capable of promoting the survival of regulatory B-cells by inhibiting apoptosis, which can result in chemotherapy resistance in various hematologic cancers (Adams and Cory, 2007).




Figure 2. A proposed pathway by which Bcl-2 and Bax interact to induce apoptosis after cell damage is detected²⁰.

1.1.5 Aims of our Study

The goals of this study were to determine the anti-proliferative and apoptotic effects of curcumin on MDA-MB 468 cells and to determine how curcumin affects the abundance of two specific apoptotic and cell cycle proteins: p53 and Bcl-2. We hypothesized a dose-dependent increase in cell death and a dose-dependent decrease in cell proliferation following curcumin treatment. We also hypothesized that curcumin would cause an up-regulation of p53 expression and a down-regulation of Bcl-2 activation (Figure 3).




Figure 3. Curcumin is hypothesized to cause DNA damage. The p53 pathway on the left begins with the activation of the kinase ATM as soon as DNA damage is detected. ATM will then phosphorylate the H2A.X protein (the double-strand break detector) causing an up-regulation of p53. The pathway on the right shows the Bcl-2 pathway. We predict curcumin to cause an increase in inactive Bcl-2 via phosphorylation, which will cause a conformational change and prevent Bax (a pro-apoptotic protein) to be bound to Bcl-2. This will allow Bax to form its own complex and activate caspases that will eventually lead to apoptosis.

Chapter II

Materials and Methods

Materials

MDA-MB-468 breast adenocarcinoma cells were obtained from the American Type Cell Culture Collection (ATCC, Manassas, VA) and cultured in Leibovitz L-15 Media (Hyclone, Logan, UT) supplemented with 10% Fetal Bovine Sera (Atlanta Biologicals, Atlanta, GA) and 1% Antibiotic/antimycotic (Invitrogen, Grand Island, NY). MDA-MB-468 were grown at 37°C in a free gas exchange with atmospheric air. Curcumin (diferuloylmethane) and dimethyl sulfoxide (DMSO) were obtained from GE Healthcare Bio-Sciences (Pittsburgh, Pennsylvania). Annexin V and Dead Cell kit, Multi-Colored DNA Damage kit, and Bcl-2 Dual Detection kit were obtained from EMD Millipore Muse® (Billerica, Massachusetts).

Cell Treatment

Curcumin was first dissolved in DMSO (Sigma-Aldrich, St. Louis, MS) as a 10 mM stock solution and subsequently diluted in L-15 culture medium. Medium containing the same amount of DMSO was used as control at concentrations not exceeding the 0.1% v/v of the culture medium. Cells were incubated with curcumin at 0, 25, 50, and 100 μ M, at three time points namely 24, 48, and 72 h. Briefly, MDA-MB-468 cells were inoculated at a density of 2×10^4 cells/well in a 12-wellplate for 72 h in 1 mL of L-15 medium. The culture supernatant was then removed and medium containing the above mentioned curcumin concentrations were added to cells and subsequently incubated for 24 h.




Figure 4. Cell treatment in 12-well plate.

Annexin V and Dead Cell Kit

This kit was used to determine percentage of cell death and proliferation. Cells were detached from the plate using a cell scraper and 100 μ L of the cell suspension was placed into microcentrifuge tubes. Next, 100 μ L of Muse Annexin V and Dead Cell Reagent was added into each tube. Samples were mixed thoroughly using a vortex at medium speed for about 3 to 5 seconds. Finally, samples were stained for 20 min. at RT and analyzed using a Muse Cell Analyzer.

Multi-Color DNA Damage Kit

This kit was used to indirectly measure p53 expression. Briefly, cells were centrifuged at 300 x g for 5 min. and washed once with 1X PBS. Following centrifugation, cells were fixed with Fixation Buffer and permeabilized with Permeabilization Buffer for 10 min. on ice and washed with PBS as before. Cells were then treated with 10 μ L of the Anti-phospho-ATM /Anti-

phospho-Histone H2A.X antibody cocktail. Samples were incubated in the dark for 30 min. at room temperature. Following incubation, cells were centrifuged at 300 x g for 5 min. and washed with 1X Assay Buffer. Cells were then resuspended in 200 μ L 1X Assay Buffer and analyzed on a Muse Cell Analyzer.

Bcl-2 Dual Detection Kit

This kit was used to measure percentage of Bcl-2 protein activation. The procedure for this assay was exactly as the Multi-Color DNA Damage Kit, except cells were fixed and permeabilized for 5 min. and an antibody cocktail consisting of a phospho-specific anti-phospho-Bcl-2 +anti-Bcl-2 was used.

Statistical Analyses

The independent variable in the experiment was the curcumin concentration. The dependent variables were percentage of cell death, cell proliferation, DNA Double-strand breaks, and Bcl-2 inactivation. Data were expressed as means \pm 1 SD. Statistical significance of differences between means was determined by 1-way ANOVA and a Least Significant Differences (LSD) Post-Hoc test. Probability values of $p < 0.05$ were considered statistically significant.



Chapter III

Results and Discussion

Figure 1. Screenshot of a software interface showing a window with a title bar and a main content area. The content area appears to be a grid or table with some text and possibly a small chart or diagram. The interface is somewhat blurry and low-resolution.

Figure 2. Screenshot of a software interface, similar to Figure 1, showing a window with a title bar and a main content area. The content area displays a different set of data or a different view of the same data, possibly a different chart or table layout.

Results




Figure 5. Results of a sample acquired using Muse® Cell Analyzer for the Multi-Color DNA Damage assay. The lower left quadrant shows percentage of negative cells that had no DNA damage. The lower right quadrant shows percentage of H2A.X- activated cells. The upper left shows percentage of ATM-activated cells, and the upper right indicates percentage of DNA double-strand breaks via dual activation of both ATM and H2A.X. This is the quadrant we were interested in since p53 expression involves the phosphorylation of both ATM and H2A.X.




Figure 6. Results of a sample acquired using Muse® Cell Analyzer for the Bcl-2 Activation Assay. The lower left quadrant shows percentage of non-expressing cells. The upper right quadrant indicates percentage of activated cells via phosphorylation, and the upper left shows percentage of inactivated cells. This is the quadrant we were interested in since inactivation of Bcl-2 allows Bax to form a complex where it can induce apoptosis in the cells via activation of a series of downstream proteins.




Figure 7. Effect of curcumin on average (± 1 S.E) death of MDA-MB 468 cells. The x-axis corresponds to the different concentrations of curcumin measured in μM (0, 25, 50, and 100) and the y-axis corresponds to the total cell death percentage. Cells were treated with the above curcumin concentrations for 24 h. Cell death was determined using the Annexin V Dead Cell kit and a Muse Cell Analyzer.




Figure 8. Effect of curcumin on average (± 1 S.E) proliferation of MDA-MB 468 cells. The x-axis corresponds to the different concentrations of curcumin measured in μM (0, 25, 50, and 100) and the y-axis corresponds to the total cell proliferation in cells/mL. Cells were treated with the above curcumin concentrations for 24 h. Proliferation was determined using the Annexin V Dead Cell kit and a Muse Cell Analyzer.




Figure 9. Effect of curcumin on average (\pm 1 S.E) p53 expression in MDA-MB 468 cells. Cells were exposed to 0, 25, 50, and 100 μ M curcumin for 24 hours to induce DNA damage, and then stained with both anti-phospho-Histone H2A.X and anti-phospho-ATM antibodies in multiplex. Indirect p53 expression was determined via percentage of double-strand breaks in DNA as a result of phosphorylation of the ATM and H2A.X proteins as seen on the y-axis. Samples were acquired using the MuseTM Cell Analyzer.




Figure 10. Effect of curcumin on average (\pm 1 S.E) expression of inactive Bcl-2 in MDA-MB 468 cells. Cells were exposed to 0, 25, 50, and 100 μ M curcumin for 24 hours to induce DNA damage, and then stained with both a phospho-specific anti-phospho-Bcl-2 and anti-Bcl-2 antibodies in multiplex. Samples were acquired using the Muse™ Cell Analyzer.

Discussion

Curcumin caused a significant dose-dependent increase in cell death at the different concentrations (1-way ANOVA, $F_{3,32}=4.119$, $P=0.014$; Figure 7) with the 100 μM concentration having a significantly higher amount of cell death compared to the control, 25 μM , and 50 μM . Curcumin was hypothesized to induce a dose-dependent decrease in cell proliferation. However, our data shows that curcumin caused a dose-dependent increase in cell proliferation when compared to the control (1-way ANOVA, $F_{3,32}=6.005$, $P=0.002$; Figure 8) which is contrary to the hypothesis. Hence, our data does not support the hypothesis. A possible explanation for this difference may be as follows. Since curcumin is derived from turmeric, a root plant containing starch, the cells could be using this starch as an additional carbon source to provide the energy needed to grow and reproduce.

Our results also suggest that curcumin did not cause an increase in p53 expression because there was no significant difference in curcumin-induced double strand breaks (DSB) in DNA (1-way ANOVA, $F_{3,32}=1.505$, $P=0.232$; Figure 9). The connection between DSB and p53 is in the pathway involving activation of ATM which detects DNA damage (Figure 1). When DNA damage occurs (curcumin-induced DSB) ATM phosphorylates the H2A.X protein which responds to DSBs. H2A.X then activates p53 to induce apoptosis or DNA repair. Therefore, an increase in DSBs should indirectly correspond to an increase in p53 expression. Furthermore, this increase in p53 expression should allow for p53 to initiate apoptosis because of a dose-dependent increase in DNA damage. Although a dose-dependent increase in DSB was shown in both the 25 μM and 50 μM curcumin treated samples relative to the control, the data was not statistically significant, $p=0.232$. Therefore, more trials need to be conducted in order to get a

significant result and prove the proposed pathway. In a study conducted by Rowe et al. in 2009, they found that following curcumin treatment, phosphorylation occurred at ATM-specific sites on the breast cancer (BRCA) 1 gene, which was consistent with the activation of a DNA damage response. In addition, phosphorylated H2A.X was detected in cells treated with curcumin, indicating that the type of damage induced by curcumin was a DNA double strand break (Rowe et al., 2009).

Curcumin did not cause a significant dose-dependent increase in inactive Bcl-2 expression as previously hypothesized (1-way ANOVA, $F_{3,20}=0.102$, $P=0.958$; Figure 10). It is known that curcumin induces apoptosis via suppression of anti-apoptotic Bcl-2, allowing the release of cytochrome c into the cytosol and activation of caspases (Anto et al., 2002; Rao et al., 2011). Inactivation (phosphorylation) of Bcl-2 allows the Bax protein to form a complex within the mitochondrial membrane that induces cell death. Although there was a slight increase in phosphorylated Bcl-2, it was not significant, $p=0.958$. Hence, more trials need to be conducted in order to see a more accurate reflection of the pathway taken by these cells to reach apoptosis.

Although there was no significant increase in p53 expression or Bcl-2 inactivation, there was a significant increase in both cell death and proliferation. Hence, future studies can look at the rates of curcumin-induced cell death versus cell growth in the cells. Also, the cells can be treated with curcumin for longer periods of time (24, 48, and 72 hours) which may result in curcumin inducing a higher percentage of DNA damage thereby decreasing the amount of cell growth. Also, curcumin's effects on p53 and Bcl-2 proteins should be studied on other carcinomas since these two oncoproteins play vital roles in cellular pathways.

References

1. Adams, J. M., and Cory, S. 2007. The Bcl-2 apoptotic switch in cancer development and therapy. *Oncogene* 26:1324-37.
2. Anto, R. J., Mukhopadhyay, A., Denning, K., and Aggarwal, B. B. 2002. Curcumin (diferuloylmethane) induces apoptosis through activation of caspase-8, BID cleavage and cytochrome c release: its suppression by ectopic expression of Bcl-2 and Bcl-xl. *Carcinogenesis* 23:143-150.
3. Breastcancer.org. What is Breast Cancer? N.p., n.d. Web. 5 May 2014.
<http://www.breastcancer.org/symptoms/understand_bc/what_is_bc>.
4. Burma, S., Chen, B., Murphy, M., Kurimasi, A., and Chen, D. 2001. ATM Phosphorylates Histone H2AX in Response to DNA Double-strand Breaks. *The Journal of Biological Chemistry* 276:42462-42467.
5. Cine, N., Limtrakul, P., Sunnetci, D., Nagy, B., and Savli, H. 2013. Effects of curcumin on global gene expression profiles in the highly invasive human breast carcinoma cell line MDA-MB 231: A gene network-based microarray analysis. *The Journal of Experimental Medicine* 5:23-27.
6. Crawford, S., and Richard, A. 2013. Magill's Medical Guide (Online Edition). Breast cancer. *Research Starters* 1:175-180.
7. Elmore, S. 2007. Apoptosis: A Review of Programmed Cell Death. *Toxicologic Pathology* 35:495-516.

8. Guillouf, C., Rosselli, F., and Krishnaraju, K. 1995. p53 involvement in control of G2 exit of the cell cycle: role in DNA damage-induced apoptosis. *Oncogene* 10:2263-2270.
9. Jemal, A., Bray, F., Center, M. M., Ferlay, J., Ward, E., and Forman, D. 2011. Global cancer statistics, *CA- A Cancer Journal for Clinicians* 61:69-90.
10. Jin, Z., and El-Deiry, W. S. 2005. Overview of cell death signaling pathways. *Cancer Biology and Therapy* 4:139-163.
11. Liu, D., and Chen, Z. 2013. The effect of curcumin on breast cancer cells. *Journal of Breast Cancer* 16:133-137.
12. Meikrantz, W. and Schlegel, R. 1995. Apoptosis and the cell cycle. *Journal of Cellular Biochemistry* 58:160-174.
13. Merritt, A. J., Allen, T. D., Potten, C.S., and Hickman, J.A. 1997. Apoptosis in small intestinal epithelial from p53-null mice, evidence for a delayed, p53-independent G2/M associated cell death after gamma-irradiation. *Oncogene* 14: 2759-2766.
14. Nagaraju, G. P., Aliya, S., Zafar, S. F., Basha, R., Diazd, R., and El-Rayes, B. F. 2012. The impact of curcumin on breast cancer. *Integrative Biology* 4:996-1007.
15. Rao, J., Xu, D. R., Zheng, F. M., Long, Z. J., Huang, S. S., Wu, X., Zhou, W. H., Huang, R. W., and Liu, Q. 2011. Curcumin reduces expression of Bcl-2, leading to apoptosis in daunorubicin-insensitive CD34+ acute myeloid leukemia cell lines and primary sorted CD34+ acute myeloid leukemia cells. *Journal of Translational Medicine* 9:1-32.
16. Rowe, D., Ozbay, T., and Nahta, R. 2009. Modulation of the BRCA1 Protein and Induction of Apoptosis in Triple Negative Breast Cancer Cell Lines by the

Polyphenolic Compound Curcumin. *Breast Cancer: Basic and Clinical Research* 3:61-75.

17. The DNA Damage Signaling Pathway. 2009. Muse™ Multi-Color DNA Damage Kit User's Guide [Internet]. Available from:
http://www.emdmillipore.com/US/en/product/,MM_NF-MCH200107#anchor_UG.
18. Wilken, R., Veena, M. S., Wang, M. B., and Srivatsan, E. S. 2011. Curcumin: A review of anti-cancer properties and therapeutic activity in head and neck squamous cell carcinoma. *Molecular Cancer* 10:1-19.
19. Watts, T. 2013. Primary Breast Cancer: What do practice nurses need to know? *Practice Nurse* 43:36-42.
20. Zhang, S. 2003. Cancer Research: School of Life Sciences, Xiamen University [Internet] Available from: <http://121.192.179.196/english/teacher/ShowArticleID=114>.

