

# The Effects of Teacher Candidates' Physical Activity Levels on Health-Related Quality of Life

Barış Gürol<sup>1</sup>, Gülsün Güven<sup>1</sup> & Dilek Yalız-Solmaz<sup>1</sup>

<sup>1</sup> Faculty of Sports Sciences, Eskişehir Technical University, Turkey

Correspondence: Barış Gürol, Faculty of Sports Sciences, Eskişehir Technical University, Eskişehir, Turkey.

Received: August 15, 2019

Accepted: September 21, 2019

Online Published: October 28, 2019

doi:10.5539/ies.v12n11p178

URL: <https://doi.org/10.5539/ies.v12n11p178>

## Abstract

The aim of this research was to determine the effects of physical activity levels of teacher candidates on the sub-dimensions of health-related quality of life. In the research among the quantitative research methods, relational survey model was used. A total of 90 teacher candidates participated in this research. The International Physical Activity Questionnaire-Short Form (IPAQ-SF) and the 36-Item Short Form Health Survey (SF-36) were used in this study. In the data analysis, "percentage, frequencies, standard deviation, mean, Product-Moment Correlation coefficients and Multiple regression" were used. According to the results, role functioning/emotional, pain and general health sub-dimensions are important predictors on physical activity levels. However, physical functioning, emotional well-being, vitality, social functioning, role functioning/physical, sub-dimensions have not an important impact on physical activity levels statistically. As a conclusion, participation in physical activity can be said to have a negative effect on emotional problems and pain, and a positive effect on general health status.

**Keywords:** physical activity, health related quality of life, teacher candidates, regression

## 1. Introduction

Inactive lifestyle increases the risk of developing many health conditions particularly cardiovascular and coronary heart diseases, diabetes mellitus, risks of certain cancers as colon and breast cancers, hypertension, osteoporosis, poor musculoskeletal and cardiorespiratory fitness, high blood pressure and elevated cholesterol levels, overweight or obesity, anxiety and depression (Cummings et al., 1985; Berlin & Colditz, 1990; Lee et al., 1991; Arroll & Beaglehole, 1992; Wannamethee & Shaper, 1992; Martinez et al., 1997; Uçkun & Çalikoğlu, 2003; Bize et al., 2007; Lahti et al., 2010; Li et al. 2016).

Regular moderate intensity physical activity decreases the risk of various chronic diseases as hypertension, coronary heart disease, cardiovascular disease, diabetes, cancer, all-cause mortality and preserves both physical and intellectual health and functions (Saxena et al., 2005; Acree et al., 2006; Blair & Morris, 2009).

Health-related quality of life (HRQL) can be described as how well his/her perceived wellbeing in intellectual, physical and social domains of health (Brown et al., 2004). The health benefits from physical activity are well-known and research on HRQL in the physical activity domain has predominately been focused on elder populations and among patients with chronic diseases (Elavsky et al., 2005; Acree et al., 2006; Bize et al., 2007; Burke et al., 2017; Pati et al., 2018; Groessl et al., 2019).

While little is known regarding HRQL and its relationship with physical activity level (Bize et al., 2007), it is stated that younger people propose that physical activity level is positively effect to HRQL (Maher et al., 2016).

However, it is not clear whether physical activity increase specific sub-dimensions of HRQL (Acree et al., 2006). The aim of this study was to reveal the effects of teacher candidates' physical activity levels on the sub-dimensions of HRQL.

## 2. Methods

### 2.1 Research Model

In the research among the quantitative research methods, relational survey model was used. The relational survey model is a research design that aims to determine the existence and degree of change between two and more variables together (Karasar, 2007).

## 2.2 Participants

A total of 110 questionnaires were distributed to teacher candidates from Department of Physical Education and Sports Teacher Training in Eskişehir Technical University and 90 of them were completed and returned, giving an overall response rate of 81.82 percent. The data were gathered during 2018-2019 Academic Year. 42.2% of the teacher candidates are women and 57.8% of the teacher candidates are male in this research. The ages differences of the participants are as follow: 42.2% of them are in 18-20 years old, 43.3% of them are in 21-23 years old and 14.4% of these teacher candidates are in 24 years old and above.

## 2.3 Data Collection Tools

### 2.3.1 The International Physical Activity Questionnaire–Short Form (IPAQ-SF)

There are various questionnaires to determine the physical activity levels (Lee et al., 2011). It was indicated that IPAQ-SF version is one of the most appropriate and widely used self-report form to assess physical activity levels (Tierney et al., 2015; Lavelle et al., 2019). IPAQ is designed to evaluate the young people/middle-aged adults and special groups' physical activity levels (Craig et al., 2003). The validity and reliability study of the scale, in Turkey in 2005 by Ozturk (2005).

It consists of seven questions to measure average daily time spent walking, sitting and participating in activity of moderate and high intensity and produces a physical activity score based on last seven days (Tierney et al., 2015; Sanda et al., 2017; Lavelle et al., 2019).

Physical activity data from the survey were converted to energy expenditure values predicated as MET scores. The total hours calculated in each physical activity levels were multiplied by the specific MET score for the activity to determine the weekly physical activity level (Hagstromer et al., 2006). Physical activity levels are determined by using MET scores as vigorous intensity physical activity level (8.0 MET), moderate intensity physical activity level (4.0 MET), walking (3.3 MET), sitting (1.5 MET) (Özüdoğru, 2013).

### 2.3.2 The 36-Item Short Form Health Survey (SF-36)

The 36-Item Short Form Health Survey (SF-36) is developed for assessing HRQL (Lins & Carvalho, 2016). The validity and reliability study of the scale, in Turkey (Koçyiğit et al., 1998).

The SF-36 consisting of 36 items assessing eight sub-dimensions: Physical Function (PF), Social Function (SF), Role Function – Physical (RFP), Role Function – Emotional (RFE), Emotional Well-being (EW), Vitality (VT), Bodily Pain (BP) and General Health Perception (GHP) (Khanna et al., 2010; Burholt & Nash, 2011; Lins & Carvalho, 2016; Matcham et al., 2016; İnci et al., 2017; Jacobsen et al., 2018).

SF-36 sub-dimensions are evaluated from 0 to 100, with a higher score which indicated better health living (Khanna et al., 2010; İnci et al., 2017). Item scores were transformed to 0-100 point scales (0=worst, 100=best) using the SF-36 algorithm (Jacobsen et al., 2018).

## 2.4 Data Analysis

Data obtained through IPAQ short form and HRQL short form-36 (SF-36), and the scores obtained from the assessment were coded and computed using the SPSS 24.0, which was also used to perform analyses. In the data analysis, “percentage, frequencies, standard deviation, mean, Product-Moment Correlation coefficients and Multiple regression” were calculated. To determine the relationship between quality of life behaviors and physical activity levels, correlation coefficients were used. Multiple regression analysis was performed to determine whether the scores are predictive of each other ( $\beta$ ).

## 3. Results

Table 1. Teacher candidates' physical activity levels

	f	%
Low Intensity Level	10	11.1
Moderate Intensity Level	31	34.4
High Intensity Level	49	54.4

As shown in Table 1, it is find out that 54.4% of teacher candidates were in high physical activity level while 34.4% of them were in moderate physical activity level and 11.1% of them were in low physical activity level.

Table 2. Teacher candidates' quality of life behaviors

	N	$\bar{X}$	SD
PF	72	89.94	14.17
RFP	72	71.67	37.70
RFE	72	57.78	42.63
VT	72	63.94	17.00
EW	72	70.22	15.19
SF	72	65.69	25.59
BP	72	73.69	19.37
GHP	72	69.00	15.65

The SF-36 consisting of 36 items assessing eight sub-dimensions: Physical Function (PF), Social Function (SF), Role Function – Physical (RFP), Role Function – Emotional (RFE), Emotional Well-being (EW), Vitality (VT), Bodily Pain (BP) and General Health Perception (GHP).

As shown in Table 2, it was examined that the highest score belong to in PF (M=89.94) and then BP (M=73.69) and RFP (M=71.67) sub-dimensions, the lowest score belong to EW (M=70.22), GHP (M=69.00), SF (M=65.69), VT (M=63.94) and RFE (M=57.78) sub-dimensions.

Table 3. Correlations between physical activity levels and quality of life sub-dimensions

	1	2	3	4	5	6	7	8	9	
PF	r	1								
	p									
	N	90								
RFP	r	.391**	1							
	p	.000								
	N	90	90							
RFE	r	.163	.488**	1						
	p	.124	.000							
	N	90	90	90						
VT	r	.118	.095	-.083	1					
	p	.270	.372	.438						
	N	90	90	90	90					
EW	r	.179	.141	-.189	.648**	1				
	p	.091	.186	.402	.000					
	N	90	90	90	90	90				
SF	r	.206	.397**	.455**	.126	.057	1			
	p	.052	.000	.000	.238	.596				
	N	90	90	90	90	90	90			
BP	r	.272**	.402**	.198	.222*	.196	.364**	1		
	p	.009	.000	.062	.035	.064	.000			
	N	90	90	90	90	90	90	90		
GHP	r	.253*	.223*	-.039	.143	.179	.097	.275**	1	
	p	.016	.035	.717	.180	.092	.361	.009		
	N	90	90	90	90	90	90	90	90	
Physical Activity Level	r	.043	-.160	-.276**	.073	.100	-.175	-.214*	.140	1
	p	.688	.131	.008	.493	.347	.098	.042	.189	
	N	90	90	90	90	90	90	90	90	90

\*p<.05; \*\*p<.01.

The SF-36 consisting of 36 items assessing eight sub-dimensions: Physical Function (PF), Social Function (SF), Role Function – Physical (RFP), Role Function – Emotional (RFE), Emotional Well-being (EW), Vitality (VT), Bodily Pain (BP) and General Health Perception (GHP).

The analysis of Table 3 demonstrates that there was a significant negative relationship between the physical

activity and RFE ( $r=-.276$ ,  $p=.008$ ), BP ( $r=-.214$ ,  $p=.042$ ) whereas the study found no significant relationship with other sub-dimensions.

Multiple regression analysis (backward) was performed to determine the factors affecting the physical activity levels of teacher candidates. Purpose of this analysis is to determine the independent variables that explain the highest dependent variable by adding all independent variable which acting on the dependent variable to the model. Firstly, all the independent variables are included in the model and then statistically insignificant variables and variables which decrease the explaining rate are removed from the model one by one. This statistical process continues until getting the highest corrected  $R^2$  value. In accordance with these explanations, model of physical activity was stated as dependent variable in Table 4.

Table 4. Regression table for teacher candidates' physical activity to predict their quality of life behaviors

	Model	$\beta$	t	p	R	Adjusted $R^2$	F	p
Physical Activity	1 (Constant)	2.076	3.592	.001	.395	.072	1.868	.076
	PF	.005	.936	.352				
	RFP	-.001	-.373	.710				
	RFE	-.003	-1.506	.136				
	VT	.002	.308	.759				
	EW	.003	.460	.647				
	SF	-.001	-.211	.834				
	BP	-.009	-2.038	.045				
	GHP	.007	1.529	.130				
	2 (Constant)	2.066	3.608	.001	.394	.083	2.153	.047
	PF	.005	.934	.353				
	RFP	-.001	-.400	.690				
	RFE	-.003	-1.687	.095				
	VT	.002	.286	.775				
	EW	.003	.476	.635				
	BP	-.009	-2.152	.034				
	GHP	.007	1.536	.128				
	3 (Constant)	2.081	3.672	.000	.393	.093	2.527	.027
	PF	.005	.934	.353				
	RFP	-.001	-.407	.685				
	RFE	-.003	-1.711	.091				
	EW	.004	.837	.405				
	BP	-.009	-2.145	.035				
	GHP	.007	1.546	.126				
	4 (Constant)	2.145	3.959	.000	.391	.102	3.029	.015
	PF	.005	.862	.391				
	RFE	-.004	-2.158	.034				
	EW	.004	.806	.423				
BP	-.009	-2.326	.022					
GHP	.007	1.513	.134					
5 (Constant)	2.316	4.652	.000	.382	.106	3.639	.009	
PF	.005	.977	.331					
RFE	-.004	-2.297	.024					
BP	-.009	-2.234	.028					
GHP	.008	1.597	.114					
6 (Constant)	2.642	7.168	.000	.370	.106	4.536	.005	
RFE	-.004	-2.184	.032					
BP	-.008	-2.089	.040					
GHP	.008	2.001	.049					

The SF-36 consisting of 36 items assessing eight sub-dimensions: Physical Function (PF), Social Function (SF), Role Function – Physical (RFP), Role Function – Emotional (RFE), Emotional Well-being (EW), Vitality (VT),

#### Bodily Pain (BP) and General Health Perception (GHP).

As shown in Table 4, SF sub-dimension was removed firstly and then VT, RFP, EW and PF sub-dimensions were removed from the model. Priority of removing the sub-dimension from the model was determined according to low explanation effect on dependent variable. RFE, BP and GHP sub-dimensions explained a large part (11 %) of the total variance in model 6 which has the highest corrected  $R^2$  value and also explaining the highest dependent variable ( $R^2=.106$ ,  $p<.05$ ). According to analysis it was found that  $F=4.536$ ,  $p=.005$ . According to the results of the analysis on the significance of regression coefficients, RFE, BP and GHP sub-dimensions were found to be significant predictors of physical activity ( $p<.05$ ).

#### 4. Discussion and Conclusions

Among the teacher candidates, the highest percentage (54.4%) was students with a high physical activity level, followed by 34.4% with a moderate physical activity level and 11.1% with a low physical activity level. According to this finding, it can be said that most of the teacher candidates are physically active people. The main reason for this is that the teacher candidates are students of the Department of Physical Education and Sports Teaching and that practical courses are included in the curriculum. Another reason for the high level of physical activity levels of teacher candidates is that they have opportunities to do physical activities. It is known that the physical activity areas have a significant effect on the participation of people (Honca & Çetinkaya, 2017; Gümüş & Özgül, 2017). However, when the literature is examined, it contradicts the findings of this study. Yahia, Wang, Rapley, and Dey (2016) reported that physical activity levels and lifestyle score stated that most of the students were not in a active lifestyle. In this research, only 7% of students indicated having a very active lifestyle. Ölçücü et al. (2015) stated in their research on university students that 36% of the students were at a sufficient physical activity level, 43% were at a low physical activity level and 21% were not physically active. Savcı et al. (2006) were obtained similar results in their study.

As our research results indicated that the highest means according to HRQL Questionnaire belongs to PF, BP and RFP sub-dimensions and the lowest means belongs to VT and RFE sub-dimensions. Our research results are alike to the research results which examined by Pekmezovic et al. (2011). Pekmezovic et al. (2011) stated that the highest means of the SF-36 scales were obtained for PF sub-dimension and the lowest means for VT sub-dimension.

In the correlation analysis, Pearson correlation coefficients representing the relationship between physical activity levels and HRQL sub-dimensions were calculated. When correlation values between physical activity levels and HRQL sub-dimensions are examined, it is observed that there is negatively a significant relationship between physical activity levels and RFE sub-dimension ( $r=.199$ ;  $p<.05$ ). Also, it has been found out that there is negatively a significant relationship between physical activity levels and BP sub-dimension ( $r=-.214$ ;  $p<.05$ ). It has been determined that there is no significant relationship between physical activity levels and PF ( $r=.043$ ;  $p>.05$ ), RFP ( $r=-.160$ ;  $p>.05$ ), VT ( $r=.073$ ;  $p>.05$ ), EW ( $r=.100$ ;  $p>.05$ ), SF ( $r=.175$ ;  $p>.05$ ) and GHP sub-dimensions ( $r=.140$ ;  $p>.05$ ).

Besides, according to another result of the study, among sub-dimensions of quality of life, only RFE, BP and GHP sub-dimensions could explain 11% of total variance of physical activity level. This result shows that RFE, BP and GHP sub-dimensions are important predictors on physical activity levels. Other sub-dimensions as PF, VT, RFP, SF, EW sub-dimensions have not an important impact on physical activity level. According to this finding, participation in physical activity can be said to have a negative effect on emotional problems (stress, anxiety, and pain), and a positive effect on general health status.

Most researchers suggest that physical activity had a positive result on quality of life (Gümüş & Işık, 2018; Hawkins & Duncan, 1991; Stewart & King, 1991; Buckelew et al., 1995; Puciato, Borysiuk, & Rozpara, 2017). Some earlier studies also indicated that there is a positive relationship between physical activity and health (Krzepota, Biernat, & Florkiewicz, 2015; Quehenberger, Cichocki, & Krajic, 2014). They may have resulted as physically active people's health status is better than inactive people (Skrzek et al., 2015). Also, researchers have documented the positive effect of physical activity as a factor enhancing the prevention or treatment of anxiety, depression or other neurological disorders (Teychenne, Ball, & Salmon, 2008; Dziubek et al., 2016; Lancer, Motta, & Lancer, 2007). For this reason, improving all kinds of infrastructure activities affecting the quality of life of teachers who will train the new generations of the society is important for educating healthy generations.

#### 5. Limitations

The current study focused on a limited number of teacher candidates ( $n=90$ ) from Department of Physical Education and Sports Teacher Training in Eskişehir Technical University.

## References

- Acree, L. S., Longfors, J., Fjeldstad, A. S., Fjeldstad, C., Schank, B., Nickel, K. J., ... & Gardner, A. W. (2006). Physical activity is related to quality of life in older adults. *Health and quality of life outcomes*, 4(1), 37. <https://doi.org/10.1186/1477-7525-4-37>
- Arroll, B., & Beaglehole, R. (1992). Does physical activity lower blood pressure? A critical review of the clinical trials. *Journal of Clinical Epidemiology*, 45, 439-447. [https://doi.org/10.1016/0895-4356\(92\)90093-3](https://doi.org/10.1016/0895-4356(92)90093-3)
- Berlin, J. A., & Colditz, G. A. (1990). A meta-analysis of physical activity in the prevention of coronary heart disease. *American Journal of Epidemiology*, 132, 612-627. <https://doi.org/10.1093/oxfordjournals.aje.a115704>
- Bize, R., Johnson, J. A., & Plotnikoff, R. C. (2007). Physical activity level and health-related quality of life in the general adult population: A systematic review. *Preventive Medicine*, 45, 401-415. <https://doi.org/10.1016/j.ypmed.2007.07.017>
- Blair, S. N., & Morris, J. N. (2009). Healthy Hearts—and the Universal Benefits of Being Physically Active: Physical Activity and Health. *Annals of Epidemiology*, 19(4), 253-256 <https://doi.org/10.1016/j.annepidem.2009.01.019>
- Brown, D. W., Brown, D. R., Heath, G. W., Balluz, L., Giles, W. H., Ford, E. S., & Mokdad, A. H. (2004). Associations between physical activity dose and health-related quality of life. *Medicine and science in sports and exercise*, 36(5), 890-896. <https://doi.org/10.1249/01.MSS.0000126778.77049.76>
- Buckelew, S. P., Murray, S. E., Hewett, J. E., Johnson, J., & Huysen, B. (1995). Self-efficacy, pain, and physical activity among fibromyalgia subjects. *Arthritis Care & Research*, 8(1), 43-50. <https://doi.org/10.1002/art.1790080110>
- Burholt, V., & Nash, P. (2011). Short Form 36 (SF-36) Health Survey Questionnaire: normative data for Wales. *Journal of Public Health*, 33(4), 587-603. <https://doi.org/10.1093/pubmed/fdr006>
- Burke, S., Wurz, A., Bradshaw, A., Saunders, S., West, M. A., & Brunet, J. (2017). Physical Activity and Quality of Life in Cancer Survivors: A Meta-Synthesis of Qualitative Research. *Cancers*, 9(5), 1-29. <https://doi.org/10.3390/cancers9050053>
- Craig, C. L., Marshall, A. L., Sjoström, M., Bauman, A. E., Booth, M. L., Ainsworth, B. E., ... Oja, P. (2003). International physical activity questionnaire: 12-country reliability and validity. *Medicine and Science in Sports and Exercise*, 35, 1381-95. <https://doi.org/10.1249/01.MSS.0000078924.61453.FB>
- Cummings, S. R., Kelsey J. L., Nevitt M. C., & O'Dowd K. J. (1985). Epidemiology of osteoporosis and osteoporotic fractures. *Epidemiological Reviews*, 7, 178-208. <https://doi.org/10.1093/oxfordjournals.epirev.a036281>
- Dziubek, W., Kowalska, J., Kuztal, M., Rogowski, L., Gołębiowski, T., Nikifur, M., ... & Woźniewski, M. (2016). The level of anxiety and depression in dialysis patients undertaking regular physical exercise training—a preliminary study. *Kidney and Blood Pressure Research*, 41(1), 86-98. <https://doi.org/10.1159/000368548>
- Elavsky, S., McAuley, E., Motl, R. W., Konopack, J. F., Marquez, D. X., Hu, L., ... Diener, E. (2005). Physical activity enhances long-term quality of life in older adults: Efficacy, esteem, and affective influences. *Annals of Behavioral Medicine*, 30(2), 138-145. [https://doi.org/10.1207/s15324796abm3002\\_6](https://doi.org/10.1207/s15324796abm3002_6)
- Farris, M. S., Kopciuk, K. A., Courneya, K. S., McGregor, S. E., Wang, Q., & Friedenreich, C. M. (2017). Associations of postdiagnosis physical activity and change from prediagnosis physical activity with quality of life in prostate cancer survivors. *Cancer Epidemiology and Prevention Biomarkers*, 26(2), 179-187. <https://doi.org/10.1158/1055-9965.EPI-16-0465>
- Groessl, E. J., Kaplan, R. M., Rejeski, W. J., Katula, J. A., Glynn, N. W., King, A. C., ... & Spring, B. (2019). Physical activity and performance impact long-term quality of life in older adults at risk for major mobility disability. *American journal of preventive medicine*, 56(1), 141-146. <https://doi.org/10.1016/j.amepre.2018.09.006>
- Gümüş, H., & Işık O. (2018). The Relationship of Physical Activity Level, Leisure Motivation and Quality of Life in Candidate Teachers. *International Journal of Progressive Education*, 14(5), 22-32. <https://doi.org/10.29329/ijpe.2018.157.3>
- Gümüş, H., & Özgül, S. A. (2017). Development of Scales for Barriers to Participation and Preference Factors in

- The Use of Recreation Area. *Journal of Human Sciences*, 14(1), 865-882. <https://doi.org/10.14687/jhs.v14i1.4448>
- Hagströmer, M., Oja, P., & Sjöström, M. (2006). The International Physical Activity Questionnaire (IPAQ): A study of concurrent and construct validity. *Public health nutrition*, 9(6), 755-762. <https://doi.org/10.1079/PHN2005898>
- Hawkins, W. E., & Duncan, T. (1991). Structural equation analysis of an exercise/sleep health practices model on quality of life of elderly persons. *Perceptual and motor skills*, 72(3), 831-836. <https://doi.org/10.2466/pms.1991.72.3.831>
- Honça, A. A., & Çetinkaya, T. (2017). Rekreasyonel Faaliyetlere Katılım Engelleri Algısı: Üniversite Öğrencileri Örneği. *International Journal of Recreation and Sports Science*, 1(1), 30-37.
- İnci, A., Çoban, M., Sarıkaya, M., & Maden, Ü. (2017). Prediyaliz Kronik Böbrek Hastalarında Kesitsel Bir Çalışma: Yaşam Kalitesi ve Etkileyen Faktörler. *Kocatepe Tıp Dergisi*, 18, 130-135. <https://doi.org/10.18229/kocatepetip.368668>
- Jacobsen, E. L., Bye, A., Aass, N., Fosså, S. D., Grotmol, K. S., Kaasa, S., ... & Hjerstad, M. J. (2018). Norwegian reference values for the Short-Form Health Survey 36: development over time. *Quality of Life Research*, 27(5), 1201-1212. <https://doi.org/10.1007/s11136-017-1684-4>
- Khanna, P. P., Ruiz, F. P., Maranian, P., & Khanna, D. (2010). Long-term therapy for chronic gout results in clinically important improvements in the health related quality of life: Short form-36 is responsive to change in chronic gout. *Rheumatology*, 50, 740-745. <https://doi.org/10.1093/rheumatology/keq346>
- Koçyiğit, H., Aydemir, Ö., & Ölmez, N. (1998) Kısa Form-36 (SF-36) nm Türkçe için güvenilirliği ve geçerliliği: Romatizmal hastalığı olan bir grup hasta ile çalışma. 34. Ulusal Psikiyatri Kongresi Bilimsel Çalışma Özet Kitabı. In S. Kültür, & L. Küey (Eds.), *İzmir Güneş Ofset* (pp. 290-291).
- Krzepota, J., Biernat, E., & Florkiewicz, B. (2015). The relationship between levels of physical activity and quality of life among students of the university of the third age. *Central European journal of public health*, 23(4), 335-339. <https://doi.org/10.21101/cejph.a4136>
- Lahti, J., Laaksonen, M., Lahelma, E., & Rahkonen, O. (2010). SF-36 total score as a single measure of health-related quality of life: Scoping review. *Preventive Medicine*, 50, 246-250. <https://doi.org/10.1016/j.ypmed.2010.02.007>
- Lancer, R., Motta, R., & Lancer, D. (2007). The effect of aerobic exercise on obsessive-compulsive disorder, anxiety, and depression: A preliminary investigation. *The Behavior Therapist*, 30, 53-62.
- Lavelle, G., Noorkoiv, M., Theis, N., Korff, T., Kilbride, C., Baltzopoulos, V., ... & Ryan, J. M. (2019). Validity of the International Physical Activity Questionnaire Short Form (IPAQ-SF) as a measure of physical activity (PA) in young people with cerebral palsy. *Physiotherapy*. <https://doi.org/10.1016/j.physio.2019.08.013>
- Lee, I., Paffenbarger, R. S., & Hsieh, C. (1991). Physical activity and risk of developing colorectal cancer among college alumni. *Journal of the National Cancer Institute*, 83, 1324-1329 <https://doi.org/10.1093/jnci/83.18.1324>
- Lee, P. H., Macfarlane, D. J., Lam, T. H., & Stewart, S. M. (2011). Validity of the international physical activity questionnaire short form (IPAQ-SF): A systematic review. *International Journal of Behavioral Nutrition and Physical Activity*, 8, 115. <https://doi.org/10.1186/1479-5868-8-115>
- Li, Y. N., Shapiro, B., Kim, J. C., Zhang, M., Porszasz, J., Bross, R., ... & Kopple, J. D. (2016). Association between quality of life and anxiety, depression, physical activity and physical performance in maintenance hemodialysis patients. *Chronic diseases and translational medicine*, 2(2), 110-119. <https://doi.org/10.1016/j.cdtm.2016.09.004>
- Lins, L., & Carvalho, F. M. (2016). SF-36 total score as a single measure of health-related quality of life: Scoping review. *SAGE Open Medicine*, 4, 1-12. <https://doi.org/10.1177/2050312116671725>
- Maher, C. A., Toohey, M., & Ferguson, M. (2016). Physical activity predicts quality of life and happiness in children and adolescents with cerebral palsy. *Disability and rehabilitation*, 38(9), 865-869. <https://doi.org/10.3109/09638288.2015.1066450>
- Martínez, M. E., Giovannucci, E., Spiegelman, D., Willett, W. C., Hunter, D. J., & Colditz, G. A. (1997). Leisure-time physical activity, body size, and colon cancer in women. *Journal of the National Cancer Institute*, 89(13), 948-955. <https://doi.org/10.1093/jnci/89.13.948>

- Matcham, F., Norton, S., Steer, S., & Hotopf, M. (2016). Usefulness of the SF-36 Health Survey in screening for depressive and anxiety disorders in rheumatoid arthritis. *BMC musculoskeletal disorders*, 17(1), 224. <https://doi.org/10.1186/s12891-016-1083-y>
- Ölçücü, B., Vatanserver, Ş., Özcan, G., Çelik, A. & Paktaş, Y. (2015). Üniversite öğrencilerinde fiziksel aktivite düzeyi ile depresyon ve anksiyete ilişkisi. *Uluslararası Türk Eğitim Bilimleri Dergisi*, 3(4), 294-303.
- Öztürk M. (2005). *Üniversitede Eğitim-Öğretim Gören Öğrencilerde Uluslararası Fiziksel Aktivite Anketinin Geçerliliği Ve Güvenirliği Ve Fiziksel Aktivite Düzeylerinin Belirlenmesi*. Yüksek Lisans Tezi. Hacettepe Üniversitesi Sağlık Bilimleri Enstitüsü.
- Özudođru, E. (2013). *Üniversite personelinin fiziksel aktivite düzeyi ile yaşam kalitesi arasındaki ilişkinin incelenmesi*. Yüksek Lisans Tezi. Mehmet Akif Ersoy Üniversitesi Eğitim Bilimleri Enstitüsü, Beden Eğitimi ve Spor Öğretimi Programı. Burdur.
- Pati, S., Swain, S., Patel, S. K., Chauhan, A. S., Panda, N., Mahapatra, P., & Pati, S. (2018). An assessment of health-related quality of life among patients with chronic obstructive pulmonary diseases attending a tertiary care hospital in Bhubaneswar City, India. *Journal of Family Medicine and Primary Care*, 7(5), 1047-1053. [https://doi.org/10.4103/jfmpc.jfmpc\\_37\\_18](https://doi.org/10.4103/jfmpc.jfmpc_37_18)
- Pekmezovic, T., Popovic, A., Tepavcevic, D. K., Gazibara, T., & Paunic, M. (2011). Factors associated with health-related quality of life among Belgrade University students. *Quality of life research*, 20(3), 391-397. <https://doi.org/10.1007/s11136-010-9754-x>
- Puciato, D., Borysiuk, Z., & Rozpara, M. (2017). Quality of life and physical activity in an older working-age population. *Clinical interventions in aging*, 12, 1627-1634. <https://doi.org/10.2147/CIA.S144045>
- Quehenberger, V., Cichocki, M., & Krajic, K. (2014). Sustainable effects of a low-threshold physical activity intervention on health-related quality of life in residential aged care. *Clinical interventions in aging*, 9, 1853-1864. <https://doi.org/10.2147/CIA.S70359>
- Sanda, B., Vistad, I., Haakstad, L. A. H., Berntsen, S., Sagedal, L. R., Lohne-Seiler, H., & Torstveit, M. K. (2017). Reliability and concurrent validity of the International Physical Activity Questionnaire short form among pregnant women. *BMC sports science, medicine and rehabilitation*, 9(1), 7. <https://doi.org/10.1186/s13102-017-0070-4>
- Savcı, S., Öztürk, M., Arıkan, H., İnal-İnce, D., & Tokgözođlu, L. (2006). Üniversite öğrencilerinin fiziksel aktivite düzeyi. *Türk Kardiyoloji Arşivi*, 34(3), 166-172.
- Saxena, S., Van Ommeren, M., Tang, K. C., & Armstrong, T. P. (2005). Mental health benefits of physical activity. *Journal of Mental Health*, 14(5), 445-451. <https://doi.org/10.1080/09638230500270776>
- Skrzek, A., Přidalová, M., Sebastjan, A., Harásková, D., Fugiel, J., Ignasiak, Z., ... & Rozek, K. (2015). Fine motor skills of the hands in Polish and Czech female senior citizens from different backgrounds. *Aging clinical and experimental research*, 27(4), 491-498. <https://doi.org/10.1007/s40520-014-0299-7>
- Stewart, A. L., & King, A. C. (1991). Evaluating the efficacy of physical activity for influencing quality-of-life outcomes in older adults. *Annals of Behavioral Medicine*, 13(3), 108-116.
- Teychenne, M., Ball, K., & Salmon, J. (2008). Physical activity and likelihood of depression in adults: A review. *Preventive medicine*, 46(5), 397-411. <https://doi.org/10.1016/j.ypmed.2008.01.009>
- Tierney, M., Fraser, A., & Kennedy, N. (2015). Criterion validity of the International Physical Activity Questionnaire Short Form (IPAQ-SF) for use in patients with rheumatoid arthritis: comparison with the SenseWear Armband. *Physiotherapy*, 101(2), 193-197. <https://doi.org/10.1016/j.physio.2014.07.005>
- Uçkun, A., & Çalıkođlu, A. S. (2003). Çocukluk çağında tip iki diyabet. *Sted*, 12(5), 174-179.
- Wannamethee, G., & Shaper, A. G. (1992). Physical activity and stroke in British middle-aged men. *British Medical Journal*, 304, 597-601. <https://doi.org/10.1136/bmj.304.6827.597>
- Yahia, N., Wang, D., Rapley, M., & Dey, R. (2016). Assessment of weight status, dietary habits and beliefs, physical activity, and nutritional knowledge among university students. *Perspectives in Public Health*, 136(4), 231-244. <https://doi.org/10.1177/1757913915609945>



**Copyrights**

Copyright for this article is retained by the author(s), with first publication rights granted to the journal.

This is an open-access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/4.0/>).