


Summer 2003

The Relationship of MMPI-A Item Effectiveness to Item Content, Diagnostic Category, and Classification Accuracy

Kathleen D. Lynch
Old Dominion University

Follow this and additional works at: https://digitalcommons.odu.edu/psychology_etds

 Part of the [Clinical Psychology Commons](#), and the [Quantitative Psychology Commons](#)

Recommended Citation

Lynch, Kathleen D.. "The Relationship of MMPI-A Item Effectiveness to Item Content, Diagnostic Category, and Classification Accuracy" (2003). Doctor of Psychology (PsyD), dissertation, Psychology, Old Dominion University, DOI: 10.25777/mwp3-hw58 https://digitalcommons.odu.edu/psychology_etds/205

This Dissertation is brought to you for free and open access by the Psychology at ODU Digital Commons. It has been accepted for inclusion in Psychology Theses & Dissertations by an authorized administrator of ODU Digital Commons. For more information, please contact digitalcommons@odu.edu.

**THE RELATIONSHIP OF MMPI-A ITEM EFFECTIVENESS TO ITEM
CONTENT, DIAGNOSTIC CATEGORY, AND CLASSIFICATION ACCURACY**

by

Kathleen D. Lynch
Virginia Consortium Program in Clinical Psychology

A Dissertation Submitted to the Faculties of

The College of William and Mary
Eastern Virginia Medical School
Norfolk State University
Old Dominion University

In Partial Fulfillment of the Requirement for the Degree of

DOCTOR OF PSYCHOLOGY

CLINICAL PSYCHOLOGY

VIRGINIA CONSORTIUM PROGRAM IN CLINICAL PSYCHOLOGY

August, 2003

Approved by:

Robert P. Archer, Ph.D. (Director)
Eastern Virginia Medical School

J.D. Ball, Ph.D. (Member)
Eastern Virginia Medical School

Richard W. Handel, Ph.D. (Member)
Eastern Virginia Medical School

Neill Watson, Ph.D. (Member)
College of William and Mary

Michelle Kelley, Ph.D. (Member)
Old Dominion University

ABSTRACT

THE RELATIONSHIP OF MMPI-A ITEM EFFECTIVENESS TO ITEM CONTENT, DIAGNOSTIC CATEGORY, AND CLASSIFICATION ACCURACY

Kathleen D. Lynch
Old Dominion University, 2000
Director: Robert P. Archer, Ph.D.

Archer, Handel, and Lynch (2001) recently compared the item endorsement frequencies for the MMPI-A normative sample against two adolescent clinical samples. Results showed that the MMPI-A contains a substantial number of items that do not show a significant difference in item endorsement frequency between normative and clinical samples. The current study extends Archer et al.'s (2001) research in three ways: 1) it examines the item endorsement frequencies of the Supplementary scales, Harris Lingoes subscales, and subtle-obvious items; 2) it examines the Basic, Content, and Supplementary scales, and Harris Lingoes subscales with two homogeneous diagnostic criterion groups (as suggested by Archer, Handel, and Lynch); and finally 3) it re-examines and recalculates Basic scale data using only those items that were shown by Archer, Handel, and Lynch (2001) to effectively discriminate between the normative and clinical populations. The mean profiles of the normative and clinical groups were contrasted based on these "revised" Basic scales using a newly acquired independent clinical sample to evaluate the extent to which profile sensitivity and specificity is affected by these scale modifications. Results demonstrated that examining the Supplementary scales and Harris-Lingoes subscales, or subtle-obvious items, or when extended to homogeneous criterion groups led to no improvement in item effectiveness from the results of Archer and his colleagues' study. However, results supported the

hypothesis in that the overall effectiveness of Basic scale discrimination increased, in terms of sensitivity, specificity, positive and negative predictive power, and hit rate, when the Basic clinical scale items were removed that did not discriminate between normative and clinical groups of adolescents.

ACKNOWLEDGMENTS

I would like to thank Bob Archer for his endless source of support, guidance and sense of humor. You are the ultimate mentor. I would also like to thank Rick Handel, who went above and beyond his duties as a committee member. Your constant involvement in this project was much appreciated. Thank you to my committee for their time and help throughout this process. Thanks to David Elkins for saving a “damsel in distress” more times than I can count. A million “thank you’s” to Denise Holloway for *everything!*

To my parents, thank you for all of your emotional and financial support, I honestly could not have done this without you. To my family, thank you for your patience and encouragement through this very long endeavor. Finally, thanks to my friends, who are constant reminders that there is a world outside of school and for helping me live it; especially, Jenny, Nicole, Katie, Jen and Jessie.

TABLE OF CONTENTS

	Page
LIST OF TABLES	ix
Chapter	
I. INTRODUCTION.....	1
Adolescent Development	3
Prevalence Data	5
Overview of the MMPI	8
History of the MMPI with Adolescents	9
Development of the MMPI-A	12
Administration and Scoring of the MMPI-A	15
Interpretation of the MMPI-A	17
Reliability and Validity	28
The Problem of Low <i>T</i> -scores	29
II. METHOD	36
Participants	36
Design and Procedure	38
Data Analysis.....	40
III. RESULTS.....	42
IV. DISCUSSION.....	79
REFERENCES.....	95
VITA.....	105

LIST OF TABLES

Table	Page
1. List of Basic Scales.....	21
2. List of Content and Supplementary Scales.....	24
3. List of Harris Lingoes Subscales.....	26
4. Percentage of MMPI-A Supplementary Scale Items Classified as Effective Based on Discrimination Performance Between the Normative Sample and Three Adolescent Clinical Samples.....	43
5. Percentage of MMPI-A Harris Lingoes Subscale Items Classified as Effective Based on Discrimination Performance Between the Normative Sample and Three Adolescent Clinical Samples	47
6. Percentage of MMPI-A Subtle-Obvious Items Classified as Effective Based on Discrimination Performance.....	52
7. Percentage of MMPI-A Basic Scale Items Classified as Effective Based on Discrimination Performance Between the Normative Sample and Three Adolescent Clinical Samples.....	54
8. Percentage of MMPI-A Content Scale Items Classified as Effective Based on Discrimination Performance Between the Normative Sample and Three Adolescent Clinical Samples.....	58
9. MMPI-A Ineffective Item Deletions with Standard MMPI-A Scale Membership	66
10. MMPI-A Validity and Basic Scale Item Totals for Original and Revised Scales .	74
11. MMPI-A Basic Scale ANOVA Results for Girls	75
12. MMPI-A Basic Scale ANOVA Results for Boys	76
13. Hit Rate, Sensitivity, Specificity, PPP, and NPP for Predicting Adolescent Membership in Normal (N= 1600) and Clinical (N= 163) Groups using Standard and Revised Scales at Two Cutting Scores by Gender	78

CHAPTER I

INTRODUCTION

The ability to assess psychological functioning is crucial to identifying adolescents with significant psychopathology, developing treatment plans, and evaluating treatment outcomes. A recent survey of practitioners has shown that the Minnesota Multiphasic Personality Inventory-Adolescent (MMPI-A) is currently the most widely used objective personality assessment instrument with adolescents (Archer & Newsom, 2000). The adolescent form of the original MMPI was developed after several researchers (e.g., Archer, 1992, 1987; Klinge, Lachar, Grisell, & Berman, 1978) documented that the original MMPI was producing extremely elevated profiles when administered to adolescents. While the development of the MMPI-A reduced or eliminated several problems associated with administering the original form of the MMPI to adolescents, the MMPI-A now has been found to produce a high frequency of Within Normal Limits Basic scale profiles for adolescents in clinical settings. Archer, Handel, and Lynch (2001) suggested that these within normal limit profiles may be produced, at least in part, because many MMPI-A items are not effective at discriminating between normative and clinical populations. These authors recently compared the item endorsement frequencies for the MMPI-A normative sample against two adolescent clinical samples. These adolescent sample results were also contrasted with item endorsement frequencies

The Publication Manual of the American Psychological Association (5th edition) was the model used for this manuscript.

obtained for the MMPI-2 normative samples and from a clinical adult psychiatric inpatient sample. Results showed that the MMPI-A contains a substantial number of items that do not show a significant difference in item endorsement frequency between normative and clinical samples. Further, the frequency of “ineffective” items was much higher for the MMPI-A in contrast to similar comparisons conducted with the MMPI-2.

It is unclear why there is a high frequency of non-discriminating items between the normative and clinical sample. One possibility is that the high frequency of non-discriminating items is due not to the characteristics of any particular clinical sample, but the high frequency of endorsement of MMPI-A items found for the normative sample. This latter hypothesis was supported in Archer et al.’s (2001) results that normal adolescents endorse many items with a higher frequency than do normal adults.

The current study extends the work of Archer, Handel, and Lynch (2001) by examining item-endorsement frequencies for the Harris-Lingoes subscales, Subtle-Obvious items, and homogeneous diagnostic criterion groups. These comparisons helped to further clarify the specific characteristics and content of items that do, and do not, discriminate in terms of the item endorsement frequencies found for normative and clinical samples of adolescents. Further, the effects of the deletion of non-discriminating items on classification accuracy of the MMPI-A Basic scales in discriminating clinical and normative samples was evaluated.

In order to evaluate adolescent response patterns on an objective personality assessment measure, it is important to first examine historical views on adolescence and how development effects psychological functioning. It is also necessary to understand the frequency with which adolescent psychopathology occurs, and how such pathology

deviates from normal development, in order to underscore the importance of accurate assessment, diagnosis, and treatment. In the following sections, adolescent development, adolescent psychopathology and an overview of the most widely used adolescent personality assessment instrument to date are examined.

Adolescent Development

Historically, normal adolescence has been viewed as a time of emotional instability, rapid shifts in mood, and conflictual relationships with authority figures (Erikson, 1956; Freud, 1958; Hall, 1916). G. Stanley Hall (1916), Anna Freud (1958) and others supported a “storm and stress” model of adolescence that suggested adolescents were likely to appear mildly, mentally ill due to the stress associated with attempting to manage physical, social, and psychological changes. Hall characterized adolescence as a time in which the young person is oscillating between contradictory tendencies. A typical adolescent, according to Hall, was emotionally labile, vacillating between euphoria and depression.

Anna Freud (1958) also shared the notion that adolescence was filled with emotional lability and periods of psychiatric illness. Freud believed that adolescents who did not display adjustment difficulties during this period were at risk to develop psychopathology later in life. Freud stated “adolescence is by its nature an interruption of peaceful growth, and...the upholding of a steady equilibrium during the adolescent process is in itself abnormal” (p. 275).

Erikson (1956) perceived adolescence as the time in which the individual moved toward identity formation. He posed the question, “how can a stage as ‘abnormal’ as adolescence be trusted to accomplish it [identity formation]?” (p.72). Erikson also

suggested that this period is a normal state of increased conflict in which neurosis and psychosis commonly occur. However, he noted that the neurosis and psychosis that ensue during adolescence are productive for the individual, unlike the neurosis and psychosis of mental illness that are seen later in life.

Although many have endorsed this psychopathological view of adolescence, others have suggested that adolescence is filled with peaceful and productive times. In a study of middle-class teenage boys, Bandura (1964) found that the adolescents in his sample were not typically in conflict with their parents over peer relations or household rules and that they typically had positive interactions with their parents. Bandura suggested that emotional lability and oppositional behaviors were an extension of pre-adolescent and childhood problems that were more easily managed in childhood. Additionally, Offer and Offer (1975) found transient episodes of mild depression and anxiety to be fairly common in their sample of adolescents; however, moderate to severe levels of psychopathology were reported in about 20% of their subjects. As will be discussed later, this percentage is consistent with findings from contemporary studies on the prevalence of adolescent psychopathology.

Ferdinand and Verhulst (1995) examined the stability of behavioral and emotional problems from adolescence to young adulthood and found psychological maladjustment to be a stable, rather than a transient phenomenon for adolescents in their sample. The authors evaluated 459 adolescents using the Child Behavior Checklist (CBC) completed by parents and eight years later with a young adult self-report (YAS). They found that those participants who exhibited significant levels of distress during adolescence (approximately 27% of their sample), also had scores in the deviant range as indicated by

the YAS at the eight-year follow up. Ferdinand and Verhulst concluded “the absence of psychopathology in adolescence, rather than the presence of behavioral or emotional problems or adolescent turmoil, should be regarded as normative” (p.1593).

Weiner and Del Gaudio (1976) offered three conclusions regarding adolescent psychopathology: (1) a significant level of psychological distress is not a normative feature of adolescence; (2) boundary lines between normal and abnormal levels of psychological distress during adolescence should be drawn with careful evaluation; (3) clinically significant psychopathology in an adolescent is unlikely to remit without intervention and will most likely continue into adulthood. Although it is important to understand that psychopathology is not a normal part of adolescence, it is still a significant problem in today’s society. The next section will attempt to explore just how prevalent psychopathology has become in children and adolescents.

Prevalence Data

Prevalence studies conducted in the past decade have suggested that the number of children and adolescents suffering from mental disorders ranges between 14 and 22 percent (Bradenburg, Friedman, & Silver, 1990; Gould, Wunsch-Hitzig, & Dohrenwend, 1981; McGee, Feehan, Williams, Partidge, Silva, & Kelly, 1990; NIMH, 1990; Rutter, Graham, Chadwick, & Yule, 1976). Although large methodological discrepancies exist between studies, to date, no investigations suggest that less than one in ten children or adolescents could be diagnosed with a mental disorder.

Gould, Wunsch-Hitzig, and Dohrenwend (1981) reviewed 25 studies conducted in the United States between 1928 and 1975 that examined the prevalence of clinical maladjustment in children and adolescents. They suggested that the rate for child and

adolescent psychiatric disturbance was probably not lower than 11.8%, which is considered a very conservative estimate. The authors stated that many of the studies they reviewed had serious methodological problems. These problems included single informant methodologies. The authors concluded that interviewing the adolescent directly could provide a more accurate estimate of adolescent psychopathology.

More recent studies have utilized more refined approaches to study the prevalence of adolescent psychopathology. Brandenburg, Friedman, and Silver (1990) used a multimethod, multistage model of identification which collected information from a variety of sources and measures. This model allowed for the identification of different levels of psychiatric disturbances. Brandenburg and colleagues examined eight studies which utilized this approach. The authors were then able to more accurately estimate the prevalence rate of child and adolescent psychopathology. They concluded that the prevalence rate was between 14% and 20%.

McGee and colleagues (1990) evaluated 943 fifteen-year-old adolescents using interviews and parental report measures. The authors found that 207 adolescents (22% of the sample) had at least one psychological disorder. The most prevalent problem was overanxious disorder, followed by non-aggressive conduct disorder, and simple phobia. Females had higher prevalence rates for all disorders with the exception of social phobia, attention deficit disorder, and aggressive conduct disorder.

In their study of 1710 high school students, Lewinsohn, Hops, Roberts, Seeley, and Andrews (1993) found that almost 10% of the adolescents met the criteria for a DSM-III-R disorder, and that more than 33% had experienced a psychological disorder at some point in their lifetime. Lewinsohn and colleagues used the Schedule for Affective

Disorders and Schizophrenia for School-Age Children to obtain these prevalence rates. Major depression had the highest lifetime prevalence rate. This was followed by anxiety disorders. As in the McGee and colleagues (1990) study, female respondents at all age levels had higher incidence rates for several disorders including unipolar depression, anxiety, eating disorders, and adjustment disorders. Male subjects had higher incidence rates of disruptive behavior disorder.

McDermott (1996) conducted a nationwide study of youths 5 to 17 years of age. The sample comprised the national normative sample of the Adjustment Scales for Children and Adolescents (ASCA). A subsample of 1400 subjects were randomly selected and then stratified to conform to U.S. Census proportions for a variety of demographic variables. McDermott examined age and gender prevalence for six core syndromes. A demarcation point of $T \geq 60$ was chosen to represent significant pathology on the ASCA, which reflected a 15% prevalence rate for the sample. Results indicated that for the hyperactive, aggressive-provocative, and aggressive-impulse syndromes, prevalence was significantly elevated among the youngest children, whereas the avoidant syndrome showed a higher incidence among the adolescent age group. For all but one syndrome, the percentage of boys exhibiting the syndrome was higher for boys than for girls.

More recently, Kessler, Avenevoli, and Merikangas (2001) reviewed the epidemiological literature on child and adolescent mood disorders. Their review suggests that major depression is common among adolescents, with up to a 25% lifetime prevalence by the end of adolescence. Kessler and colleagues state that retrospective data suggests 50% of adolescents experiencing depression and 90% of adolescents

experiencing mania will have reoccurrences in adulthood. The authors state that these reoccurrences are likely mediated by adverse role transitions (e.g., truncated educational achievement and teenage childbearing) that typically occur before the time of initial treatment.

Hyman (2001) reviewed evolving NIMH priorities for research on pediatric mood disorders. Ultimately, over the next decade, NIMH will be striving to understand the etiology of disorders such as depression and ADHD but also attempting to develop cures for these disorders, and most importantly, to prevent new onset of these disorders. Until a cure or preventative measures are developed, effective use of diagnostic assessment measures not only aids in designing treatment interventions for adolescents, but also serves to identify those adolescents at increased risk for several forms of psychiatric disorders during childhood. The next section attempts to explore the development of the MMPI-A, one of the most widely used diagnostic assessment tools for adolescents.

Overview of the MMPI

In the 1930's and 40's Stark Hathaway and J.C. McKinley were searching for a more efficient and reliable way to arrive at appropriate psychodiagnostic labels. They collected a large pool of potential inventory items from textbooks and earlier published scales. Then, appropriate diagnostic criterion groups were selected for patient populations. The Minnesota "normals" consisted of visitors and patients to the University of Minnesota Hospital. The clinical group was comprised of psychiatric patients at the University of Minnesota Hospital. Clinical participants were divided into subgroups determined by their clinical diagnosis. Only those with a clear diagnosis were included. The different subgroups formed were hypochondriasis, depression, hysteria, psychopathic

deviate, paranoia, psychasthenia, schizophrenia, and hypomania. The potential items were then administered to the Minnesota "normals" and the clinical groups. An item analysis was conducted for each of the clinical groups to identify which items differentiated between the clinical group and a group of "normal" persons. Those 504 items that were identified as discriminating between groups were retained. Using this criterion keying method, Hathaway and McKinley (1943) developed the original form of the MMPI. This form originally contained eight basic clinical scales that measured the presence of a variety of psychological disorders and traits and included hypochondriasis (scale 1), depression (scale 2), hysteria (scale 3), psychopathic deviant (scale 4), paranoia (scale 6), anxiety (scale 7), schizophrenia (scale 8), and mania (scale 9). Two more scales, the Masculinity-Femininity scale (scale 5) and the Social Introversion-Extroversion scale (scale 0) were added later and used to identify personal preferences and personality traits, rather than psychological disturbance. The MMPI also includes four validity scales which aid in the detection of invalid protocols due to aberrant or deceptive responding to test items. These include the Cannot Say scale (scale ?), which is the number of items the responder did not answer; the *L* or lie scale, which may be elevated when a responder consciously or unconsciously attempts to present oneself in a favorable manner; the *F* scale, which contains infrequently endorsed items; and the *K* scale, which measures the responders level of psychological defensiveness. Other scales such as the Supplementary, Content, and Harris-Lingoes (1955) scales and subscales have also been developed to allow for refinement of interpretation and research purposes.

History of the MMPI with Adolescents

While the original instrument was intended for use with individuals 16 years and

older, Dahlstrom, Welsh, and Dahlstrom (1972) stated that the MMPI could be used with "bright children as young as 12" (p.21). In fact, use of the MMPI with adolescents appears to have been more widespread than originally intended by Hathaway and McKinley. Both Capwell (1945a) and Monachesi (1948, 1950) found that the MMPI accurately discriminates between delinquent and non-delinquent adolescents based on the elevations of the Pd scale and that these differences were maintained over time (Capwell, 1945b).

Although the MMPI was used with adolescents, the first systematic effort aimed at collecting a large amount of data on adolescents using the MMPI was known as the "statewide sample" and was undertaken by Hathaway and Monachesi (1953, 1961, 1963). The statewide sample included approximately 15,000 Minnesota adolescents tested between 1948 and 1954. Participants also completed intelligence measures and a vocational interest inventory. Additionally, participants' teachers were asked to report on which students they thought were likely to have psychiatric or legal problems. A subsample was re-tested three years later in an effort to identify and predict longitudinally the personality variables related to the onset of delinquency. Monachesi and Hathaway (1969) stated that higher scores on the *Pd*, *Sc*, and *Ma* scales were predictive of higher rates of delinquent behavior in their sample of adolescents. Archer (1992, 1997) concluded that the Hathaway and Monachesi findings were important in several ways. Most importantly, the studies demonstrated that the MMPI could predict delinquent behavior. Additionally, the studies provided information regarding differences in item endorsements between males versus females, adults versus adolescents, and longitudinal test-retest differences in item endorsement for individuals in middle versus

late adolescence. Hathaway and Monachesi also provided clinical correlate data for their sample of adolescents for both high and low scores for each of the ten standard clinical scales. These early efforts provided the data for the development of two sets of adolescent norms for the MMPI.

Marks and Briggs (1972), Gottesman, Hanson, Kroeker, and Briggs (1987), and Colligan and Offord (1989) all developed adolescent norms for use with the original form of the MMPI. However the Marks and Briggs and the Gottesman norms included the Hathaway and Monachesi data set which was extremely dated and did not represent contemporary adolescent response patterns (Archer & Krishnamurthy, 1996).

Additionally, Archer (1987) found that the use of these norms produced an inflated number of false-negative MMPI profiles for adolescents evaluated in clinical settings, a problem that continues to be seen today, even with the revision of the instrument.

Although Colligan and Offord used a more contemporary sample of adolescents tested in the 1980's, their norms were criticized due to the geographically and ethnically restricted sample from which the norms were developed (Archer & Krishnamurthy, 1996).

Several studies have examined the differences that exist on an adolescents' profile when scored on each of the three adolescent normative sets developed for the MMPI. Some studies (Archer, Pancoast, & Klinefelter, 1989; Klinefelter, Pancoast, Archer, & Pruitt, 1990) indicate that when an adolescent's MMPI is scored on all three normative sets, large differences are observed in the profiles. The lowest *T*-scores are produced by the Colligan and Offord (1989) norms. In addition, major differences occur when attempting to derive code-types from the three scoring systems. Archer (1987) suggests that clinicians use caution in applying codetype descriptors developed on one set of

adolescent norms to an MMPI profile scored on a different set of adolescent norms.

Research also has indicated that when an adolescent's MMPI profile is scored on both adult and adolescent norms, the profile is more elevated overall when the adult norms are used (Archer, 1984; Ehrenworth & Archer, 1985; Klinge, Lachar, Grissell, & Berman, 1978; Klinge & Strauss, 1976; Lachar, Klinge, & Grissell, 1976; Marks, Seeman, & Haller, 1974). These Basic scale profiles have been particularly elevated on scales *F*, *4*, and *8*. It appears that both normative and clinical samples of adolescents more frequently endorse items that indicate the presence of psychopathology than do their adult counterparts. This results in profiles that overemphasize pathological features (Archer, 1984).

In addition to the aforementioned problems in using the MMPI with adolescents, several more obvious problems exist. Archer, Maruish, Imhof, and Piotrowski (1991) found that the length of the test, the relatively high reading level of test items, and the outdated and sometimes inappropriate language used in the test items were the most commonly cited problems clinicians mentioned related to the use of the MMPI with an adolescent population. Additionally, the content of many items was thought to be inappropriate for adolescents and did not address some of the specific issues adolescents face, such as drug use, school related problems, and eating disorders (Butcher et al., 1992). Due to the many problems in using the MMPI with an adolescent population, the University of Minnesota appointed a project committee to assess the need for an adolescent form of the MMPI.

Development of the MMPI-A

In 1989, the MMPI Adolescent Project Committee convened to develop an

adolescent form of the MMPI. The committee established goals to develop contemporary norms based on a national representative sample, shorten the length of the instrument, retain the standard validity and clinical scales to ensure continuity with the original test instrument, modify and improve existing scales, and standardize and refine assessment practices with adolescents (Butcher et al., 1992).

Form TX

An experimental test booklet (labeled MMPI Form TX) was created for the development of the MMPI-A (Butcher et al., 1992). This form consisted of the 550 original MMPI items and 154 new items. Approximately 13 percent of the original MMPI items were reworded or modified to eliminate awkward wording and outdated expressions or sexist language. Also, the 16 repeated items from the MMPI were dropped. Modifications that occurred in the MMPI-A did not result in significant changes in response patterns for these items (Archer & Gordon, 1994). Form TX was used in both adolescent normative and clinical data collection along with a biographical information form and a Life Events form. These were administered in order to obtain data on family structure, parental occupation, residence, and family history. The Life Events form was developed to assess stressful events that had occurred in the six months prior to the testing situation.

Sample

The normative sample of the MMPI-A included junior high and high school students from California, Minnesota, New York, North Carolina, Ohio, Pennsylvania, Virginia, and Washington (Butcher, et al., 1992). These eight states were chosen to obtain a wide range of diversity in geographical location, ethnic background, and rural-urban

residence. Students were each paid ten dollars for their participation (with the exception of those students from New York) and data collection was conducted in the schools (with groups of 5 to 100 students). Two criteria were developed in order to ensure that only complete and valid data were included in the normative sample. First, only participants who completed all three measures were included in the sample. Second, if the participant left more than 35 items blank, or they had an *F* scale raw score greater than 25, the data from that participant was not included in the sample. The final MMPI-A normative sample was comprised of 1620 participants (805 boys and 815 girls).

The clinical sample included a smaller and less ethnically diverse sample of 420 boys and 293 girls from a variety of treatment settings around the Minneapolis area. The settings included were inpatient alcohol and drug treatment units, inpatient mental health facilities, day-treatment programs, and a special school program. These subjects ranged in age from 14 to 18 years, and were currently enrolled in school in grades 7 through 12. These subjects completed all three measures, Form TX, the Biographical Information form, and the Life Events form. Additionally, data on the clinical sample of adolescents were collected using the Child Behavior Checklist (Achenbach & Edelbrock, 1983), the Devereux Adolescent Behavior Rating Scale (Spivack, Haimes, & Spotts, 1967), and a review of hospital records. These additional measures provided behavioral ratings from parents and treatment staff that supplemented self-report data in the creation of scale descriptors.

The current form of the MMPI-A has been in use for almost a decade. Today, the MMPI-A is one of the most widely used assessment instruments with adolescents (Archer & Newsom, 2000). Due to the frequency with which the MMPI-A is administered, it is

important that the test be re-examined to ensure that the test is providing an accurate diagnostic picture for clinicians.

Administration and Scoring of the MMPI-A

Qualifications of Test Users

As with any testing instrument, it is important that the individual administering the test should have sufficient background in test theory and test construction, as well as an understanding of the instrument itself. Although the MMPI-A manual states that the MMPI-A may be administered by carefully trained support staff under close supervision, the responsibility of an appropriate administration ultimately lies with the clinician.

Archer (1992) recommends that at a minimum the test user should have completed graduate-level coursework in psychological testing, adolescent development, personality, psychopathology, and psychodiagnosis.

Age Criteria

Although the MMPI-A was essentially created for use with individuals' ages 14 to 18 years, it may be selectively administered to 12- and 13- year-old adolescents who have the cognitive and social maturity that would enable them to read and understand the items (Archer, 1992). Adolescents who are 18 years of age are eligible to take either the MMPI-2 or the MMPI-A. Eighteen-year-old high school students who are still living at home and are still dependent on parents should be administered the MMPI-A. An 18-year-old individual who is living independently (e.g., married or holds a full-time job) should be administered the MMPI-2.

Reading Requirements

It is essential that the test-taker have an adequate level of reading comprehension

in order to take the MMPI-A. The majority of the items were determined by Butcher and colleagues (1992) to be at the fifth- to seventh-grade reading level. Audiotaped versions of the MMPI-A are available for adolescents who have literacy problems. Adolescents with limited intelligence, reading disorders, or speak English as a second language are examples of individuals who may have difficulties taking the MMPI-A.

Testing Environment

The test-taker should have adequate privacy and supervision during the testing session. Examiners may respond to questions about the meaning of words using a dictionary definition or clarifying idioms. It is important for the test-taker to have a quiet environment free from distractions with comfortable seating and adequate lighting. Additionally, breaks from testing are permitted for adolescents who become fatigued during testing (Butcher et al., 1992).

Materials

The MMPI-A is available in a booklet, audiocassette, and computer version. In the booklet format, items are presented in either a hard or soft cover, reusable, 478-item test booklet and a separate answer sheet. Subjects fill in the circle marked "T" if the item is true or mostly true and "F" if the item is false or mostly false. The audiocassette version can be used in either the individual or group setting (Butcher et al., 1992). The audiocassette version may be useful for the visually impaired, as well as for those adolescents with significant reading difficulties. Individuals can also take the MMPI-A on a computer using software provided through National Computer Systems (NCS).

Scoring

Answer sheets should be carefully examined upon completion for items left

unanswered and items endorsed in both the true and false direction (Archer, 1992). If there are a substantial number of items left blank, the individual may be asked to complete as many of those items as possible. Raw scores for all validity, clinical, and Supplementary scales are obtained through the use of scoring templates or a computer-scoring program. Butcher et al. (1992) note that raw score values are then converted to either linear *T*-scores (scales *VRIN*, *TRIN*, *F1*, *F2*, *F*, *L*, *K*, *5*, *0*, *MAC-R*, *ACK*, *PRO*, *IMM*, *R*, and *A*) or uniform *T*-scores (scales *1*, *2*, *3*, *4*, *6*, *7*, *8*, *9*, and all 15 Content scales).

Interpretation of the MMPI-A

Assessment of the Validity of Individual Protocols

The MMPI-A contains scales that provide information regarding the test-takers tendency to answer the items in an accurate and consistent manner: Cannot Say (?), *L* (Lie), *F* (Frequency), *K* (Defensiveness), *VRIN* (Variable Response Inconsistency), and *TRIN* (True Response Inconsistency).

Cannot say (?). The Cannot Say (?) scale consists of the total number of items that have been omitted or that have been answered in both the true and false direction. A profile is considered invalid with more than 30 items omitted, and therefore uninterpretable. Archer (1997) suggests that adolescents who omit more than 30 items be asked to complete the remaining items or to retake the entire test if their reading level is consistent with this task.

The L (Lie) scale. The MMPI-A *L* scale consists of 14-items designed to detect naïve attempts of individuals to portray themselves in a favorable or overly moralistic light. Individuals endorsing items on this scale are denying minor flaws or weaknesses.

MMPI-A profiles with elevated *L* scale scores ($T \geq 65$) suggest a conscious or unconscious effort to appear moral, ethical, and without common foibles.

The F (Frequency) scale and the F1 and F2 subscales. The *F* scale consists of 66 items that were endorsed in the deviant direction by no more than 20% of the MMPI-A normative sample. Those individuals with high scores on this scale are endorsing a wide variety of strange and unusual experiences either due to the presence of psychopathology, a random response style, or a response style in which the individual is exaggerating symptoms. The *F1* subscale consists of the first 33 items that make up the *F* scale, and the *F2* scale consists of the second 33 *F* scale items. The MMPI-A manual (Butcher et al., 1992) suggests that comparison of the *T*-score values for these two subscales can provide information on the consistency of the individual's responding throughout the test. However, recent data by Archer, Handel, Lynch, and Elkins (2002) indicate that comparison of *F1* and *F2* subscale *T*-score values is of limited effectiveness in detecting random response patterns in the latter half of the test booklet.

The K (Defensiveness) scale. The *K* scale attempts to identify individuals who display a significant degree of psychopathology, but produce profiles within normal limits. This 30-item scale can be considered a measure of defensiveness against psychopathology. In the MMPI-2, there is a *K*-correction procedure which was not carried over to the MMPI-A because *K*-weights developed for the MMPI-A were not effective at improving classification accuracy for adolescent subgroups when compared to the standard non-*K* corrected norms (Alperin, Archer, & Coates, 1996). MMPI-A profiles with elevated *T*-scores (≥ 65) may represent a defensive test-taking attitude in the adolescent.

The VRIN (Variable Response Inconsistency) and TRIN (True Response Inconsistency) scales. These scales indicate the tendency of an individual to respond to items in an inconsistent or contradictory manner. The *VRIN* scale is made up of pairs of items that are either similar or opposite in content. For each pair of items that is answered inconsistently, the test-taker receives a point on the *VRIN* scale. The *TRIN* scale is made up of pairs that are opposite in content only. One point is added to the *TRIN* scale raw score if the test-taker answers true to both items in a particular pair. One point is subtracted from the *TRIN* scale raw score if the test-taker answers false to both items in the pair. Therefore, a very high raw score on this scale represents a tendency to answer indiscriminately true to the items and a very low score indicates a tendency to answer indiscriminately false to the items. The *VRIN* scale is helpful in identifying inconsistent profiles that represent either acquiescent or nonacquiescent response patterns. Archer (1992) suggests that profiles containing *VRIN* or *TRIN* *T*-scores ≥ 80 should be considered invalid.

If the profile is invalid, Greene (2000) suggests that the MMPI-A be readministered if the clinician determines that the profile is unacceptable due to inconsistency of responding. However, Greene suggests that the instrument not be readministered if it is deemed invalid due to accuracy problems. When a profile is deemed reliable, the clinician should proceed with interpretation of the clinical scales.

The Basic Clinical Scales

The MMPI contains 10 Basic clinical scales, as well as Content, Supplementary, and a variety of other special scales that may provide interpretive information in regard to the MMPI-A respondent (see Table 1). In most cases, interpretations are limited to

clinical scales with *T*-score elevations ≥ 65 (Butcher et al., 1992). Scales with *T*-score elevations between 60 and 64, that fall into what test developers call the “shaded” or gray zone”, are considered moderately elevated, making interpretations more difficult. Butcher and colleagues (1992) suggest examiners should exercise caution when interpreting scales with *T*-scores in this range.

Code Type Interpretation

Using the highest pattern of elevation on clinical scales, codetypes can be developed to classify profiles. Although there is not yet any clinical codetype correlate data available from studies using the MMPI-A beyond single scale correlates, Archer (1992, 1997) provided codetype correlate descriptions for 29 code types based on studies of adolescent populations using the MMPI.

Table 1

List of Basic Scales

Scale Number	Abbreviation	Scale Name
Scale 1	<i>Hs</i>	Hypochondriasis
Scale 2	<i>D</i>	Depression
Scale 3	<i>Hy</i>	Hysteria
Scale 4	<i>Pd</i>	Psychopathic Deviate
Scale 5	<i>Mf</i>	Masculinity-Femininity
Scale 6	<i>Pa</i>	Paranoia
Scale 7	<i>Pt</i>	Psychasthenia
Scale 8	<i>Sc</i>	Schizophrenia
Scale 9	<i>Ma</i>	Hypomania
Scale 0	<i>Si</i>	Social Introversion

Content and Supplementary Scales

The Content scales were created by adapting those MMPI-2 Content scales that were appropriate for adaptation to the MMPI-A. Next, items were added or deleted to improve psychometric properties. Eleven of the 15 Content MMPI-A scales overlap with the MMPI-2 and four scales are unique to the MMPI-A (see Table 2): Alienation (*A-aln*), Low Aspiration (*A-las*), School Problems (*A-sch*), and Conduct Problems (*A-con*). Uniform *T*-score transformation procedures are used when converting MMPI-A Content scale raw score totals to *T*-score values. The Content scales appear to have relatively high internal reliability with alpha coefficient values ranging from .55 to .83 (Archer, 1997). Additionally, Content scales can only be interpreted when all 478-items of the MMPI-A are administered. Content scales can be considered helpful in supplementing the information received from Basic scales but should not be used independently of the Basic scales.

There are six Supplementary scales, three of which were taken from the MMPI-2 (MacAndrew Alcoholism scale, *MAC-R*; Welsh's Anxiety and Repression scales, *A*, and *R*, respectively) and three new scales (see Table 2). The Immaturity (*IMM*) scale, is a scale measuring immaturity as it relates to interpersonal style, cognitive complexity, self-awareness, and a variety of other factors (Archer, Pancoast, & Gordon, 1994). Items for this scale were chosen using a multistage procedure using both rational and statistical criteria. The Alcohol-Drug Problem (*ACK*) scale was developed to assess an adolescent's willingness to acknowledge alcohol or drug use-related symptoms. The 13 items were initially selected based on the rational judgement that item content was related to drug use, and then later refined based on statistical criteria. The Alcohol-Drug Proneness

(*PRO*) scale consists of 36 items that were empirically selected based on item endorsement differences found between adolescents in alcohol and drug treatment programs and adolescents in inpatient psychiatric facilities (Archer, 1992; Weed, Butcher, & Williams, 1994). In contrast to the Content scales, the raw scores for all Supplementary scales are converted to *T*-score values based on linear *T*-score transformation procedures. Consistent with the Content scales, all 478- items must be administered and the results from the Supplementary scales should be used to supplement and refine interpretations of the MMPI-A Basic scales.

Table 2

List of Content and Supplementary Scales

Content Scales	Supplementary Scales
Anxiety (<i>A-anx</i>)	Anxiety (<i>A</i>)
Obsessiveness (<i>A-obs</i>)	Repression (<i>R</i>)
Depression (<i>A-dep</i>)	MacAndrew Alcoholism (<i>MAC-R</i>)
Health Concerns (<i>A-hea</i>)	Alcohol/Drug Problem Proneness (<i>PRO</i>)
Bizarre Mentation (<i>A-biz</i>)	Alcohol/Drug Problem Acknowledgment (<i>ACK</i>)
Anger (<i>A-ang</i>)	Immaturity (<i>IMM</i>)
Cynicism (<i>A-cyn</i>)	
Alienation (<i>A-aln</i>)	
Conduct Problems (<i>A-con</i>)	
Low Self-Esteem (<i>A-lse</i>)	
Low Aspirations (<i>A-las</i>)	
Social Discomfort (<i>A-sod</i>)	
Family Problems (<i>A-fam</i>)	
School Problems (<i>A-sch</i>)	
Negative Treatment Indicators (<i>A-trt</i>)	

Harris Lingoies and Si Subscales

Six of the Basic clinical scales have subscales that allow for a more specific focus in item content areas. Harris and Lingoies (1955) created subscales for these six MMPI clinical scales (2,3,4,6,8,9) which were carried over to the MMPI-A (see Table 3). The scales were developed by examining the content of items within a clinical scale and grouping items together which appeared to reflect a similar trait. Archer (1997) cautions that the subscales should only be used to supplement or refine interpretation due to the lack of validity data on these measures. Additionally, there are three Scale 0 subscales created for the MMPI and carried over to the MMPI-A (Ben-Porath, Hostetler, Butcher, & Graham, 1989) to help differentiate the nature of social isolation.

Table 3

List of Harris Lingoes and Si Subscales

Harris Lingoes Scales		Si Subscales	
<i>D1</i>	Subjective Depression	<i>Si1</i>	Shyness/Self-Consciousness
<i>D2</i>	Psychomotor Retardation	<i>Si2</i>	Social Avoidance
<i>D3</i>	Physical Malfunctioning	<i>Si3</i>	Self/Other Alienation
<i>D4</i>	Mental Dullness		
<i>D5</i>	Brooding		
<i>Hy1</i>	Denial of Social Anxiety		
<i>Hy2</i>	Need for Affection		
<i>Hy3</i>	Lassitude-Malaise		
<i>Hy4</i>	Somatic Complaints		
<i>Hy5</i>	Inhibition of Aggression		
<i>Pd1</i>	Familial Discord		
<i>Pd2</i>	Authority Problems		
<i>Pd3</i>	Social Imperturbability		
<i>Pd4</i>	Social Alienation		
<i>Pd5</i>	Self-alienation		
<i>Pa1</i>	Poignancy		
<i>Pa3</i>	Naiveté		

Table 3 (continued)

List of Harris Lingoes and Si Subscales

Harris Lingoes Scales

- Sc1* Social Alienation
- Sc2* Emotional Alienation
- Sc3* Lack of Ego Mastery, Cognitive
- Sc4* Lack of Ego Mastery, Conative
- Sc5* Lack of Ego Mastery, Defective Inhibition
- Sc6* Bizarre Sensory Experiences
- Ma1* Amorality
- Ma2* Psychomotor Acceleration
- Ma3* Imperturbability
- Ma4* Ego Inflation
-

Structural Summary

Archer and Krishnamurthy (1994) created the Structural Summary approach to the MMPI-A interpretation based on factor-analytic research conducted by Archer, Belevich, and Elkins (1994). The Structural Summary identifies eight scale-level factors in the MMPI-A: General Maladjustment, Immaturity, Disinhibition/Excitatory Potential, Social Discomfort, Health Concerns, Naiveté, Familial Alienation, and Psychoticism. The Structural Summary provides an organized approach to viewing the information provided by the 69 Clinical, Content, Supplementary scales and subscales.

Reliability and Validity

The MMPI-A manual reports one-week, test-retest reliability correlations obtained from a subsample of normative subjects (Butcher et al., 1992). The clinical scales appear to have an adequate range of test-retest correlations (.65 to .84). However, the coefficients for the validity scores are lower (.49 to .75). The internal consistency coefficients (coefficient alphas) for the MMPI-A Basic validity and clinical scales are below .80 for both boys and girls for nine of the 15 scales in the normative sample, and ten of the 15 scales for the clinical sample. Although Black (1994) states that most of these coefficients in the "low to moderate" range should be considered insufficient, the MMPI-A manual states that most of the coefficients for the MMPI-A are in the low to moderate range but that some scales demonstrated strong internal consistencies despite the fact that most of the scales were derived through empirical item selection. The MMPI Content scales, however, have generally higher alpha coefficients than their Basic scale counterparts because of the development method used for these latter scales (Butcher et al., 1992). The alpha coefficients for the Content scales fall mostly in the .70 to .80 range.

Test-retest coefficients for the Content scales range from .62 to .82.

Factor analysis of the MMPI-A Basic clinical scales was conducted by Butcher and colleagues (1992) using the normative sample; It indicated that a four-factor solution was appropriate for both boys and girls: (a) a general anxiety or maladjustment factor marked by high loadings on most scales; (b) an overcontrol or repression factor marked by high loadings on scales *L*, *K*, and *9*; (c) a third factor largely defined by scale *0*; and (d) a fourth factor defined almost entirely by scale *5*. Expanding on this work, Archer, Belevich, and Elkin's (1994) identified eight factors in an analysis of the 69 scales and subscales of the MMPI-A.

Studies evaluating the concurrent validity of the MMPI-A have found moderate relationships between the MMPI-A and other self-report measures. However, there is still limited information regarding concurrent validity of the MMPI-A. Krishnamurthy, Archer, and House (1996) found that there were limited relationships between Rorschach variables and MMPI-A variables. The two instruments should be viewed as providing different types of information and not to confirm or disconfirm the data provided in each measure.

The Problem of Low T-Scores

As previously discussed, adolescent respondents on the MMPI endorse significantly more unusual items on the MMPI than do their adult counterparts, particularly items that are scored on Scales *F*, *4*, and *8* (Archer, 1984; Klinge & Strauss, 1976; Marks, Seeman, & Haller, 1974). Because of the high rate of endorsement of items on scale *F* and *8*, it is difficult to construct a critical item list for adolescents (Archer & Jacobson, 1993). It is still not fully understood why adolescents tend to more frequently

endorse unusual items.

Some researchers (Archer, 1992, 1997) suggest that the phenomenon is related to the reading level required for the MMPI and the MMPI-A. If an adolescent is unable to read at the sixth or seventh grade level, they may provide inaccurate, and therefore invalid MMPI profiles due to an inability to accurately read and comprehend MMPI items. A second explanation for the overall higher *T*-score patterns seen in adolescents may be due to the nature of adolescence as a stormy time of life (Archer, 1984). In contrast to adults, adolescents tend to report more unusual symptoms that are consistent with the views of adolescence as a turbulent stage of development (Hall, 1916; Freud, 1958). Finally, adolescents may tend to be more honest or candid than adults when responding to items on the MMPI, thereby elevating clinical scale *T*-scores. Butcher et al. (1992) have found that adolescents typically endorse fewer *K* scale items than do adults, suggesting they may be less defensive when responding to items.

MMPI researchers have attempted to address the problem of elevated adolescent MMPI profiles (when scored on adult norms) by developing adolescent norm sets for the MMPI (Colligan & Offord, 1989; Gottesman et al., 1987; Marks & Briggs, 1972), and later, the MMPI-A (Butcher et al., 1992). However, by interpreting adolescent responses scored on adolescent norms, the profiles then produce subclinical elevations, even for adolescent inpatient psychiatric settings. Ehrenworth and Archer (1985) then recommended using a *T*-score value of ≥ 65 for clinical range elevations when interpreting adolescent MMPI profiles on adolescent norms. This aided in reducing the frequency of within-normal-limits profiles and increased the sensitivity in accurately identifying profiles by normative versus clinical samples of adolescents (Archer, 1987).

When the MMPI-A was published, the authors decided to use a range of *T*-score values, or a “shaded” zone, in place of a demarcation line value that separates clinical-range from normal-range elevations. *T*-scores < 60 are classified in the normal-range of elevation, and *T*-scores ≥ 65 are clearly in the clinical-range elevation. Butcher et al. (1992) suggest that *T*-scores falling in the “shaded” zone between 60 and 65 are considered marginally elevated, and should be interpreted with caution. Archer (1997) suggests that scores in this range reflect adolescents who may exhibit some, but not all, of the characteristics measured by a particular scale. Additionally, Alperin, Archer, and Coates (1996) examined the possibility that the development of an MMPI-A *K*-correction factor could increase test sensitivity. However, their results did not show any systematic improvements in MMPI-A accuracy in identifying normative and clinical samples when *K*-correction factors were used. Finally, Fontaine, Archer, Elkins and Johansen (2001) indicated that a reduction in the *T*-score criterion used to define a “clinical range elevation” did not produce a more effective balance between test sensitivity and specificity (i.e., that lowering the clinical range elevation *T*-score criterion from 65 to 60 did not improve classification performance).

Regardless of the steps that have been taken to improve the problem of low *T*-scores, the MMPI-A continues to produce a high frequency of within normal limits Basic scale profiles for individuals with substantial psychopathology. In fact, 30.4% of boys and 29.1% of girls receiving mental health services had profiles containing no clinical scale *T*-score values ≥ 65 (Archer, 1997). As stated previously, Archer (1984, 1987) suggested that this problem might be related to the observation that many normal adolescents experience psychological turbulence and distress making it difficult to

distinguish between normative and clinical populations. If this were accurate, one would expect to find fewer differences in the frequency of endorsement of MMPI-A items between groups of normative and clinical adolescents than found in similar comparisons for adults on the MMPI-2. Archer and Jacobson (1993) examined the item endorsement frequencies for the Koss-Butcher (1973) critical items in the MMPI-2 normative sample and in a clinical comparison group of 232 male and 191 female adult inpatients.

Additionally, Archer and Jacobson's study examined the item endorsement frequencies for the same Koss-Butcher items for the MMPI-A normative sample and the clinical sample of 420 boys and 293 girls reported in the MMPI-A manual. Results indicated that most critical items showed large frequency differences between normative and clinical adult samples but that the critical item endorsement frequencies often were equivalent for normative and clinical samples of adolescents.

Using a similar methodology, Archer, Handel, and Lynch (2001) compared the item endorsement frequencies for the MMPI-A normative sample against two adolescent clinical samples and contrasted these results with the item endorsement frequencies for the MMPI-2 normative sample and a clinical sample of adult psychiatric inpatients. Of the two MMPI-A clinical samples, the first consisted of those adolescents recruited from Minnesota treatment facilities. The findings for this group were presented in the MMPI-A manual (Butcher et al., 1992). The second clinical sample consisted of adolescents evaluated upon admission to several North Carolina inpatient psychiatric facilities.

The frequency of "true" responses was calculated for each of the items that appear in the MMPI Basic clinical scales and in the eleven Content scales held commonly between the two MMPI forms, the Harris-Lingoes and the Supplementary scales were not

examined in this study. The item endorsement frequencies for the MMPI-2 and the MMPI-A were examined separately. An item was considered to show a significantly higher rate of endorsement in the clinical group if the difference in endorsement frequency between the clinical group and the normative sample was at least twice the standard error of the proportions of true responses of the two groups under examination. The test for the significance of difference between two independent proportions is expressed as a Z-score, with an absolute value of ≥ 2 indicating that the percentage difference was twice or greater the standard error of the independent proportions. This was the criterion employed by Hathaway and McKinley (1943) (as noted by Greene, 2000) in their original selection of items of the MMPI Basic scales. This procedure was utilized by Hathaway and McKinley to select the initial items for scale membership, and items were selected based on the additional requirements that the criterion group response frequency was at least 10% for the selected item.

Results from the above study showed that the MMPI-A contains a substantial number of items that do not show a significant difference in item endorsement frequency between normative and clinical samples. Further, MMPI-A Basic and Content scales generally showed a much lower percentage of effective items than did the corresponding scales for the MMPI-2. Out of the 567 items that appear on the MMPI-2, 83% met the criterion as effective items for both men and women. However for the clinical sample reported in the MMPI-A manual, only 56% of the 478 MMPI-A items met the criterion required as effective for boys and only 49% of total items met the effectiveness criterion for girls. In the independent clinical sample, only 54% met the criterion as effective for boys and only 45% of items were classified as effective for girls.

The percentage of effective items was higher for the MMPI-2 comparisons for nine of the ten Basic scales, with similar levels of effectiveness for the *Pd* scale items across adolescent and adult comparisons. In fact, the *Pd* scale was the only MMPI-A Basic scale for which the majority of items were considered to meet the criterion to be classified as effective.

The Content scales for the MMPI-2 also appeared to have a higher frequency of effective items than the MMPI-A. For the MMPI-A Content scales, Family Problems (*A-fam*) was the only MMPI-A Content scale for which the majority of items were classified as effective in comparisons for both samples. The lowest percentage of effective items occurred for the MMPI-A Social Discomfort (*A-sod*) scale.

It is suggested that the high frequency of non-discriminating items between adolescent normative and clinical groups does not reflect the unique characteristics of the clinical adolescent samples, but the high frequency of endorsement of MMPI-A items found for the normative sample. These findings also suggest that the item content of scales created by empirical keying methodologies with adult populations may not be directly generalizable to adolescent respondents. The authors suggest that an investigation of MMPI-A item effectiveness in a variety of separate diagnostic groups would provide more specific information regarding item discrimination. The authors also observed that if the majority of items do not effectively discriminate between normative and clinical populations, it is possible that the 478-item form could be reduced to only those items that do effectively discriminate between normative and clinical populations.

The current study is designed to further investigate the issue of MMPI-A item endorsement frequency between normative and clinical populations. Specifically, this

study has three purposes: 1) To extend the Archer, Handel and Lynch (2001) study by examining the item endorsement frequencies of the Supplementary scales, Harris Lingo's subscales, and subtle-obvious items; 2) To examine the Basic, Content, and Supplementary scales, and Harris Lingo's subscales with two homogeneous diagnostic criterion groups (as suggested by Archer, Handel, and Lynch), and finally; 3) To re-examine and recalculate Basic scale data using only those items that prove to discriminate effectively between the normative and clinical populations. The mean profile of the normative and clinical groups was contrasted based on the "revised" Basic scales with an independent clinical sample to evaluate the extent to which profile sensitivity and specificity is affected by these scale modifications. The hypotheses are as follows:

1) The frequency of effective items would not be significantly related to membership on specific Supplementary and Harris Lingo's scales. The frequency of item effectiveness in the Basic scales would be substantially higher among obvious items in contrast to subtle items.

2) The specific content of non-discriminating items would vary as a function of the diagnostic groups used in contrast to normative groups, (i.e., the frequency of effective items would increase when item content and diagnostic group are related.)

3) The overall rate of discrimination would increase, in terms of sensitivity, when the Basic clinical scale items are removed that did not discriminate between normative and clinical groups of adolescents, (i.e., when the Basic scales are revised based only on responses to retained items and Basic scale clinical profiles recalculated for normative and clinical groups.).

CHAPTER II

METHOD

Participants

Participants for this study consisted of norms for a group of normative subjects, and a clinical sample as reported in the MMPI-A manual, and two samples of adolescents evaluated with the MMPI-A while receiving clinical services. The MMPI-A normative sample was used as the control group in this study, and consisted of 1620 adolescents (805 males and 815 females), between grades 7 and 12, inclusive. Participants ranged in age from 14 through 18, inclusive, mean age of 15.5 ($SD= 1.17$) for boys and 15.6 ($SD= 1.10$) for girls. The MMPI-A normative sample is described in detail in Archer (1997) and in the MMPI-A manual by Butcher et al. (1992). The MMPI-A normative sample will be referred to as the normative sample. The MMPI-A manual clinical sample consisted of 420 boys and 293 girls, aged 14 through 18, with grade placements of 7 through 12, inclusive. This clinical sample was recruited from treatment facilities in the Minneapolis area, including inpatient alcohol and drug treatment centers ($N= 299$ boys, 163 girls), inpatient mental health facilities ($N= 67$ boys, 96 girls), day-treatment programs ($N= 13$ boys, 24 girls), and a special school program ($N= 41$ boys, 10 girls). Diagnoses are not available for this clinical sample. This sample is referred to as the manual-clinical sample.

The first additional clinical sample for the current study used to test the diagnostic homogeneity hypothesis were selected from 271 boys and 160 girls between the ages of 14 through 18, inclusive, who were evaluated with the MMPI-A upon their admission to adolescent units in inpatient psychiatric treatment facilities in North Carolina ($N=90$),

Texas ($N=202$), and Virginia ($N=139$). This clinical sample will be designated as the inpatient clinical sample. The diagnostic breakdown (based on DSM-III-R and DSM-IV criteria) for the combined inpatient sample, based on principle diagnosis as provided in the admission medical record was as follows: Conduct Disorder = 192 (44.5%), Dysthymic disorders = 63 (14.6%); missing = 42 (9.8%); Major Depression = 34 (7.9%); other = 30 (6.9%); Depressive disorder NOS = 16 (3.7%); Disruptive Behavior disorder = 15 (3.5%); Oppositional Defiant Disorder = 13 (3.0%); Adjustment disorders = 12 (2.8%); Bipolar disorders = 10 (2.0%); and Sexual disorders = 4 (.9%). The first diagnostic subsample from this inpatient sample are those participants diagnosed with conduct disorder and are referred to as the conduct disorder sample. The second diagnostic subsample from this inpatient sample are comprised of those participants diagnosed with depression and dysthymia and are referred to as the depression sample.

The second clinical sample in this study, used to calculate classification characteristics for standard and revised Basic MMPI-A scales, consisted of 90 boys and 73 girls between grades 7 and 12 inclusive, who received outpatient evaluations or outpatient therapy services at the Hampton-Newport News Community Services Board (CSB). Participants ranged in age from 13 to 18, inclusive. The mean age was 15.7 ($SD=1.29$) for boys and 15.7 ($SD=1.07$) for girls. The diagnostic breakdown (based on DSM-IV criteria) was Depressive Disorders ($N=14$), Conduct Disorder/Oppositional Defiant Disorder ($N=21$), Bipolar I Disorder ($N=3$), Substance Abuse ($N=2$), Adjustment Disorder ($N=5$), Attention Deficit/Hyperactivity Disorder ($N=5$), Learning Disorders ($N=2$), Child Abuse/Neglect ($N=4$), Missing ($N=107$). This data collection project is not yet completed; therefore, many of the diagnoses are missing for this data set. This sample

is referred to as the independent outpatient clinical sample.

Design and Procedure

The first purpose of this study was to evaluate the item discrimination effectiveness of MMPI-A items when examined in relation to the Supplementary scales, Harris Lingoes subscales, and subtle versus obvious Basic scale items for the normative versus manual clinical sample. Differences in item endorsement frequencies were evaluated by using the Test for the Significance of Difference between two independent proportions expressed as a Z-score, with an absolute value of ≥ 2 indicating that the percentage difference was twice or greater the standard error of the independent proportions. This criterion was used to define a significant difference in item endorsement, and as noted earlier, is consistent with the primary criterion employed by Hathaway and McKinley (1940) in their selection of the original items for the MMPI Basic scales. The subtle and obvious components of the MMPI-A Basic scales were identified by consulting the MMPI-2 (Butcher et al., 1989, 2001) manual for the item composition of subtle and obvious components of the Basic scales, and then extrapolating this information to all of those Basic scales items retained within the MMPI-A Basic scale structure.

A second purpose of the present study was to evaluate the frequency of item effectiveness for Basic, Content, and Supplementary scales and Harris Lingoes subscales in relation to membership in two homogeneous diagnostic groups. Participants for specific diagnostic groups were selected from the first clinical sample based on their membership in the conduct disorders diagnostic group ($N=192$) and in the depression group ($N=113$), that consisted of the combined dysthymic disorders ($N=63$), major

depression ($N=34$), and depressive disorders NOS ($N=16$) subgroups.

The third and final purpose of this study was to examine the rate of classification accuracy on an independent outpatient clinical sample after those items previously determined as ineffective were removed from Basic scale membership. Items were considered ineffective, and subsequently removed from the scales, if they met the “ineffective” criteria (Z score absolute value of < 2) for the manual clinical sample and an independent sample used in Archer et al. (2001). Scales 2, 4, and 9 were required to meet additional criteria in that they also had to be ineffective for the homogeneous criterion group that corresponds to the scale (i.e., scale 2- Depression; scales 4 and 9, Conduct Disorder). The uniform T -scores utilized to evaluate the revised Basic scales, that is, the Basic scale compositions following the deletion of non-discriminating items, was based upon the derivation of Uniform T -score values from the MMPI-A normative sample, calculated separately by gender. A software package for the derivation of Uniform T -scores for the MMPI-2 Scales (Hoeglund & Tellegen, 1998) was utilized for this purpose. Uniform T -scores were developed to provide a raw score to a T -score transformation procedure that produced relatively comparable percentile values across scales for a given T -score value (Hoeglund & Tellegen, 1998). The Uniform T -score transformation process consisted of three stages. The goal of the first stage was to derive from raw score frequencies the corresponding linear T -scores for each scale. The second stage was to derive for each scale the regression weights and intercept values for the Uniform T -score equation ($UT = B_0 + B_1X + B_2D^2 + B_3D^3$). Finally, the third stage involved the development of a table with all possible raw scores and the corresponding Uniform T -scores for each raw score value.

Data Analysis

The test for significance of difference between two independent proportions was used as means of evaluating item endorsement frequency differences. This test is expressed as a Z-score, and a criterion of an absolute value of ≥ 2 was utilized to signify a significant difference in endorsement frequencies. Two-tailed Chi square tests, with Yates correction, were used in order to evaluate differences between endorsement of subtle versus obvious items; gender, homogeneous versus manual clinical samples, and Harris Lingoes subscales versus Basic scales. The Yates continuity correction is designed to yield more accurate chi-square approximations under conditions in which expected frequencies are limited and there is only one degree of freedom. However, Howell (1992) argues that Chi square approximations with limited frequencies are not completely accurate with or without a chi square approximation. To evaluate *T*-score differences for the original versus revised scales on the independent clinical sample, a series of univariate ANOVA's were performed. To protect against the inflation of the alpha level that occurs as a product of conducting multiple tests on related data sources, a MANOVA was used to evaluate the overall group effect on the basic scales. Further, due to the number of tests conducted, alpha level was also adjusted to $p < .01$ for subsequent univariate tests for individual scales.

Finally, overall hit-rate, sensitivity, specificity, positive predictive power and negative predictive power was used to evaluate the effects of the deletion of non-discriminating items from the Basic scales on the level of accurate prediction of adolescents status in either the normative or clinical groups. Hit rate, in the current study, is the ability of the Basic scale profile to accurately identify true positives and true

negatives (calculated by adding the true positives and the true negatives and dividing by the total number of subjects). Sensitivity is the percentage of participants whose protocols indicate the presence of clinical status who in fact were from the clinical sample. Specificity is the percentage of participants whose within normal limits protocols accurately indicate the absence of pathology because, in fact, they were from the normative group. Positive predictive power is defined as the probability that an elevated score is being produced by a protocol from the clinical sample. Negative predictive power is the probability that a within normal limits protocol was produced by a participant in the normative group.

CHAPTER III

RESULTS

Results will be examined in three parts. First, the endorsement frequencies for the manual's normative and clinical sample are presented for the Supplementary scales, Harris Lingoes subscales, and subtle-obvious items. Next, the endorsement frequencies for the homogeneous criterion groups (i.e., depression and conduct disorder samples) are examined on the Basic, Content, and Supplementary scales and Harris Lingoes subscales. Finally, overall hit-rate, sensitivity, specificity, positive predictive power and negative predictive power are examined for the revised Basic scales in contrast to the standard MMPI-A Basic scales for the independent clinical sample.

The first objective of this study was to extend Archer, Handel, and Lynch's (2001) study by examining the item endorsement frequencies of the Supplementary scales, the Harris Lingoes subscales, and subtle versus obvious Basic scale items for the normative and the manual clinical sample. There are six Supplementary scales on the MMPI-A (*MAC-R*, *ACK*, *PRO*, *IMM*, *A* and *R*). As shown in Table 4, the right hand columns illustrate the percentage of effective items by gender for the normative versus manual clinical sample. There were no significant gender differences in item endorsement frequency on any of the Supplementary scales. *MAC-R*, *ACK*, and *PRO* scales all contained more than 60% effective items (except *MAC-R* for girls, 49%) where as *IMM*, *A*, and *R*, all contained less than 50% effective items.

Table 4

Percentage of MMPI-A Supplementary Scale Items Classified as Effective Based on Discrimination Performance Between the Normative Sample and Two Adolescent Clinical Samples

	MMPI-A Percent Effective Items Normative/ Depression Sample		MMPI-A Percent Effective Items Normative/Conduct Disorder Sample		MMPI-A Percent Effective Items Normative/Manual Clinical Sample	
	Boys	Girls	Boys	Girls	Boys	Girls
<i>MAC-R</i>	27% (13/49)	27% (13/49)	53% (26/49)	31% (15/49)	65% (32/49)	49% (24/49)
<i>ACK</i>	31% (4/13)	46% (6/13)	69% (9/13)	31% (4/13)	62% (8/13)	62% (8/13)
<i>PRO</i>	36% (13/36)	56% (20/36)	58% (21/36)	42% (15/36)	94% (34/36)	89% (32/36)
<i>IMM</i>	26% (11/43)	42% (18/43)	35% (15/43)	30% (13/43)	44% (19/43)	44% (19/43)
<i>A</i>	26% (9/35)	34% (12/35)	29% (10/35)	03% (1/35)	20% (7/35)	20% (7/35)
<i>R</i>	12% (4/33)	18% (6/33)	9% (3/33)	12% (4/33)	09% (3/33)	21% (7/33)

Table 4 (continued)

Note. Percent figures reflect the total percentage of items within each scale that showed significant differences in item endorsement frequencies between the normative and clinical sample, i.e., items classified as "effective" in terms of discrimination between samples. The first number within each parentheses indicates the number of "effective" items within the scale, and the second number reflects the total number of items in the scale.

There are 28 Harris Lingo (HL) subscales for seven of the ten Basic scales. As shown in Table 5, the right hand columns illustrate the percentage of effective items using the normative versus manual clinical sample by gender for each of the Harris Lingo subscales. There were no significant gender differences in item endorsement frequency for any of the HL subscales. On the HL *D* subscale, the percentage of effective items was highest on *D4* (Mental Dullness) for girls (67%, 10/15 effective items) and for boys (80%, 12/15). The percentage of effective items was lowest on *D2* (Psychomotor Retardation) for girls (7%, 1/14 items) and for boys (21% 3/14). Of the HL *Hy* subscales, only *Hy3* (Lassitude-Malaise) for girls had more than 50% of the items classified as effective. *Hy4* (Somatic Complaints) had only one effective item out of 17 for both boys and girls. The HL *Pd* subscales contained the highest percentage of effective items of all the Harris Lingo subscales; all of the *Pd* subscales had more than half of their items classified as effective. On HL *Pa*, only *Pa1* (Persecutory ideas) was above 50% effectiveness and only for girls. None of the 9 items on *Pa3* (Naivete) distinguished between the normative and clinical samples. Similarly, on HL *Sc*, only one subscale (*Sc2*, Emotional Alienation) was above 50 % item effectiveness, and that result was found only for girls. There were no items on either *Sc5* (Lack of Ego Mastery, Defective Inhibition) or *Sc6* (Bizarre Sensory Experiences) that distinguished between the normative and clinical samples. For HL *Ma*, only *Ma1* (Amorality) had more than 50% classified as effective and this was only for boys. All the other *Ma* scales were below 50% effectiveness.

Comparing the HL subscales against their corresponding parents scales, none performed significantly better than the parent scales. However, on two occasions (*Hy3*

($\chi^2(1, N=60) = 7.59, p < .01$) and *Sc6* ($\chi^2(1, N=77) = 6.26, p < .01$), the parent scales performed significantly better than the corresponding HL subscale.

Table 5

Percentage of MMPI-A Harris Lingoes Subscale Items Classified as Effective Based on Discrimination Performance Between the Normative Sample and Three Adolescent Clinical Samples

Scales	MMPI-A Percent Effective Items Normative/ Depressed Sample		MMPI-A Percent Effective Items Normative/Conduct Disorder Sample		MMPI-A Percent Effective Items Normative/Manual Clinical Sample	
	Boys	Girls	Boys	Girls	Boys	Girls
<i>D1</i>	50% (16/32)	63% (20/32)	44% (14/32)	19% (6/32)	34% (11/32)	50% (16/32)
<i>D2</i>	14% (2/14)	21% (3/14)	21% (3/14)	07% (1/14)	21% (3/14)	07% (1/14)
<i>D3</i>	64% (7/11)	64% (7/11)	46% (5/11)	09% (1/11)	27% (3/11)	46% (5/11)
<i>D4</i>	67% (10/15)	80% (12/15)	47% (7/15)	20% (3/15)	33% (5/15)	67% (10/15)
<i>D5</i>	50% (5/10)	50% (5/10)	05% (5/10)	20% (2/10)	50% (5/10)	50% (5/10)
<i>Hy1</i>	0% (0/6)	0% (0/6)	33% (2/6)	33% (2/6)	50% (3/6)	33% (2/6)
<i>Hy2</i>	0% (0/12)	08% (1/12)	42% (5/12)	17% (2/12)	25% (3/12)	08% (1/12)

Table 5 (continued)

Scales	MMPI-A Percent Effective Items Normative/ Depressed Sample		MMPI-A Percent Effective Items Normative/Conduct Disorder Sample		MMPI-A Percent Effective Items Normative/Manual Clinical Sample	
	Boys	Girls	Boys	Girls	Boys	Girls
<i>Hy3</i>	80% (12/15)	73% (11/15)	53% (8/15)	20% (3/15)	40% (6/15)	67% (10/15)
<i>Hy4</i>	35% (6/17)	53% (9/17)	06% (1/17)	0% (0/17)	06% (1/17)	06% (1/17)
<i>Hy5</i>	0% (0/7)	0% (0/7)	14% (1/7)	0% (0/7)	0% (0/7)	29% (2/7)
<i>Pd1</i>	56% (5/9)	33% (3/9)	67% (6/9)	11% (1/9)	78% (7/9)	78% (7/9)
<i>Pd2</i>	50% (4/8)	63% (5/8)	75% (6/8)	88% (7/8)	75% (6/8)	88% (7/8)
<i>Pd3</i>	0% (0/6)	17% (1/6)	33% (2/6)	33% (2/6)	50% (3/6)	50% (3/6)
<i>Pd4</i>	42% (5/12)	58% (7/12)	67% (8/12)	42% (5/12)	58% (7/12)	75% (9/12)
<i>Pd5</i>	58% (7/12)	67% (8/12)	92% (11/12)	50% (6/12)	75% (9/12)	83% (10/12)
<i>Pa1</i>	41% (7/17)	59% (10/17)	59% (10/17)	41% (7/17)	41% (7/17)	53% (9/17)

Table 5 (continued)

Scales	MMPI-A Percent Effective Items Normative/ Depressed Sample		MMPI-A Percent Effective Items Normative/Conduct Disorder Sample		MMPI-A Percent Effective Items Normative/Manual Clinical Sample	
	Boys	Girls	Boys	Girls	Boys	Girls
<i>Pa2</i>	22% (2/9)	33% (3/9)	33% (3/9)	0% (0/9)	33% (3/9)	33% (3/9)
<i>Pa3</i>	0% (0/9)	0% (0/9)	0% (0/9)	0% (0/9)	0% (0/9)	0% (0/9)
<i>Sc1</i>	33% (7/21)	48% (10/21)	48% (10/21)	19% (4/21)	38% (8/21)	19% (4/21)
<i>Sc2</i>	46% (5/11)	73% (8/11)	46% (5/11)	27% (3/11)	27% (3/11)	55% (6/11)
<i>Sc3</i>	50% (5/10)	60% (6/10)	50% (5/10)	10% (1/10)	10% (1/10)	30% (3/10)
<i>Sc4</i>	64% (9/14)	57% (8/14)	43% (6/14)	14% (2/14)	21% (3/14)	29% (4/14)
<i>Sc5</i>	27% (3/11)	09% (1/11)	18% (2/11)	0% (0/11)	0% (0/11)	0% (0/11)
<i>Sc6</i>	30% (6/20)	15% (3/20)	15% (3/20)	0% (0/20)	0% (0/20)	0% (0/20)
<i>Ma1</i>	0% (0/6)	0% (0/6)	100% (6/6)	67% (4/6)	83% (5/6)	33% (2/6)
<i>Ma2</i>	09% (1/11)	0% (0/11)	09% (1/11)	0% (0/11)	27% (3/11)	0% (0/11)

Table 5 (continued)

Scales	MMPI-A Percent Effective Items Normative/ Depressed Sample		MMPI-A Percent Effective Items Normative/Conduct Disorder Sample		MMPI-A Percent Effective Items Normative/Manual Clinical Sample	
<i>Ma3</i>	25% (2/8)	25% (2/8)	38% (3/8)	13% (1/8)	38% (3/8)	25% (2/8)
<i>Ma4</i>	0% (0/9)	33% (3/9)	44% (4/9)	22% (2/9)	22% (2/9)	11% (1/9)

Note. Percentage figures reflect the total percentage of items within each scale that showed significant differences in item endorsement frequencies between the normative and clinical sample, (i.e., items classified as “effective” in terms of discrimination between samples). The first number within each parentheses indicates the number of “effective” items within the scale; the second number reflects the total number of items in the scale.

Although the MMPI-A does not have specific subtle-obvious scales (Butcher et al., 1992), each item on the Weiner-Harmon Subtle-Obvious Subscales used from the MMPI-2 was matched with a corresponding MMPI-A item. There were no significant differences between subtle and obvious item endorsement effectiveness for any of the five scales as evaluated in Chi-square analyses (see Table 6).

Table 6

Percentage of MMPI-A Subtle-Obvious Items Classified as Effective Based on
Discrimination Performance

Scale	Subtle Items	Obvious Items	χ^2 Value
<i>D</i>	78% (14/18)	79% (31/39)	.02
<i>Hy</i>	44% (12/27)	64% (21/33)	1.50
<i>Pd</i>	68% (15/22)	96% (26/27)	5.11
<i>Pa</i>	35% (6/17)	74% (17/23)	4.49
<i>Ma</i>	83% (19/23)	61% (14/23)	1.72

Note. All Chi square values produced alpha levels $p > .05$.

The second purpose of this study was to evaluate the frequency of item effectiveness for the Basic, Content, Supplementary scales, and Harris Lingoes subscales in relation to two homogeneous diagnostic criterion groups (i.e., depression and conduct disorder). Results will be examined under the heading of each diagnosis.

Depression sample

The results found for the frequency of effective items produced in comparisons of the normative and depressed clinical sample for Basic scales are shown in the left hand side of Table 7. For the Depression scale, there were no significant differences in frequency of effective items between the depression sample and the manual clinical sample for both boys and girls. As shown in the left column of Table 7, there were only two scales that demonstrated over 50% effectiveness, *Hs* and *Pd* (both results restricted to girls). Not surprisingly, given its classification as a “non-clinical” scale, *Mf* performed the most poorly (7% effective items), with only 3 of 44 items effectively distinguishing between the normative sample and the depression sample for either gender.

Table 7

Percentage of MMPI-A Basic Scale Items Classified as Effective Based on Discrimination Performance Between the Normative Sample and Three Adolescent Clinical Samples

Scale	MMPI-A Percent Effective Items Normative/ Depression Sample		MMPI-A Percent Effective Items Normative/Conduct Disorder Sample		MMPI-A Percent Effective Items Normative/Manual Clinical Sample	
	Boys	Girls	Boys	Girls	Boys	Girls
<i>Hs (1)</i>	31% (10/32)	53% (17/32)	13% (4/32)	06% (2/32)	03% (1/32)	13% (4/32)
<i>D (2)</i>	39% (22/57)	47% (27/57)	58% (33/57)	16% (9/57)	28% (16/57)	40% (23/57)
<i>Hy (3)</i>	30% (18/60)	35% (21/60)	22% (13/60)	10% (6/60)	25% (15/60)	28% (15/60)
<i>Pd (4)</i>	43% (21/49)	53% (26/49)	63% (31/49)	41% (20/49)	63% (31/49)	71% (35/49)
<i>Mf (5)</i>	07% (3/44)	07% (3/44)	11% (5/44)	07% (3/44)	16% (7/44)	25% (11/44)
<i>Pa (6)</i>	23% (9/40)	35% (14/40)	38% (15/40)	22% (9/40)	30% (12/40)	33% (13/40)
<i>Pt (7)</i>	31% (15/48)	40% (19/48)	31% (15/48)	08% (4/48)	23% (11/48)	29% (14/48)

Table 7 (continued)

Scale	MMPI-A Percent Effective Items Normative/ Depression Sample		MMPI-A Percent Effective Items Normative/Conduct Disorder Sample		MMPI-A Percent Effective Items Normative/Manual Clinical Sample	
	Boys	Girls	Boys	Girls	Boys	Girls
<i>Sc (8)</i>	38% (29/77)	40% (31/77)	31% (24/77)	14% (11/77)	18% (14/77)	30% (23/77)
<i>Ma (9)</i>	13% (06/46)	15% (7/46)	44% (20/46)	22% (10/46)	35% (16/46)	17% (8/46)
<i>Si (0)</i>	10% (6/62)	15% (9/62)	24% (15/62)	05% (3/62)	11% (7/62)	19% (12/62)

Note. Percent figures reflect the total percentage of items within each scale that showed significant differences in item endorsement frequencies between the normative and clinical sample, i.e., items classified as “effective” in terms of discrimination between samples. The first number within each parentheses indicates the number of “effective” items within the scale, and the second number reflects the total number of items in the scale.

There were no significant differences in item endorsement frequency between boys and girls in the normative and the depression sample on any of the 10 scales. There was a significant difference between the boys depressed group and the boys from the manual clinical sample for *Sc* ($\chi^2(1, N = 77) = 6.32, p < .01$). There were more items in the depression sample that correctly discriminated between the normative and clinical population than for the manual clinical sample. Additionally, on *Hs*, there was a significant difference in the frequency of effective items between the depression group and the manual clinical sample for both boys ($\chi^2(1, N = 32) = 7.02, p < .01$) and girls ($\chi^2(1, N = 32) = 10.20, p < .01$). There were significantly more items that were able to effectively discriminate between the normative and the depression sample in these comparisons.

The data related to the item effectiveness found in normative versus the depression clinical sample for the Content scales is shown in the left hand side of Table 8. There were no significant differences in item endorsement frequency between boys and girls in the depression sample and the normative sample for any of the Content scales, as evaluated by Chi square analyses. The *A-dep* (Depression) was the only Content scale that had 50% or more effective items for both boys and girls, however, it did not differ significantly from the effective item frequency found for the manual clinical sample. *A-ang* (Anger) was the only Content scale that showed a significant difference in frequency of item effectiveness between the depression group and the manual clinical sample (only for boys) ($\chi^2(1, N = 17) = 17.17, p < .01$). There were significantly more effective items produced in the comparisons for boys between the

normative and manual clinical sample, in contrast to the normative-depressed sample comparisons.

Table 8

Percentage of MMPI-A Content Scale Items Classified as Effective Based on Discrimination Performance Between the Normative Sample and Three Adolescent Clinical Samples.

Scale	MMPI-A Percent Effective Items Normative/ Depression Sample		MMPI-A Percent Effective Items Normative/Conduct Disorder Sample		MMPI-A Percent Effective Items Normative/Manual Clinical Sample	
	Boys	Girls	Boys	Girls	Boys	Girls
<i>A-anx</i>	33% (7/21)	52% (11/21)	38% (8/21)	14% (3/21)	29% (6/21)	38% (8/21)
<i>A-obs</i>	07% (1/15)	13% (2/15)	20% (3/15)	07% (1/15)	20% (3/15)	20% (3/15)
<i>A-dep</i>	50% (13/26)	62% (16/26)	50% (13/26)	23% (6/26)	50% (13/26)	50% (13/26)
<i>A-hea</i>	27% (10/37)	41% (15/37)	14% (5/37)	08% (3/37)	08% (3/37)	14% (5/37)
<i>A-aln</i>	25% (5/20)	45% (9/20)	25% (5/20)	10% (2/20)	Not Available	Not Available
<i>A-biz</i>	16% (3/19)	21% (4/19)	26% (5/19)	11% (2/19)	11% (2/19)	21% (4/19)
<i>A-ang</i>	06% (1/17)	41% (7/17)	35% (6/17)	18% (3/17)	82% (14/17)	41% (7/17)

Table 8 (continued)

Scale	MMPI-A Percent Effective Items Normative/ Depression Sample		MMPI-A Percent Effective Items Normative/Conduct Disorder Sample		MMPI-A Percent Effective Items Normative/Manual Clinical Sample	
	Boys	Girls	Boys	Girls	Boys	Girls
<i>A-cyn</i>	09% (2/22)	23% (5/22)	68% (15/22)	14% (3/22)	32% (7/22)	27% (6/22)
<i>A-con</i>	17% (4/23)	30% (7/23)	70% (16/23)	35% (8/23)	Not Available	Not Available
<i>A-lse</i>	39% (7/18)	56% (10/18)	22% (4/18)	0% (0/18)	28% (5/18)	44% (8/18)
<i>A-las</i>	13% (2/16)	38% (6/16)	19% (3/16)	13% (2/16)	Not Available	Not Available
<i>A-sod</i>	13% (3/24)	17% (4/24)	25% (6/24)	0% (0/24)	4% (1/24)	4% (1/24)
<i>A-fam</i>	43% (15/35)	57% (20/35)	46% (16/35)	29% (10/35)	69% (24/35)	66% (23/35)
<i>A-sch</i>	40% (8/20)	65% (13/20)	55% (11/20)	40% (8/20)	Not Available	Not Available
<i>A-trt</i>	15% (4/26)	27% (7/26)	35% (9/26)	04% (1/26)	23% (6/26)	31% (8/26)

Note. Percent figures reflect the total percentage of items within each scale that showed significant differences in item endorsement frequencies between the normative and clinical sample, (i.e., items classified as "effective" in terms of discrimination between

samples). The first number within each parentheses indicates the number of “effective” items within the scale; the second number reflects the total number of items in the scale.

The results found for the frequency of effective items produced in comparisons of the normative and depressed clinical sample for Supplementary scales are shown in the left hand columns of Table 4. There were no significant gender differences, as evaluated by Chi Square analyses, in item endorsement frequency within the depression group for any of the Supplementary scales. However, on *PRO* Supplementary scale, there were significant differences in frequency of effective items for both boys ($\chi^2 (1, N= 36) = 24.51, p < .01$) and girls ($\chi^2 (1, N= 36) = 8.37, p < .01$) between the depression group and the manual clinical sample. In both comparisons, there were higher frequencies of effective items in the normative versus the manual clinical sample comparison. There were also significant differences for boys between the depression group and the manual clinical sample for *MAC-R* ($\chi^2 (1, N= 49) = 13.31, p < .01$). There was a significantly higher frequency of effective items in the manual clinical versus normative sample comparison, in contrast to the depression versus normative sample comparison.

The findings generated by the comparisons of frequency of effective items in the normative versus depressed sample for the Harris Lingoes subscales are shown in the left columns of Table 5. There were no significant gender differences in the frequency of effective items for the depression group versus the normative sample comparisons on any of the Harris Lingoes subscales. There was a significant difference between the depression group and the manual clinical sample in frequency of item effectiveness for girls on *Hy4* ($\chi^2 (1, N= 17) = 6.94, p < .01$). There were more effective items in normative sample versus the depression sample comparisons than were found for comparisons between manual clinical and normative samples.

Conduct Disorder sample

The frequency of effective items found in normative versus conduct disorder sample comparisons for the Basic scales is shown in the middle columns of Table 7.

There were significant differences in item endorsement frequency between boys and girls on 3 of the 10 scales, D (χ^2 (1, $N=57$) = 19.94, $p < .01$), Pt (χ^2 (1, $N=48$) = 6.65, $p < .01$), and Si (χ^2 (1, $N=46$) = 7.86, $p < .01$). Specifically, on all three scales, the frequency of occurrence of items that effectively discriminated between normative and conduct disorder samples was higher for boys than girls.

There were significant differences in frequency of effective items on D between the conduct disorder sample and the manual clinical sample for both boys (χ^2 (1, $N=57$) = 9.16, $p < .01$) and girls (χ^2 (1, $N=57$) = 7.34, $p < .01$), however these differences were in opposite directions. Comparisons between the normative and conduct disorder samples for boys show a significantly higher frequency of effective items than comparisons for the normative and the manual clinical samples. Comparisons between the normative and manual clinical samples for girls show a significantly higher frequency of effective items than comparisons for the normative and the conduct disorder samples. There was also a significant difference in frequency of effective items on scale Pd for girls, between the conduct disorder sample and the manual clinical sample (χ^2 (1, $N=49$) = 8.12, $p < .01$). There were more effective items in the manual clinical sample versus the normative sample comparisons than were found between the conduct disorder and the normative sample comparisons.

The frequency of effective items found in normative versus conduct disorder sample comparisons for the Content scales is shown in the middle column of Table 8.

There were significant differences in the frequency of effective items found for comparisons between boys and girls on *A-cyn* ($\chi^2 (1, N= 22) = 11.37, p < .01$) and *A-trt* ($\chi^2 (1, N= 26) = 6.06, p < .01$). The frequency of effective items produced by comparisons of the normative and the conduct disorder clinical sample was higher for boys than for girls on both *A-cyn* and *A-trt*. There were also significant differences in the frequency of effective items between the girls in the conduct disorder sample and girls in the manual clinical sample for *A-lse* ($\chi^2 (1, N= 18) = 7.87, p < .01$) and *A-fam* ($\chi^2 (1, N= 35) = 8.25, p < .01$). There were more effective items in the manual clinical sample versus the normative sample comparisons than were found for comparisons between the conduct disorder and the normative samples.

The frequency of effective items found in normative versus conduct disorder sample comparisons for the Supplementary scales is shown in the middle column of Table 4. There was a significant gender difference in the frequency of effective items for the Welsh's Anxiety (*A*) scale ($\chi^2 (1, N= 35) = 6.90, p < .01$). Boys showed a higher frequency of items that effectively discriminated between normative and conduct disorder samples than girls. There was also a significant difference for the *PRO* scale between the frequency of effective items produced in comparisons of the normative sample and the conduct disorder sample, versus the normative sample and manual clinical sample. This difference was significant for both boys ($\chi^2 (1, N= 36) = 11.08, p < .01$) and girls ($\chi^2 (1, N= 36) = 15.68, p < .01$). A significantly higher frequency of effective items was produced by the normative-manual clinical sample comparisons (in contrast to the normative-conduct disorder clinical sample).

The frequency of effective items found in normative versus conduct disorder sample comparisons for the Harris Lingoes subscales is shown in the middle column of Table 5. There were no significant gender differences in the frequency of occurrence of items that effectively discriminated between the normative and conduct disorder samples. When comparing the frequency of effective items between the conduct disorder sample and the manual clinical sample, there were no significant differences for both genders.

The third purpose of this study was to evaluate the predictive accuracy of revised MMPI-A Basic scales, constructed by deleting ineffective items from scale membership and recalculating gender specific Uniform *T*-scores for raw score values produced by the revised and shortened scales on an independent clinical sample. Items were considered ineffective, and subsequently removed from the scales, if they met the “ineffective” criteria (*Z* score absolute value of < 2) for the manual clinical sample and an independent sample used in Archer et al. (2001). Scales 2, 4, and 9 were required to meet additional criteria in that they also had to be ineffective for the homogeneous criterion group that corresponds to the scale (i.e., scale 2- Depression; scales 4 and 9, Conduct Disorder). Table 9 presents the list of deleted items and their scale membership. Additionally, Table 10 presents MMPI-A validity and Basic scale item totals for the original and revised scales. MANOVAs were also performed separately by gender and both were found to be significant at the .01 level (Girls: $F(8, 137) = 5.14, p < .001$ and Boys: $F(8, 171) = 2.72, p < .01$). ANOVAs were performed separately for each scale by gender to assess for significance of difference between the standard and revised scales. For girls, all eight scales showed a significant difference ($p < .01$) between standard and revised scales (see

Table 11). For boys, five of the eight scales showed a significant difference ($p < .01$) between standard and revised scales (see Table 12).

Table 9

MMPI-A Ineffective Item Deletions with Standard MMPI-A Scale Membership

Item	Item Content	Scale membership
3	I wake up fresh and rested most mornings.	1, 3, 7, <i>dep</i>
4	I seldom worry about my health.	2, 7
7	I like to read newspaper articles on crime.	<i>MAC-R</i> , 3, <i>R</i>
8	My hands and feet are usually warm enough.	1, 3
14	I work under a great deal of tension.	2, 9, <i>anx</i>
15	Once in a while I think about things too bad to talk about.	6, 7, 8, <i>L</i>
21	At times I have fits of laughing and crying that I cannot control.	6, 7, 8, 9
23	I feel that it is certainly best to keep my mouth shut when I am in trouble.	3, 5
34	At times I feel like smashing things.	2, <i>K</i> , <i>R</i> , <i>ang</i>
*35	I have periods of days, weeks, or months where I couldn't take care of things because I couldn't get going.	2, 7, 8, <i>dep</i> , <i>A</i>
41	Once a week or oftener I suddenly feel hot all over for no real reason.	3, 8, <i>hea</i>
*42	I am in just as good physical health as most of my friends.	1, 2, 3, <i>hea</i> , <i>R</i>
*43	I prefer to pass people I know but have not seen for a long time, unless they speak to me first.	2, 8, <i>sod</i>

Table 9 (Continued)

Item	Item Content	Scale membership
44	I am almost never bothered by pains over my heart or in my chest.	1, 3, <i>hea</i>
45	Most anytime I would rather sit and daydream than do anything else.	8, <i>IMM</i>
46	I am a very sociable person.	2, 0, <i>sod</i> , <i>MAC-R</i>
47	I have often had to take orders from someone who did not know as much as I did.	9, <i>cyn</i>
50	Parts of my body often feel like they are burning, tingling, or "going to sleep."	1, <i>hea</i>
52	I sometimes keep on at a thing until others lose their patience with me.	2, 9, <i>obs</i>
58	I am an important person.	9, <i>lse</i>
67	I am easily downed in an argument.	4, 0, <i>lse</i>
77	I think most people would lie to get ahead.	3, 6, <i>cyn</i>
81	At times I have a strong urge to do something harmful or shocking.	8, 9, <i>ACK</i>
85	My hardest battles are within myself.	4, 7
93	There seems to be a fullness in my head or nose most of the time.	1, <i>hea</i>
96	I have never done anything dangerous for the thrill of it.	6, 9, 0, <i>con</i>

Table 9 (Continued)

Item	Item Content	Scale membership
100	Most people are honest chiefly because they are afraid of being caught.	5, 6, 0, <i>cyn</i>
102	My speech is the same as always (not faster or slower, no slurring or hoarseness).	8, 9, 0
107	Most people will use somewhat unfair means to get what they want.	3, 6, 0, <i>K, cyn</i>
111	Often I can't understand why I have been so irritable and grouchy.	3, <i>K, ang</i>
118	I often wonder what hidden reason another person may have for doing something nice for me.	3, <i>cyn</i>
121	Criticism or scolding hurts me terribly.	2, 0, <i>K, A</i>
128	At times I feel like picking a fist fight with somebody.	2, <i>IMM, R,</i> <i>ang</i>
129	I have often lost out on things because I couldn't make up my mind soon enough.	3, 0, <i>A, obs</i>
146	I do not tire quickly.	1, 3
150	What others think of me does not bother me.	3, 4, <i>K</i>
152	I have never had a fainting spell.	3, <i>hea</i>
159	I am worried about sex.	3, 5- <i>m, 5-f, 8,</i> <i>MAC-R</i>

Table 9 (Continued)

Item	Item Content	Scale membership
160	I find it hard to make talk when I meet new people.	3, 4, 9, 0, K, <i>sod</i>
166	I can read a long while without tiring my eyes.	1, 3, <i>sch</i>
168	I have very few headaches.	1, 3, <i>hea</i>
169	My hands have not become clumsy or awkward.	5, 8
170	I like to study and read about things that I am working at.	7, <i>las, IMM</i>
171	Sometimes, when embarrassed, I break out in a sweat which annoys me greatly.	2, <i>R</i>
185	I frequently find myself worrying about something.	5, 7, K, <i>anx,</i> <i>obs</i>
189	It is not hard for me to ask help from my friends even though I cannot return the favor.	9
194	Some of my family have habits that bother and annoy me very much.	5, 9, <i>fam</i>
196	I hardly ever notice my heart pounding and I am seldom short of breath.	1, 3, <i>anx</i>
201	I get mad easily and then get over it soon.	3, K, <i>ang</i>
202	I have been quite independent and free from family rule.	4, <i>MAC-R</i>
204	My relatives are nearly all in sympathy with me.	4

Table 9 (Continued)

Item	Item Content	Scale membership
205	I have periods of such great restlessness that I cannot sit long in a chair.	3, 7, 8, 9
209	I believe I am no more nervous than most others.	2, <i>anx</i>
211	My way of doing things is apt to be misunderstood by others.	2, <i>cyn, aln</i>
216	I can be friendly with people who do things which I consider wrong.	3
218	I have difficulty starting to do things.	2, 8, <i>las, A, IMM</i>
225	It is safer to trust nobody.	3, <i>cyn</i>
227	When in a group of people I have trouble thinking of the right things to talk about.	3, 4, 9, 0, <i>K, A, aln</i>
229	When I leave home I do not worry about whether the door is locked and the windows are closed.	2
231	I have numbness in one or more places on my skin.	1, 8, <i>hea</i>
233	My eyesight is as good as it has been for years.	1, 3, <i>hea</i>
239	I do not often notice my ears ringing or buzzing.	1, 6, 8, 0, <i>hea, R</i>
243	Once in a while I laugh at a dirty joke.	2, <i>L</i>

Table 9 (Continued)

Item	Item Content	Scale membership
248	I am likely not to speak to people unless they speak to me.	3, 0, <i>sod</i>
251	I wish I were not bothered by thoughts about sex.	5- <i>m</i> , 5- <i>f</i> , 8
253	I think that I feel more intensely than most people do.	5, 6
257	In school I find it very hard to talk in front of the class.	7, 0, <i>sch</i>
258	I love my mother, or (if your mother is dead) I loved my mother.	8, F2, <i>fam</i>
260	I get all the sympathy I should.	8, <i>aln</i>
262	I seem to make friends about as quickly as others do.	8, 0, MAC-R, <i>sod</i>
263	A person who leaves valuable property unprotected is about as much to blame when it is stolen as the one who steals it.	6, <i>cyn</i>
264	I dislike having people around me.	8, F2, <i>sod</i>
265	I think nearly anyone would tell a lie to keep out of trouble.	6, 0, K, <i>cyn</i>
266	I am more sensitive than most other people.	6, 7
267	Most people inwardly dislike putting themselves out to help other people.	6, <i>cyn</i>
268	Many of my dreams are about sex.	8, MAC-R
270	I am easily embarrassed.	7, 0, A
272	I have never been in love with anyone.	8, PRO
273	I am afraid of using a knife or anything very sharp or pointed.	8, F2

Table 9 (Continued)

Item	Item Content	Scale membership
274	I almost never dream.	7
275	I have never been paralyzed or had any unusual weakness of any of my muscles.	8, <i>hea</i>
276	Sometimes my voice leaves me or changes even though I have no cold.	8, 0
277	My mother or father often makes me obey even when I think it is unreasonable.	6, <i>R, fam</i>
278	Peculiar odors come to me at times.	8, <i>biz</i>
281	I feel anxiety about something or someone almost all the time.	7, <i>A, anx</i>
284	Sometimes I become so excited that I find it hard to get to sleep.	7
287	At times I hear so well it bothers me.	6, 8
290	Often I cross the street in order to not meet someone I see.	7, <i>A, sod</i>
291	I often feel as if things are not real.	8, <i>A, biz</i>
293	I have a habit of counting things that are not important such as bulbs on electric signs, and so forth.	7, <i>obs</i>
295	I tend to be on my guard with people who are somewhat more friendly than I had expected.	6, <i>cyn</i>
296	I have strange and peculiar thoughts.	7, 8, <i>biz</i>
299	I hear strange things when I am alone.	8, <i>biz</i>

Table 9 (Continued)

Item	Item Content	Scale membership
300	I have been afraid of things or people that I knew could not hurt me.	7, 8
301	I have no dread of going into a room by myself where other people have already gathered and are talking.	7, 0
303	Sometimes I enjoy hurting the persons I love.	8, F2, <i>fam</i>
307	Bad words, often terrible words, come into my mind and I cannot get rid of them.	7, IMM, <i>obs</i>
308	Sometimes some unimportant thought will run through my mind and bother me for days.	7, 0, A, <i>obs</i>
309	Almost everyday something happens to frighten me.	7, 8, F2
310	I usually have to stop and think before I act in even small matters.	7, A, <i>obs</i>
311	I am inclined to take things hard.	7, <i>dep</i>
315	Someone has control over my mind.	6, F2, <i>biz</i>
332	At one or more times in my life I felt that someone was making me do things by hypnotizing me.	6, 8, F2, <i>biz</i>

Note. Items with an asterisk next to them were dropped on all listed scales except scale 2. The item was able to distinguish between the normative sample and the homogenous depression sample on all other listed scales except scale 2. Therefore, these items were not completely dropped from the test.

Table 10

MMPI-A Validity and Basic Scale Item Totals for Original and Revised Scales

Scale	Standard	Deleted	Revised
<i>L</i>	14	2	12
<i>F</i>	66	7	59
<i>FI</i>	33	0	33
<i>F2</i>	33	7	26
<i>K</i>	30	10	20
<i>1-Hs</i>	32	13	19
<i>2-D</i>	57	12	45
<i>3-Hy</i>	60	27	33
<i>4-Pd</i>	49	8	41
<i>5-Mf</i>	44	8	36
<i>6-Pa</i>	40	17	23
<i>7-Pt</i>	48	25	23
<i>8-Sc</i>	77	33	44
<i>9-Ma</i>	46	13	33
<i>0-Si</i>	62	19	43

Note. Items were not deleted directly from scales *L*, *F*, *FI*, *F2*, *K*, *Mf*, and *Si* because these scales were not evaluated in terms of item effectiveness. However deletions of items from the remaining Basic scales would result in removal of items for the former scales due to item overlap in scale membership and these subsequent effects are shown in this table.

Table 11

MMPI-A Basic Scale ANOVA Results for Girls

Scale	MMPI-A Standard Basic Scales			MMPI-A Revised Basic Scales			F
	Mean	(SD)	α level	Mean	(SD)	α level	
<i>Hs (1)</i>	51.2	(12.5)	.79	54.1	(11.8)	.74	33.1***
<i>D (2)</i>	54.3	(9.8)	.66	55.6	(10.3)	.66	10.3**
<i>Hy (3)</i>	49.7	(11.5)	.55	56.3	(12.0)	.57	105.6***
<i>Pd (4)</i>	61.3	(10.5)	.68	62.6	(10.4)	.61	15.5***
<i>Pa (6)</i>	54.5	(11.2)	.59	60.2	(11.1)	.70	111.1***
<i>Pt (7)</i>	48.4	(11.5)	.86	52.4	(11.9)	.81	95.7***
<i>Sc (8)</i>	53.3	(13.1)	.89	56.9	(13.3)	.82	76.4***
<i>Ma (9)</i>	55.2	(10.7)	.61	57.2	(11.5)	.55	12.7***

Note. * $p < .05$, ** $p < .01$, *** $p < .001$.

Table 12

MMPI-A Basic Scale ANOVA Results for Boys

Scale	MMPI-A Standard Basic Scales			MMPI-A Revised Basic Scales			F
	Mean	(SD)	α level	Mean	(SD)	α level	
<i>Hs (1)</i>	52.6	(9.4)	.78	56.1	(10.5)	.71	59.1***
<i>D (2)</i>	58.3	(9.3)	.65	58.9	(9.2)	.63	2.5
<i>Hy (3)</i>	54.8	(9.7)	.63	59.4	(9.5)	.61	73.5***
<i>Pd (4)</i>	60.3	(11.3)	.63	61.1	(11.5)	.55	5.7*
<i>Pa (6)</i>	53.3	(11.3)	.57	56.0	(10.9)	.67	13.6***
<i>Pt (7)</i>	47.9	(10.3)	.84	49.7	(10.6)	.76	4.2*
<i>Sc (8)</i>	50.6	(11.5)	.88	54.3	(12.0)	.80	103.6***
<i>Ma (9)</i>	50.6	(9.9)	.61	53.9	(11.4)	.53	54.1***

Note. * $p < .05$, ** $p < .01$, *** $p < .001$.

Table 13 shows the percentages for hit rate, sensitivity, specificity, positive predictive power and negative predictive power are given for boys and girls at two different cut-off scores for the standard and revised Basic scales. The revised scales demonstrated an increase in hit rate, sensitivity, specificity, positive predictive power and negative predictive power (see Table 13). Further, the overall prediction accuracy found for both the standard and revised scales was consistently higher for predictions based on the $T \geq 65$ criterion in contrast to the $T \geq 60$ criterion. Specifically, using the $T \geq 65$, all classification indices increased (i.e., the ability of the Basic scales to correctly identify clinical respondents and normal respondents increased when the clinical designation was set at $T \geq 65$). Specifically, there was a decrease in sensitivity for the revised scales for boys, and a decrease in specificity in the revised scales for females. However, as expected, sensitivity was generally lower than specificity for a cut-off of $T \geq 65$, and sensitivity was generally higher than specificity for a cut off of $T \geq 60$.

Table 13

Hit Rate, Sensitivity, Specificity, PPP, and NPP for Predicting Adolescent Membership in Normal (N= 1600) and Clinical (N= 163) Groups using Standard and Revised Scales at Two Cutting Scores by Gender

Gender	Scale	Cut-off Score (T \geq)	Hit Rate	Sensitivity	Specificity	PPP	NPP
Boys	Standard	65	.66	.57	.66	.16	.93
	Revised	65	.68	.63	.69	.19	.94
	Standard	60	.54	.78	.52	.16	.95
	Revised	60	.61	.76	.59	.18	.96
Girls	Standard	65	.66	.52	.68	.13	.94
	Revised	65	.72	.63	.73	.18	.96
	Standard	60	.56	.69	.55	.12	.95
	Revised	60	.57	.84	.54	.14	.97

Note: PPP= Positive Predictive Power, NPP= Negative Predictive Power.

CHAPTER IV

DISCUSSION

The discussion of findings will be organized into three sections, corresponding with the three primary objectives of the study. The first objective of this study was to extend Archer, Handel, and Lynch's (2001) investigation of item effectiveness within the MMPI-A Basic and Content scales to item-effectiveness evaluations for the Supplementary scales, Harris Lingoes subscales, and subtle versus obvious items. The purpose of this extension was to evaluate if there was an increase in the frequency of occurrence of effective items in scales and subscales that were more homogeneous than the Basic clinical scales, in terms of item content. It was hypothesized that this research extension of Archer et al.'s (2001) study to include additional MMPI-A scales and subscales would not significantly effect the frequency of effective items in contrast to those frequencies established for the Basic and Content Scales.

In terms of supplementary scale findings, using the manual clinical sample as the criterion group, the three substance abuse scales (*MAC-R*, *ACK*, and *PRO*) demonstrated a relatively high frequency of effective items, ranging from 62% of the item composition for the *ACK* scale (for both boys and girls) to 94% of the item composition for the *PRO* scale (for boys). While these findings are impressive, results are also consistent with the observation that of the 420 boys and 293 girls included in the manual clinical sample, 299 of those boys and 163 of those girls were evaluated in inpatient alcohol and drug treatment facilities (Butcher et al., 1992). Additionally, higher scores on the substance abuse scales may have been a product of the process by which these MMPI-A scales were developed. The *MAC-R* scale was derived from the *MAC* scale on the original form

of the MMPI by contrasting the item responses from 300 adult male alcoholics with 300 adult male psychiatric patients (MacAndrew, 1965). The items that showed the greatest difference between the substance abusing group and the psychiatric group were selected for scale membership. Forty-five of the forty-nine items from *MAC* were retained for the MMPI-A, and four new items were added to the scale (Butcher et al., 1992). The *PRO* scale was developed for the MMPI-A in a similar fashion to the *MAC* scale. *PRO* scale items were empirically selected based on item endorsement differences found between adolescents in alcohol and drug treatment programs and adolescents receiving inpatient psychiatric services (Weed, Butcher, & Williams, 1994). These adolescents in the alcohol and drug treatment settings were the same adolescents used to develop the clinical sample data reported in the MMPI-A manual and used in our study as one of the primary clinical groups to evaluate item effectiveness. Because the clinical group used to create the *PRO* scale heavily overlapped with the clinical group we used to determine item effectiveness, it would be expected that the *PRO* scale would have the highest percentage of effective items for boys (94%) and girls (89%). However, previous research has also shown that the MMPI-A substance abuse scales are effective with independent substance abuse samples. Michucci (2002), for example, was able to accurately classify approximately 90% of substance abuse cases on at least one of the scales. Gantner, Graham, and Archer (1992) examined three different samples (residential substance abusers, psychiatric inpatients, and high school students) and results indicated that the highest degree of discrimination on the *MAC* scale occurred between substance abusing adolescents and normal adolescents. Additionally, other studies have found positive correlations between the substance abuse scales and therapist ratings of adolescents substance-abusing

behaviors (Gallucci, 1997a). These findings, in conjunction with numerous other studies, support the utility of the MMPI-A substance abuse scales (e.g., Aharoni, 1999; Gallucci, 1997b; Gantner, Graham & Archer, 1992; Stein & Graham, 2001; Walfish, Massey, & Krone, 1990; Weed, Butcher, & Williams, 1994).

Since each of the MMPI-A Basic Scales typically contain several content areas, and it is possible that item effectiveness may vary as a function of content area, this hypothesis was tested using the Harris Lingo's subscales. These subscales were developed by Harris and Lingo's (1955) for the MMPI to identify the major item content areas. The items scored in each of the Basic scales were examined and grouped into subscales that reflected a single trait or content area and then given a name that was thought to be descriptive of this trait or attitude. The Harris-Lingo's subscales were carried over to the MMPI-A because few of the original items used to construct the scales were deleted during the development of the MMPI-A.

In general, current findings provide little evidence that the frequency of item effectiveness was related to the homogeneity of content areas, at least as those areas were defined in the Harris Lingo's subscale (see Table 5). Of the 28 Harris Lingo's subscales (56 tabulations done separately by gender), only 13 subscales showed item effectiveness above 50% when comparing the normative versus clinical manual sample. Not surprisingly, HL *Pd* subscales contained the highest percentages of effective items, similar to the performance of the *Pd* parent scale among the Basic scales.

Comparing the Harris Lingo's subscales (see right side of Table 5) with their Basic clinical scale counterparts (see right side of Table 7), there were no HL subscales that contained a significantly higher frequency of effective items when contrasted to their

“parent” Basic scale. Indeed, two parent scales (Basic scales) had a frequency of item effectiveness that was significantly higher than the corresponding Harris Lingoes subscales (i.e. *Hy3* vs. *Hy* and *Sc6* vs. *Sc*), while all other remaining comparisons were not significantly different. Thus, there appears to be no difference in item effectiveness when the scales are more homogeneous in terms of content.

Further, it was possible that the frequency of item effectiveness varied as a function of obviousness or subtlety of MMPI-A items. Although the MMPI-A does not contain a set of subtle-obvious items, Weiner and Harmon subtle-obvious items developed for the MMPI were utilized in this study. The Weiner-Harmon Subtle-Obvious scales on the MMPI were developed by dividing MMPI items from five scales into two groups (Weiner, 1948). One group consisted of items that were relatively easy to detect as indicating disturbance (obvious) and the other group consisted of items that were difficult to detect as indicating disturbance (subtle). These scales were developed to assess the frequency with which respondents endorse obvious or subtle items, thus allowing test-givers to assess the test-taking attitude of a respondent. Weiner and Harmon assumed those respondents who endorse more obvious items and less subtle items, may be over-reporting or magnifying their problems, whereas the converse is true for respondents underreporting their problems. However, there have been mixed findings regarding the utility of these scales on the MMPI and on the MMPI-2. Some studies demonstrate that the usefulness of the Weiner-Harmon scales in determining profile accuracy is limited (e.g., Bagby, Buis, & Nicholson, 1995; Boone, 1994; Herkov, Archer, & Gordon, 1991; Timbrook, Graham, Keiller, & Watts, 1993; Weed, Ben-Porath, & Butcher, 1990), where as others have found that these scales are useful in accurately

identifying individuals who over-report or under-report symptomatology (e.g., Brems & Johnson, 1991; Hsu, Santelli, & Hsu, 1989; Schretlen, 1988). In the current study, the obvious item component on the Basic clinical scales did not show a significant difference in frequency of item effectiveness when contrasted with the frequencies of effective items found for subtle item components within each of these basic scales. Specifically, there was no significant difference between subtle and obvious item endorsement effectiveness for the Depression, Hysteria, Psychopathic Deviate, Paranoia and Mania Scales. Thus, overall findings provide no support for the possibility that obvious items would produce a higher frequency of effective items than subtle items. This finding is consistent with the decision of the MMPI-A steering committee to discourage clinical use of these subscales by not including a subtle-obvious profile sheet or norms for the MMPI-A (Archer, 1997; Butcher et al., 1992).

Overall, there has been no improvement in item effectiveness from the results of Archer and his colleagues' study when examining the Supplementary scales and Harris-Lingoes subscales, or subtle-obvious items. Because the MMPI-A scales have not been useful in helping to understand the item effectiveness issue, a logical step was to examine item effectiveness in relationship to various diagnostic groups.

The second broad purpose of the study was to examine frequency of item effectiveness among the Basic, Content, and Supplementary scales, and Harris Lingoes subscales with reference to two homogeneous diagnostic criterion groups. This hypothesis proposed that the frequency of non-discriminating items within specific scales would vary as a function of the diagnostic groups used in contrast to normative groups, i.e., the frequency of effective items would increase when the normative sample was

contrasted with homogeneous groups that were specifically related to the purpose of the scale (e.g., depressed patients used to determine item effectiveness for the *D* Basic scale).

Although this hypothesis proposed that the frequency of effective items in the Depression scale (*D*) would increase when normative samples were compared with specific depression samples, this was not supported. Specifically, there was no significant item effectiveness difference between the manual clinical sample frequency and the homogeneous depression sample frequency. The depression sample contrasts only produced a higher frequency of effective items than the general clinical sample on scales *Hs* (for both genders) and scale *Sc* (for boys), that is, Basic scales that are not directly related to the depression construct.

Similar comparisons regarding the frequency of item effectiveness were made for the Conduct disorder group on Psychopathic Deviate scale (*Pd*). One would expect the frequency of effective items to be higher on the *Pd* scales for evaluations that contrasted the endorsement frequency of the normative sample with that of conduct disordered adolescents rather than a more heterogeneous clinical sample. However, the results of the current study did not produce evidence that the frequency of item effectiveness for the *Pd* scale improved as a result of using the more homogeneous and construct relevant clinical sample.

In contrast to the results obtained when the normative sample was compared to the homogeneous depression sample for scale 2, the comparison of item endorsement frequencies between the normative and the conduct disorder sample produced a higher frequency of effective items for boys than the normative-manual sample for scale 2. Although it was expected that boys in the conduct disorder group would also endorse

symptoms of depression, it is interesting that there would be a significant difference between the normative-conduct samples on the *D* scale but no significant difference between the normative-depression samples when compared to the normative-manual clinical sample. Given that these results may be due to chance, it is suggested that these findings be replicated on other independent clinical samples.

Further, on the Content scales (see Table 8), the depression sample did not show a higher frequency of effective items than the general manual clinical sample on the conceptually related scales. Although statistical comparisons were unable to be made between *A-con* for the conduct disorder group and the manual clinical sample, boys had a 70% frequency of item effectiveness, a relatively high rate, compared to other content scales. However, it is unknown if this scale performs better with the conduct disorder group than the general manual clinical sample. Among the 15 MMPI-A Content scales, the only significant difference is that the manual clinical sample has a higher frequency of effective items than the depression sample for boys on *A-ang*, and in the conduct disorder sample for girls on *A-lse* and *A-fam*. In both the depression and conduct disorder sample, results are not consistent with hypotheses in that significant differences do not reflect apriori predictions.

As might be expected, given the prior pattern of results, none of the Supplementary scales or Harris Lingoes subscales performed significantly better in comparisons using the homogeneous diagnostic groups rather than the manual clinical sample. However, when examining Supplementary scale results for the manual clinical sample there was a statistical difference between the manual clinical sample and the two homogeneous diagnostic groups, especially on the *PRO* subscale. Specifically, the

Supplementary substance abuse scales (especially *PRO*) performed better (in terms of a higher frequency of effective items) in contrast with the manual clinical sample rather than the homogeneous samples. As stated before, this may be due to the fact that the manual sample was largely composed in part from a residential substance abuse treatment sample, all of whom were used as the criterion group in the development of the *PRO* scale (Weed, Butcher, & Williams, 1994).

The overall conclusion regarding the second purpose of this study is that there are no stable or reliable improvements in item endorsement effectiveness when using homogeneous clinical samples in contrast to the general clinical sample. It does not appear that the diagnostic specificity of the sample systematically increases item effectiveness for scales selected based on constructs related to the diagnostic criterion group. However, it may be argued that in the case of the Supplementary scales, the substance abuse scales performed at a significantly higher rate of effectiveness for the manual clinical sample (largely collected at substance abuse settings) than any of the other samples because we (unintentionally) included a clinical sample with a substance abuse component- the very criterion group used to construct the *PRO* scale as stated by Weed et al. (1994). Thus, at least for the *PRO* findings, results may be artificially inflated. This is illustrated by the relatively increased frequency of item effectiveness for the *PRO* scale versus the *MAC-R* and *ACK* scale.

A remaining question for future research focuses on why the substance abuse largely homogeneous sample produced a higher frequency of effective items within their conceptually related scales (i.e., *MAC-R*, *ACK*, and *PRO*) than the depression or the conduct disorder samples for their conceptually related scales, (i.e., Basic scales *D* and

Pd, respectively). One possible explanation might be that substance abuse constitutes a more specific and reliable diagnosis in contrast to Depression and Conduct disorder. However, there is no persuasive evidence to support this argument in the literature. Although there are many potentially relevant diagnostic groups that could have been used in this study, only two homogeneous samples were utilized. Future research should examine various disorders on several scales, e.g., anxiety disorders and scale 7; school problems and *A-sch*; psychosomatic symptoms and scales 1, 3, and *A-hea*; psychotic disorders and scale 8 and *A-biz*, etc. Using additional samples will help to define areas, if any exist, in which specific diagnostic groups produce higher frequencies of effective items.

Despite the impressive substance abuse scale item effectiveness findings, there are still numerous non-discriminating items within all the scales on the MMPI-A, both when examined in terms of homogeneous diagnostic groups and in terms of a general sample. Additionally, there appears to be no consistent or reliable gender differences. The only exception is for the conduct disordered group, where there were more effective items for boys than girls on three Basic scales (*D*, *Pt*, and *Si*), two content scales (*A-cyn* and *A-trt*), and one Harris Lingoes subscale (*A*). This finding is consistent with the perspective of the DSM-IV (1994) authors who note "Conduct Disorder, especially the Childhood-Onset Type, is much more common in males" (p. 88). However, non-discriminating items appear to persist regardless of item content, sample, and gender. This problem may exist for several reasons explored in detail below.

The decision to carry over the Basic scales from the MMPI to the MMPI-A without re-establishing item composition through the use of criterion groups may have

failed to be sensitive to relatively unique factors that play a part in adolescent development. Generalizing MMPI results based on adults to adolescents (for example, scale construction and composition) may have contributed to the inability of many of items to distinguish between a normative and clinical population. This idea is partially supported by the Archer et al. (2001) findings that there were many more items on the MMPI-2 scales that were able to discriminate between normative and clinical samples than found for MMPI-A counterpart scales. However, scales that were developed solely for the use of the MMPI-A (*A-ahn*, *A-las*, *A-sch*, *A-con*, *ACK*, *PRO*, *IMM*), did not appear to perform that much better than those carried over from the MMPI.

Additionally, the relatively higher frequency of ineffective items may be created, in part, by lower base rates in adolescents for certain psychiatric disorders and symptoms found among the standard MMPI Basic scales (e.g., Schizophrenia). For example, none of the adolescents in our clinical sample were diagnosed with schizophrenia. However, many of the adolescents in this sample were diagnosed with Depression or Conduct Disorder, and there were still few items within these scales that were able to discriminate between clinical and normative group endorsement frequencies. Further research should attempt to examine those disorders that are frequently diagnosed in adolescence, extending this examination of item effectiveness to new homogeneous samples (e.g., Anxiety disorders), as well as replicating item-effectiveness results using additional independent Depression, Conduct disorder, and Substance abuse samples.

Another possible explanation for current findings is that “normal” adolescents tend to endorse more symptoms of psychological disturbance than do “normal” adults, tending to blunt the discrimination achievable between normal and clinical adolescent

groups. Support for this idea was found in the study by Archer and his colleagues (2001) that demonstrated mean raw score values for the majority of the Basic Scales in the normative groups was higher for the MMPI-A than the MMPI-2. This further underscores the observation that methodologies used to develop scales for adults may not prove appropriate for adolescent inventories. There is substantial literature supporting the salient differences between adolescents and adults in their responses to objective personality inventories, including the MMPI (e.g., Archer 1984, 1987). Further, normal adolescence has been viewed as a time of emotional instability, rapid shifts in mood, and conflictual relationships with authority figures (Erikson, 1956; Freud, 1958; Hall, 1916). Offer and Offer (1975) found transient episodes of mild depression and anxiety to be fairly common in their surveys of normal adolescents.

The results shown in the current study may also have been produced, in part, by limitations or confounding influences inherent in the MMPI-A normative sample. Current literature indicates that the prevalence and incidence of psychopathology in adolescents ranges from 14-22% (Bradenburg, Friedman, & Silver, 1990; Gould, Wunsch-Hitzig, & Dohrenwend, 1981; McGee, Feehan, Williams, Partidge, Silva, & Kelly, 1990; NIMH, 1990; Rutter, Graham, Chadwick, & Yule, 1976). This raises an important question regarding the normative sample used in the development of the MMPI-A. That is, the sample was not screened for the occurrence of psychological disorders. The MMPI-A normative sample may have included adolescents endorsing items similar to the clinical populations, because they were, in fact, clinical respondents. While data on this issue is not directly available, the normative sample did complete a Life Events form that contained an item asking respondents if they had "been referred to a therapist or

counselor” in the past six months. Of the 1620 respondents in the normative sample, 84% ($n=1360$) responded “no” to this question and 15% ($n= 256$) responded yes (four respondents did not answer the question). Interestingly, this incidence is consistent with estimates of the national incidence of adolescent psychopathology (Bradenburg, et al., 1990; Gould, et al., 1981; McGee, et al., 1990; NIMH, 1990; Rutter, et al., 1976). An important extension to this study is to explore the implications of a normative group that contains respondents recently referred to counseling or therapy. Removing those respondents who answered “yes” to the question “Have you been referred to counseling or therapy?” from the normative sample and comparing the frequency of effective items on the revised normative and clinical groups on several independent samples would help in understanding if the normative sample is indeed part of the problem in producing ineffective items. This extension, while logistically relatively easy to perform, would aid in clarifying the problem of item-ineffectiveness. If, in fact, a substantial difference was found between the current normative sample and a revised normative sample, procedures similar to Colligan and Offord (1989) screening out those individuals diagnosed with medical or psychological problems or who report they have been referred for therapy) could and should be utilized in collecting a truly normative sample for future research in this area.

In addition to limitations of the MMPI-A normative sample, there were differences between the MMPI-A manual clinical sample and those independent clinical samples used in this study. The MMPI-A clinical sample presented in the test manual included inpatient adolescents, but also contained adolescents from day treatment and special school programs for emotionally disturbed children. In contrast, the independent

homogeneous clinical samples used for this study for comparison purposes exclusively contained adolescents whose psychiatric symptomatology required treatment in inpatient treatment facilities. Although those differences did not appear to significantly effect current results, it would be methodologically more desirable to have clinically similar groups for comparisons, if at all possible, because one could argue that the normative sample versus the manual-clinical sample comparisons might have produced higher rates of ineffective items (because the clinical sample population may be closer to the normative sample, in terms of psychological functioning, than a completely independent inpatient population).

The third purpose of this study was to re-examine and recalculate Basic scale data using only those items that effectively discriminated between the normative and clinical populations. The mean profile of the normative and clinical groups was contrasted based on the "revised" Basic scales with an independent clinical sample (CSB) to evaluate the extent to which profile sensitivity and specificity was effected by these scale modifications. Results supported the hypothesis in that the overall effectiveness of Basic scale discrimination increased, in terms of sensitivity, specificity, positive and negative predictive power and hit rate, when the Basic clinical scale items were removed that did not discriminate between normative and clinical groups of adolescents. Hit rate, sensitivity, specificity, positive predictive power (PPP) and negative predictive power (NPP) improved for both genders at a $T \geq 65$. While most classification scores improved at $T \geq 60$ it was not as effective for sensitivity and specificity as a cut-off of $T \geq 65$. Fontaine, Archer, Elkins, and Johansen (2001) demonstrated that a T -score cut-off of 65 resulted in higher levels of accurate classification overall while minimizing the

misclassification of both clinical and normal cases. This study indicated that a reduction in the T -score criterion used to define a “clinical range elevation” did not produce a more effective balance between test sensitivity and specificity (i.e., that lowering the clinical range elevation T -score criterion from 65 to 60 did not improve classification performance). However, a unique feature on the MMPI-A is the “shaded” zone between T scores in the 60 to 64 range alerting the test-user to approaching psychopathology. For this reason, classification scores were evaluated at both $T \geq 65$ and $T \geq 60$. All classification percentages were above .50 except for PPP (defined as the probability that an elevated score is being produced by a protocol from the clinical sample). One explanation for low PPP values is that 15% of the normative sample reported being referred for counseling or therapy, i.e., these adolescents may not have been “normal”. Thus, there were many respondents (potentially 256 of 1620) that were misclassified as “normal” but “accurately” produced elevated protocols, thus, dramatically reducing PPP. That also may be the reason that sensitivity (the percentage of participants whose protocols indicate the presence of clinical status who in fact were from the clinical sample) is lower than specificity (the percentage of participants whose within normal limits protocols accurately indicate the absence of pathology because, in fact, they were from the normative group) when $T \geq 65$. Future research should attempt to find a truly “normal” sample, as discussed earlier, in order to improve PPP. However, regardless of the limitations of PPP, there is still substantial evidence that removal of the ineffective items results in an overall increase in prediction accuracy.

In addition to improving hit rate performance, deleting “ineffective” items would also serve another important purpose. Archer, Maruish, Imhof, and Piotrowski (1991)

conducted a practitioner survey concerning the use of the MMPI with adolescents. Results demonstrated of the 124 respondents, 49% ($n= 61$ respondents) indicated that the primary disadvantage of using the MMPI with adolescents was the length/administration time of the test, followed by concerns regarding poor/outdated norms (20%) and reading level too high (18%). The MMPI-A, while shorter (478 items) than the original version of the MMPI (566 items), is still a lengthy instrument and demands a good deal of concentration and attention. While attempts have been made to shorten the test (i.e., Archer, Tirrell, & Elkins, 2001), short forms tend to be generally less reliable than their short form counterparts. In the current study, essentially shortened Basic scales (by over 100 items) produced results showing small increases in overall hit rate. Before applying the current revised basic clinical scales in clinical settings, however, many questions require evaluation. For example, when deleting items from a scale, does the scale still retain its original "meaning" in terms of extra-test correlates? Related to this point, would the code-type patterns produced by revised scales retain correlate patterns established for specific code-types with the standard instrument? An important future research direction is to examine the revised MMPI-A Basic scales with external correlate data such as diagnoses, self-report forms (e.g., Youth Self Report), and parent-report forms (Child Behavior Checklist and Personality Inventory for Children-Adolescent version). Another limitation of the current study is that items were only evaluated in terms of effectiveness on clinical scales and not on validity scales. For this reason, ineffective items should not be dropped from the test until the effects of such deletions on MMPI-A validity scales are assessed. Because the strength and usefulness of the validity scales distinguish the

MMPI-A from many other self report measures, future research focusing on the deletion of non-discriminating items on *L,F,K*, *VRIN*, and *TRIN* is warranted.

REFERENCES

- Achenbach, T. M., & Edelbrock, C. S. (1983). *Manual of the Child Behavior Checklist and revised child behavior profile*. Burlington: University of Vermont.
- Aharoni, D. M. (1999). The effectiveness of the MMPI-A in the assessment of adolescent substance abuse. *Dissertation Abstracts International*, 60, 6B. (UMI No. 9502196).
- Alperin, J. J., Archer, R. P., & Coates, G. D. (1996). Development of the effects of an MMPI-A K-correction procedure. *Journal of Personality Assessment*, 67, 155-168.
- American Psychiatric Association. (1994). *Diagnostic and statistical manual of mental disorders* (4th ed.). Washington DC: American Psychiatric Association.
- Archer, R. P. (1984). The use of the MMPI with adolescents: A review of salient issues. *Clinical Psychology Review*, 4, 241-251.
- Archer, R. P. (1987). *Using the MMPI with adolescents*. Hillsdale, NJ: Lawrence Erlbaum Associates.
- Archer, R. P. (1992). *MMPI-A: Assessing Adolescent Psychopathology*. Hillsdale, NJ: Lawrence Erlbaum Associates.
- Archer, R. P. (1997). *MMPI-A: Assessing adolescent psychopathology* (2nd ed.). Mahwah, NJ: Lawrence Erlbaum.
- Archer, R. P., Belevich, J. K. S., & Elkins, D. E. (1994). Item-level and scale-level factor structures of the MMPI-A. *Journal of Personality Assessment*, 62, 332-345.

- Archer, R. P., & Gordon, R. A. (1994). Psychometric stability of MMPI-A item modifications. *Journal of Personality Assessment, 62*, 416-426.
- Archer, R. P., Handel, R. W., Lynch, K. D. (2001). The effectiveness of MMPI-A items in discriminating between normative and clinical samples. *Journal of Personality Assessment, 77*, 420-435.
- Archer, R. P., Handel, R. W., Lynch, K. D., Elkins, D. E. (2002). MMPI-A validity scale uses and limitations in detecting varying levels of random responding. *Journal of Personality Assessment, 78*, 417-431.
- Archer, R. P., & Jacobson, J. M. (1993). Are critical items "critical" for the MMPI-A? *Journal of Personality Assessment, 61*, 547-556.
- Archer, R. P., & Krishnamurthy, R. (1994). A structural summary approach for the MMPI-A: Development and empirical correlates. *Journal of Personality Assessment, 63*, 554-573.
- Archer, R. P., & Krishnamurthy, R. (1996). The Minnesota Multiphasic Personality Inventory- Adolescent (MMPI-A). In C. S. Newmark (Ed.), *Major psychological assessment instruments* (pp. 59-107). Boston: Allyn and Bacon.
- Archer, R. P., Maruish, M., Imhof, E. A., & Piotrowski, C. (1991). Psychological test usage with adolescent clients: 1990 survey findings. *Professional Psychology: Research and Practice, 22*, 247-252.
- Archer, R. P., & Newsom, C. R. (2000). Psychological test usage with adolescent clients: Survey update. *Assessment, 7*, 227-236.

- Archer, R. P., Pancoast, D. L., & Gordon, R. A. (1994). The development of the MMPI-A Immaturity (IMM) scale: Findings for normal and clinical samples. *Journal of Personality Assessment, 62*, 145-156.
- Archer, R. P., Pancoast, D. L. & Klinefelter, D. (1989). A comparison of MMPI codetypes produced by traditional and recent adolescent norms. *Psychological Assessment: A Journal of Consulting and Clinical Psychology, 1*, 23-29.
- Archer, R. P., Tirrell, C. A., & Elkins, D. E. (2001). An evaluation of an MMPI-A short form: Implications for adaptive testing. *Journal of Personality Assessment, 76*, 76-89.
- Bagby, R. M., Buis, T. N., Nicholson, R. A. (1995). Relative effectiveness of the standard validity scales in detecting fake-bad and fake-good responding: Replication and extension. *Psychological Assessment, 7*, 84-92.
- Bandura, A. (1964). The stormy decade: Fact of fiction? *Psychology in the School, 1*, 224-231.
- Ben-Porath, Y. S., Hostetler, K., Butcher, J. N., Graham, J. R., (1989). New subscales for the MMPI-2 Social Introversion (Si) scale. *Psychological Assessment: A Journal of Consulting and Clinical Psychology, 1*, 169-174.
- Black, K. (1994). A critical review of the MMPI-A. *Child Assessment News, 4*, 9-12.
- Boone, D. (1994). Reliability of the MMPI-2 subtle and obvious scales with psychiatric inpatients. *Journal of Personality Assessment, 62*, 346-351.
- Bradenburg, N. A., Friedman, R. M., & Silver, S. E. (1990). The epidemiology of childhood psychiatric disorders: Prevalence findings from recent studies. *Journal of the American Academy of Child and Adolescent Psychiatry, 29*, 76-83.

- Brems, C., & Johnson, M. E. (1991). Subtle-obvious scales of the MMPI: Indicators of profile validity in a psychiatric population. *Journal of Personality Assessment*, 56, 536-544.
- Butcher, J. N., Dahlstrom, W. G., Graham, J. R., Tellegen, A., Kaemmer, B. (1989). *MMPI-2 Manual for Administration and Scoring*. Minneapolis, MN: University of Minnesota Press.
- Butcher, J. N., Graham, J. R., Ben-Poarth, Y. S., Tellegen, A., Dahlstrom, W. G., & Kaemmer, B. (2001). *Minnesota Multiphasic Personality Inventory-2. Manual for administration, scoring, and interpretation. Revised edition*. Minneapolis: University of Minnesota Press.
- Butcher, J. N., Williams, C. L., Graham, J. R., Archer, R. P., Tellegen, A., Ben-Poarth, Y.S., & Kaemmer, B. (1992). *MMPI-A (Minnesota Multiphasic Personality Inventory - Adolescent): Manual for Administration, Scoring, and Interpretation*. Minneapolis, MN: University of Minnesota Press.
- Capwell, D. F. (1945a). Personality patterns of adolescent girls. I. Girls who show improvement in IQ. *Journal of Applied Psychology*, 29, 212-228.
- Capwell, D. F. (1945b). Personality patterns of adolescent girls. II. Delinquents and non-delinquents. *Journal of Applied Psychology*, 29, 284-297.
- Colligan, R. C., & Offord, K. P. (1989). The aging MMPI: Contemporary norms for contemporary teenagers. *Mayo Clinic Proceedings*, 64, 3-27.
- Dahlstrom, W. G., Welsh, G. S., & Dahlstrom, L. E. (1972). *An MMPI handbook: Vol. 1 Clinical interpretation* (rev. ed.). Minneapolis: University of Minnesota Press.

- Ehrenworth, N. V., & Archer, R. P. (1985). A comparison of clinical accuracy ratings of interpretive approaches for adolescent MMPI responses. *Journal of Personality Assessment, 49*, 413-421.
- Erikson, E. H. (1956). The concept of ego identity. *The Journal of the American Psychoanalytic Association, 4*, 56-121.
- Ferdinand, R. F., & Verhulst, F. C. (1995). Psychopathology from adolescence into young adulthood: An 8-year follow-up study. *American Journal of Psychiatry, 152*, 1586-1594.
- Fontaine, J. L., Archer, R. P., Elkins, D. E., & Johansen, J. (2001). The effects of MMPI-A T-score elevation on classification accuracy of normal and clinical adolescent samples. *Journal of Personality Assessment, 76*, 264-281.
- Freud, A. (1958). Adolescence. *Psychoanalytic Study of the Child, 13*, 255-278.
- Gallucci, N. T. (1997a). Correlates of MMPI-A substance abuse scales. *Assessment, 4*, 87-94.
- Gallucci, N. T. (1997b). On the identification of patterns of substance abuse with the MMPI-A. *Psychological Assessment, 9*, 224-232.
- Gantner, A. B., Graham, J. R., & Archer, R. P. (1992). Usefulness of the MAC scale in differentiating adolescents in normal, psychiatric, and substance abuse settings. *Psychological Assessment, 4*, 133-137.
- Gottesman, I. I., Hanson, D. R., Kroeker, T. A., & Briggs, P. F. (1987). New MMPI normative data and power-transformed T-score tables for the Hathaway-Monachesi Minnesota cohort of 14,019 fifteen-year-old and 3,674 eighteen-year-

- olds. In R. P. Archer, *Using the MMPI with adolescents* (pp. 241-297). Hillsdale, NJ: Lawrence Erlbaum Associates.
- Gould, M. S. Wunsch- Hitzig, R., & Dohrenwend, B. (1981). Estimating the prevalence of childhood psychopathology: A critical review. *Journal of the American Academy of Child Psychiatry*, 20, 462-476.
- Greene, R. L. (2000). *The MMPI-2: An interpretive manual (2nd ed.)*. Boston, MA: Allyn & Bacon.
- Hall, G. S. (1916). *Adolescence*. New York: Appelton.
- Harris, R. E., & Lingo, J. C. (1955). *Subscales for the MMPI: An aid to profile interpretation*. Department of Psychiatry, University of California School of Medicine and the Langley Porter Clinic, mimeographed materials.
- Hathaway, S. R., & McKinley, J. C. (1943). *The Minnesota Multiphasic Personality Inventory* (rev. ed.). Minneapolis: University of Minnesota Press.
- Hathaway, S. R., & Monachesi, E. D. (Eds.). (1953). *Analyzing and predicting juvenile delinquency with the MMPI*. Minneapolis: University of Minnesota Press.
- Hathaway, S. R., & Monachesi, E. D. (1961). *An atlas of juvenile MMPI profiles*. Minneapolis: University of Minnesota Press.
- Hathaway, S. R., & Monachesi, E. D. (1963). *Adolescent personality and behavior: MMPI patterns of normal, delinquent, dropout, and other outcomes*. Minneapolis: University of Minnesota Press.
- Herkov, M. J., Archer, R. P., & Gordon, R. A. (1991). MMPI response sets among adolescents: An evaluation of the limitations of the subtle-obvious subscales.

- Psychological Assessment: A Journal of Consulting and Clinical Psychology*, 3, 424-426.
- Hoeglund, T. & Tellegan A. (1998). *A software package for the derivation of Uniform T scores for MMPI-2 scales*. Minneapolis: Regents of the University of Minnesota.
- Howell, D. C. (1992). *Statistical methods for Psychology*. Belmont, CA: Duxbury Press.
- Hsu, L. M., Santelli, J., & Hsu, J. (1989). Faking detection validity and incremental validity of response latencies to MMPI subtle and obvious items. *Journal of Personality Assessment*, 53, 278-295.
- Hyman, S. E. (2001). Mood disorders in children and adolescents: An NIMH perspective. *Biological Psychiatry*, 49, 962-969.
- Kessler, R. C., Avenevoli, S., & Merikangas, K. R. (2001). Mood disorders in children and adolescents: An epidemiological perspective. *Biological Psychiatry*, 49, 1002-1014.
- Klinefelter, D., Pancoast, D. L., Archer, R. P., & Pruitt, D. L. (1990). Recent adolescent MMPI norms: T-score elevation comparisons to Marks and Briggs. *Journal of Personality Assessment*, 54, 379-389.
- Klinge, V. & Lachar, D., Grissell, J., & Berman, W. (1978). Effects of scoring norms on psychiatric drug users' and nonusers' MMPI profiles. *Adolescence*, 13, 1-11.
- Klinge, V., & Strauss, M. E. (1976). Effects of scoring norms on adolescent psychiatric patients' MMPI profiles. *Journal of Personality Assessment*, 40, 13-17.
- Koss, M. T., & Butcher, J. N. (1973). A comparison of psychiatric patients' self-report with other sources of clinical information. *Journal of Research in Personality*, 7, 225-236.

- Krishnamurthy, R., Archer, R. P., & House, J. J. (1996). The MMPI-A and Rorschach: A failure to establish convergent validity. *Assessment*, 3, 179-191.
- Lachar, D., Klinge, V., & Grissell, J. L. (1976). Relative accuracy of automated MMPI narratives generate from adult norm and adolescent norm profiles. *Journal of Consulting and Clinical Psychology*, 44, 20-24.
- Lewinsohn, P. M., Hops, H. Roberts, R. E., Seeley, J. R., & Andrews, J. A. (1993). Adolescent psychopathology: I. Prevalence and incidence of depression and other DSM-III-R disorders in high school students. *Journal of Abnormal Psychology*, 102, 133-144.
- MacAndrew, C. (1965). The differentiation of male alcoholic outpatients from non-alcoholic psychiatric outpatients by means of the MMPI. *Quarterly Journal of Studies on Alcohol*, 26, 238-246.
- Marks, P. A., & Briggs, P. F. (1972). Adolescent norm tables for the MMPI. In W. G. Dahlstrom, G. S. Welsh, & L. E. Dahlstrom, *An MMPI Handbook: Vol. 1. Clinical Interpretation (rev. ed., pp. 388-399)*. Minneapolis: University of Minnesota Press.
- Marks, P. A., Seeman, W., & Haller, D. L. (1974). *The actuarial use of the MMPI with adolescents and adults*. Baltimore: Williams & Wilkins.
- McDermott, P. A. (1996). A nationwide study of developmental and gender prevalence for psychopathology in childhood and adolescence. *Journal of Abnormal Child Psychology*, 24, 53-66.

- McGee, R., Feehan, M., Williams, S., Partridge, F., Silva, P. A., & Kelly, J. (1990).
DSM-III disorders in a large sample of adolescents. *Journal of the American
Academy of Child and Adolescent Psychiatry, 29*, 611-619.
- Michucci, J. (2002). Accuracy of MMPI-A scales, ACK, MAC-R, and PRO in detecting
comorbid substance abuse among psychiatric inpatients. *Assessment, 9*, 111-122.
- Monachesi, E. D. (1948). Some personality characteristics of delinquents and non-
delinquents. *Journal of Criminal Law and Criminology, 38*, 487-500.
- Monachesi, E. D. (1950). Personality characteristics of institutionalized and non-
institutionalized male delinquents. *Journal of Criminal Law and Criminology,
41*, 167-179.
- Monachesi, E. D., & Hathaway, S. R. (1969). The personality of delinquents. In
J. N. Butcher (Ed.). *MMPI: Research developments and clinical applications* (pp.
207-219). Minneapolis: University of Minnesota Press.
- National Institute of Mental Health (NIMH). (1990). *National plan for research on child
and adolescent mental disorders* (DHHS Publication No. ADM 90-1683).
Washington, DC: U. S. Government Printing Office.
- Offer D., & Offer, J. B. (1975). *From teenager to young manhood*. New York: Basic
Books.
- Rutter, M., Graham, P., Chadwick, O. F. D., & Yule, W. (1976). Adolescent turmoil: Fact
or fiction? *Journal of Child Psychology and Psychiatry, 17*, 35-56.
- Schretlen, D. J. (1988). The use of psychological tests to identify malingered symptoms
of mental disorders. *Clinical Psychology Review, 8*, 451-456.

- Spivack, G., Haimes, P. E., & Spotts, J. (1967). *Devereux Adolescent Behavior (DAB) Rating Scale Manual*. Devon, PA: The Devereux Foundation.
- Stein, L. A., & Graham, J. R. (2001). Use of the MMPI-A to detect substance abuse in a juvenile correctional setting. *Journal of Personality Assessment*, 77, 508-523.
- Timbrook, R. E., Graham, J. R., Keiller, S. W., & Watts, D. (1993). Comparison of the Wiener-Harmon Subtle Obvious Scales and the standard validity scales in detecting valid and invalid MMPI-2 profiles. *Psychological Assessment*, 5, 53-61.
- Walfish, S., Massey, R., & Krone, A. (1990). MMPI profiles of cocaine-addicted individuals in residential treatment: Implications for practical treatment planning. *Journal of Substance Abuse Treatment*, 7, 151-154.
- Weed, N. C., Ben-Porath, Y. S., & Butcher, J. N. (1990). Failure of Wiener and Harmon MMPI subtle scales as personality descriptors and as validity indicators. *Psychological Assessment: A Journal of Consulting and Clinical Psychology*, 2, 281-285.
- Weed, N. C., Butcher, J. N., & Williams, C. L. (1994). Development of MMPI-A alcohol/drug problem scales. *Journal of Studies on Alcohol*, 55, 296-302.
- Weiner, I. B. (1948) Subtle and obvious keys for the MMPI. *Journal of Consulting Psychology*, 12, 164-170.
- Weiner, I. B., & Del Gaudio, A. C. (1976). Psychopathology in adolescence: An epidemiological study. *Archives of General Psychiatry*, 33, 187-193.

VITA

Kathleen D. Lynch, M.A.

1416 Hampton Boulevard, Apt. 6B
Norfolk, Virginia 23517
(757) 640-1405

VCPCP
Pembroke Two/Suite 301
287 Independence Blvd.
Virginia Beach, VA 23462
(757) 518-2550

Kathleen D. Lynch attended the College of Charleston from 1993-1997 and received a B.S. in Psychology. She then attended Western Carolina University (1997-1999) for her masters in Clinical Psychology. She completed her thesis "Parental Predictions of Children's Interests and Cognitive Performance" under the supervision of Bruce Henderson, Ph.D. and her internship in Washington D.C. working with children. Kathleen was then accepted to the Virginia Consortium Program in Clinical Psychology (VCPCP) in 1999. She trained at several sites including the Hampton Veterans Affairs Medical Center, Virginia Beach Public Schools, Suffolk Department of Social Services, and Hampton Roads Neuropsychology Clinic. Kathleen taught Introductory Psychology and Developmental Psychology at Norfolk State University and held a position as a research assistant for Robert P. Archer, Ph.D. at Eastern Virginia Medical School (EVMS). Her third year concentration was in Neuropsychology at Eastern State Hospital and The Neuropsychology Center at EVMS. In July of 2003, she will finish her internship at Eastern Virginia Medical School and graduate with her Psy.D. in Clinical Psychology. In September 2003, Kathleen will be a Post-Doctoral Fellow in Clinical Neuropsychology at Eastern Virginia Medical School and Eastern State Hospital. Kathleen also works for Virginia Beach Psychiatric Center as a consulting clinician to area Emergency Rooms.