### IOWA STATE UNIVERSITY Digital Repository

**Retrospective Theses and Dissertations** 

Iowa State University Capstones, Theses and Dissertations

1984

## The effects of silica, nitrogen, and phosphorus limitation on the biochemical composition of Cyclotella meneghiniana Kütz: an experimental analysis

David Fred Millie Iowa State University

Follow this and additional works at: https://lib.dr.iastate.edu/rtd Part of the <u>Botany Commons</u>

### **Recommended** Citation

Millie, David Fred, "The effects of silica, nitrogen, and phosphorus limitation on the biochemical composition of Cyclotella meneghiniana Kütz: an experimental analysis " (1984). *Retrospective Theses and Dissertations*. 9013. https://lib.dr.iastate.edu/rtd/9013

This Dissertation is brought to you for free and open access by the Iowa State University Capstones, Theses and Dissertations at Iowa State University Digital Repository. It has been accepted for inclusion in Retrospective Theses and Dissertations by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digrep@iastate.edu.



### **INFORMATION TO USERS**

This reproduction was made from a copy of a document sent to us for microfilming. While the most advanced technology has been used to photograph and reproduce this document, the quality of the reproduction is heavily dependent upon the quality of the material submitted.

The following explanation of techniques is provided to help clarify markings or notations which may appear on this reproduction.

- 1. The sign or "target" for pages apparently lacking from the document photographed is "Missing Page(s)". If it was possible to obtain the missing page(s) or section, they are spliced into the film along with adjacent pages. This may have necessitated cutting through an image and duplicating adjacent pages to assure complete continuity.
- 2. When an image on the film is obliterated with a round black mark, it is an indication of either blurred copy because of movement during exposure, duplicate copy, or copyrighted materials that should not have been filmed. For blurred pages, a good image of the page can be found in the adjacent frame. If copyrighted materials were deleted, a target note will appear listing the pages in the adjacent frame.
- 3. When a map, drawing or chart, etc., is part of the material being photographed, a definite method of "sectioning" the material has been followed. It is customary to begin filming at the upper left hand corner of a large sheet and to continue from left to right in equal sections with small overlaps. If necessary, sectioning is continued again-beginning below the first row and continuing on until complete.
- 4. For illustrations that cannot be satisfactorily reproduced by xerographic means, photographic prints can be purchased at additional cost and inserted into your xerographic copy. These prints are available upon request from the Dissertations Customer Services Department.
- 5. Some pages in any document may have indistinct print. In all cases the best available copy has been filmed.



. . -

Millie, David Fred

THE EFFECTS OF SILICA, NITROGEN, AND PHOSPHORUS LIMITATION ON THE BIOCHEMICAL COMPOSITION OF CYCLOTELLA MENEGHINIANA KUTZ: AN EXPERIMENTAL ANALYSIS

Iowa State University

Рн. Л. 1984

University Microfilms International 300 N. Zeeb Road, Ann Arbor, MI 48106

# Copyright 1984 by

Millie, David Fred

All Rights Reserved

.

# The effects of silica, nitrogen, and phosphorus limitation on the biochemical composition of <u>Cyclotella meneghiniana</u> Kütz.:

An experimental analysis

by

### David Fred Millie

### A Dissertation Submitted to the Graduate Faculty in Partial Fulfillment of the Requirements for the Degree of DOCTOR OF PHILOSOPHY

Department: Botany Major: Botany (Aquatic Plant Biology)

Approved:

Signature was redacted for privacy.

### In Charge of Major Work

Signature was redacted for privacy.

### For the Major Department

Signature was redacted for privacy.

For the Graduate College

Iowa State University Ames, Iowa

### 1984

Copyright C David Fred Millie, 1984. All rights reserved.

### © 1984

DAVID FRED MILLIE

All Rights Reserved

### TABLE OF CONTENTS

	Page
ABBREVIATIONS	iii
INTRODUCTION	1
LITERATURE REVIEW	4
METHODS	14
Experimental Design	16
Sample Analysis	22
Statistical Analysis	25
RESULTS	28
DISCUSSION	50
BIOTECHNICAL PERSPECTIVE	62
SUMMARY	65
LITERATURE CITED	67
ACKNOWLEDGMENTS	77
APPENDIX: MEAN SQUARE VALUES OF THE FACTORIAL ANALYSIS AND THE ANALYSIS OF VARIANCE	79

### ABBREVIATIONS

ANOVA	-	analysis of variance
ATP	-	adenosine triphosphate
Ch1/Ph	-	chlorophyll/phaeophytin
DNA	-	deoxyribonucleic acid
FAMEs	-	fatty acid methyl esters
FAS	-	fatty acid synthetase
gj	-	gigajoule
ha	-	hectare
L/G	-	lipid/glucan
LSD	-	least significant difference
m]	-	milliliter
Ν	-	nitrogen
Р	-	phosphorus
P/G	-	protein/glucan
Si	-	silica
Si(OH) <sub>4</sub>	-	silicic acid
yr	-	year

#### INTRODUCTION

The biochemical composition of algae has been studied extensively. This information is useful to researchers analyzing the suitability of algae as food sources in aquatic food chains. In addition, biochemical information may aid in the characterization of the physiological condition of natural algal populations and may provide phylogenetic implications (Darley, 1977).

In recent years, the concern over limited energy sources has prompted researchers to investigate algal biochemistry and physiology in a technological perspective. Some algae, particularly diatoms, accumulate lipid as a storage product. This storage product, coupled with the high photosynthetic efficiency of the algal cell, has prompted researchers to consider algae as a potential energy source. The high-energy lipid, when harvested, could be used as is or converted to hydrocarbon fuels (Bergeron <u>et al.</u>, 1983). The high-energy content of the lipid-enriched cells may also allow algae to serve as a potential food source.

The effect of nutrient concentration on algal biochemical composition is not well-documented. Researchers have reported that nutrient limitation generally causes a decrease in protein and photosynthetic pigments and an increase in lipid and carbohydrate storage products (Darley, 1977). However, most researchers have used batch cultures where algal cells are limited to a finite amount of nutrients. In batch cultures, cells progressively modify the nutrient concentration of the medium and exhibit exponential, linear, and stationary growth phases.

Any physiological measurements made on such a constantly changing population cannot be identified with a particular growth stage or nutrient concentration (Weete, 1980).

Continuous culture systems maintain a constant rate of nutrient supply coupled with a constant rate of cell removal. Under these conditions, cultures achieve a steady state where cell density, growth rate, and mean biochemical composition are constant. Changes in the rate of nutrient supply will cause changes in the growth rate, biochemical composition, and physiological state of the cells. Physiological measurements made on steady-state cells at different rates of nutrient supply would characterize a population at specific points along a growth curve and/or nutrient gradient.

Little information is available on the effect of nutrient limitation on the biochemical composition of algae under steady state conditions. Richardson <u>et al</u>. (1969) analyzed the amount of cellular lipid and chlorophyll of two green algae in response to increasing nitrogen concentration in a chemostat. Myklestad (1974), using continuous culture techniques, reported the amount of cellular carbohydrate and protein of a marine diatom in response to increasing nitrogen and phosphorus concentrations. To my knowledge, the amount of cellular lipid of diatoms under steady state conditions along a nutrient gradient has not been investigated.

The intent of this study was to analyze the effects of nutrient starvation and nutrient stress on the amount of cellular lipid, fatty acid, glucan, protein, and chlorophyll of a diatom. In this study,

the term "nutrient-starved cultures" refers to batch cultures in which the nutrient under investigation is deleted from the growth medium. The term "nutrient-stressed cultures" refers to semi-continuous cultures in which the nutrient under investigation is present within the growth medium, but at lower concentrations than necessary for maximum rates of growth. Silica (Si), nitrogen (N), and phosphorus (P) were chosen for investigation because other researchers have shown these nutrients to frequently limit diatom growth. The centric diatom, <u>Cyclotella</u> <u>meneghiniana</u> Kütz., was selected for study because it can be easily cultured in defined media and produces large amounts of lipid.

#### LITERATURE REVIEW

The importance of Si, N, and P in algal biochemistry and physiology is well-documented. Diatoms readily absorb Si as silicic acid  $(Si(OH)_4)$ and incorporate it in a polycondensed form as the main structural component of the cell wall (Werner, 1977). The amount of Si of diatoms typically comprises 10 to 30% of the cell's dry weight (Paasche, 1980), but may change with variations in the cell's growth rate, life cycle, structural features, and amount of surface area as well as the Si(OH)<sub>4</sub> concentration available for uptake (Eppley, 1977; Werner, 1977). Researchers have indicated that diatoms do not accumulate excess Si early in their development for later use in cell division (Darley, 1969; Paasche, 1980). However, Kilham <u>et al</u>. (1977) reported <u>Diatoma elongatum</u> can store up to 4.4 times more Si than needed for cell function.

The role of Si in diatom physiology is not well-understood. Besides its obvious function as a structural component, Si is also considered to be a required element for certain physiological functions. Metabolic Si has been reported to affect the citric acid cycle (between acetyl coenzyme A and 2-oxoglutarate), ATP oxidative phosphorylation, protein synthesis, respiration, chrysolaminaran utilization, chlorophyll synthesis, and DNA synthesis (Coombs <u>et al</u>., 1967b; Darley and Volcani, 1969; Azam <u>et al</u>., 1974; Werner, 1977; Sullivan and Volcani, 1981).

The role of N in algal physiology has been extensively studied (Morris, 1974; McCarthy, 1980). N is often reported to be the limiting nutrient in marine systems. Along with carbon, hydrogen, and oxygen, N

comprises a major portion of the dry matter of algal cells (Syrett, 1981). It is an essential constituent of important molecules such as amino acids, nucleic acids, and chlorophyll.

Algae have been reported to absorb N in the form of nitrate, nitrite, ammonia and urea (Lewin and Guillard, 1963; McCarthy, 1980). Amino acids have also been reported to be a N source for several diatoms (Hellebust and Lewin, 1977). The preferred order of N utilization appears to proceed from ammonia to urea to nitrate in both natural and culture systems (McCarthy, 1980). Morris (1974) stated that cellular growth rate is usually the same with either ammonia or nitrate as the N source. Paasche (1971), however, reported that <u>Dunaliella tertiolecta</u>, grew 30% faster with ammonia as the N source than with nitrate. Nitrate and nitrite must undergo an energy-requiring reduction prior to their incorporation into glutamate and amino acids (Syrett, 1962; Bidwell, 1974; Morris, 1974). Therefore, cells which can readily assimilate reduced forms of N would be expected to have increased growth rates because energy could then be utilized for cell synthesis rather than for cell maintenance.

The role of P in algal physiology has also been studied extensively (Kuhl, 1974; Nalewajko and Lean, 1980). Researchers often report P to be the nutrient responsible for increased algal productivity in freshwater systems. It is essential to all life forms as it functions in storage and transfer of metabolic energy (adenosine triphosphate) and is required for the synthesis of nucleotides, phospholipids, and other phosphorylated compounds.

Algae assimilate P in an inorganic soluble form such as orthophosphate (Wetzel, 1975). In some algae, organically-bound P appears to be the preferred form for P uptake (Nalewajko and Lean, 1980). Unlike N, however, P does not undergo any valency change during its assimilation and within the cell it forms polyphosphates. Many algae, when provided with sufficient P, can store excess P as polyphosphate bodies in the cytoplasm or vacuole. Stevenson and Stoermer (1982) reported these P "sinks" in algal cells to be related to the P concentration in the water. Tilman and Kilham (1976) reported that the diatom <u>Asterionella formosa</u> can store up to 82 times more P than needed for cell maintenance. These P "sinks" have great physiological and ecological significance for they allow the cell to continue normal metabolic functions even when P in the environment is limited (Kuhl, 1974).

Researchers have proposed that optimum cell nutrient ratios be used to indicate the limiting role of algal nutrients. The optimum cell nutrient ratio is the ratio at which transition from limitation by one nutrient to another takes place (Rhee and Gotham, 1980). Redfield (1958) reported a N:P ratio of approximately 15:1 for algal cells. However, cellular N:P ratios have been reported to range from 5.1 to 25 depending upon the taxon and the environmental conditions to which the cells are exposed (Parsons <u>et al</u>., 1961; Davis, 1976; McCarthy, 1980; Rhee, 1978; Rhee and Gotham, 1980). Generally optimum cellular N:P ratios appear to be high in green algae and low in diatoms (Rhee and Gotham, 1980).

Little information exists describing optimum cellular P:Si ratios

for diatoms. Parsons <u>et al</u>. (1961) reported P:Si ratios in <u>Skeletonema</u> <u>costatum</u> and <u>Coscinodiscus</u> sp. to be 0.19 and 0.02, respectively. Davis (1976), in a study of Si-limited cultures of <u>Skeletonema</u> <u>costatum</u> exposed to various light intensities, reported values which ranged from 0.27 to 1.2.

Whether lipid or carbohydrate is the photosynthetic reserve product of diatoms has long intrigued algal researchers. For many years, lipid (oil) was considered to be the only storage product of diatoms. The discovery that diatoms accumulated a  $\beta$ -1,3 linked glucyopyranoside during photosynthesis led some researchers to believe that carbohydrate was the main storage product. Most likely, diatoms can store both lipid and carbohydrate as photosynthetic reserve products with the exact abundance of each product dependent upon the physiological state of the cells (Barker, 1935; Fogg, 1956; Badour and Gergis, 1965).

Carbohydrate in diatoms is present primarily as reserve polysaccharides and cell wall constituents (Fogg, 1953). The principal monosaccharides are glucose, galactose, mannose, ribose, xylose, rhamnose, and fucose with glucose representing the greatest percent of total carbohydrate (Darley, 1977). Chrysolaminaran (leucosin) is the principal polysaccharide of diatoms and chrysophytes. It was first isolated from diatoms by von Stosch (1951). Beattie <u>et al</u>. (1961) and Ford and Percival (1965) reported that chrysolaminaran consists of approximately 12  $\beta$ -1,3 and  $\beta$ -1,6 linked glucose units in a ratio of 11:1. Chrysolaminaran is similar to laminaran, the photosynthetic reserve product of brown algae, but possesses a smaller ratio of  $\beta$ -1,3 to

 $\beta$ -1,6 linkages (11:1 vs. 15:1), a smaller glucose chain length (12 vs. 16), and lacks the terminal mannitol residues.

Chrysolaminaran has been identified as a photosynthetic reserve product in diatoms and chrysophytes. It was the radioactive product formed during photosynthesis in chrysophytes after exposure to  ${}^{14}\text{CO}_2$ (Kauss, 1962). Coombs and Volcani (1968) noted that the "main labelled soluble carbohydrate" after  ${}^{14}\text{CO}_2$  uptake by <u>Navicula pelliculosa</u> was a glucan. Handa (1969), in an analysis of carbohydrate metabolism, reported that the  $\beta$ -1,3 glucan of <u>Skeletonema costatum</u> was consumed during the dark.

Chrysolaminaran has been reported to accumulate in membrane-bound vesicles outside of, and directly abutting, the chloroplast (Duke and Reimann, 1977; Lee, 1980). Although chrysolaminaran can comprise 15 to 20% of the mass of the cell (Meeuse, 1962), it has rarely been mentioned in electron microscope studies of diatoms (Duke and Reimann, 1977). This is because chrysolaminaran is dissolved during the fixation and embedding process.

Lipids are broadly defined as materials which are soluble in organic solvents and are essentially insoluble in water (Stryer, 1981). The lipid composition of diatoms resembles that of green algae and higher plants. Lipid components of diatoms include triglycerides, sulfoquinavosyl diglyceride, digalactosyl diglyceride, monogalactosyl diglyceride, lecithin, phosphatidyl inositol, phosphatidyl glycerol, phosphatidyl ethanolamine, brassicosterol, clionosterol, and chondrillosterol (Low, 1955; Kates and Volcani, 1966; Kates and Volcani, 1968;

Tornabene et al., 1974).

Fatty acids are esterfied with glycerol (or other alcohols) to form the principal building blocks of polar and nonpolar lipids (Fisher and Schwarzenbach, 1978). Polar lipids, such as phospholipids, sulfolipids, and galactosyl lipids, are the major constituents of cell and organelle membranes. Most non-polar lipids are triglycerides (Otsuka and Morimura, 1966). Non-polar lipids serve as a photosynthetic reserve product (Fisher and Schwarzenbach, 1978) and accumulate in membrane-bound lipid "bodies" (Darley, 1977). Lipid "bodies" have been found in the chloroplast near the pyrenoids, in the central vacuole, and scattered throughout the cytoplasm (Drum, 1963; Stoermer <u>et al</u>., 1964; Stoermer <u>et al</u>., 1965; Crawford, 1973; Dawson, 1973; Duke and Reimann, 1977).

Fatty acid composition varies among algal groups. Diatoms characteristically possess fatty acids that have carbon chain lengths ranging from 12 to 22 (Ackman and Tocher, 1968; Wood, 1974). Diatoms generally possess large amounts of tetradecanoic ( $C_{14:0}$ ), hexadecanoic ( $C_{16:0}$ ), hexadecenoic ( $C_{16:1}$ ), eicosatetraenoic ( $C_{20:4}$ ) and eicosapentanaenoic ( $C_{20:5}$ ) and small amounts of hexadecadienoic ( $C_{16:2}$ ), hexadecatrienoic ( $C_{16:3}$ ), octadecenoic ( $C_{18:1}$ ), octadecadienoic ( $C_{18:2}$ ), and ocatdecatrienoic ( $C_{18:3}$ ) acids (Ackman <u>et al</u>., 1964; Brockerhoff <u>et al</u>., 1964; Kates and Volcani, 1966; Ackman and Tocher, 1968; Kates and Volcani, 1968; Pugh, 1971; DeMort <u>et al</u>., 1972; Opute, 1974a; Tornabene <u>et al</u>., 1974; Fisher and Schwarzenbach, 1978).

Several researchers have indicated that the biochemical composition

of algae is variable and dependent on the cellular growth stage. In exponentially growing populations, the main biochemical "product" is protein (Olive <u>et al.</u>, 1969). Darley (1977) stated that in "nutrient sufficient cells, protein content almost always exceeds carbohydrate content, which, in turn, usually exceeds the lipid content." Morris (1981) reported the amount of cellular protein in exponentially-growing diatom cultures ranged from 17 to 35% of the cell dry weight. The amount of cellular carbohydrate and lipid ranged from 4 to 20% and 2 to 7%, respectively.

Researchers have reported algal fatty acid composition to vary as cells age. Collyer and Fogg (1955) reported the amount of fatty acid per culture increased or decreased with age depending upon the taxon. Pugh (1971) reported an increase in fatty acids that have carbon chain lengths of 18 as cells age. Dodecanoic acid ( $C_{12:0}$ ) decreased with cell senescence. Ackman <u>et al</u>. (1964) noted that tetradecanoic acid ( $C_{14:0}$ ) increased and eicosapentaenoic acid ( $C_{20:5}$ ) decreased as cells age.

The amount of cellular carbohydrate and lipid is greater in senescent cells than in actively-growing cells (Badour and Gergis, 1965; Handa, 1969; Opute, 1974b; Conover, 1975). The amount of cellular protein remains constant (Badour and Gergis, 1965; Handa, 1969) or decreases (Myklestad and Haug, 1972; Conover, 1975) with cell senescence. The amount of cellular chlorophyll content is lower in senescent cells than in young, actively-growing cells (Antia <u>et al.</u>, 1963; Ackman and Tocher, 1968).

Algal physiology and biochemistry have been reported to be affected

by the Si, N, and P concentrations of the growth medium. Diatom cell division is inhibited as Si is depleted from the medium (Eppley, 1977). In a Si-starved culture of <u>Navicula pelliculosa</u>, cell division stopped but cellular morphological development continued until the culture consisted mostly of biprotoplastic cells (two daughter protoplasts contained within a parent frustule, each surrounded by a new plasmalemma and separated by an intercellular space). In these cells, the rate of carbon and P uptake was lower than that in Si-replete conditions (Coombs et al., 1967a,b; Coombs and Volcani, 1968).

Diatoms show a gradual decrease in major metabolic activity as Si becomes limiting (Darley, 1977). Protein, chlorophyll, carbohydrate, and nucleic acid synthesis are partially, or completely, inhibited. Lipid synthesis increases as Si becomes limited (Coombs <u>et al.</u>, 1967a,b; Healey <u>et al.</u>, 1967; Coombs and Volcani, 1968). Darley (1977) noted a 60 to 70% reduction in glycolysis when <u>Cyclotella cryptica</u> cells were transferred to Si-deficient media. The rate of fatty acid synthesis, however, increased by greater than 100%.

When Si is introduced into Si-starved cultures, a reduction in cell lipid synthesis and an increase in cell carbohydrate, protein, and DNA synthesis occurs. Repletion of Si also causes an increase in cell nucleosidetriphosphate and a resumption of cell division (Coombs <u>et al.</u>, 1967a; Coombs and Volcani, 1968; Darley, 1969; Darley, 1977).

The effects of N limitation on algal cells have been well-documented. Eppley and Renger (1974) and Opute (1974b) observed a reduction in photosynthetic efficiency when actively-growing cells were exposed to N

limitation. The amount of cellular lipid of diatoms and green algae also increased as N becomes limiting (Fogg, 1956; Badour and Gergis, 1965; Richardson <u>et al</u>., 1969; Werner, 1977; Shifrin and Chisholm, 1981; Rosen, 1982; Bergeron <u>et al</u>., 1983). Collyer and Fogg (1955) stated that accumulation of lipid is not associated with low cellular N in red and blue-green algae. However, de Vasconcelos and Fay (1974) reported increased lipid content in a blue-green alga during N starvation.

The amount of fatty acids in algal cells has been reported to increase as N becomes limiting (Zhukova <u>et al.</u>, 1968; Klyachko-Gurvich <u>et al.</u>, 1969). Cells exposed to N limitation predominantly possess large concentrations of saturated or monounsaturated fatty acids that have carbon chain lengths of 16 and 18. Cells exposed to high concentrations of N predominantly possess polyunsaturated fatty acids that have carbon chain lengths of 16 and 18 (Ackman <u>et al.</u>, 1964; Matsuka <u>et al.</u>, 1966; Klyachko-Gurvich <u>et al.</u>, 1969; DeMort <u>et al.</u>, 1972; Pohl, 1974).

The amount of cellular carbohydrate in diatoms and green-algae has also been reported to increase as N becomes limiting (Matsuka <u>et</u> <u>al.</u>, 1966; Klyachko-Gurvich <u>et al.</u>, 1969; Myklestad and Haug, 1972; Werner, 1977). The amount of cellular protein and chlorophyll has been observed to decrease as N becomes limiting (Fogg, 1956; Matsuka <u>et al.</u>, 1966; Thomas and Dodson, 1972; Eppley and Renger, 1974; Morris <u>et al.</u>, 1974; Myklestad, 1974; Syrett, 1981). Syrett (1981), however, reported a study of <u>Thalassiosira</u> in which N deprivation caused a decrease in

photosynthetic rate but little change in the amount of cellular chlorophyll.

The biochemical composition of algae is also affected by P limitation. Researchers observed an accumulation of cellular lipid and carbohyrate only after all cellular P reserves had been utilized (Kuhl, 1974; Werner, 1977; Rosen, 1982). Rosen (1982) reported a decrease in the amount of cellular chlorophyll of <u>Cyclotella meneghiniana</u> as P becomes limiting. Spoehr and Milner (1949) observed no biochemical differences between P-stressed and P-sufficient cells of <u>Chlorella</u>.

#### METHODS

Cells of <u>Cyclotella meneghiniana</u> Kütz. were collected from the Des Moines River (Boone County, IA) on 18 May, 1982. Several cells were isolated into soil water extract (Nichols, 1973) using a micropipette. A clonal culture was obtained by washing cells through 4 distilled water baths and reisolating a single cell into a defined medium (Table 1). This clone was used in all subsequent cultures. Diatoms were grown in 350 ml of culture media in 500 ml, polycarbonate, Erlenmeyer flasks. Cultures were maintained in a Percival Growth Chamber (Boone, IA) at  $18 \pm 2^{\circ}$ C. Six parallel 40-watt cool white fluorescent lamps alternated with 6 34-watt warm white incandescent lamps provided approximately 450 foot candles of illumination (approximately 0.1 mMoles photons  $(M^2)^{-1}$ .  $day^{-1}$ ). Cultures were exposed to a 16:8 hr light:dark cycle in an attempt to obtain synchronously-dividing cultures and thus minimize cell variability (Cook, 1961). Diatom cultures were shaken daily to keep cells suspended in the medium.

<u>Cyclotella</u> cultures were unialgal, but not axenic. To minimize further contamination of the cultures, sterile techniques and equipment were utilized throughout the study. Glass, double-distilled water was used in the preparation of the culture media. Media solutions minus vitamins were prepared for maintenance cultures by autoclaving the solution 15 minutes at  $250^{\circ}$ C. Vitamins were added aseptically with a Gelman 0.2 µm mesh Acrodisc filter assembly attached to a syringe. For nutrient-starvation experiments, the treatment media were filtered through a Gelman 0.2 µm mesh Metricel membrane filter into a sterilized flask.

Nutrient	mg/liter	Nutrient	mg/liter
Macronutrients:		Micronutrients:	
CaCl <sub>2</sub> ·2H <sub>2</sub> O	36.76	Na <sub>2</sub> •EDTA	4.36
MgS0 <sub>4</sub> •7H <sub>2</sub> 0	36.97	FeC13•6H20	3.15
NaHCO3	12.60	CuS0 <sub>4</sub> • 5H <sub>2</sub> 0	0.01
K <sub>2</sub> HPO4 <sup>a,b</sup>	8.71	ZnS04•7H20	0.022
NaNO3 <sup>C,d</sup>	85.01	CoC1 <sub>2</sub> •6H <sub>2</sub> 0	0.01
NaSi0 <sub>3</sub> •9H <sub>2</sub> 0 <sup>e,f</sup>	28.42	$MnC1_2 \cdot 4H_20$	0.18
NaCl	584.50	Na2MoO4 • 2H2O	0.006
к <sub>2</sub> нсо <sub>3</sub> д	6.91	H <sub>3</sub> B0 <sub>3</sub>	1.0
Vitamins:		Buffer:	
Thiamine•HC1	0.1	Tris(hydroxymeth	iyl)
Biotin	0.0005	aminomethane	500.00
Cyanocobalamin	0.0005		

Table 1. Composition of culture medium: modified from Guillard and Lorenzen (1972) and Rosen (1982)

<sup>a</sup>Omitted from P-deficient cultures in nutrient starvation experiments.

<sup>b</sup>Used at 1/10th strength in P-limited media in nutrient stressed experiments.

<sup>C</sup>Omitted from N-deficient cultures in nutrient starvation experiments.

<sup>d</sup>Used at 1/10th strength in N-limited media in nutrient stressed experiments.

<sup>e</sup>Omitted from Si-deficient cultures in nutrient starvation experiments.

<sup>f</sup>Used at 1/10th strength in Si-limited media in nutrient stressed experiments.

<sup>g</sup>Used in P-deficient cultures.

Treatment media in nutrient-stressed experiments were prepared and autoclaved in 2.5 liter polycarbonate bottles. Vitamin concentrations in these bottles were increased 2.5-fold over normal strength to overcome degradation of vitamins during autoclaving.

### Experimental Design

Diatom cultures were subjected to Si-, N-, and P-starvation and to Si-, N-, and P-stressed experiments. In nutrient-starvation experiments, diatom cells were removed from nutrient-replete batch cultures and resuspended in nutrient media with either  $NaSiO_3 \cdot 9H_2O$ ,  $NaNO_3$ , or  $K_2HPO_4$  omitted. Since the only potassium source normally in the medium was  $K_2HPO_4$ , P-deficient media would have lacked potassium also. Therefore, an appropriate amount of potassium was added to the P-deficient media in the form of  $K_2HCO_3$  (Table 1).

In nutrient-starvation experiments, a  $2^3$  factorial experimental design in fractional replication (Cochran and Cox, 1957) was used. In this type of design, the three nutrients in various combinations produced 8 possible nutrient treatments (Table 2). However, due to the number of variables analyzed per flask and the number of flasks considered, all 8 treatments could not be analyzed collectively. Rather, only half of the total treatments were considered at one time. Each pair of 4 treatments (hereafter referred to as a half-replicate) was selected to minimize experimental "confounding" of nutrient effects (Table 2). The 2 halfreplicates were then put together to make a single replication of the factorial design.

In a single replication of the  $2^3$  factorial design, each nutrient

is indicated by a plus (+) or minus (-). Half-replicates
of an asterisk (*)

Table 2. Nutrient combinations in media treatments for nutrient-

<b>-</b>	Nutrient			
Ireatment	Silica	Nitrogen	Phosphorus	
]	-	-		
2*	+	-	-	
3*	-	+	-	
4	+	+	-	
5*	-	-	+	
6	+	-	+	
7	-	+	+	
8*	+	+	+	

effect was an average over 4 combinations of other factors and had, in effect, 4-fold replication (Cochran and Cox, 1957). Variable means for nutrient deficient effects were calculated using treatments in which the nutrient under consideration was omitted. Variable means for nutrient-replete effects were calculated using treatments in which the nutrient under consideration was present.

A schematic representation of the experimental procedure used in nutrient-starvation experiments is presented in Fig. 1. For each halfreplicate, 10 culture flasks containing nutrient sufficient media were inoculated with equal amounts of cells from a maintenance culture. Cultures were placed in the growth chamber and allowed to grow until cell density reached approximately 14,000 cells/ml. At this density, cell division was still occurring, but at a much reduced rate than cells of cultures undergoing exponential growth.

Two of the flasks were then analyzed. The day in which experimental analysis began was designated Day O. The remaining 8 flasks were randomly divided into 4 pairs and each pair was assigned to 1 of the 4 possible media treatments in the half-replicate considered. The entire volume of each flask was centrifuged at approximately 3,000 rpm for 4 minutes. Centrifuged cells were washed 4 times in their assigned treatment medium and resuspended in flasks containing the appropriate treatment medium. Cultures were replaced in the growth chamber. Cultures were analyzed 3 and 9 days after transfer to treatment media, a time sequence used by Shifrin and Chisholm (1981). On day 3, 1 flask from each of the 4 treatment pairs was removed for analysis. On day 9,

Fig. 1. Schematic representation of the experimental procedure used for each treatment in nutrientstarvation experiments

ı



### DAY 3

DAY 9







the four remaining flasks were analyzed. For the nutrient-starvation experiment, a single replication of the factorial design was completed.

In nutrient-stressed experiments, semi-continuous culture techniques were used. Cultures were diluted by removing a certain percentage of the culture at regular intervals (hereafter referred to as dilution rate) and replacing it with an equal amount of fresh media. Cell densities of semi-continuous cultures are inversely proportional to the dilution rate. A net decrease in cell density occurs when dilution rate is greater than growth rate. If dilution rate exceeds growth rate for a sufficient period of time, the cell population "washes out" of the culture flask. A net increase in cell density occurs when growth rate is greater than dilution rate. Cell density increases until dilution rate equals growth rate. Consequently, a steady-state cell density is achieved. At steady state, the population possesses a constant growth rate and constant mean biochemical characteristics. High dilution rates produce steady state populations possessing low cell densities and, therefore, a high nutrient supply per cell. Low dilution rates, in turn, produce steady-state populations possessing high cell density and a low nutrient supply per cell. By varying the dilution rate, the culture can be "held" at various points along its growth curve (Fogg, 1965). This allows analysis of the physiological and biochemical condition of steady-state populations along a nutrient gradient.

The molar ratio of N:P:Si was 1.0:0.05:0.1 in media used for maintenance cultures. For nutrient-stressed experiments, culture media were prepared with the nutrient under investigation at 1/10th normal

strength (Table 1). Preliminary experimentation indicated that cultures in media with Si, N, and P at 1/10th normal strength were nutrient limited. Cell densities of cultures for all dilution rates were less than similarly diluted cultures using normal strength media. In this manner, the N:P:Si molar ratio was 1.0:0.05:0.01 in Si-stressed media, 0.1:0.05:0.1 in N-stressed media, and 1.0:0.005:0.1 in P-stressed media.

In each of the nutrient-stressed experiments, cells from a maintenance culture growing exponentially were centrifuged and washed 3 times in the appropriate medium. Aliquots of the washed cell suspension were resuspended in 9 flasks containing nutrient stressed media. Cultures were placed in the growth chamber. Cells within the flasks were allowed to divide without interruption for 3 or 4 days, after which the dilution process was initiated. The 9 flasks were randomly divided into 3 groups and a daily dilution rate of 10%, 25%, or 50% assigned to the groups. Dilutions were continued until a steady state cell density was determined by microscopic counting. When steady state conditions were reached, the daily supply of limiting nutrient per cell for flasks within a dilution rate was calculated (Table 3). Flasks were then analyzed.

### Sample Analysis

Diatom cultures were analyzed for the amount of cellular lipid, glucan, protein, chlorophyll, and phaeophytin. Samples were obtained by centrifuging a known volume of the cultures at approximately 2,000 rpm for 3 minutes. Examination of centrifuged material by phase contrast microscopy revealed some bacteria, but their biomass was insignificant in comparison to that of diatom cells. Samples for glucan

Limiting nutrient	Dilution rate (%·day <sup>-1</sup> )	Supply rate (mg x 10-7.cell- <sup>1</sup> .day <sup>-1</sup> )
	10	0.140 (± 0.009)
Silica	25	0.449 (± 0.051)
	50	1.178 (± 0.147)
	10	0.126 (± 0.012)
Nitrogen	25	0.868 (± 0.012)
	50	3.475 (± 0.137)
	10	0 013 (+ 0 003)
	10	$0.013 (\pm 0.002)$
Phosphorus	25	$0.064 (\pm 0.014)$
	50	0.434 (± 0.019)

Table 3. Supply of limiting nutrients per cell per day for dilution rates of media in nutrient-stressed experiments. Means and standard errors (in parentheses) for replicate flasks are presented analysis were obtained by filtering a known volume of culture directly onto Whatman 934-AH glass microfibre filters. Culture cell densities were determined by microscopic counting techniques utilizing a Palmer-Maloney chamber (Guillard, 1973).

Total lipid was extracted from diatom samples by aqueous alkaline hydrolysis (modified from Hammond <u>et al.</u>, 1981). Diatom cells were refluxed for approximately  $1\frac{1}{2}$  hours in 10 ml of 12% alcoholic KOH. The digested cells were cooled, adjusted to pH 3 with 1N HCl, and diluted with 50 ml of distilled water. Ten ml of hexane was added to the reflux flask. The solution was mixed thoroughly and filtered through a Whatman GF/C glass microfibre filter to remove diatom frustules. The reflux flask was rewashed with 5 ml of hexane and the filtering repeated. The filtered solution was transferred to a separatory funnel fitted with a teflon stopcock. The hexane layer was allowed to separate, removed from the funnel, and transferred to a tared glass vial. The hexane was evaporated under N gas and the resulting residue and vial were weighed. Total lipid was expressed gravimetrically as lipid per cell.

Total glucan was extracted and determined spectrophotometrically by the sulfuric acid-phenol method presented by Myklestad (1978). Total protein was determined by the "Lowry Method" (Lowry <u>et al.</u>, 1951) using bovine serum albumin as a standard. Chlorophyll and phaeophytin values were determined by spectrophotometric methods (Richards with Thompson, 1952) using the equations of Parsons and Strickland (1963). Values for glucan, protein, chlorophyll, and phaeophytin were expressed on a per cell basis. Chlorophyll/phaeophytin (Chl/Ph), protein/glucan (P/G),

and lipid/glucan (L/G) ratios were also calculated.

The fatty acid composition of lipid residues from cells grown in nutrient-stressed experiments was analyzed. The lipid residues for replicate samples within a supply rate were resuspended in hexane, pooled, and re-evaporated under N gas. The fatty acids were converted to fatty acid methyl esters (FAMEs) by adding 1 ml of 3%  $\rm H_2SO_4$  in methanol to the residue and heating in a teflon-sealed vial at  $55^{\circ}$ C for 12 hours. After heating, 3 ml of distilled water and 1 ml of hexane were added to the vial. The solution was mixed and centrifuged. The hexane layer was removed and washed 3 or 5 times with distilled water to remove traces of methanol. The hexane was evaporated to near dryness to concentrate the FAMEs. One  $\mu$ l of sample was injected into a Beckman GC-5 gas chromatograph equipped with a 183 x 0.3 cm stainless steel column packed with 15% EGSSX on 100/120 mesh Chromosorb P (Applied Science Laboratories, State College, PA). The carrier gas was N at a flow rate of 50 ml/minute. The column temperature varied between 139<sup>0</sup> and 176°C, depending upon the date of analysis. Retention peaks of sample FAMEs were compared to standard FAME mixtures (Supelco, Inc., Bellefonte, PA) for identification. The fatty acid samples from Pstressed experiments were inadvertently destroyed in preparation.

### Statistical Analysis

For nutrient-starvation experiments, values of biochemical variables and the calculated ratios for flasks on days 3 and 9 were compared by a factorial analysis using a split-plot, randomized block design

(Cochran and Cox, 1957). Since only cells that had been cultured in nutrient-sufficient media were used as inoculum for the experimental flasks, no comparison to nutrient-deficient treatments could be made for day 0. Therefore, values for this initial day of experimentation were not included in the statistical analysis. The sources of variation investigated in the factorial analysis are presented in Table 4. Halfreplicates were considered blocks. The significance of nutrient effects (Si, N, and P) and their 2-factor interactions (Si·N, Si·P, and N·P) were assessed by a Half-Normal Probability Plot (Daniel, 1959).

For each nutrient-stressed experiment, values of biochemical variables and the calculated ratios were compared in terms of nutrient supply rate by an Analysis of Variance (ANOVA) (Snedecor and Cochran, 1980). The sources of variation investigated in the ANOVA are presented in Table 5. The effect of nutrient supply rate was subdivided to analyze for deviation from linearity. The significance between specific pairs of variable means was analyzed by a Least Significant Difference (LSD) analysis (Snedecor and Cochran, 1980).

Table 4.	Sources of variation with their corresponding degrees of	
	freedom investigated in the Factorial Analysis performed on	
	variables of nutrient-starvation experiments	

Source of variation	Degrees of freedom
Fractional replicate	1
Silica	(3)
Nitrogen	i
Phosphorus	1
Error (A) <sup>a</sup>	3
Days	1
Interaction	(3)
Days.Silica	1
Days•Nitrogen	1
Days·Phosphorus	1
Error (B) <sup>D</sup>	4

a(Silica·Nitrogen + Silica·Phosphorus + Nitrogen·Phosphorus). b(Days·Error (A) + Days·Fractional replicate).

Table 5. Sources of variation and their corresponding degrees of freedom investigated in the Analysis of Variance performed on variables of nutrient-stressed experiments

Source of variation	Degrees of freedom
Nutrient supply rate	(2)
Linear	1
Quadratic	1
Error (replicates)	6 <sup>a</sup>

<sup>a</sup>Value expressed is based on 3 replicates per nutrient supply rate.
## RESULTS

The biochemical composition of the cells varied little throughout the nutrient-starvation experiments. The amount of cellular lipid (Fig. 2) generally ranged from 2.0 to 4.0 x  $10^{-3} \ \mu g \cdot cell^{-1}$ . The amount of cellular lipid initially appeared to increase in cells grown in Nsufficient media and decrease in N-deficient media. The amount of cellular glucan remained fairly constant (approximately 0.6 to 0.8 x  $10^{-3} \ \mu g \cdot cell^{-1}$ ) in all treatments (Fig. 3). The amount of cellular protein varied greatly between Si-sufficient and Si-deficient treatments (Fig. 4). In Si-deficient treatments, the amount of cellular protein increased from 3.1 to 5.3 x  $10^{-1} \ \mu g \cdot cell^{-1}$  between days 0 and 3. The amount of cellular protein then remained constant until day 9. The amount of cellular protein in N and P treatments remained fairly constant (3.0 to 4.0 x  $10^{-1} \ \mu g \cdot cell^{-1}$ ) throughout the study. The amount of cellular chlorophyll varied little (1.0 to  $1.25 \times 10^{-5} \ \mu g \cdot cell^{-1}$ ) throughout all nutrient treatments (Fig. 5).

The calculated biochemical ratios also varied little throughout the nutrient-starvation experiments. Chl/Ph ratios remained fairly constant (approximately 1.7 to 2.5) over all treatments (Table 6). L/G ratios (Table 7) ranged from 1.5 to 7.0 with the majority of values approximately 5.0. P/G ratios ranged from 300 to 1900 among all nutrient treatments (Table 8).

A factorial analysis using a split-plot, randomized block design was used to analyze the effects of half-replicates and nutrient effects on

Fig. 2. Changes in the amount of cellular lipid in silica, nitrogen, and phosphorus-sufficient (solid lines) and deficient (dashed lines) cultures through time. Values represent mean of four replicate flasks. Bars indicate standard error of the means



Fig. 3. Changes in the amount of cellular glucan in silica, nitrogen, and phosphorus-sufficient (solid lines) and deficient (dashed lines) cultures through time. Values represent mean of four replicate flasks. Bars indicate standard error of the means

.



Fig. 4. Changes in the amount of cellular protein in silica, nitrogen, and phosphorus-sufficient (solid lines) and deficient (dashed lines) cultures through time. Values represent mean of four replicate flasks. Bars indicate standard error of the means



Fig. 5. Changes in the amount of cellular chlorophyll in silica, nitrogen, and phosphorus-sufficient (solid lines) and deficient (dashed lines) cultures through time. Values represent mean of four replicate flasks. Bars indicate standard error of the means



Nutriont	Nutrient	Day			
nucrienc	availability	0	3	9	
Silica	+ -	2.46	2.08 1.84	2.33 2.16	
Nitrogen	+ -	2.46	1.68 2.24	2.49 2.00	
Phosphorus	+ -	2.46	2.34 1.57	1.66 2.82	

Table 6. Chlorophyll/phaeophytin ratios for replicate flasks in nutrient-starvation experiments. Means of replicate flasks are

Table 7. Lipid/glucan ratios for replicate flasks in nutrient-starvation experiments. Means of replicate flasks are presented

Nutrient	Nutrient availability	Day			
		0	3	9	
Silica	+ -	5.13	5.06 4.20	1.51 6.40	
Nitrogen	+ -	5.13	6.97 2.29	2.37 5.55	
Phosphorus	+ -	5.13	4.46 4.80	2.62 5.29	

Table 8. Protein/glucan ratios for replicate flasks in nutrientstarvation experiments. Means of replicate flasks are presented

Nutniont	Nutrient availability	Day			
Nucrienc		0	3	9	
Silica	+ -	377.88	303.27 527.46	255.16 1926.31	
Nitrogen	+ -	377.88	361.21 434.52	400.92 1780.55	
Phosphorus	+ -	377.88	386.32 444.42	440.69 1740.78	

biochemical variables and their calculated ratios between days 3 and 9 (Appendix Table A1). The Half-Normal Probability Plot indicated no significant effect of 2-factor nutrient interactions on the biochemical composition of the cells. Therefore, only the effects of individual nutrients were considered in the factorial analysis.

In this factorial analysis, the fractional-replicate effect measured the variability of values between fractional replicates. This effect, in theory, measured the reproducibility of values between halfreplicates. Only the values for the amount of cellular protein were significantly different between the 2 half-replicates (p<0.05). The limiting-nutrient effect indicated the variability between the mean of values on days 3 and 9 in nutrient-sufficient treatments and the mean of values on days 3 and 9 in nutrient-deficient treatments. Only Si had a significant effect on a biochemical variable. A significant increase in the amount of cellular protein occurred with Si deficiency. The effect of days measured the variability between the mean of values on day 3 and the mean of values on day 9. Interaction effects measured the variability between the difference of nutrient-deficient and nutrient-sufficient treatments on day 3 and this difference on day 9. Neither day nor day.nutrient interactions significantly affected any biochemical variable investigated.

The biochemical composition of the cells varied in response to nutrient supply rate (dilution rate) within and between nutrient-stressed experiments (Figs. 6 to 8). An ANOVA was used to measure the effect of supply rate on the biochemical composition of the cells (Appendix Table

Fig. 6. The amount of cellular lipid, glucan, protein, and chlorophyll in silica-stressed cultures as a function of silica supply rate. Values represent mean of replicate flasks. Bars indicate standard error of the means



Fig. 7. The amount of cellular lipid, glucan, protein, and chlorophyll in nitrogen-stressed cultures as a function of nitrogen supply rate. Values represent mean of replicate flasks. Bars indicate standard error of the means

.



Fig. 8. The amount of cellular lipid, glucan, protein, and chlorophyll in phosphorus-stressed cultures as a function of phosphorus supply rate. Values represent mean of replicate flasks. Bars indicate standard error of the means

-



A2). A LSD analysis was used to test the significance between specific pairs of variable means within an experiment (Table 9).

The LSD analysis indicated that increasing Si supply rate caused a significant decrease in the amounts of cellular lipid and protein and cellular L/G ratios. However, the ANOVA revealed no significant effects of Si supply rate on any variable except the L/G ratio (p<0.05). Cellular L/G ratios decreased from 2.6 to 1.2 between 0.140 and 0.449 x  $10^{-7}$  mg Si·cell<sup>-1</sup>·day<sup>-1</sup> but remained constant between 0.449 and 1.178 x  $10^{-7}$  mg Si·cell<sup>-1</sup>·day<sup>-1</sup>.

The LSD analysis indicated that increasing N supply rate caused significant increases in the amount of cellular lipid, glucan, protein, and chlorophyll. The ANOVA revealed significant effects of N supply rate on all of these variables except chlorophyll (p<0.001). The amount of cellular lipid remained constant (approximately 3.4 x  $10^{-3} \,\mu\text{g}\cdot\text{cell}^{-1}$ ) between 0.126 and 0.868 x  $10^{-7} \,\text{mg N}\cdot\text{cell}^{-1}\cdot\text{day}^{-1}$  but increased to 9.6 x  $10^{-3} \,\mu\text{g}$  at 3.475 x  $10^{-7} \,\text{mg N}\cdot\text{cell}^{-1}\cdot\text{day}^{-1}$ . The amount of cellular glucan remained constant (approximately 1.1 x  $10^{-3} \,\mu\text{g}\cdot\text{cell}^{-1}$ ) between 0.126 and 0.868 x  $10^{-7} \,\text{mg N}\cdot\text{cell}^{-1}\cdot\text{day}^{-1}$ . The amount of cellular glucan remained constant (approximately 1.1 x  $10^{-3} \,\mu\text{g}\cdot\text{cell}^{-1}$ ) between 0.126 and 0.868 x  $10^{-7} \,\text{mg N}\cdot\text{cell}^{-1}\cdot\text{day}^{-1}$  but increased to 3.3 x  $10^{-3} \,\mu\text{g}$  at 3.475 x  $10^{-7} \,\text{mg N}\cdot\text{cell}^{-1}\cdot\text{day}^{-1}$  but increased from 1.95 to 4.11 x  $10^{-1} \,\mu\text{g}\cdot\text{cell}^{-1}$  between 0.126 and 0.868 x  $10^{-7} \,\text{mg}$  N·cell<sup>-1</sup> day<sup>-1</sup> to 6.7 x  $10^{-1} \,\mu\text{g}\cdot\text{cell}^{-1}$  at 3.475 x  $10^{-7} \,\text{mg N}\cdot\text{cell}^{-1}\cdot\text{day}^{-1}$ .

The LSD analysis indicated that significant increases in the amount of cellular lipid and glucan occurred with increasing P supply rate. The ANOVA also indicated significant increases in the amount of cellular lipid (p<0.001) and glucan (p<0.001). The amount of cellular lipid

Supply Limiting rate		Cellular composition			<u>Cellular ratio</u>			
nutrient	(mg x 10-7. ce11-1.day-1)	Lipid (µg x 10-3)	Glucan (µg x 10-3)	Protein (µg x 10-3)	Chloro- phyll (µgxl0 <sup>-3</sup> )	Chlorophyll/ Phaeophytin	Protein/ Glucan	Lipid/ Glucan
	0.140	8.62 <sup>a</sup>	3.33	2.13 <sup>a</sup>	0.95	No data	68.4	2.6 <sup>a,b</sup>
Silica	0.449	5.37 <sup>a</sup>	4.79	1.17 <sup>a</sup>	1.05	No data	28.1	1.2 <sup>a</sup>
	1.178	6.60	5.45	1.33	0.94	No data	30.4	1.3 <sup>b</sup>
	0.126	2.10 <sup>a</sup>	0.66 <sup>a</sup>	1.95 <sup>a</sup>	0.91 <sup>a</sup>	7.1	298.7	3.1
Nitrogen	0.868	4.67 <sup>b</sup>	1.58 <sup>b</sup>	4.11 <sup>a,b</sup>	1.20 <sup>a</sup>	5.1	263.4	3.0
-	3.475	9.63 <sup>a,b</sup>	3.33 <sup>a,b</sup>	6.71 <sup>a,b</sup>	1.03	5.3	211.7	3.0
	0.013	2.59 <sup>a</sup>	0.72 <sup>a</sup>	2.56	0.75	No data	363.6	3.9
Phosphorus	s 0.064	2.68 <sup>b</sup>	1.09 <sup>b</sup>	3.73	1.04	No data	363.6	2.6
	0.434	7.35 <sup>a,b</sup>	3.49 <sup>a,b</sup>	4.24	1.05	No data	121.8	2.1

Table 9. Mean values of the cellular biochemical composition and the cellular biochemical ratios for nutrient supply rates in nutrient-stressed experiments

 $^{a,b}$ Variable means within a nutrient category with similar superscripts are significantly different (p<0.05).

.

remained constant (approximately 2.63 x  $10^{-4} \ \mu g \cdot cell^{-1}$ ) between 0.013 and 0.064 x  $10^{-7} \ mg \ P \cdot cell^{-1} \cdot day^{-1}$ . The amount of cellular glucan remained constant (approximately 0.91 x  $10^{-3} \ \mu g \cdot cell^{-1}$ ) between 0.013 and 0.064 x  $10^{-7} \ mg \ P \cdot cell^{-1} \cdot day^{-1}$  but increased to 3.5 x  $10^{-3} \ \mu g \cdot cell^{-1}$ at 0.434 x  $10^{-7} \ mg \ P \cdot cell^{-1} \cdot day^{-1}$ .

The fatty acids of lipid extracted in N-stressed and Si-stressed experiments were tetradecanoic  $(C_{14:0})$ , pentadecanoic  $(C_{15:0})$ , hexadecanoic  $(C_{16:0})$ , heptadecanoic  $(C_{17:0})$ , heptadecanoic-branched  $(C_{17:branched})$ , octadecanoic  $(C_{18:0})$ , octadecenoic  $(C_{18:1})$ , octadecadienoic  $(C_{18:2})$ , octadecatrienoic  $(C_{18:3})$ , eicosanoic  $(C_{20:0})$ , eicosenoic  $(C_{20:1})$ , eicosatetraenoic  $(C_{20:4})$ , docosadienoic  $(C_{22:2})$ , docosatetraenoic  $(C_{22:4})$ , and tetracosanoic  $(C_{24:0})$ . Although FAME samples were washed several times with distilled water, there was, probably, a retention of methanol in the sample. This caused a gradual slope off the solvent peak during gas chromatographic analysis which did not allow the area under the FAME retention peaks to be measured. Therefore, the relative percentage of each FAME in the lipid sample could not be calculated. Rather, only the presence or absence of a FAME could be determined.

The fatty acid composition of the extracted lipid differed little among nutrient supply rates within an experiment but was distinctly different between N-stressed and Si-stressed experiments (Tables 10 and 11). In the N-stressed experiment, carbon chain length of fatty acids ranged from 14 to 24. Except for the absence of  $C_{15:0}$ ,  $C_{17:0}$ , and  $C_{24:0}$ at 0.126 x 10<sup>-7</sup> mg N·cell<sup>-1</sup>·day<sup>-1</sup> and  $C_{20:4}$  and  $C_{24:0}$  at 0.868 x 10<sup>-7</sup>

	Supply rate (x 10 <sup>-7</sup> mg N·cell <sup>-1</sup> ·day <sup>-1</sup> )				
rally acto	0.126	0.868	3.475		
14:0 <sup>a</sup>	+	+			
15:0		+			
16:0	+	+	+		
17:branched	+	+			
17:0		+			
18:0	+	+	+		
18:1	+	+	+		
18:2	+	+	+		
20:0	+	+			
20:1 (or 2)	+	+	+		
20:4 (or 5)	+		+		
22:2	+	+	+		
22:4 ?	+	+	+		
24:0 ?			+		

Table 10. Fatty acid composition of <u>Cyclotella meneghiniana</u> for supply rates of nitrogen-stressed cultures. Presence of a fatty acid is indicated by a plus (+)

<sup>a</sup>Shorthand notation for carbon chain length:number of double bonds.

Table 11. Fatty acid composition of <u>Cyclotella meneghiniana</u> for supply rates of silica-stressed cultures. Presence of a fatty acid is indicated by a plus (+)

	Supply rate (x 10 <sup>-7</sup> mg Si·cell <sup>-1</sup> ·day <sup>-1</sup> )				
Fatty acid	0.140	0.449	1.178		
14:0 <sup>a</sup>	+	+	+		
15:0	+				
16:0	+	+	+		
17:0	+	+	+		
18:0	+	÷	+		
18:1	+	+	+		
18:2	+	+			

<sup>a</sup>Shorthand notation for carbon chain length:number of double bonds.

mg P.cell<sup>-1</sup>.day<sup>-1</sup>, all of the aforementioned fatty acids were observed at these two nutrient supply rates. At 3.475 x  $10^{-7}$  mg N.cell<sup>-1</sup>.day<sup>-1</sup>, fatty acids with a low carbon chain length (e.g. C<sub>14</sub> to C<sub>17</sub>) were, for the most part, absent.

In contrast, the fatty acids in the Si-stressed experiment were limited to low carbon chain lengths ( $C_{14}$  to  $C_{18}$ ). In addition, there was a preponderance of FAMEs with retention times less than that associated with  $C_{14}$ . These peaks were, most likely, FAMEs with carbon chain lengths of 10 to 13. Since the standard samples used did not possess FAMEs with carbon chain lengths less than 14, no definite identification of these fatty acids could be made. Fatty acids at 0.140 and 0.449 x 10<sup>-7</sup> mg Si·cell<sup>-1</sup> ·day<sup>-1</sup> ranged from  $C_{14:0}$  to  $C_{18:2}$ .  $C_{15:0}$  was absent in the extracted lipid at 0.449 x 10<sup>-7</sup> mg Si·cell<sup>-1</sup>·day<sup>-1</sup>. At 1.178 x 10<sup>-7</sup> mg Si·cell<sup>-1</sup> ·day<sup>-1</sup>,  $C_{15:0}$  and  $C_{18:2}$  were absent from the extracted lipid.

## DISCUSSION

Researchers have reported that algae accumulate storage products when exposed to nutrient limitation (Fogg, 1956; Coombs <u>et al.</u>, 1967a,b; Darley, 1977; Shifrin and Chisholm, 1981). However, in this study, accumulation of storage products by <u>C. meneghiniana</u> varied with the type and degree of nutrient limitation. In addition, the statistical significance of nutrient-limitation effects on the biochemical composition of the cells varied with the culture design used.

A reduction in the amount of cellular carbohydrate, protein, and chlorophyll and an increase in the amount of cellular lipid has been reported to occur in diatoms with increasing Si limitation (Werner, 1977). In this study, the LSD analysis indicated some increases in the amount of cellular lipid and protein as Si becomes limiting. However, Snedecor and Cochran (1980) suggested that the LSD analysis should be used to claim significance between variable means only if the "F test" is significant. Therefore, the LSD analysis was considered applicable only if the ANOVA indicated a significant difference between variable means among nutrient supply rates. The ANOVA indicated that only L/G ratios were significantly different among Si supply rates.

The significant decrease in cell L/G ratios with increasing Si supply rate indicated a shift from glucan to lipid accumulation as Si becomes limiting. While the ANOVA did not indicate a significant difference in the amount of cellular lipid and/or glucan among Si supply rates, it is possible that the shift from glucan to lipid accumulation occurs at a particular Si concentration. In the nutrient-stressed

experiments, the use of only 3 distinct supply rates may have restricted the ANOVA from redefining lipid differences above and below this threshold concentration. In this study, the threshold concentration appeared to be between 0.140 and 0.449 x  $10^{-7}$  mg Si·cell<sup>-1</sup>·day<sup>-1</sup>. Future studies should investigate cellular lipid accumulation over this range.

The amount of cellular lipid of <u>C</u>. <u>meneghiniana</u> significantly increased with increasing N supply rate. This is in contrast to results from researchers who observed lipid accumulation with N limitation (Fogg, 1956; Badour and Gergis, 1965; Richardson <u>et al.</u>, 1969; Werner, 1977; Shifrin and Chisholm, 1981). However, several researchers have observed results similar to those reported in this study. Collyer and Fogg (1955) stated that N concentration had no direct effect on fat accumulation in certain algae. Fogg (1956) stated that when fat-producing, N-deficient cells of <u>Navicula pelliculosa</u> underwent N repletion, fat accumulation continued. Badour and Gergis (1965) noted fat accumulation in Nitzschia sp. at times when N was not deficient.

The amount of cellular glucan of <u>C</u>. <u>meneghiniana</u> significantly increased with increasing N supply rate. In contrast, Myklestad and Haug (1972) observed a decrease in carbohydrate in <u>Chaetoceros affinis</u> var. <u>willei</u> with increasing N concentration. In this study, no significant differences in the amount of cellular glucan were observed between N-sufficient and N-deficient batch cultures.

Because of the essential role of N in amino acids and chlorophyll, N limitation might be expected to cause a decrease in the amount of

cellular protein and chlorophyll. However, no significant differences in the amount of cellular protein or chlorophyll were noted between N-sufficient and N-deficient batch cultures. However, decreasing N supply rate in the nutrient-stressed experiment did cause a significant decrease in cellular protein.

Decreasing P supply rate caused a significant reduction in the amount of cellular lipid and glucan. This contrasts with reports from several researchers (Kuhl, 1974; Werner, 1977; Rosen, 1982) who observed an accumulation of lipid and carbohydrate in algal cells as P becomes limiting. Rosen (1982) also noted a decrease in the amount of cellular chlorophyll of <u>C</u>. <u>meneghiniana</u> as P becomes limiting. In this study, a significant decrease in the amount of cellular chlorophyll as P becomes limiting was not observed.

Chloroplast ultrastructure has been reported to be differentially affected by Si, N, and P limitation (Butler and Simon, 1971; Duke and Reimann, 1977; Hurkman, 1979; Rosen, 1982). It might be expected that the ratio of cellular chlorophyll to its breakdown product, phaeophytin, would decrease as nutrients become limiting. However, no significant differences in the ratios were noted between nutrient-sufficient and nutrient-limited cultures. This observation agrees with results of a previous study dealing with the effects of nutrient deficiency on <u>C</u>. <u>meneghiniana</u> (Rosen, 1982). Rosen stated that "If adjustments to light intensity and nutrient deficiency were occurring through the degradation of chlorophyll, they did not appear to be through the creation of phaeophytin." In addition, Marker et al. (1980) noted that phaeophorbid,

rather than phaeophytin, is the major naturally-occurring breakdown product of chlorophyll and little is known of its spectral character-istics.

Negative phaeophytin values were calculated for several replicate cultures within supply rates of the N-stressed and P-stressed experiments. These values, in turn, produced negative Chl/Ph ratios which were omitted from further analysis. The negative values might have occurred because of improper acidification procedures prior to spectrophotometric readings or to extremely variable absorbance readings. Cell densities of cultures in the nutrient-stressed experiments were much less than cell densities of cultures in the nutrient-starvation experiments. The low chlorophyll concentrations resulting from the low cell densities produced spectrophotometric readings consistently below 0.1 absorbance. Since absorbance readings are relatively inaccurate at levels below 0.1 (A.P.H.A. et al., 1971), extreme variability among replicate samples could have occurred. Therefore, if these values were substituted into the defined equations for determination of chlorophyll and phaeophytin concentrations, highly inaccurate values could have resulted.

Myklestad and Haug (1972) considered the protein/carbohydrate ratio to be a valuable indicator of the physiological state of the cell. In rapidly-growing cells, most cellular carbon is incorporated into protein. In times of stress, more cellular carbon is incorporated into storage products (lipid and carbohydrate) and protein synthesis is retarded (Cook, 1961; Mylkestad and Haug, 1972; Weete, 1980). Therefore,

high P/G ratios would be expected in healthy, nutrient-sufficient cells and low P/G ratios in stressed, nutrient-limited cells (Myklestad and Haug, 1972; Sakshaug and Myklestad, 1973; Myklestad, 1974). In this study, no significant differences in P/G ratios were noted in any experiment.

Carbon chain lengths of fatty acids generally ranged from 14 to 24. This is in agreement with results of previous studies on diatom fatty acids (Ackman and Tocher, 1968; Wood, 1974; Fisher and Schwarzenbach, 1978). No distinct differences in fatty acid composition were noted among supply rates within Si-stressed and N-stressed cultures. However, distinct differences in fatty acid composition were observed between Sistressed and N-stressed cultures. The cultures exposed to Si stress did not possess fatty acids that have carbon chain lengths greater than 18. The cultures exposed to Si stress also possessed fatty acids that have carbon chain lengths less than 14. The cultures exposed to N stress did possess fatty acids that have carbon chain lengths greater than 18. In addition, a greater amount of unsaturated acids was observed in Nstressed cultures than in Si-stressed cultures.

The synthesis of long chain fatty acids from  $CO_2$  involves the sequential addition of 2 carbons to the growing fatty acid chain utilizing the enzyme complex, Fatty Acid Synthetase (FAS). This elongation process stops at the formation of hexadecanoic acid ( $C_{16:0}$ ) (Stryer, 1981). Subsequent elongation of the fatty acid chain into longer carbon chain lengths and/or unsaturation involves several enzymatic systems besides FAS. The FAS or the additional enzyme systems may have been

blocked, in some manner, by Si stress, thereby preventing formation of long carbon chain, saturated fatty acids.

Algae are reported to possess large amounts of unsaturated fatty acids (Otsuka and Morimura, 1966; DeMort et al., 1972; MacCarthy and Patterson, 1974a, b; Opute, 1974a; Pohl, 1974; Fisher and Schwarzenbach, 1978). Since long chain unsaturated acids are believed to be involved in photosynthetic reactions and are essential components of diatom cell membranes (Kates and Volcani, 1966; Opute, 1974a; Tornabene et al., 1974), it appears that Si stress caused a greater disruption of C. meneghiniana's lipid composition than did N stress. In addition, a predominance of fatty acids that have carbon chain lengths less than 14 has also been observed in marine diatoms exposed to Si stress (Larry Raymond, Solar Energy Research Institute, personal communication). However, the current study did not quantitate the fatty acids of the extracted lipids and the mere presence or absence of an acid may not provide a realistic appraisal of the actual fatty acid dynamics. In addition, researchers (Ackman and Tocher, 1968; Orcutt and Patterson, 1974) have proposed that changes in the fatty acid composition of a cell reflects changes in the type of lipid. Therefore, "it is really necessary to consider the fatty acids of each individual group of lipids and their changes, rather than the fatty acids in toto, before a proper interpretation of nutrient-linked changes in the fatty acid composition of a cell can be made" (Pugh, 1971).

The exact reason for the biochemical response of <u>C</u>. <u>meneghiniana</u> to Si, N, and P limitation is not known. Coombs and Volcani (1968)

reported researchers who suggested that Si limitation leads to a reduction in Si-containing compounds essential to the metabolism of the cell. However, several researchers (Coombs et al., 1967a; Darley, 1969; Darley, 1977) theorized that the biochemical response to Si limitation results from cessation of growth prior to cell division. Cells of cultures in exponential growth would develop into mature cells or progress to a biprotoplastic stage before Si deficiency inhibited their development. Cellular constituents would accumulate in the biprotoplastic cells since these cells have more of their division cycle to complete than do the older, mature cells (Darley, 1969). A biochemical response associated with Si limitation may, therefore, simply "reflect the percentage of morphologically arrested cells in the culture" (Coombs et al., 1967a). Darley and Volcani (1969) supported this hypothesis as they observed a decrease in the overall energy metabolism and net biosynthetic capacity with the presence of biprotoplastic cells in Si-deficient, diatom cultures. This hypothesis might explain the increase in cellular protein in Si-deficient batch cultures of C. meneghiniana.

Researchers have proposed several theories to explain the effects of N and P concentration on lipid accumulation in algae. Fogg (1956) theorized that N deficiency caused an increase in the proportion of fats to other organic constituents within a cell as a result of the hydrolysis and subsequent loss from the cell of materials other than lipids. De Vasconcelos and Fay (1974) and Rosen (1982) suggested that the formation of lipid droplets in algal chloroplasts results from the degradation

of the chloroplasts' thylakoid membranes. The formation of these lipid droplets would, of course, cause increases in the total amount of cellular lipid. Atkinson (1965) suggested that in N-deficient cells, an increase of ATP occurs due to the inhibition of protein synthesis. The concentration of AMP would decrease and the reduced activity of isocitric dehydrogenase would cause an accumulation of citric acid. Since citrate is the allosteric activator of acetyl CoA carboxylase (Stryer, 1981), the conversion of acetyl CoA to fatty acids would be enhanced. On the other hand, increased photosynthesis caused by increasing N and P supply rates might cause increases in the citric acid intermediates of C. meneghiniana. Beardall et al. (1976) noted photosynthetic fixation of carbon in 2 marine diatoms by reactions similar to those of  $\rm C_{A}$  plants. More than 70% of the carbon fixed in these 2 taxa was in the form of amino acids or as intermediates of the citric acid cycle (including citric acid). Increased citric acid concentrations would cause an increase in the activity of acetyl CoA carboxylase and, subsequently, an increase in fatty acid synthesis. If C. meneghiniana possesses a photosynthetic mechanism similar to these 2 diatoms, the increase in the amount of cellular lipid with increasing N and P concentrations might be explained.

In addition, N stress could cause a reduction in the cells' protein synthesis and, consequently, in enzymatic activity. Collyer and Fogg (1955) speculated "that N deficiency brings about a change in the proportions of enzyme systems in favor of those concerned in fat synthesis." It may be that the enzymatic systems involved in fatty acid synthesis

are of low "priority" in the metabolism of <u>C</u>. <u>meneghiniana</u>. As N becomes increasingly limited, the low "priority" enzymatic systems might be inhibited while systems more essential for cell survival remain operative. Since the amount of cellular protein appeared to be positively correlated with the amount of cellular lipid of <u>C</u>. <u>meneghiniana</u>, this hypothesis might explain the small amounts of cellular lipid at low N concentrations.

Several factors could have contributed to differences in results between this study and others. The extraction techniques used in this study were chosen for their precision, ease of use, and applicability to algal samples. For example, the "Lowry Method" was used for protein determination because of its ease of use and universal acceptance. However, Coombs <u>et al</u>. (1967a) noted that protein determination by the "Lowry Method" produced values 30% lower than the "Kjeldahl Method." In addition, most researchers have used the chloroform/methanol extraction technique for lipid analysis. Only a few researchers (Otsuka and Morimura, 1966; Fisher and Schwarzenbach, 1978) have used alcoholic KOH to extract algal lipids. The alcoholic KOH techniques used in this study have been shown to be more effective in extracting lipids than chloroform/methanol techniques (Moon and Hammond, 1978). Therefore, differences in protein and lipid results between this study and others may have resulted from the extraction techniques that were used.

No direct comparisons between data in this study and others can be made. The biochemical data in this study were expressed on a "per cell" basis. Other researchers have almost exclusively expressed biochemical

data on a "per dry weight" basis. Data expressed on a dry weight basis are, however, difficult to interpret in diatoms "because of the presence of the silicious cell wall which is a highly variable percentage of the dry weight in different species" (Darley, 1977). Therefore, since data in this study were intended only for comparisons within and between experimental treatments and not for comparisons between studies, expression of biochemical data on a "per cell" basis was judged to be satisfactory.

Most researchers investigating nutrient-limitation effects on algal biochemical composition have analyzed nutrient-sufficient cells resuspended in nutrient-deficient media. The biochemical evaluation of such cells does not take into account the physiological history of the cells. Many algae can carry on normal physiological functions in nutrient-limited conditions as long as sufficient nutrient "pools" are maintained within the cells. For example, diatoms have been reported to store cytoplasmic "pools" of Si and P (Coombs <u>et al</u>., 1967a; Azam <u>et al</u>., 1974; Tilman and Kilham, 1976; Kilham <u>et al</u>., 1977; Paasche, 1980; Sullivan and Volcani, 1981). Therefore, the biochemical response of a culture during the initial period of Si or P limitation might simply be a response to the utilization of the cells' cytoplasmic "pools" rather than the nutrient concentration of the growth medium.

To minimize the variability caused by the luxury consumption of nutrients by diatoms in batch-starvation experiments, Kilham <u>et al</u>. (1977) suggested a 5-day, Si-starvation period and a 3-week, P-starvation period of cells prior to performing batch-growth experiments. In

this study, the biochemical composition of <u>C</u>. <u>meneghiniana</u> was analyzed 3 and 9 days after resuspension of nutrient-sufficient cells in nutrient-deficient media. No consideration was given for luxury consumption of nutrients by <u>C</u>. <u>meneghiniana</u> which may account for some of the differences in results between the nutrient-starvation and nutrient-stressed experiments.

Most researchers investigating diatom lipid production have used marine taxa or a species of the freshwater genera, <u>Nitzschia</u> and <u>Navicula</u>, as a test organism. Physiological characteristics of <u>C</u>. <u>meneghiniana</u> could have caused differences in results between this study and others. For example, Lowe (1974) reported <u>C</u>. <u>meneghiniana</u> to be a "facultative N heterotroph." Therefore, <u>C</u>. <u>meneghiniana</u> might possess different nutrient assimilation mechanisms than taxa used in other studies.

To my knowledge, this is the first study to investigate the amount of cellular lipid of a diatom at steady state along nutrient gradients. The semi-continuous culture technique maintained constant cell growth rates by maintaining a constant rate of nutrient supply to the cells. Changes in the rate of nutrient supply cause changes in the growth rate and the physiological state of the cell. Biochemical measurements made on cells of semi-continuous cultures in different physiological states characterized the population at specific points along a nutrient gradient. On the other hand, the batch culture technique did not maintain constant growth rate or similar physiological states of the cells. Biochemical measurements made on cells of batch cultures cannot be

identified with a particular nutrient concentration. Therefore, results of the nutrient-stressed experiments were more reliable than results of the nutrient-starvation experiments. Future studies should use continuous culture techniques for accurate biochemical and physiological analyses of algae.

## **BIOTECHNICAL PERSPECTIVE**

The application of mass algal cultures to biotechnology has attracted scientific attention since the late 1940s. Mass algal cultures have been proposed for use in wastewater treatment, bioregenerative life support systems, chemical production, water renovation and recycling, and, most recently, hydrocarbon fuel production (Anonymous, 1978; Goldman, 1979; Goldman, 1980; Bergeron <u>et al.</u>, 1983). The latter usage has received much attention in the last several years. However, the U.S. Research and Development Administration projected that for biomass "fuels" to have potential in energy production, they must provide 5 to 10% of the total U.S. energy needs (Goldman and Ryther, 1977).

If <u>C</u>. <u>meneghiniana</u> was mass cultured as a biotechnological energy source, the data presented in nutrient-stressed experiments could be used to estimate "conceivable" production of lipid, glucan, and protein. The amount of cellular lipid, glucan, and protein was greatest in Nstressed cultures with the greatest N supply rate. The following hypothetical yields of lipid, glucan, and protein production were estimated using this data set.

Assume that <u>C</u>. <u>meneghiniana</u> could achieve a uniform growth of 25,000 cells·ml<sup>-1</sup> in a 1 hectare (ha) by 50 cm pond. Since the harvesting of unicellular algae is difficult (Goldman and Ryther, 1977), a sampling efficiency of 75% is assumed. An extraction efficiency of 90% (Shifrin and Chisholm, 1980) is also assumed. Using these assumptions, hypothetical yields of 0.78 metric tons lipid·ha<sup>-1</sup>, 0.31 metric tons

glucan·ha<sup>-1</sup>, and 61.87 metric tons protein·ha<sup>-1</sup> would result. If harvesting could be completed 3 times per year (yr), the hypothetical yield would be 2.33 metric tons lipid·ha<sup>-1</sup>·yr<sup>-1</sup>, 0.94 metric tons glucan·ha<sup>-1</sup>·yr<sup>-1</sup>, and 185.61 metric tons protein·ha<sup>-1</sup>·yr<sup>-1</sup>. Using the caloric values of 15 gigajoules (gj) per metric ton carbohydrate and protein and 40 gj per metric ton lipid (Dubinsky <u>et al</u>., 1979), the total energy yield would be 2,891 gj·ha<sup>-1</sup>·yr<sup>-1</sup>.

The importance and applicability of such an energy yield is difficult to interpret. Based on the projected biomass cost target (i.e., gross income) of 0.95 to 1.40 dollars  $\cdot gj^{-1}$  (Lipinsky, 1978), mass cultures of <u>C</u>. <u>meneghiniana</u> could produce a gross income of 2,602 to 4,047 dollars  $\cdot ha^{-1} \cdot yr^{-1}$ . This value is substantially higher than the yearly gross income calculated for algal biomass by Dubinsky <u>et al</u>. (1979) and approximately 5 to 8 times greater than that projected for plant energy "farms" (Lipinsky, 1978). However, these estimated gross incomes do not take into account any harvesting, production, or material costs. Consequently, no comparisons between algal and higher plant net incomes can be made.

The success of algal mass cultures in biotechnology is dependent upon the selection and utilization of taxa which produce optimal yields concurrent with maximum growth. Researchers have indicated that algae produce optimal biochemical yields under conditions that are not conducive to optimal growth. The data generated from nutrient-stressed experiments indicated that <u>C</u>. <u>meneghiniana</u> produced greater biochemical yields with decreasing N stress. The ability of a taxon to produce
high biochemical yields under optimal growth conditions does, therefore, exist. This may be an occurrence specific to a particular taxon or a particular strain of a taxon.

The challenge for scientists in biotechnology is to select organisms capable of producing the desired compounds under specific growth conditions. Such organisms could possibly be obtained in several ways. Certain taxa in Israeli mass algal cultures are the "preferred" taxa with large yields of glycerin and beta carotene (Anonymous, 1978). Additional studies may allow researchers to identify other taxa that can produce large quantities of desired biochemical compounds (e.g., octadecatrienoic acid, a fatty acid of high commercial value). In addition, some researchers (Hall, 1979; Bergeron et al., 1983) have proposed that manipulation of a taxon's genetic structure has great promise in biotechnology. The data from the nutrient-stressed experiments with C. meneghiniana indicate that taxa can produce large amounts of desired compounds under low-stress growth conditions. Genetic engineering would be best used on these "preferred" taxa. Therefore, a thorough screening for these "preferred" taxa should be completed before manipulation of a taxon's gene structure is used to obtain large biochemical yields.

## SUMMARY

- (1) The statistical significance of nutrient-limitation effects on the biochemical composition of the cells varied between batch and semi-continuous culture experiments. In batch cultures, Si deficiency caused a significant increase (300%) in the amount of cellular protein. Cellular protein was also significantly different between fractional replicates. In semi-continuous cultures, cellular L/G ratios significantly decreased (50%) with increasing Si supply rate. The amount of cellular lipid, glucan, and protein significantly increased (300%) with increasing N supply rate. The amount of cellular lipid and glucan significantly increased (300 and 400%, respectively) with increasing P supply rate.
- (2) Fatty acids of lipid extracted from cells of semi-continuous cultures generally possessed carbon chain lengths of 14 to 24. Carbon chain lengths of fatty acids from cells in Si-stressed cultures ranged from less than 14 to 18. No distinct differences in diatom fatty acid composition were observed among supply rates within Si-stressed and N-stressed cultures.
- (3) Greater amounts of unsaturated fatty acids were found in lipid extracted from cells of N-stressed cultures than in cells of Sistressed cultures.
- (4) Discrepancies between results of this study and others were attributed to culture design, extraction techniques, the scaling factor used for expression of biochemical content, and the physiological characteristics of <u>C</u>. meneghiniana.

- (5) An estimated yield for a hypothetical mass culture of <u>C</u>.
  <u>meneghiniana</u> was calculated to be 2.33 metric tons lipid.ha<sup>-1</sup>.yr<sup>-1</sup>,
  0.94 metric tons glucan.ha<sup>-1</sup>.yr<sup>-1</sup>, and 185.61 metric tons protein.
  - $ha^{-1} \cdot yr^{-1}$ . The total energy yield derived from such a biochemical yield would be 2,891 gj $\cdot$ ha<sup>-1</sup> $\cdot$ yr<sup>-1</sup>.

## LITERATURE CITED

- Ackman, R. G., and C. S. Tocher. 1968. Marine phytoplankter fatty acids. J. Fish. Res. Bd. Can. 25:1603-1620.
- Ackman, R. G., P. M. Jangaard, R. J. Hoyle, and H. Brockerhoff. 1964. Origin of marine fatty acids. I. Analysis of the fatty acids produced by the diatom <u>Skeletonema costatum</u>. J. Fish. Res. Bd. Can. 21:747-756.
- Ackman, R. G., C. S. Tocher, and J. McLachlan. 1968. Marine phytoplankter fatty acids. J. Fish. Res. Bd. Can. 21:1603-1620.
- American Public Health Association, American Water Works Association, and Water Pollution Control Federation. 1971. Standard methods for the examination of water and wastewater. 13th ed. American Public Health Association, New York. 874 pp.
- Anonymous. 1978. Algae may yield bumper crop. Chemical Week 122(18): 42.
- Antia, N. J., C. D. McAllister, T. R. Parsons, K. Stephens, and J. D. H. Strickland. 1963. Further measurements of primary production using a large-volume plastic sphere. Limnol. Oceangr. 8:166-183.
- Atkinson, D. E. 1965. Biological feedback control at the molecular level. Science 150:851-857.
- Azam, F., B. B. Hemmingsen, and B. E. Volcani. 1974. Role of silicon in diatom metabolism. V. Silicic acid transport and metabolism in heterotrophic diatom <u>Nitzschia alba</u>. Archiv. Mikrobiol. 97:103-114.
- Badour, S. S., and M. S. Gergis. 1965. Cell division and fat accumulation in <u>Nitzschia</u> sp. grown in continuously illuminated cultures. Archiv. Mikrobiol. 51:94-102.
- Barker, H. A. 1935. Photosynthesis in diatoms. Archiv. Mikrobiol. 6:141-156.
- Beardall, J., D. Mukerji, H. E. Glover, and I. Morris. 1976. The path of carbon in photosynthesis by marine phytoplankton. J. Phycol. 12:409-417.
- Beattie, A., E. L. Hirst, and E. Percival. 1961. Studies on the metabolism of the Chrysophyceae; Comparative structural investigations on leucosin (chrysolaminaran) separated from diatoms and laminaran from the brown algae. Biochem. J. 79:531-537.

- Bergeron, P. W., R. E. Corder, A. M. Hill, H. Lindsey, and M. Z. Lowenstein. 1983. SERI biomass program annual technical report: 1982. Solar Energy Research Institute Special Report for the U. S. Department of Energy; Contract No. EG-77-C-01-4042. 82 pp.
- Bidwell, R. G. S. 1974. Plant physiology. Macmillan Publishing Co., Inc., New York. 643 pp.
- Brockerhoff, H., M. Yurkowski, R. J. Hoyle, and R. G. Ackman. 1964. Fatty acid distribution in lipids of marine plankton. J. Fish. Res. Bd. Can. 21:1379-1384.
- Butler, R. D., and E. W. Simon. 1971. Ultrastructural aspects of senescence in plants. Advances in Gerontological Research. 3: 73-129.
- Cochran, W. G., and G. M. Cox. 1957. Experimental designs. 2nd ed. John Wiley & Sons, Inc., New York. 611 pp.
- Collyer, D. M., and G. E. Fogg. 1955. Studies on fat accumulation by algae. J. Exp. Bot. 6:256-275.
- Conover, S. A. M. 1975. Partitioning of nitrogen and carbon in cultures of the marine diatom <u>Thalassiosira fluviatilis</u> supplied with nitrate, ammonium, or urea. Mar. Biol. 32:231-246.
- Cook, J. R. 1961. Euglena gracilis in synchronous division. II. Biosynthetic rates over the life cycle. Biol. Bull. 121:277-289.
- Coombs, J., and B. E. Volcani. 1968. Studies on the biochemistry and fine structure of silica shell formation in diatoms: Siliconinduced metabolic transients in Navicula pelliculosa (Breb.) Hilse. Planta (Berl.) 80:264-279.
- Coombs, J., W. M. Darley, O. Holm-Hansen, and B. E. Volcani. 1967a. Studies on the biochemistry and fine structure of silica shell formation in diatoms: Chemical composition on <u>Navicula pelliculosa</u> during silicon-starvation synchrony. Plant Physiol. 42:1601-1606.
- Coombs, J., C. Spanis, and B. E. Volcani. 1967b. Studies on the biochemistry and fine structure of silica shell formation in diatoms: Photosynthesis and respiration in silicon-starvation synchrony of Navicula pelliculosa. Plant Physiol. 42:1607-1611.
- Crawford, R. M. 1973. The protoplasmic ultrastructure of the vegetative cell of <u>Melosira varians</u> C. A. Agardh. J. Phycol. 9:50-61.

- Daniel, C. 1959. Use of half-normal plots in interpreting factorial two-level experiments. Technometrics 1:311-341.
- Darley, W. M. 1969. Silicon and the division cycle of the diatom <u>Navicula pelliculosa</u> and <u>Cylindrotheca fusiformis</u>. Proc. N. Am. Paleon. Conf. 1969:944-1009.
- Darley, W. M. 1977. Biochemical composition. Pages 198-223 in D. Werner (ed.). The biology of diatoms. University of California Press, Berkeley.
- Darley, W. M., and B. E. Volcani. 1969. Role of silicon in diatom metabolism. Exp. Cell Res. 58:334-342.
- Davis, C. O. 1976. Continuous culture of marine diatoms under silicate limitation. II. Effect of light intensity on growth and nutrient uptake of Skeletonema costatum. J. Phycol. 12:291-300.
- Dawson, P. A. 1973. Observations on the structure of some forms of <u>Gomphonema parvulum</u> Kutz. II. The internal organization. J. Phycol. 9:165-175.
- DeMort, C. L., R. Lowry, I. Tinsley, and H. K. Phinney. 1972. The biochemical analysis of some estuarine phytoplankton species. I. Fatty acid composition. J. Phycol. 8:211-216.
- de Vasconcelos, L., and P. Fay. 1974. Nitrogen metabolism and ultrastructure in <u>Anabaena cylindrica</u>. I. The effect of nitrogen starvation. Arch. Microbiol. 96:271-279.
- Drum, R. W. 1963. The cytoplasmic fine structure of the diatom <u>Nitschia palea</u>. J. Cell Biol. 18:429-440.
- Dubinsky, Z., T. Berner, and S. Aaronson. 1979. Potential of large scale algal culture for biomass and lipid production in arid lands. Biotech. Bioeng. Symp. 8:51-68.
- Duke, E. L., and B. E. F. Reimann. 1977. The ultrastructure of the diatom cell. Pages 65-109 in D. Werner (ed.). The biology of diatoms. University of California Press, Berkeley.
- Eppley, R. W. 1977. The growth and culture of diatoms. Pages 24-64 in D. Werner (ed.). The biology of diatoms. University of California Press, Berkeley.
- Eppley, R. W., and E. H. Renger. 1974. Nitrogen metabolism of an oceanic diatom in nitrogen-limited continuous culture. J. Phycol. 10:15-23.

- Fisher, N. S., and R. P. Schwarzenbach. 1978. Fatty acid dynamics in <u>Thalassiosira pseudonana</u> (Bacillariophyceae): Implications for physiological ecology. J. Phycol. 14:143-150.
- Fogg, G. E. 1953. The metabolism of algae. Methuen and Company, Ltd., London. 149 pp.
- Fogg, G. E. 1956. Photosynthesis and formation of fats in a diatom. Ann. Bot. 20:265-285.
- Fogg, G. E. 1965. Algal cultures and phytoplankton ecology. University of Wisconsin Press, Madison. 126 pp.
- Ford, C. W., and E. Percival. 1965. The carbohydrates of <u>Phaeodactylum</u> <u>tricornatum</u>. I. Preliminary examination of the organism, and characterization of low molecular weight material and a glucan. J. Chem. Soc. (London): 7035-7041.
- Goldman, J. C. 1979. Outdoor algal mass cultures. I. Applications. Water Research 13:1-19.
- Goldman, J. C. 1980. Physiological aspects in algal mass cultures. Pages 343-359 in G. Shelef and C. J. Soeder (eds.). Algae biomass. Elsevier/North Holland Biomedical Press, Amsterdam.
- Goldman, J. C., and J. H. Ryther. 1977. Mass production of algae: Bioengineering aspects. Pages 367-378 in A. Mitsui, S. Miyachi, A. San Pietro, and S. Tamura (eds.). Biological solar energy conversion. Academic Press, New York.
- Guillard, R. L. 1973. Division rates. Pages 289-312 in J. R. Stein (ed.). Handbook of phycological methods; Culture methods and growth measurements. Cambridge University Press, Cambridge.
- Guillard, R. L., and C. J. Lorenzen. 1972. Yellow-green algae with chlorophyllide C. J. Phycol. 8:10-14.
- Hall, D. O. 1979. Solar energy use through biology--Past, present, and future. Solar Energy 22:307-328.
- Hammond, E. G., B. A. Glatz, Y. Choi, and M. T. Teasdale. 1981. Oil production by <u>Candida curvata</u> and extraction, composition, and properties of the oil. Pages 171-187 in E. H. Pride, L. H. Princen and K. D. Mukherjee (eds.). New sources of fats and oils. Am. Oil Chem. Soc. Monogr. No. 9. Champaign.
- Handa, N. 1969. Carbohydrate metabolism in the marine diatom Skeletonema costatum. Mar. Biol. 4:208-214.

- Healey, F. P., J. Coombs, and B. E. Volcani. 1967. Changes in pigment content of the diatom <u>Navicula pelliculosa</u> (Breb.) Hilse in siliconstarvation synchrony. Archiv Mikrobiol. 59:131-142.
- Hellebust, J. A., and J. Lewin. 1977. Heterotrophic nutrition. Pages 169-197 in D. Werner (ed.). The biology of diatoms. University of California Press, Berkeley.
- Hurkman, W. J. 1979. Ultrastructural changes of chloroplasts in attached and detached, aging primary wheat leaves. Am. J. Bot. 66:64-70.
- Kates, M., and B. E. Volcani. 1966. Lipid components of diatoms. Biochim. Biophys. Acta. 116:264-278.
- Kates, M., and B. E. Volcani. 1968. Studies on the biochemistry and fine structure of silica shell formation in diatoms; Lipid components of the cell walls. Z. Pflangenphysiol. Bd. 60:19-29.
- Kauss, H. 1962. Der einfluss von thiamin- und biotin-mangel auf den stoffwechsel von <u>Ochromonas malhamensis</u>. Votr. Gesamtbeg. Bot. 1:129-132.
- Kilham, S. S., C. L. Kott, and D. Tilman. 1977. Phosphate and silicate kinetics for the Lake Michigan diatom <u>Diatoma</u> <u>elongatum</u>. J. Great Lakes Res. 3:93-99.
- Klyachko-Gurvich, G. L., T. S. Zhukova, M. G. Vladimirova, and T. A. Kurnosova. 1969. Comparative characterization of the growth and direction of biosynthesis of various strains of <u>Chlorella</u> under nitrogen deficiency conditions. III. Synthesis of fatty acids. Sov. Plant Physiol. 16:168-171.
- Kuhl, A. 1974. Phosphorus. Pages 636-654 in W. D. P. Stewart (ed.). Algal physiology and biochemistry. University of California Press, Berkeley.
- Lee, R. E. 1980. Phycology. Cambridge University Press, Cambridge. 478 pp.
- Lewin, J. C., and R. R. L. Guillard. 1963. Diatoms. Ann. Rev. Microbiol. 17:373-414.
- Lipinsky, E. S. 1978. Fuels from biomass: Integration with food and materials systems. Science 199:644-651.
- Low, E. M. 1955. Studies on some chemical constituents of diatoms. J. Mar. Res. 14:199-204.

- Lowe, R. L. 1974. Environmental requirements and pollution tolerance of freshwater diatoms. Environmental Monitoring Series. EPA-670/ 4-74-005. U. S. Environmental Protection Agency, Cincinnati. 334 pp.
- Lowry, O. H., N. J. Rosebrough, A. L. Farr, and R. J. Randall. 1951. Protein measurement with the Folin phenol reagent. J. Biol. Chem. 193:265-275.
- MacCarthy, J. J., and G. W. Patterson. 1974a. Effects of cation levels of the nutrient medium on the biochemistry of <u>Chlorella</u>. I. Concentration series. Plant Physiol. 54:192-132.
- MacCarthy, J. J., and G. W. Patterson. 1974b. Effects of cation levels of the nutrient medium on the biochemistry of <u>Chlorella</u>. II. Factorial experiment. Plant Physiol. 54:133-135.
- Marker, A. F. H., E. A. Nusch, H. Rai, and B. Reimann. 1980. The measurement of photosynthetic pigments in freshwaters and standardization of methods: Conclusions and recommendations. Ergebn. Limnol. 14:91-106.
- Matsuka, M., H. Otsuka, and E. Hase. 1966. Changes in contents of carbohydrate and fatty acid in the cells of Chlorella <u>protothecoides</u> during the processes of de- and re-generation of chloroplasts. Plant Cell Physiol. 7:651-662.
- McCarthy, J. J. 1980. Nitrogen. Pages 191-233 in I. Morris (ed.). The physiological ecology of phytoplankton. University of California Press, Berkeley.
- Meeuse, B. J. D. 1962. Storage products. Pages 289-314 in R. A. Lewin (ed.). Physiology and biochemistry of algae. Academic Press, New York.
- Moon, N. J., and E. G. Hammond. 1978. Oil production by fermentation of lactose and the effect of temperature on the fatty acid composition. J. Am. Oil Chem. Soc. 55:683-688.
- Morris, I. 1974. Nitrogen assimilation and protein synthesis. Pages 583-609 in W. D. P. Stewart (ed.). Algal physiology and biochemistry. University of California Press, Berkeley.
- Morris, I. 1981. Photosynthetic products, physiological state, and phytoplankton growth. Pages 83-102 in T. Platt (ed.). Physiological bases of phytoplankton ecology. Can. Bull. Fish. Aquat. Sci. 210.

- Morris, I., H. E. Glover, and C. S. Yentsch. 1974. Products of photosynthesis by marine phytoplankton: The effect of environmental factors on the relative rates of protein synthesis. Mar. Biol. 27:1-9.
- Myklestad, S. 1974. Production of carbohydrates by marine planktonic diatoms. I. Comparison of nine different species in culture. J. Exp. Mar. Biol. Ecol. 15:261-274.
- Myklestad, S. 1978. β 1,3-glucans in diatoms and brown seaweeds. Pages 133-142 <u>in</u> J. A. Hellebust and J. S. Craigie (eds.). Handbook of phycological methods; Physiological and biochemical methods. Cambridge University Press, Cambridge.
- Myklestad, S., and A. Haug. 1972. Production of carbohydrates by the marine diatom <u>Chaetoceros affinis</u> var. <u>Willei</u> (Grun.) Hustedt. I. Effect of concentration of nutrients in the culture medium. J. Exp. Mar. Biol. Ecol. 9:125-136.
- Nalewajko, C., and D. R. S. Lean. 1980. Phosphorus. Pages 235-258 <u>in</u> I. Morris (ed.). The physiological ecology of phytoplankton. University of California Press, Berkeley.
- Nichols, H. W. 1973. Growth media-freshwater. Pages 7-24 in J. R. Stein (ed.). Handbook of phycological methods; Culture methods and growth measurements. Cambridge University Press, Cambridge.
- Olive, J. H., D. M. Benton, and J. Kishler. 1969. Distribution of C-14 in products of photosynthesis and its relationship to phytoplankton composition and rate of photosynthesis. Ecology 50:380-386.
- Opute, F. I. 1974a. Lipid and fatty acid composition of diatoms. J. Exp. Bot. 25:823-835.
- Opute, F. I. 1974b. Studies on fat accumulation in <u>Nitzschia palea</u>. Ann. Bot. 38:889-902.
- Orcutt, D. M., and G. W. Patterson. 1974. Effect of light intensity upon lipid composition of <u>Nitzschia</u> <u>closterium</u> (<u>Cylindrotheca</u> <u>fusiformis</u>). Lipids 9:1000-1003.
- Otsuka, H., and Y. Morimura. 1966. Change of fatty acid composition of <u>Chlorella ellipsoidea</u> during its life cycle. Plant Cell Physiol. 7:663-670.
- Paasche, E. 1971. Effect of ammonia and nitrate on growth, photosynthesis and ribulose-diphosphate carboxylase content of <u>Dunaliella</u> <u>tertiolecta</u>. Physiologia Planta 25:294-299.

- Paasche, E. 1980. Silicon. Pages 259-284 in I. Morris (ed.). The physiological ecology of photoplankton. University of California Press, Berkeley.
- Parsons, T. R., K. Stephens, and J. D. H. Strickland. 1961. On the chemical composition of eleven species of marine phytoplankters. J. Fish. Res. Bd. Can. 18:1001-1016.
- Parsons, T. R., and J. D. Strickland. 1963. Discussion of spectrophotometric determination of marine-plant pigments, with revised equations of ascertaining chlorophylls and carotenoids. J. Mar. Res. 21:155-163.
- Pohl, P. 1974. Control of unsaturated fatty acid biosynthesis in unicellular algae by the nitrogen content of the medium and by the wavelength of light. J. Am. Oil Chem. Soc. 51:521A.
- Pugh, P. R. 1971. Changes in the fatty acid composition of <u>Coscinodis</u>-<u>cus eccentricus</u> with culture-age and salinity. Mar. Biol. 11:118-124.
- Redfield, A. C. 1958. The biological control of chemical factors in the environment. Am. Sci. 46:205-221.
- Rhee, G-Y. 1978. Effects of N:P atomic ratios and nitrate limitation on algal growth, cell composition, and nitrate uptake. Limnol. Oceanogr. 23:10-25.
- Rhee, G-Y., and I. J. Gotham. 1980. Optimum N:P ratios and coexistence of planktonic algae. J. Phycol. 16:486-489.
- Richards, F. A., with T. G. Thompson. 1952. The estimation and characterization of plankton populations by pigment analyses: II. A spectrophotometric method for the estimation of plankton pigments. J. Mar. Res. 11:156-171.
- Richardson, B., D. M. Orcutt, H. A. Schwertner, C. L. Martinez, and H. E. Wickline. 1969. Effects of nitrogen limitation on the growth and composition of unicellular algae in continuous culture. Appl. Microbiol. 18:245-250.
- Rosen, B. H. 1982. Physiological and ultrastructural responses to light intensity and nutrient deficiency in the planktonic diatom <u>Cyclotella meneghiniana</u> Kütz. Ph.D. thesis. Bowling Green State University, Bowling Green, Ohio. 136 pp.

- Sakshaug, E., and S. Myklestad. 1973. Studies on the phytoplankton ecology of the Trondheimsfjord. III. Dynamics of phytoplankton blooms in relation to environmental factors, bioassay experiments, and parameters for the physiological state of the populations. J. Exp. Mar. Biol. Ecol. 11:157-188.
- Shifrin, N. S., and S. W. Chisholm. 1980. Phytoplankton lipids: Environmental influences on production and possible commercial applications. Pages 627-645 in G. Shelef and C. J. Soeder (eds.). Algae biomass. Elsevier/North Holland Press, Amsterdam.
- Shifrin, N. I., and S. W. Chisholm. 1981. Phytoplankton lipids: Interspecific differences and effects of nitrate, silicate, and light-dark cycles. J. Phycol. 17:374-384.
- Snedecor, G. W., and W. G. Cochran. 1980. Statistical methods. 7th ed. The Iowa State University Press, Ames. 507 pp.
- Spoehr, H. A., and H. W. Milner. 1949. The chemical composition of <u>Chlorella</u>; Effect of environmental conditions. Plant Physiol. 24:120-149.
- Stevenson, R. J., and E. F. Stoermer. 1982. Luxury consumption of phosphorus by five <u>Cladophora</u> epiphytes in Lake Huron. Trans. Am. Microsc. Soc. 101:151-161.
- Stoermer, E. F., H. S. Pankratz, and R. W. Drum. 1964. The fine structure of <u>Mastogloia grevillei</u> Wm. Smith. Protoplasma 59:1-13.
- Stoermer, E. F., H. S. Pankratz, and C. C. Bowen. 1965. Fine structure of the diatom <u>Amphipleura pellucida</u>. II. Cytoplasmic fine structure and frustule formation. Am. J. Bot. 52:1067-1078.
- Stryer, L. 1981. Biochemistry. 2nd ed. W. H. Freeman and Company, San Francisco. 949 pp.
- Sullivan, C. W., and B. E. Volcani. 1981. Silicon in the cellular metabolism of diatoms. Pages 15-42 in T. L. Simpson and B. E. Volcani (eds.). Silicon and siliceous structures in biological systems. Springer-Verlag, Inc., New York.
- Syrett, P. J. 1962. Nitrogen assimilation. Pages 171-188 in R. A. Lewin (ed.). Physiology and biochemistry of algae. Academic Press, New York.
- Syrett, P. J. 1981. Nitrogen metabolism by microalgae. Pages 182-210 in T. Platt (ed.). Physiological bases of phytoplankton ecology. Can. Bull. Fish. Aquat. Sci. 210.

- Tilman, D., and S. S. Kilham. 1976. Phosphate and silicate growth and uptake kinetics of the diatoms <u>Asterionella formosa</u> and <u>Cyclotella</u> <u>meneghiniana</u> in batch and semi-continuous culture. J. Phycol. 12:375-383.
- Titman, D., and P. Kilham. 1976. Sinking in freshwater phytoplankton: Some ecological implications of cell nutrient status and physical mixing processes. Limnol. Oceanogr. 21:409-417.
- Thomas, W. H., and A. N. Dodson. 1972. On nitrogen deficiency in tropical pacific oceanic phytoplankton. II. Photosynthetic and cellular characteristics of a chemostat-grown diatom. Limnol. Oceangr. 17:515-523.
- Tornabene, T. G., M. Kates, and B. E. Volcani. 1974. Sterols, aliphatic hydrocarbons, and fatty acids of a non-photosynthetic diatom, Nitzschia alba. Lipids 9:279-284.
- von Stosch, H. A. 1951. Uber das leukosin, den reservestaff der Chrysophyten. Naturwissenschaften 38:192-193.
- Weete, J. D. 1980. Lipid biochemistry of fungi and other organisms. Plenum Press, New York. 388 pp.
- Werner, D. 1977. Silicate metabolism. Pages 110-149 <u>in</u> D. Werner (ed.). The biology of diatoms. University of California Press, Berkeley.
- Wetzel, R. G. 1975. Limnology. W. B. Saunders Co., Philadelphia. 743 pp.
- Wood, B. J. B. 1974. Fatty acids and saponifiable lipids. Pages 236-265 <u>in</u> W. D. P. Stewart (ed.). Algal physiology and biochemistry. University of California Press, Berkeley.
- Zhukova, T. S., G. L. Klyachko-Gurvich, M. G. Vladimirova, and T. A. Kurnosova. 1968. Comparative characterization of the growth and direction of biosynthesis of various strains of <u>Chlorella</u> under conditions of nitrogen starvation. Formation of carbohydrates and lipids. Sov. Plant Physiol. 16:96-101.

## ACKNOWLEDGMENTS

This research was supported, in part, by grants from Sigma Xi, the Department of Botany, Iowa State University, and the Graduate College, Iowa State University.

I dedicate this study to my parents, Fred and Aileen Millie, as an expression of my appreciation for their unyielding encouragement, guidance, support, and love. Their installment of pride and confidence in me was, at times, the only belief I had left to "hold onto." Without them, I could never have completed this personal and professional goal. I owe them so much that a mere "thanks" seems trivial.

I am grateful to Dr. John D. Dodd for bringing me to Iowa and teaching me to look "five years down the road." I extend deep appreciation to Dr. Lois H. Tiffany for taking on the responsibility of serving as my major professor upon Dr. Dodd's retirement. Without Dr. "T's" continual friendship, support, and guidance, my dream probably would not have materialized. I thank Drs. Roger Bachmann, Ronald Coolbaugh, William Crumpton, David Czarnecki, Tom Elthon, Earl Hammond, Paul Hinz, David Klarer, Barry Rosen, Mike Starr, Lois Tiffany, and Arnold van der Valk for providing technical assistance, advice, and/or equipment during this study. Carolyn Taylor provided valuable advice while typing the manuscript.

I thank Kelly Applegate, Tim Bliss, Ellen Bluth, Mark Buchheim, Dr. Mike and Sue Crum, Dr. David Czarnecki, Denise Farrar, Dawn Fortner, John Gamble, Maria Garrison, Ron and Diann Getman, Al Hancock, Gary Hartman, Cheryl Jacobsen, Jim and Mari Kemis, Jane Kletschke, Dr.

George Knaphus, Ken Kortge, Jeff Love, Stan and Kay Maas, Steve Mahoney, Becky Miller, Mark and Luann Pisarik, Pam Rypkema, Steve and Renee Schaaf, Neil Shaw, Judy Shearer, Audrey Wacha, Lynette Walsh, Dr. Jim and Beth Wee, and Gale West for making my time spent in Iowa very rewarding. Extra appreciation is extended to Stan and Kay for acting as my "adopted parents" and for providing me with enough Scotch to forget the tensions and frustrations of life. These people have made me realize that Iowa really is "a place to grow."

Last, but not least, I extend appreciation to Karen Getman for providing me assistance and friendship during my doctoral program. I have included this as a reminder to her that I do, indeed, "believe in miracles." Just remember, Karen, "No matter how bad things get, if you persevere, you will survive. . . and sometimes even win!"

"...Don't you understand? When you give up your dream, you die...."

Michael Nouri, Flashdance (1983)

## APPENDIX: MEAN SQUARE VALUES OF THE FACTORIAL ANALYSIS AND THE ANALYSIS OF VARIANCE

Source of	Variable								
variation	Lipid	Glucan	Protein	Chloro- C phyll P	hlorophyll/ haeophytin	Protein/ Glucan	Lipid/ Glucan		
Fractional replicate	8.60	7.60x10-2	6.88x10 <sup>4</sup> *	1.21x10 <sup>-6</sup>	3.58	8.86x10 <sup>5</sup>	6.41x10 <sup>1</sup>		
Limiting nutrients		_			_		_		
Silica	9.17	$2.67 \times 10^{-1}$	6.87x10 <sup>5</sup> *	1.43x10 <sup>-4</sup>	$1.68 \times 10^{-1}$	3.59x10 <sup>6</sup>	1.62x10 <sup>1</sup>		
Nitrogen	8.78	$5.89 \times 10^{-3}$	4.40x10 <sup>4</sup>	9.03x10 <sup>-7</sup>	$3.97 \times 10^{-3}$	2.01x10 <sup>6</sup>	2.25		
Phosphorus	7.01x10 <sup>-1</sup>	$1.24 \times 10^{-2}$	2.33x10 <sup>4</sup>	8.10x10 <sup>-7</sup>	1.56x10 <sup>-1</sup>	1.84x10 <sup>6</sup>	9.14		
Error (A)	2.43	$5.59 \times 10^{-2}$	3.28x10 <sup>4</sup>	3.92x10 <sup>-5</sup>	4.50×10	1.66x10 <sup>b</sup>	9.48		
Days	1.12x10 <sup>1</sup>	9.17x10 <sup>-2</sup>	2.68x10 <sup>3</sup>	1.16x10 <sup>-5</sup>	3.21x10 <sup>-1</sup>	1.82x10 <sup>6</sup>	1.83		
Interaction	_	-	-	_					
Days•Silica	1.56x10 <sup>1</sup>	$1.61 \times 10^{-1}$	6.60x10 <sup>3</sup>	2.03x10 <sup>-7</sup>	$4.36 \times 10^{-3}$	2.09x10 <sup>6</sup>	3.30x10		
Days•Nitrogen	6.39	2.43x10 <sup>-2</sup>	1.54x10 <sup>4</sup>	6.25x10 <sup>-8</sup>	1.08	1.80x10 <sup>6</sup>	6.17x10 <sup>1</sup>		
Days.Phosphorus	6.38x10 <sup>-2</sup>	$2.05 \times 10^{-2}$	3.33x10 <sup>2</sup>	2.11x10 <sup>-5</sup>	3.76	$1.54 \times 10^{6}$	5.43		
Error (b)	4.02	3.78x10 <sup>-2</sup>	8.89x10 <sup>3</sup>	$2.05 \times 10^{-5}$	1.49	1.59x10 <sup>6</sup>	1.98x10 <sup>1</sup>		

Table Al. Mean square values for the sources of variation investigated in the factorial analysis of nutrient-starvation experiments

\*p<0.05.

Experi- ment	Source of variation	Variable							
		Lipid	Glucan	Protein	Chloro- phyll	Chloro- phyll/ Phaeo- phytin	Protein/ Glucan	Lipid/ Glucan	
Silica	Supply rate Linear Quadratic Error	8.09x10 <sup>-6</sup> 4.01x10 <sup>-6</sup> 1.22x10 <sup>-5</sup> 4.37x10 <sup>-6</sup>	3.55x10 <sup>-6</sup> 6.22x10 <sup>-6</sup> 8.70x10 <sup>-7</sup> 1.43x10 <sup>-6</sup>	5.23x10 <sup>-3</sup> 4.78x10 <sup>-3</sup> 5.67x10 <sup>-3</sup> 8.25x10 <sup>-4</sup>	9.51x10-3 3.67x10 <sup>-5</sup> 1.90x10 <sup>-2</sup> 5.23x10 <sup>-2</sup>	  	1.03x10 <sup>3</sup> 1.16x10 <sup>3</sup> 8.87x10 <sup>2</sup> 1.74x10 <sup>2</sup>	1.75* 2.04* 1.47 3.44x10 <sup>-1</sup>	
Nitrogen	Supply rate Linear Quadratic Error	4.36x10 <sup>-5***</sup> 8.72x10 <sup>-5***</sup> 1.00x10 <sup>-7</sup> 1.24x10 <sup>-6</sup>	5.40x10 <sup>-6***</sup> 1.08x10 <sup>-5***</sup> 1.00x10 <sup>-8</sup> 1.90x10 <sup>-7</sup>	1.70x10 <sup>-1***</sup> 3.30x10 <sup>-1***</sup> 2.08x10 <sup>-3</sup> 4.31x10 <sup>-3</sup>	5.26x10 <sup>-2</sup> 1.40x10 <sup>-2</sup> 9.12x10 <sup>-2</sup> 1.24x10 <sup>-2</sup>	3.34 4.31 2.36 5.38	5.73x10 <sup>3</sup> 1.14x10 <sup>4</sup> 1.02x10 <sup>1</sup> 2.49x10 <sup>3</sup>	1.36x10 <sup>-2</sup> 1.37x10 <sup>-2</sup> 1.34x10 <sup>-2</sup> 1.45	
Phosphorus	Supply rate Linear Quadratic Error	1.48x10 <sup>-5*</sup> 2.58x10 <sup>-5**</sup> 3.77x10 <sup>-6</sup> 9.10x10 <sup>-7</sup>	4.51x10 <sup>-6***</sup> 8.43x10 <sup>-6***</sup> 5.80x10 <sup>-7</sup> 4.00x10 <sup>-8</sup>	1.49x10 <sup>-2</sup> 2.59x10 <sup>-2</sup> 3.76x10 <sup>-3</sup> 4.13x10 <sup>-3</sup>	6.03×10 <sup>-2</sup> 7.72×10 <sup>-2</sup> 4.35×10 <sup>-2</sup> 1.37×10 <sup>-2</sup>	  	3.89x10 <sup>4</sup> 6.72x10 <sup>4</sup> 1.08x10 <sup>4</sup> 1.02x10 <sup>4</sup>	1.80 3.06 5.44×10 <sup>-1</sup> 3.61	

Table A2. Mean square values for the sources of variation investigated in the Analysis of Variance for nutrient-stressed experiments

\*p<0.05. \*\*p<0.01. \*\*\*p<0.001.