

11-1997

## Thermal Properties of Starch from 62 Exotic Maize (*Zea mays* L.) Lines Grown in Two Locations

K.Y. Ng

*Iowa State University*

Linda M. Pollak

*United States Department of Agriculture*

Susan A. Duvick

*United States Department of Agriculture*

Pamela J. White

*Iowa State University, pjwhite@iastate.edu*

Follow this and additional works at: [http://lib.dr.iastate.edu/fshn\\_ag\\_pubs](http://lib.dr.iastate.edu/fshn_ag_pubs)

 Part of the [Agricultural Education Commons](#), [Food Biotechnology Commons](#), [Food Processing Commons](#), and the [Human and Clinical Nutrition Commons](#)

The complete bibliographic information for this item can be found at [http://lib.dr.iastate.edu/fshn\\_ag\\_pubs/130](http://lib.dr.iastate.edu/fshn_ag_pubs/130). For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

This Article is brought to you for free and open access by the Food Science and Human Nutrition at Iowa State University Digital Repository. It has been accepted for inclusion in Food Science and Human Nutrition Publications by an authorized administrator of Iowa State University Digital Repository. For more information, please contact [digirep@iastate.edu](mailto:digirep@iastate.edu).

---

# Thermal Properties of Starch from 62 Exotic Maize (*Zea mays* L.) Lines Grown in Two Locations

## Abstract

The variability in thermal properties among 62 S<sub>3</sub> lines derived from a high-yielding exotic corn (*Zea mays*) population, Antigua 1 (PI 484990), was evaluated by differential scanning calorimetry (DSC). The S<sub>3</sub> lines were grown in Puerto Rico (1990–1991) and Georgia (1994). Separate single-kernel starch extractions for five kernels (five replicates) from each line grown in each location were performed, and the starch was analyzed. The DSC values reported included gelatinization onset ( $T_{oG}$ ), range ( $R_G$ ), enthalpy ( $\Delta H_G$ ), and peak height index (PHI) and retrogradation onset ( $T_{oR}$ ), range ( $R_R$ ), enthalpy ( $\Delta H_R$ ), and percent retrogradation (%R) (an indication of the stability of gelatinized starch after storing at 4°C for 7 days). Significant differences ( $P < 0.05$ ) were found among the 62 lines of Antigua 1 for  $T_{oG}$ ,  $R_G$ , and PHI and highly significant differences ( $P < 0.01$ ) were found for  $\Delta H_G$ . The starches from plants grown in Georgia (1994) had significantly ( $P < 0.05$ ) greater  $T_{oG}$ ,  $\Delta H_G$ , and PHI but a significantly lower  $R_G$  than those from Puerto Rico (1990–1991). These data suggest that the starch from plants grown in Georgia (1994) might have a greater degree of crystallinity than that from Puerto Rico (1990–1991). None of the retrogradation values were significantly different among starches of the 62 lines of Antigua 1 and the starches from plants grown in the two locations.

## Disciplines

Agricultural Education | Food Biotechnology | Food Processing | Food Science | Human and Clinical Nutrition

## Comments

This article is from *Cereal Chemistry*, November 1997, 74(6); 837-841. Doi: <http://dx.doi.org/10.1094/CCHEM.1997.74.6.837>.

## Rights

Works produced by employees of the U.S. Government as part of their official duties are not copyrighted within the U.S. The content of this document is not copyrighted.

## Thermal Properties of Starch from 62 Exotic Maize (*Zea mays* L.) Lines Grown in Two Locations<sup>1</sup>

K.-Y. NG,<sup>2</sup> L. M. POLLAK,<sup>3</sup> S. A. DUVICK,<sup>3</sup> and P. J. WHITE<sup>2,4</sup>

### ABSTRACT

Cereal Chem. 74(6):837–841

The variability in thermal properties among 62 *S*<sub>3</sub> lines derived from a high-yielding exotic corn (*Zea mays*) population, Antigua 1 (PI 484990), was evaluated by differential scanning calorimetry (DSC). The *S*<sub>3</sub> lines were grown in Puerto Rico (1990–1991) and Georgia (1994). Separate single-kernel starch extractions for five kernels (five replicates) from each line grown in each location were performed, and the starch was analyzed. The DSC values reported included gelatinization onset ( $T_{oG}$ ), range ( $R_G$ ), enthalpy ( $\Delta H_G$ ), and peak height index (PHI) and retrogradation onset ( $T_{oR}$ ), range ( $R_R$ ), enthalpy ( $\Delta H_R$ ), and percent retrogradation (%*R*) (an indication of the stability of gelatinized starch after storing at

4°C for 7 days). Significant differences ( $P < 0.05$ ) were found among the 62 lines of Antigua 1 for  $T_{oG}$ ,  $R_G$ , and PHI and highly significant differences ( $P < 0.01$ ) were found for  $\Delta H_G$ . The starches from plants grown in Georgia (1994) had significantly ( $P < 0.05$ ) greater  $T_{oG}$ ,  $\Delta H_G$ , and PHI but a significantly lower  $R_G$  than those from Puerto Rico (1990–1991). These data suggest that the starch from plants grown in Georgia (1994) might have a greater degree of crystallinity than that from Puerto Rico (1990–1991). None of the retrogradation values were significantly different among starches of the 62 lines of Antigua 1 and the starches from plants grown in the two locations.

The importance of starch from completely “natural” sources has grown dramatically in the food industry during recent years. The introduction of starches containing single-, double-, and triple-mutant combinations with properties similar to chemically modified starches has resulted in a number of patents (Wurzburg and Ferguson 1984; Friedman et al 1988a–g; Friedman et al 1989a, b; Mauro et al 1991; Furscik and Mauro 1991a, b; Furscik and Deboer 1991a, b; Zallie et al 1984; Pearlstein and Ulrich 1994; White et al 1994; Hauber et al 1996).

Screening germplasm is important because the corn hybrids grown in the United States use only about 5% of the total germplasm available worldwide (Goodman 1985). This limited use of germplasm can translate into increased genetic vulnerability to insects, diseases, and stress and eventually affects agronomic performance. Tracy (1990) stressed the importance of exotic maize germplasm as an excellent source of improved quality and, possibly, improved agronomic performance in corn because “much of the maize grown outside of the United States is consumed directly by humans and has undergone centuries of selection for flavors, aromas and textures.”

Differential scanning calorimetry (DSC) has been used to evaluate thermal properties of starches since Stevens and Elton (1971) first reported the DSC properties of cereal starches. A differential scanning calorimeter is easy to operate and requires only a small sample size. In addition, it is relatively rapid compared with more traditional methods of studying starch gelatinization, making it useful for screening maize germplasm (Campbell et al 1995). Several studies have used DSC to screen corn germplasm. White et al (1990) reported differences in thermal properties revealed by DSC among and within five open-pollinated corn populations. The greatest differences were observed for gelatini-

zation onset ( $T_o$ ), range ( $R_G$ ), and total enthalpy ( $\Delta H$ ). Campbell et al (1995) reported significant differences in DSC properties among a set of exotic and domestic inbred corn lines. Starches from exotic lines generally had lower  $T_o$ , peak temperature ( $T_p$ ), and  $\Delta H$ . Li et al (1994) reported large variations in thermal properties among and within 35 tropical and subtropical maize populations. The means of  $T_o$ ,  $T_p$ , peak conclusion ( $T_c$ ), and  $\Delta H$  were 66.3°C (64.3–69.6°C range), 71.0°C (70.1–73.9°C range), and 77.8°C (76.8–79.6°C range), and 2.5 cal/g (2.0 to 2.9 cal/g range), respectively.

Properties of endosperm starches are affected by environmental factors during the development of plants. Ferguson and Zuber (1962) reported a negative relationship between amylose synthesis and temperature in maize. Asaoka et al (1984) examined the effect of environmental temperature on properties of rice starch. A higher environmental temperature seemed to increase the amount of long B chains and decrease that of short chains of amylopectin in rice starch. Shi et al (1994) studied the effect of temperature during grain filling on starches from several wheat cultivars in growth chambers. Elevated temperatures caused low test weight and shriveled kernels, reduced starch accumulation and starch granule size, and deformed starch granules. In addition, the starch lipid levels increased markedly, amylose content increased slightly, and gelatinization temperatures of starches were greater. White et al (1991) found that starches from corn grown under tropical conditions rather than temperate conditions gave an elevated and narrow gelatinization temperature range; after retrogradation, there was an elevated onset temperature.

Our objectives were to study the thermal properties of 62 *S*<sub>3</sub> lines of an exotic corn population and to examine the effect of growing location on the thermal properties of the same 62 lines.

### MATERIALS AND METHODS

#### Plant Material

Antigua 1 (PI 484990) is a native corn (*Zea mays* L.) accession with yellow semident grain that originated from the Caribbean Island of Antigua. Development of inbred lines proceeded as follows. The first self-pollination was done during the winter near Homestead, FL, in 1989–1990, and the second was done during the summer near Isabela, Puerto Rico, in 1990. Sixty-two lines (*S*<sub>3</sub> generation) of Antigua 1 were developed during the winter near Isabela, Puerto Rico, in 1990–1991 (planting date December 17,

<sup>1</sup>Journal Paper J-17440, Iowa Agriculture and Home Economics Experiment Station, Ames, IA. Projects 3396 and 3082.

<sup>2</sup>Graduate student and professor, respectively, Department of Food Science and Human Nutrition and Center for Crops Utilization Research, Iowa State University, Ames, IA 50011.

<sup>3</sup>Research geneticist and biologist, respectively, USDA-ARS, Corn Insects and Crop Genetics Research Unit, Department of Agronomy, Iowa State University, Ames, IA 50011.

<sup>4</sup>Corresponding author. E-mail: pjwhite@iastate.edu Phone: 515/294-3011. Fax: 515/294-8181.

1990) and near Tifton, GA, during the summer of 1994 (planting date April 5, 1994), using the same  $S_2$  plants grown in Puerto Rico in 1990–1991. The average monthly temperatures and precipitation of the growing seasons for  $S_3$  development in Puerto Rico and Georgia are shown in Table I. Grain filling occurred between January and February in Puerto Rico and between June and July in Georgia. The field experiments were planted in a completely randomized design. One random ear in each row was selected for DSC analysis. Ears were dried at 38°C for 48 hr to a moisture content of  $\approx 13\%$ . All samples were stored at 4°C in 40% rh until analyzed.

### Starch Extraction

Single-kernel starch isolations were conducted on five random kernels from each ear of each line according to the method of White et al (1990), with slight modifications (Krieger et al 1997). A single kernel was degermed by hand, and the pericarp was removed. The endosperm was put in a 50-ml centrifuge tube, and 5 ml of distilled water was added. A Tekmar tissue homogenizer (600 W, Ultra-Turrax T25, Cincinnati, OH) at  $2,052 \times g$  (17,000 rpm) was used to blend the endosperm and distilled water for 25 sec in the centrifuge tube. The filtering and drying processes were conducted as described by White et al (1990). After extraction, samples were dried at room temperature (22–24°C) with a fan circulating air over the starch and then stored at room temperature in air-tight vials until evaluation by DSC.

TABLE I  
Average Monthly Temperatures (°C) and Total Precipitation Near Isabela, Puerto Rico (1990–1991) and Tifton, GA (1994)

Location Month	Temperature			Precipitation (cm)
	Average Maximum	Average Minimum	Average	
Puerto Rico <sup>a</sup>				
November	30.9	21.7	26.3	10.3
December	28.9	20.2	24.6	15.2
January	28.9	19.7	24.3	6.6
February	28.8	19.9	24.4	15.8
March	29.1	20.0	24.6	6.5
Georgia <sup>b</sup>				
April	27.2	12.3	19.8	6.7
May	28.7	15.0	21.9	4.0
June	31.6	20.4	26.0	23.5
July	31.6	20.9	26.3	19.2
August	31.4	20.1	25.8	17.4

<sup>a</sup> National Climatic Data Center (1990, 1991). Grain-filling occurred during January and February.

<sup>b</sup> National Climatic Data Center (1994). Grain-filling occurred during June and July.




Fig. 1. Effects of growing location on gelatinization onset temperature ( $T_{oG}$ ) of 62  $S_3$  corn lines of Antigua 1.

### Differential Scanning Calorimetry

For DSC analysis, a Perkin-Elmer DSC-7 (Norwalk, CT) equipped with a thermal analysis data station was used. Approximately 4 mg of starch dwb was weighed in an aluminum pan, and 8  $\mu$ L of distilled water was added. Samples were heated from 30 to 110°C at a rate of 10°C/min. The starch gelatinization peaks and the peak caused by melting the amylose-lipid complex were completed at 110°C. Further heating provided no additional information about the starches. For all the DSC analyses, an empty pan with 8 mg of distilled water was used as a reference. The DSC parameters recorded included onset ( $T_{oG}$ ), range ( $R_G$ ), enthalpy ( $\Delta H_G$ ), and peak height index (PHI). Onset ( $T_{oG}$ ), peak ( $T_{pG}$ ), and enthalpy ( $\Delta H_G$ ) were computed directly by the DSC software. At the water level used, the DSC endotherms were essentially symmetrical, which allowed the total gelatinization range to be computed as  $2(T_{pG} - T_{oG})$  according to the method of Krueger et al (1987). The PHI, which is the ratio  $\Delta H_G / (T_{pG} - T_{oG})$ , was calculated to allow quantitative evaluation of variations in peak shape (Krueger et al 1987). To determine retrogradation characteristics, starch samples used for gelatinization were stored for 7 days at 4°C and rescanned on the DSC from 30 to 90°C at the same heating rate as was used for the gelatinization run. By this method, we determined onset temperature ( $T_{oR}$ ), range ( $R_R$ ), enthalpy ( $\Delta H_R$ ), and starch gel retrogradation (%R), as described by White et al (1989). Further heating above 90°C provided no additional changes on the DSC thermograms. The %R is a measure of the tendency of the gelatinized starch to retrograde after storage. The lower the %R, the less tendency there is to retrograde, and the more stable is the gelatinized starch paste. All DSC values reported are the average of five scans, one from each single-kernel starch extraction. Enthalpies were calculated on a starch dwb.

### Statistical Analysis

A completely randomized design was used to investigate the DSC properties of the starch among 62 lines of the population Antigua 1 and between the two locations. Analysis of variance (ANOVA) and least significant difference (LSD) analyses ( $\alpha = 0.05$ ) were computed by the Statistical Analysis System (SAS Institute, Cary, NC).

## RESULTS AND DISCUSSION

### Variations Among Lines

There were significant differences ( $P < 0.05$ ) for  $T_{oG}$ ,  $R_G$ , and PHI and highly significant differences ( $P < 0.01$ ) for  $\Delta H_G$  among the 62 Antigua 1 lines. Individual values are not shown (Ng




Fig. 2. Effects of growing location on gelatinization range ( $R_G$ ) of 62  $S_3$  corn lines of Antigua 1.

1997). However, scatter plots of the values for the 62 lines of these DSC parameters are shown in Figs. 1–4. None of the retrogradation characteristics, however, were significantly different among the lines. In each generation of self-pollination, the percentage of heterozygotes decreases by 50%. These corn kernels were S<sub>3</sub> lines, so 87.5% of genes within a line would be expected to be fixed. The fact that the lines are fixed for different genes likely accounts for the differences in thermal properties observed in this research. Antigua 1-20 had the lowest T<sub>oG</sub>, 66.1°C, whereas Antigua 1-56 had the highest T<sub>oG</sub>, 69.7°C when averaged from both locations (data not shown). For R<sub>G</sub>, Antigua 1-56 had the narrowest R<sub>G</sub>, 5.7°C, whereas Antigua 1-20 had the widest R<sub>G</sub>, 10.7°C. The ΔH<sub>G</sub> ranged from 2.63 cal/g for Antigua 1-20 and 3.35 cal/g for Antigua 1-57. For PHI, Antigua 1-20 had the lowest value, 0.51, whereas Antigua 1-57 had the highest value, 1.17 (Antigua 1-56 had a similar high PHI of 1.15). The respective values of T<sub>oR</sub>, R<sub>R</sub>, ΔH<sub>R</sub>, and %R ranged from 39.4°C (Antigua 1-27) to 43.4°C (Antigua 1-55), 19.0°C (Antigua 1-51) to 24.2°C (Antigua 1-80), 1.32 cal/g (Antigua 1-55) to 1.67 cal/g (Antigua 1-25), and 41.4% (Antigua 1-55) to 55.8% (Antigua 1-67).

Li et al (1994) evaluated the intrapopulation variability of thermal properties within 35 tropical maize populations. Significant

variations were found within all populations for T<sub>oG</sub>, T<sub>p</sub>, and T<sub>c</sub>, except for T<sub>c</sub> in two populations (Tuson Bai III and Tuson × (Mo44[2] × Tuson Cub) 75% F<sub>3</sub>). Thirty of thirty-five populations showed significant variation for R<sub>G</sub>, and only six populations showed significant variation for ΔH<sub>G</sub>. The maize population Yungueño showed the greatest variation for T<sub>oG</sub> (64.1–68.2°C), R<sub>G</sub> (11.0–16.8°C), and ΔH<sub>G</sub> (2.1–3.8 cal/g). White et al (1990) also evaluated the variability of thermal properties within the same population with 20 plants from a Corn Belt population, Weekly. The T<sub>oG</sub>, R<sub>G</sub>, and ΔH<sub>G</sub> ranged from 59.6 to 64.7°C, 10.5 to 19.5°C, and 1.83 to 2.83 cal/g, respectively. A comparison of these values to those found in the current study on Antigua 1 lines (66.1–69.7°C for T<sub>oG</sub>, 5.7–10.7°C for R<sub>G</sub>, and 2.63–3.45 cal/g for ΔH<sub>G</sub>, respectively) (data not shown) showed greater variation in populations Yungueño and Weekly than in Antigua 1. In the current study, the fact that only the gelatinization parameters and none of the retrogradation parameters were different among the Antigua 1 lines suggests that the native states of the starches, such as the degree of crystallinity, were different. Krueger et al (1987) reported that T<sub>o</sub> and ΔH increased, but the range decreased in dent corn lines that had undergone an annealing treatment (heating starch samples in water at a subgelatinization temperature). They




Fig. 3. Effects of growing location on gelatinization enthalpy (ΔH<sub>G</sub>) of 62 S<sub>3</sub> corn lines of Antigua 1.

Fig. 4. Effects of growing location on peak height index (PHI) of 62 S<sub>3</sub> corn lines of Antigua 1.

TABLE II  
Means and Ranges of Gelatinization and Retrogradation Values for 62 S<sub>3</sub> lines of Antigua 1 Grown at Two Locations Determined by Differential Scanning Calorimetry (DSC)

Location	Gelatinization				Retrogradation			
	T <sub>oG</sub> <sup>a</sup>	R <sub>G</sub> <sup>b</sup>	ΔH <sub>G</sub> <sup>c</sup>	PHI <sup>d</sup>	T <sub>oR</sub> <sup>e</sup>	R <sub>R</sub> <sup>f</sup>	ΔH <sub>R</sub> <sup>g</sup>	%R <sup>h</sup>
Puerto Rico (1990–1991)								
Mean	67.2	8.4	3.04	0.76	41.1	21.3	1.49	49.1
Min	65.0	5.0	2.66	0.43	37.6	18.6	1.14	40.1
Max	69.6	12.7	3.33	1.28	44.3	26.8	1.69	56.8
Georgia (1994)								
Mean	69.2	6.9	3.15	0.96	41.4	21.4	1.52	48.5
Min	66.8	4.6	2.60	0.56	38.2	18.6	1.26	38.5
Max	71.7	10.9	3.45	1.47	44.8	24.5	1.80	58.5
LSD <sub>0.05</sub> <sup>i</sup>	10.4	10.5	0.04	0.05	10.4	10.5	0.04	11.3

<sup>a</sup> Gelatinization onset temperature (°C).

<sup>b</sup> Gelatinization range (°C).

<sup>c</sup> Enthalpy of gelatinization (cal/g).

<sup>d</sup> Peak height index = ΔH<sub>G</sub> / (T<sub>pG</sub> - T<sub>oG</sub>), as described by Krueger et al (1987).

<sup>e</sup> Retrogradation onset temperature (°C).

<sup>f</sup> Retrogradation range (°C).

<sup>g</sup> Enthalpy of retrogradation (cal/g).

<sup>h</sup> %R = (ΔH<sub>G</sub> / ΔH<sub>R</sub>) × 100.

<sup>i</sup> Least significant difference (α = 0.05).



proposed that the shifts were attributed to an increase in crystallinity of the starch molecules that occurred during annealing. Marchant and Blanshard (1978) proposed that annealing allows partial melting of some crystallites and a general realignment of starch chains in the amorphous phase, leading to an increase in hydrogen bonding. White et al (1990) attributed significant differences among five open-pollinated corn cultivars for  $T_{oG}$ , range, and  $\Delta H$  to the differences in the native alignment and hydrogen bonding of the starch molecules. Perhaps, some variations in the gelatinization characteristics among the lines in the current study were a result of different size distributions of starch granules. In a study by Knutson et al (1982), gelatinization temperature and  $\Delta H$  of amylo maize increased as granule size decreased; however, these trends were not observed in normal dent corn. Knutson et al (1982) also noted that larger starch granules gelatinized over a smaller range. Possibly, the range and PHI values in the current study might be useful in predicting starch granule size.

Starches with narrow gelatinization ranges would make production more efficient by making gelatinization more rapid (Hegenbart 1996). Antigua 1-56 and Antigua 1-5 had narrow gelatinization ranges of 5.7 and 5.8°C (data not shown), respectively. Li et al (1994) reported gelatinization ranges of 35 tropical and subtropical maize populations that varied between 9.8 and 13.0°C. The gelatinization ranges of starches from five open-pollinated corn populations studied by White et al (1990) varied between 8.7 and 16.4°C. The Antigua 1 lines may be worth further investigation for this and other functional properties that could be useful in possible industrial applications.

#### Variations Between Locations

The average values and ranges of the DSC parameters ( $T_{oG}$ ,  $R_G$ ,  $\Delta H_G$ , PHI,  $T_{oR}$ ,  $R_R$ ,  $\Delta H_R$ , and %R) of starch from corn grown at two locations are presented in Table II. The LSD ( $\alpha = 0.05$ ) for each parameter also is listed. The differences for  $T_{oG}$ ,  $R_G$ ,  $\Delta H_G$ , and PHI between locations were highly significant ( $P < 0.01$ ). The effects of growing location on  $T_{oG}$ ,  $R_G$ ,  $\Delta H_G$ , and PHI for all 62 lines are illustrated in Figs. 1–4. The  $T_{oG}$ ,  $\Delta H_G$ , and PHI in starches from corn grown in Puerto Rico (1990–1991) were significantly lower ( $P < 0.05$ ) than for corn grown in Georgia (1994). The  $R_G$ , on the other hand, was significantly greater for starches from corn grown in Puerto Rico than from corn grown in Georgia. As noted for differences among the 62 Antigua 1 lines, there were differences in gelatinization properties, but not in retrogradation properties, between the starches grown in the two locations. Again, the data suggest differences in the native state of starches from corn grown in the two locations. Starches from plants grown in Georgia, which had higher  $T_{oG}$ ,  $\Delta H_G$ , and PHI and narrower  $R_G$  (results similar to the effect of annealing), may have a higher degree of molecular order than those from Puerto Rico, as suggested by Krueger et al (1987). Perhaps part of the variation in the gelatinization properties of starches may have been a result of partial annealing due to the higher temperatures and precipitation in Georgia than in Puerto Rico during the grain-filling period (June and July in Georgia and January and February in Puerto Rico, respectively [Table I]).

Increased temperatures during grain-filling cause a higher gelatinization temperature of starch from barley, wheat, and maize (Tester et al 1991, Shi et al 1994, Lu et al 1996). The increase in  $T_{oG}$  of starches from corn grown in Georgia in the current study, where higher temperatures occurred during the grain-filling period, agreed with those studies. Tester et al (1991) attributed the increase in gelatinization temperature of starch to differences in the perfection of amylopectin crystallinity, which increased with the temperature at which barley was grown. Lu et al (1996) reported that the increase in  $T_{oG}$  of starches from corn grown at higher temperatures correlated with an increase in chain length of the medium branch-chain fractions of amylopectin. They did not report information on the retrogradation properties of the starches.

Variations in starch thermal properties of corn between the two locations studied in this research might have been a combination of many factors, including location effects, year differences, and genotype by environment interactions. Possibly, the longer storage period of the Puerto Rico corn than of the Georgia corn affected these properties, although the seed was kept in controlled environments. Further experiments might be necessary to isolate these factors, if possible, to determine their effects on starch thermal properties. The cause of the geographic differences and the relations between thermal starch properties, structural variations in the starch, and production characteristics of the corn may be of future interest.

#### CONCLUSIONS

Significant differences were found among the 62  $S_3$  lines of one corn population, Antigua 1, when averaged over two locations for the gelatinization properties ( $T_{oG}$ ,  $R_G$ ,  $\Delta H_G$ , and PHI) but not for retrogradation characteristics. Highly significant differences ( $P < 0.01$ ) were observed for the gelatinization properties of the lines grown in two locations, but not for the retrogradation properties. These differences in starch thermal properties suggest that there may be differences in native alignment and hydrogen bonding within the starch molecules.

#### ACKNOWLEDGMENTS

We thank N. Burkhart for her assistance with starch isolation.

#### LITERATURE CITED

- Asaoka, M., Okuno, K., Sugimoto, Y., Kawakami, J., and Fuwa, H. 1984. Effect of environment temperature during development of rice plants on some properties of endosperm starch. *Starch Staerke* 36:189-193.
- Campbell, M. R., Pollak, L. M., and White, P. J. 1995. Genetic variation for starch thermal and functional properties among nonmutant maize inbreds. *Cereal Chem.* 72:281-286.
- Ferguson, V. L., and Zuber, M. S. 1962. Influence of environment on amylose content of maize endosperm. *Crop Sci.* 2:209-211.
- Friedman, R. B., Gottneid, D. J., Faron, E. J., Pustek, F. J., and Katz, F. R. 1988a. Food stuffs containing starch of an *amylose-extender dull* genotype. U.S. patent 4,790,997.
- Friedman, R. B., Gottneid, D. J., Faron, E. J., Pustek, F. J., and Katz, F. R. 1988b. Food stuffs containing starch of a *dull sugary-2* genotype. U.S. patent 4,792,458.
- Friedman, R. B., Gottneid, D. J., Faron, E. J., Pustek, F. J., and Katz, F. R. 1988c. Starch of a *wxsh1* genotype and products produced therefrom. U.S. patent 4,767,849.
- Friedman, R. B., Gottneid, D. J., Faron, E. J., Pustek, F. J., and Katz, F. R. 1988d. Starch of the *duh* genotype and products produced therefrom. U.S. patent 4,774,328.
- Friedman, R. B., Gottneid, D. J., Faron, E. J., Pustek, F. J., and Katz, F. R. 1988e. Food stuffs containing starch of a *dull waxy* genotype. U.S. patent 4,789,557.
- Friedman, R. B., Gottneid, D. J., Faron, E. J., Pustek, F. J., and Katz, F. R. 1988f. Starch of the *wxfl1* genotype and products produced therefrom. U.S. patent 4,789,738.
- Friedman, R. B., Gottneid, D. J., Faron, E. J., Pustek, F. J., and Katz, F. R. 1988g. Novel starch and products produced therefrom. U.S. patent 4,770,710.
- Friedman, R. B., Gottneid, D. J., Faron, E. J., Pustek, F. J., and Katz, F. R. 1989a. Food stuffs containing starch of an *amylose extender sugary-2* genotype. U.S. patent 4,798,735.
- Friedman, R. B., Gottneid, D. J., Faron, E. J., Pustek, F. J., and Katz, F. R. 1989b. Food stuffs containing starch of a *waxy shrunken-2* genotype. U.S. patent 4,801,470.
- Furcsik, S. L., and Deboer, E. 1991a. *duh* Batter starch for deep fat fried food. PCT international patent publication WO 9101651.
- Furcsik, S. L., and Deboer, E. 1991b. *aedu* Batter starch for deep fat fried food. PCT international patent publication WO 9101652.
- Furcsik, S. L., and Mauro, D. J. 1991a. Starch jelly candy. U.S. patent 5,035,912.
- Furcsik, S. L., and Mauro, D. J. 1991b. Starch jelly candy. European

- patent 0462693.
- Goodman, M. M. 1985. Exotic maize germplasm: Status, prospects, and remedies. *Iowa State J. Res.* 59:497-527.
- Hauber, R., Friedman, R., and Katz, F. 1996. Foodstuffs containing a *waxy waxy amylose extender* starch. U.S. Patent 5,576,048.
- Hegenbart, S. 1996. Understanding starch functionality. *Food Prod. Design* January:23-34.
- Knutson, C. A., Khoo, U., Cluskey, J. E., and Inglett, G. E. 1982. Variation in enzyme digestibility and gelatinization behavior of corn starch granule fractions. *Cereal Chem.* 59:512-515.
- Krieger, K. M., Duvick, S. A., Pollak, L. M., and White, P. J. 1997. Thermal properties of corn starch extracted with different blending methods: Microblender and homogenizer. *Cereal Chem.* 74:553-555.
- Krueger, B. R., Knutson, C. A., Inglett, G. E., and Walker, C. E. 1987. A differential scanning calorimetry study on the effect of annealing on gelatinization behavior of corn starch. *J. Food Sci.* 52:715-718.
- Li, J., Berke, T. G., and Glover, D. V. 1994. Variation for thermal properties of starch in tropical maize germplasm. *Cereal Chem.* 71:87-90.
- Lu, T.-J., Jane, J.-L., Keeling, P. L., and Singletary, G. W. 1996. Effects of ear developmental temperature on fine structure of maize starch. *Carbohydrate Res.* 282:157-170.
- Marchant, J. L., and Blanshard, J. M. V. 1978. Studies of the dynamics of the gelatinization of starch granules employing a small angle light scattering system. *Starch Staerke* 30:257-264.
- Mauro, D. J., Furscik, S. L., Katz, F. R., Faron, E. J., Gottneid, D. J., and Pustek, F. J. 1991. Food stuff containing *aewx* starch. U.S. patent 5,009,911.
- National Climatic Data Center. 1990. Climatological Data: Puerto Rico and Virgin Islands. Vol. 36. National Climatic Data Center: Asheville, NC.
- National Climatic Data Center. 1991. Climatological Data: Puerto Rico and Virgin Islands. Vol. 37. National Climatic Data Center: Asheville, NC.
- National Climatic Data Center. 1994. Georgia Monthly Climatological Data. National Climatic Data Center: Asheville, NC.
- Ng, K.-Y. 1997. Growth development and growing location effects on thermal properties of starch from mutant and exotic corn. MS thesis Iowa State University: Ames, IA.
- Pearlstein, R. W., and Ulrich, J. F. 1994. Starch and grain with a novel genotype. PCT international patent publication WO 9422291.
- Shi, Y. C., Seib, P. A., and Bernardin, J. E. 1994. Effects of temperature during grain-filling on starches from six wheat cultivars. *Cereal Chem.* 71:369-383.
- Stevens, D. J., and Elton, G. A. H. 1971. Thermal properties of the starch/water system. *Starch Staerke* 23:8-11.
- Tester, R. F., South, J. B., Morrison, W. R., and Ellis, R. P. 1991. The effects of ambient temperature during the grain-filling period on the composition and properties of starch from four barley genotypes. *J. Cereal Sci.* 13:113-127.
- Tracy, W. F. 1990. Potential contributions of five exotic maize populations to sweet corn improvement. *Crop Sci.* 30:918-923.
- White, P. J., Abbas, I., and Johnson, L. 1989. Freeze-thaw stability and refrigerated-storage retrogradation of starches. *Starch Staerke* 41:176-180.
- White, P. J., Abbas, I. R., Pollak, L. M., and Johnson, L. A. 1990. Intra- and interpopulation variability of thermal properties of maize starch. *Cereal Chem.* 67:70-73.
- White, P. J., Pollak, L. M., and Burkhart, S. 1991. Thermal properties of starches from corn grown in temperate and tropical environments. (Abstr.) *Cereal Foods World* 36:704.
- White, P. J., Pollak, L. M., and Johnson, L. A. 1994. Starch thickened acidic foodstuffs and method of preparation. U.S. patent 5,365,655.
- Wurzburg, O. B., and Ferguson, V. L. 1984. Starch thickener characterized by improved low-temperature stability. U.S. patent 4,428,972.
- Zallie, J., Trimble, R., and Bell, H. 1984. Bread containing *wxsu2* genotype starch as an anti-stalent. U.S. patent 4,615,888.

[Received June 2, 1997. Accepted August 27, 1997.]