

# Tissue-nonspecific alkaline phosphatase and plasma cell membrane glycoprotein-1 are central antagonistic regulators of bone mineralization

Lovisa Hessel<sup>\*†</sup>, Kristen A. Johnson<sup>‡</sup>, H. Clarke Anderson<sup>§</sup>, Sonoko Narisawa<sup>\*</sup>, Adnan Sali<sup>¶</sup>, James W. Goding<sup>¶</sup>, Robert Terkeltaub<sup>‡</sup>, and José Luis Millán<sup>\*¶</sup>

<sup>\*</sup>The Burnham Institute, 10901 North Torrey Pines Road, La Jolla, CA 92037; <sup>†</sup>Medical Biosciences, Medical Genetics, Umeå University, S-901 85 Umeå, Sweden; <sup>‡</sup>Veterans Administration Medical Center/University of California, 3350 La Jolla Village Drive, San Diego, CA 92161; <sup>§</sup>Department of Pathology and Laboratory Medicine, University of Kansas Medical Center, 3901 Rainbow Boulevard, Kansas City, KS 66160; and <sup>¶</sup>Department of Pathology and Immunology, Monash University Medical School, Commercial Road, Prahran, Victoria 3181, Australia

Edited by Victor A. McKusick, Johns Hopkins University School of Medicine, Baltimore, MD, and approved May 20, 2002 (received for review February 1, 2002)

**Osteoblasts mineralize bone matrix by promoting hydroxyapatite crystal formation and growth in the interior of membrane-limited matrix vesicles (MVs) and by propagating the crystals onto the collagenous extracellular matrix. Two osteoblast proteins, tissue-nonspecific alkaline phosphatase (TNAP) and plasma cell membrane glycoprotein-1 (PC-1) are involved in this process. Mutations in the TNAP gene result in the inborn error of metabolism known as hypophosphatasia, characterized by poorly mineralized bones, spontaneous fractures, and elevated extracellular concentrations of inorganic pyrophosphate (PP<sub>i</sub>). PP<sub>i</sub> suppresses the formation and growth of hydroxyapatite crystals. PP<sub>i</sub> is produced by the nucleoside triphosphate pyrophosphohydrolase activity of a family of isozymes, with PC-1 being the only member present in MVs. Mice with spontaneous mutations in the PC-1 gene have hypermineralization abnormalities that include osteoarthritis and ossification of the posterior longitudinal ligament of the spine. Here, we show the respective correction of bone mineralization abnormalities in knockout mice null for both the TNAP (*Akp2*) and PC-1 (*Enpp1*) genes. Each allele of *Akp2* and *Enpp1* has a measurable influence on mineralization status *in vivo*. *Ex vivo* experiments using cultured double-knockout osteoblasts and their MVs demonstrate normalization of PP<sub>i</sub> content and mineral deposition. Our data provide evidence that TNAP and PC-1 are key regulators of the extracellular PP<sub>i</sub> concentrations required for controlled bone mineralization. Our results suggest that inhibiting PC-1 function may be a viable therapeutic strategy for hypophosphatasia. Conversely, interfering with TNAP activity may correct pathological hyperostification because of PP<sub>i</sub> insufficiency.**

In the process of bone formation, osteoblasts mineralize the matrix by promoting the seeding of basic calcium phosphate crystals of hydroxyapatite in the sheltered interior of shed membrane-limited matrix vesicles (MVs) and by propagating hydroxyapatite mineral onto the collagenous extracellular matrix (osteoid; refs. 1 and 2). Tissue-nonspecific alkaline phosphatase (TNAP), an isozyme of a family of four homologous human alkaline phosphatase genes (3), plays a role in bone matrix mineralization. Deactivating mutations in the TNAP gene causes the inborn error of metabolism known as hypophosphatasia (4), characterized by poorly mineralized cartilage (rickets) and bones (osteomalacia), spontaneous bone fractures, and elevated extracellular inorganic pyrophosphate (PP<sub>i</sub>) concentrations (5). The severity and expressivity of hypophosphatasia depends on the nature of the TNAP mutation (6). TNAP is present in MVs (7), and it has been proposed that the inorganic phosphate (P<sub>i</sub>)-generating activity of TNAP is required to generate the P<sub>i</sub> needed for hydroxyapatite crystallization (8–10). However, the ability of TNAP to hydrolyze PP<sub>i</sub> also has been hypothesized to be important to promote osteoblastic mineralization (11, 12), because PP<sub>i</sub> suppresses the formation and growth of hydroxy-

apatite crystals (13). In fact, heritable extracellular PP<sub>i</sub> deficiencies are models of ectopic calcification such as ankylosing spinal hyperostosis and pathologic soft-tissue ossification (14–16). PP<sub>i</sub> is produced by the nucleoside triphosphate pyrophosphohydrolase (NTPPPH) activity of a family of isozymes that include PC-1, B10/PDNP3, and autotaxin (17–19). However, PC-1 seems to be the only NTPPPH present in MVs (20). TNAP knockout (KO) mice (21–23) recapitulate the heritable metabolic disease hypophosphatasia (5), whereas PC-1-null mice display hypermineralization abnormalities similar to cartilage calcification in osteoarthritis (14) and ossification of the posterior longitudinal ligament of the spine (15).

We previously identified PC-1 as the likely NTPPPH isozyme to act on the same pathway with TNAP as antagonistic regulators of extracellular PP<sub>i</sub> concentrations (20). In this paper, we have tested the hypothesis that bone abnormalities caused by the lack of TNAP could be counterbalanced by the removal of PC-1 and vice versa. We show that bone mineralization in double-KO mice lacking both TNAP and PC-1 is essentially normal, providing evidence that TNAP and PC-1 are key regulators of bone mineralization by determining the normal steady-state levels of PP<sub>i</sub>. Our work suggests that TNAP and PC-1 may be useful therapeutic targets for the treatment of bone mineralization abnormalities.

## Materials and Methods

***Akp2* and *Enpp1* KO Mice.** The generation and characterization of the *Akp2* KO mice has been reported (22, 23). These *Akp2* KO mice were hybrids of C57BL/6 × 129/J mouse strains. The generation of the *Enpp1* KO mice has been reported briefly (24). These *Enpp1* KO mice were hybrids of C57BL/6 × 129/SvTerJ mouse strains. *Akp2/Enpp1* double-heterozygous mice were rederived into a C57BL/6 background, and the colony was maintained by brother–sister mating.

**Tissue Preparation and Morphological Analysis.** Whole-mount skeletal preparations were prepared by removing the skin and viscera of the mice before they were immersed in 100% ethanol for 1 week and in acetone for 1 week. Specimens were transferred into a 100% ethanol solution containing 0.01% Alizarin Red S (Sigma), 0.015% Alcian Blue 8GX (Sigma), and 0.5% acetic acid and were incubated for 3 weeks. They were destained with 1% (vol/vol) KOH/50% (vol/vol) glycerol solution. Cleared specimens were stored in 100% glycerol.

The lumbar spines of 10-day-old mice were fixed in PBS

This paper was submitted directly (Track II) to the PNAS office.

Abbreviations: MV, matrix vesicle; TNAP, tissue-nonspecific alkaline phosphatase; P<sub>i</sub>, inorganic phosphate; PP<sub>i</sub>, inorganic pyrophosphate; KO, knockout.

¶To whom reprint requests should be addressed. E-mail: millan@burnham.org.

containing 10% (vol/vol) formalin for 5 days. After washes in PBS-based sucrose solutions, samples were embedded in optimal cutting temperature (OCT) compound. Sections (16  $\mu\text{m}$ ) were stained with the von Kossa procedure, as described (25). The femurs of 10-day-old mice were fixed in 2.5% (vol/vol) glutaraldehyde in neutral 0.1 M cacodylate buffer for 24 h. Sections (1  $\mu\text{m}$ ) were stained with hematoxylin/eosin.

**Isolation of Primary Osteoblasts and Matrix Vesicles.** Calvarial bones from 3-day-old pups were incubated three times for 10 min in a solution containing 4 mM EDTA, 137 mM NaCl, 2.7 mM KCl, 3 mM  $\text{NaH}_2\text{PO}_4$  at pH 7.2. Specimens underwent seven 10-min digestions with EDTA-free buffer containing 180  $\mu\text{g}/\text{ml}$  collagenase type 2 (Worthington). The last five fractions were pooled and seeded at  $3\text{--}4 \times 10^4$  cells per  $\text{cm}^2$  in  $\alpha$ -MEM (GIBCO/BRL) containing 10% (vol/vol) heat-inactivated FCS. When the primary calvarial cultures were confluent, the complete  $\alpha$ -MEM media was supplemented with L-ascorbic acid (50  $\mu\text{g}/\text{ml}$ ) every third day and with either  $\beta$ -glycerophosphate (2.5 mM; ref. 26) or sodium phosphate (2.5 mM; ref. 18). The pericellular MV fraction was collected by collagenase digestion for 2 h at 37°C and sequential centrifugation, as described (20).

**PP<sub>i</sub> and Mineralization Assays.** PP<sub>i</sub> concentrations were determined by differential adsorption on activated charcoal of UDP-D-[6-<sup>3</sup>H]glucose (Amersham Pharmacia) from the reaction product 6-phospho [6-<sup>3</sup>H]gluconate, as described (19).

To quantify matrix calcification by the cultured calvarial osteoblasts, we used Alizarin Red S binding assay as described (27). In brief, calvarial osteoblasts (5,000 cells per well) were plated in a volume of 0.1 ml of medium in a flat bottom 96-well plate (Falcon). At each time point, the medium was removed, and 0.05 ml of diluted Alizarin Red S [0.5% (vol/vol) Alizarin Red S, pH 5.0] was added and incubated at 23°C for 10 min. The plates were washed four times with PBS before the addition of 0.1 ml of 10% (wt/vol) cetylpyridinium chloride for 10 min to release the remaining calcium-bound Alizarin Red S. The solution was collected, diluted at a ratio of 1:10 and read at OD<sub>570</sub> on a SpectraMAX microplate reader (Molecular Devices).

## Results

**Rescue from Bone Abnormalities *in Vivo*.** We observed clear evidence of rescue from rickets and osteomalacia in *Akp2/Enpp1* double-KO mice. The unmineralized areas in the calvaria (shown by the lack of Alizarin Red staining) and the absence of secondary ossification centers in the hind limb phalanges characteristic of the *Akp2* KO mice have been corrected in the *Akp2/Enpp1* double-KO mice (Fig. 1 *a–f*). We also found correction in the length and histological appearance of the growth plates of the double-KO mice (Fig. 1 *g–i*). *Akp2* KO mice show a characteristic thickening of the hypertrophic zone of the growth plates in the tibia (23). We observed a restoration of normal growth-plate thickness and cellularity in the *Akp2/Enpp1* double-KO mice. Interestingly, double-homozygous mice live up to 25 day without vitamin B6 administration (21, 23, 25), whereas we have never observed an *Akp2* KO mouse surviving longer than 14 days in our colony ( $n = 325$ ) in the absence of vitamin B6 supplementation.

We examined the lumbar spine of all genotypes and found differences in the degree of mineralization in the vertebral apophyses. Fig. 2*a* shows a low and a high magnification view of the apophyses of each genotype. We then counted and determined the ratio of the number of ossification centers containing visible mineral deposits to those containing no visible deposits (Fig. 2*b*). Ten-day-old *Enpp1* KO mice have precipitated mineral in every lumbar vertebral apophyses (100%), whereas in the corresponding areas of age-matched *Akp2*-null mice, mineral deposits are virtually absent (6%). More than half the lumbar

vertebrae of the *Akp2/Enpp1* double-KO mice had mineral deposits, a ratio comparable to that found in the wild-type (WT) mice. Each functional allele of *Akp2* and *Enpp1* contributes to the mineralization status of the spine (Fig. 2*b*). Note that even carriers (heterozygotes) of *Akp2*- and *Enpp1*-null mutations are affected.


**Correction of Mineralization and PP<sub>i</sub> Levels *ex Vivo*.** We then examined the ability of *Akp2/Enpp1* double-KO primary osteoblasts to mineralize bone nodules *in vitro*. Similar to the *Akp2* KO primary osteoblasts, the *Akp2/Enpp1* double-KO osteoblasts cultured with  $\beta$ -glycerophosphate as the principal phosphate source showed an absence of mineralization (not shown). However, incubating primary osteoblasts with sodium phosphate as the free phosphate source, a condition that better mimics the *in vivo* environment, allows for the detection of differences in the mineralization ability of osteoblasts of differing genotypes (Fig. 3*a*). *Akp2* KO osteoblasts cultured for 21 days precipitated significantly less mineral than WT osteoblasts, whereas osteoblasts of *Enpp1* KO mice precipitated significantly more mineral than control cells. A normal mineralization capacity was restored in the *Akp2/Enpp1* double-KO osteoblasts. Because inorganic pyrophosphate is a potent inhibitor of mineralization, we also quantified the amount of PP<sub>i</sub> in the MVs isolated from the cultured osteoblasts (Fig. 3*b*). MVs from *Akp2* KO osteoblasts had increased amounts of PP<sub>i</sub>, whereas those from the *Enpp1* KO osteoblasts had decreased levels of PP<sub>i</sub>. The MVs derived from the *Akp2/Enpp1* double-KO osteoblasts exhibited PP<sub>i</sub> concentrations that were comparable to those of WT MVs.

## Discussion


It has been proposed that the activity of TNAP is required to generate the P<sub>i</sub> needed for hydroxyapatite crystallization (8–10), and ATP and PP<sub>i</sub> have been explored as potential substrates for this effect (11, 28). However, the ability of TNAP to hydrolyze PP<sub>i</sub> also has been hypothesized to be important to promote calcification by hydrolyzing this potent inhibitor of mineralization (11–13, 29, 30). Our data provide evidence that in osteoblasts, TNAP and PC-1 are central, directly antagonistic regulators of the concentration of the matrix calcification inhibitor PP<sub>i</sub>. Restoration of extracellular PP<sub>i</sub> to normal levels in *Akp2/Enpp1* double-KO mice, in conjunction with correction of the hypomineralization resulting from the TNAP deficiency, points to the key role of PP<sub>i</sub> in this phenotypic rescue (Fig. 4). Our data support the hypothesis that the extracellular PP<sub>i</sub> pool largely determines the rate of hydroxyapatite crystal formation in bone tissue. PC-1 is a major contributor to the extracellular PP<sub>i</sub> pool, whereas the action of TNAP restricts PP<sub>i</sub> concentration and also contributes to the extracellular P<sub>i</sub> pool available for the formation of hydroxyapatite crystals. The concerted action of TNAP and PC-1 ensures that the optimal concentration of PP<sub>i</sub> is achieved under normal conditions. In hypophosphatasia, the PP<sub>i</sub> pool increases because of lack of TNAP's inorganic pyrophosphatase activity, and rickets/osteomalacia ensue as PP<sub>i</sub>'s inhibition of hydroxyapatite deposition is augmented. In PC-1 deficiency, the extracellular PP<sub>i</sub> pool decreases because of reduced production of PP<sub>i</sub>, and hypermineralization takes place as PP<sub>i</sub> inhibition of hydroxyapatite deposition is relaxed. The simultaneous deletion of TNAP and PC-1 correct the levels of extracellular PP<sub>i</sub>, and normal mineralization is re-established.

Our data allow us to conclude that TNAP's primary function in mineralizing tissues is to act in concert with PC-1 to fine tune PP<sub>i</sub> concentrations to maintain steady-state levels of PP<sub>i</sub> adequate for controlled mineralization. Additionally, through hydrolysis of PP<sub>i</sub> and nucleoside triphosphates, TNAP also may contribute to the pool of P<sub>i</sub> available for deposition as hydroxyapatite. However, the fact that the TNAP-null mice, albeit inefficiently, still deposit bone mineral argues that the P<sub>i</sub> needed






**Fig. 1.** Correction of skeletal mineralization and growth-plate structure in *Akp2/Enpp1* double-KO mice. Whole-mount skeletal preparations of the calvaria (a–c) and hind limb phalanges (d–f) show skeletal correction in *Akp2/Enpp1* double-KO mice. The tissues were stained with Alizarin Red that detects bone mineral and Alcian Blue that stains unmineralized osteoid at pH 5.5. WT, calvaria, and phalanges of a 20-day-old WT mouse (a and d). At 3 weeks of age, the *Enpp1* KO mice (not shown), do not show any evidence of hypermineralization in the calvaria or phalanges, and their whole-mount preparations are indistinguishable from those of WT mice. *Akp2* KO, calvaria and phalanges of a 20-day-old *Akp2* KO mouse (b and e). Note the absence of secondary ossification centers in the phalanges; *Akp2/Enpp1* KO, calvaria and phalanges of a 20-day-old *Akp2/Enpp1* double-KO mouse displaying correction of the mineralization defects (c and f). (g–i) Nondecalfied sections of tibial growth plates of 10-day-old mice stained with hematoxylin/eosin. The WT tibial growth plate (g) shows normal proliferative zone cells at the top. Lightly stained hypertrophic zone cells are seen in the middle of the field, and the metaphysis is present at the bottom. In contrast, the tibial growth plate from *Akp2* KO mice (h) shows a characteristic thickening, especially of the hypertrophic zone, as seen (23). Note the restoration of normal growth-plate thickness and cellularity in the tibial growth plate of *Akp2/Enpp1* double-KO mice (i). Bars in the figure indicate the thickness of the hypertrophic zone of the growth plate. (Magnification  $\times 310$ .)



**Fig. 2.** Nondecalcified lumbar spine sections stained by the von Kossa procedure to visualize phosphate deposits. (a) Low ( $\times 16$ ) and high ( $\times 240$ ) magnification view of the 10-day-old vertebral apophyses of WT, *Akp2* KO, *Enpp1* KO, and *Akp2/Enpp1* double-KO mice. Arrows point to mineralized secondary ossification centers in sample sections at low magnification. Note the absence of mineral deposits in sections from the *Akp2* KO mice. \*, relative position of the nucleus pulposus as a point of reference in the high magnification panels. (b) Graph plotting the contribution of each *Akp2* and *Enpp1* allele to the mineralization status of the lumbar vertebrae. The percentage of mineralized apophyses is plotted as a function of the *Akp2* and *Enpp1* genotypes.

for deposition is supplied by other, more major, sources, e.g., intestinal absorption (31). The bone nodule mineralization assay using  $\beta$ -glycerophosphate as the phosphate source reveals a lack of *in vitro* mineralization by *Akp2* KO primary osteoblasts, as we have reported (26), as well as by the *Akp2/Enpp1* double-KO osteoblasts (data not shown). The explanation is trivial, however, because free  $P_i$  cannot be produced from this organic compound in the absence of TNAP. However, the fact that MVs from patients with hypophosphatasia contain hydroxyapatite mineral in their interior (32) indicates that TNAP is not required for the initial deposition of bone mineral, and that phosphated organic compounds are not likely crucial substrates for TNAP's role in bone mineralization. A variation of the *in vitro* mineralization assay that uses free inorganic phosphate as the principal phosphate source (18) better mimics the physiological environment where there is an excess of free phosphate resulting from intestinal absorption (31). This assay has allowed us to detect




**Fig. 3.** Primary *Akp2/Enpp1* double-KO osteoblasts show a correction in their mineralization capacity and normalized  $PP_i$  concentrations in MVs. (a) *Akp2* KO osteoblasts show poor mineralization after 21 days in culture, whereas the cultures of *Enpp1* KO osteoblasts have a higher mineralization capacity compared with WT osteoblasts. Remarkably, the *Akp2/Enpp1* double-KO and the WT cultures display indistinguishable mineralization properties ( $n = 7$  mice each genotype, in replicates of 4). \*,  $P < 0.05$  compared with single gene KO osteoblasts. (b) The histogram displays the concentration of  $PP_i$  in the MVs of each genotype. *Akp2/Enpp1* double-KO MVs show  $PP_i$  levels that are equivalent to those in WT vesicles. In contrast, MVs from single KO osteoblasts show either an elevation (*Akp2* KO) or a reduction (*Enpp1* KO) in their  $PP_i$  concentration ( $n = 5$  mice, in replicates of 3) relative to TTPP mice. \*,  $P < 0.05$  compared with WT MVs.

differences in the ability of primary osteoblasts of different genotypes to mineralize. Importantly, we found that the restoration of *in vitro* mineralization correlates perfectly with the *in vivo* correction of mineralization abnormalities in the *Akp2/Enpp1* double-KO mice and with the normalized levels of inorganic pyrophosphate resulting in the *Akp2/Enpp1* double-KO MVs. Thus, a  $P_i$ -generating role for TNAP seems to be secondary to its inorganic pyrophosphatase function in bone tissue, in agreement with previous indications (29, 30).

To date, there is no effective medical treatment for hypophosphatasia (5). Attempts to treat the disease by using oral admin-





**Fig. 4.** Schematic representation of the counterregulatory functions of TNAP and PC-1 in modulating extracellular PP<sub>i</sub> concentrations. (NTPs, nucleoside triphosphates). In the bone matrix, PC-1 is the major producer of PP<sub>i</sub> (thick arrows), which in turn has an inhibitory effect on hydroxyapatite deposition. TNAP has a positive influence on mineralization primarily by controlling the size of the inhibitory pool of PP<sub>i</sub> through its inorganic pyrophosphatase activity. TNAP also generates P<sub>i</sub> by using NTPs and PP<sub>i</sub> as substrates, but other more major sources of P<sub>i</sub> (thick arrow), e.g., intestinal absorption, are likely to contribute the bulk of the P<sub>i</sub> needed for hydroxyapatite deposition.

istration of Mg<sup>2+</sup>, Zn<sup>2+</sup>, or P<sub>i</sub> supplementation have been unsuccessful, as have been the administration of parathyroid hormone to attempt to stimulate TNAP activity (33–35). Infusing hypophosphatasia patients with normal serum (36), Paget's serum (37), and purified alkaline phosphatase (38) also have had disappointing results. Our data identify PC-1 as a useful target for interference in the treatment of hypophosphatasia. By de-

leting the *Enpp1* gene, we have restored PP<sub>i</sub> concentrations to a normal level and corrected osteomalacia. We fully expect that *in vivo* administration of inhibitors of PC-1 activity, e.g., reactive blue-2 or pyridoxal phosphate-6-azophenyl-2',4'-disulphonic acid 4-sodium (39), could be equally successful in treating rickets and osteomalacia in hypophosphatasia patients.

PC-1-deficient mice represent a model of certain human diseases in which there is uncontrolled ossification of ligamentous tissues leading to loss of spine and peripheral joint mobility. One of these diseases is ossification of the posterior longitudinal ligament (OPLL; refs. 14 and 15). OPLL is a common disorder in Asia, particularly in Japan, where the incidence is 1.5% among people over 50 years of age. Polymorphisms of the PC-1 gene, of currently unknown functional significance, have been reported in Japanese subjects with OPLL (40, 41). The PC-1 KO mice are also an accepted animal model for diffuse idiopathic skeletal hyperostosis and ankylosing spinal hyperostosis (refs. 42 and 43). Treatments for enhanced ossification, except for surgery, are not well established. There is a need for drugs that target the action of osteoblasts by affecting their ability to ossify their surrounding matrix. Our results suggest the possible utility of interfering with TNAP activity as a therapeutic strategy for the treatment of pathologic cartilage calcification and hyperostosis due to local and systemic PC-1 deficiency.

This work was supported by National Institutes of Health Grants CA 42595, DE 12889, AR 47908, AG O7996, AR 40770, and AR 47347, and by grants from the Department of Veterans Affairs, Monash University, and from the National Health and Medical Research Council of Australia.

1. Anderson, H. C. (1995) *Clin. Orthop. Relat. Res.* **314**, 266–280.
2. Boskey, A. L. (1996) *Connect. Tissue Res.* **35**, 357–363.
3. Henthorn, P., Millán, J. L. & Leboy, P. (1999) in *Principles of Bone Biology*, eds. Seibel, M. J., Robins, S. P. & Bilezikian, J. P. (Academic, San Diego), pp. 127–136.
4. Henthorn, P. S. & Whyte, M. P. (1992) *Clin. Chem.* **38**, 2501–2505.
5. Whyte, M. P. (2001) in *The Metabolic and Molecular Bases of Inherited Disease*, eds. Scriver, C. R., Beaudet, A. L., Sly, W. S., Valle, D., Childs, B., Kinzler, K. W. & Vogelstein, B. (McGraw-Hill, New York), pp. 5313–5329.
6. Zurutuza, L., Muller, F., Gibrat, J. F., Taillandier, A., Simon-Bouy, B., Serre, J. L. & Mornet, A. (1999) *Hum. Mol. Genet.* **8**, 1039–1046.
7. Hsu, H. H. & Anderson, H. C. (1978) *Proc. Natl. Acad. Sci. USA* **75**, 3805–3808.
8. Robison, R. (1923) *Biochem. J.* **17**, 286–293.
9. Majeska, R. J. & Wuthier, R. E. (1975) *Biochim. Biophys. Acta* **391**, 51–60.
10. Fallon, M. D., Whyte, M. P. & Teitelbaum, S. L. (1980) *Lab. Invest.* **43**, 489–494.
11. Moss, D. W., Eaton, R. H., Smith, J. K. & Whitby, L. G. (1967) *Biochem. J.* **102**, 53–57.
12. Rezende, A., Pizauro, J., Ciancaglini, P. & Leone, F. (1994) *Biochem. J.* **301**, 517–522.
13. Meyer, J. L. (1984) *Arch. Biochem. Biophys.* **231**, 1–8.
14. Sakamoto, M., Hosoda, Y., Kojimihara, K., Yamazaki, T. & Yoshimura, Y. (1994) *Pathol. Int.* **44**, 420–427.
15. Okawa, A., Nakamura, I., Goto, S., Moriya, H., Nakamura, Y. & Ikegawa, S. (1998) *Nat. Genet.* **19**, 271–273.
16. Ho, A. M., Johnson, M. D. & Kingsley, D. M. (2000) *Science* **289**, 265–270.
17. Terkeltaub, R., Rosenbach, M., Fong, F. & Goding, J. (1994) *Arthritis Rheum.* **37**, 934–941.
18. Johnson, K., Moffa, A., Chen, Y., Pritzker, K., Goding, J. & Terkeltaub, R. (1999) *J. Bone Miner. Res.* **14**, 883–892.
19. Johnson, K., Vaingankar, S., Chen, Y., Moffa, A., Goldring, M., Sano, K., Jin-Hua, P., Sali, A., Goding, J. & Terkeltaub, R. (1999) *Arthritis Rheum.* **42**, 1986–1997.
20. Johnson, K. A., Hessle, L., Vaingankar, S., Wennberg, C., Mauro, S., Narisawa, S., Goding, J. W., Sano, K., Millán, J. L. & Terkeltaub, R. (2000) *Am. J. Physiol.* **279**, R1365–R1377.
21. Waymire, K. G., Mahuren, J. D., Jaje, J. M., Guilarte, T. R., Coburn, S. P. & MacGregor, G. R. (1995) *Nat. Genet.* **11**, 45–51.
22. Narisawa, S., Fröhlander, N. & Millán, J. L. (1997) *Dev. Dyn.* **208**, 432–446.
23. Fedde, K. N., Blair, L., Silverstein, J., Coburn, S. P., Ryan, L. M., Weinstein, R. S., Waymire, K., Narisawa, S., Millán, J. L., MacGregor, G. R. & Whyte, M. P. (1999) *J. Bone Miner. Res.* **14**, 2015–2026.

24. Sali, A., Favalaro, J. M., Terkeltaub, R. & Goding, J. W. (1999) in *Ecto-ATPases and Related Ectonucleotidases*, eds. Vanduffel, L. & Lemmens, R. (Shaker, Maastricht, The Netherlands), pp. 267–282.
25. Narisawa, S., Wennberg, C. & Millán, J. L. (2001) *J. Pathol.* **193**, 125–133.
26. Wennberg, C., Hessle, L., Lundberg, P., Mauro, S., Narisawa, S., Lerner, U. H. & Millán, J. L. (2000) *J. Bone Miner. Res.* **15**, 1879–1888.
27. Johnson, K., Hashimoto, S., Lotz, M., Pritzker, K., Goding, J. & Terkeltaub, R. (2001) *Arthritis Rheum.* **44**, 1071–1081.
28. Hsu, H. H. & Anderson, H. C. (1977) *Biochim. Biophys. Acta* **500**, 162–172.
29. Caswell, A. M., Whyte, M. P. & Russell, R. G. (1991) *Crit. Rev. Clin. Lab. Sci.* **28**, 175–232.
30. Whyte, M. P. (1994) *Endocr. Rev.* **15**, 439–461.
31. Drezner, M. K. (1996) in *Principles of Bone Biology*, eds. Bilezikian, J. P., Raisz, L. G. & Rodan, G. A. (Academic, San Diego), pp. 263–276.
32. Anderson, H. C., Hsu, H. H., Morris, D. C., Fedde, K. N. & Whyte, M. P. (1997) *Am. J. Pathol.* **151**, 1555–1561.
33. Bongiovanni, A. M., Album, M. M., Root, A. W., Hope, J. W., Marino, J. & Spencer, D. M. (1968) *Am. J. Med. Sci.* **255**, 163–170.
34. Whyte, M. P., Teitelbaum, S. L., Murphy, W. A., Bergfeld, M. A. & Avioli, L. V. (1979) *Medicine (Baltimore)* **58**, 329–347.
35. Whyte, M. P., Magill, H. L., Fallon, M. D. & Herrod, H. G. (1986) *J. Pediatr.* **108**, 82–88.
36. Whyte, M. P., Mahuren, J. D., Fedde, K. N., Cole, F. S., McCabe, E. R. & Coburn, S. P. (1988) *J. Clin. Invest.* **81**, 1234–1239.
37. Whyte, M. P., McAlister, W. H., Patton, L. S., Magill, H. L., Fallon, M. D., Lorentz, W. B. & Herrod, H. G. (1984) *J. Pediatr.* **105**, 926–933.
38. Whyte, M. P., Habib, D., Coburn, S. P., Tecklenburg, F., Ryan, L., Fedde, K. N. & Stinson, R. A. (1992) *J. Bone Miner. Res.* **7**, Suppl., S155.
39. Grobden, B., Claes, P., Roymans, D., Esmans, E. L., Van Onckelen, H. & Slegers, H. (2000) *Br. J. Pharmacol.* **130**, 139–145.
40. Nakamura, I., Ikegawa, S., Okawa, A., Okuda, S., Koshizuka, Y., Kawaguchi, H., Nakamura, K., Koyama, T., Goto, S., Toguchida, J., et al. (1999) *Hum. Genet.* **104**, 492–497.
41. Koshizuka, Y., Kawaguchi, H., Ogata, N., Ikeda, T., Mabuchi, A., Seichi, A., Nakamura, Y., Nakamura, K. & Ikegawa, S. (2002) *J. Bone Miner. Res.* **17**, 138–144.
42. Okawa, A., Goto, S. & Moriya, H. (1999) *Calcif. Tissue Int.* **64**, 239–247.
43. Rutsch, F., Vaingankar, S., Johnson, K., Goldfine, I., Maddux, B., Schauerte, P., Kalhoff, H., Sano, K., Boisvert, W. A., Superti-Furga, A. & Terkeltaub, R. (2001) *Am. J. Pathol.* **158**, 543–554.