

10-1993

Toxicity of *Fusarium proliferatum*-fermented nixtamalized corn-based diets fed to rats: Effect of nutritional status

Suzanne Hendrich

Iowa State University, shendric@iastate.edu

Kathrine A. Miller

Iowa State University


Terrance M. Wilson

Iowa State University

Patricia A. Murphy

Iowa State University, pmurphy@iastate.edu

Follow this and additional works at: http://lib.dr.iastate.edu/fshn_ag_pubs

 Part of the [Food Science Commons](#), [Human and Clinical Nutrition Commons](#), [Plant Pathology Commons](#), and the [Veterinary Pathology and Pathobiology Commons](#)

The complete bibliographic information for this item can be found at http://lib.dr.iastate.edu/fshn_ag_pubs/105. For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

This Article is brought to you for free and open access by the Food Science and Human Nutrition at Iowa State University Digital Repository. It has been accepted for inclusion in Food Science and Human Nutrition Publications by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

Toxicity of *Fusarium proliferatum*-fermented nixtamalized corn-based diets fed to rats: Effect of nutritional status

Abstract

Male F344/ N rats were fed corn, nixtamalized corn, *Fusarium proliferatum*-fermented (FP) corn to provide 50 mg/kg fumonisin B1 (FB1), or nixtamalized FP corn (NFP), with and without supplemented nutrients. Nixtamalization of FP corn almost completely hydrolyzed FB1 and produced 7-10 mg/kg hydrolyzed fumonisin (HFB1). Relative liver weight was significantly increased in nutrient-supplemented rats fed FP or NFP corn. Plasma cholesterol increased in rats fed FP or NFP corn. Plasma glutamate pyruvate transaminase was elevated in all rats fed FP corn and in nutrient-supplemented rats fed NFP corn. All rats fed FP corn and nutrient-supplemented rats fed NFP corn exhibited hepatocellular adenomas, whereas unsupplemented rats fed NFP corn had few adenomas. HFB1 and other uncharacterized *F. proliferatum* mycotoxins produced by nixtamalization were less toxic than were toxicants, including FB1, present in FP corn. Improvement in nutritional status due to nixtamalization stimulated NFP corn toxicity.

Keywords

National Veterinary Services Laboratory

Disciplines

Food Science | Human and Clinical Nutrition | Plant Pathology | Veterinary Pathology and Pathobiology

Comments

Reprinted with permission from *Journal of Agricultural and Food Chemistry*. 41(10):1649-1654. doi: [10.1021/jf00034a025](https://doi.org/10.1021/jf00034a025). Copyright 1993 American Chemical Society.

Rights

One-time permission is granted only for the use specified in your request. No additional uses are granted (such as derivative works or other editions).

Toxicity of *Fusarium proliferatum*-Fermented Nixtamalized Corn-Based Diets Fed to Rats: Effect of Nutritional Status

Suzanne Hendrich,[†] Kathrine A. Miller,[†] Terrance M. Wilson,[‡] and Patricia A. Murphy^{*†}

Food Science and Human Nutrition, Iowa State University, Ames, Iowa 50011, and National Veterinary Services Laboratory, Ames, Iowa 50010

Male F344/N rats were fed corn, nixtamalized corn, *Fusarium proliferatum*-fermented (FP) corn to provide 50 mg/kg fumonisin B₁ (FB₁), or nixtamalized FP corn (NFP), with and without supplemented nutrients. Nixtamalization of FP corn almost completely hydrolyzed FB₁ and produced 7-10 mg/kg hydrolyzed fumonisin (HFB₁). Relative liver weight was significantly increased in nutrient-supplemented rats fed FP or NFP corn. Plasma cholesterol increased in rats fed FP or NFP corn. Plasma glutamate-pyruvate transaminase was elevated in all rats fed FP corn and in nutrient-supplemented rats fed NFP corn. All rats fed FP corn and nutrient-supplemented rats fed NFP corn exhibited hepatocellular adenomas, whereas unsupplemented rats fed NFP corn had few adenomas. HFB₁ and other uncharacterized *F. proliferatum* mycotoxins produced by nixtamalization were less toxic than were toxicants, including FB₁, present in FP corn. Improvement in nutritional status due to nixtamalization stimulated NFP corn toxicity.

INTRODUCTION

Fumonisin (FB) are recently discovered carcinogenic mycotoxins (Bezuidenhout et al., 1988; Gelderblom et al., 1991) produced by the commonly occurring corn fungi *Fusarium moniliforme* and *Fusarium proliferatum*. *F. moniliforme*-inoculated corn was hepatocarcinogenic in rats (Wilson et al., 1985). Isolated fumonisins were hepatocarcinogenic in rats (Gelderblom et al., 1988, 1991) and caused equine leukoencephalomalacia (Wilson et al., 1992; Kellerman et al., 1990) and porcine pulmonary edema (Harrison et al., 1990). The occurrence of *F. moniliforme* containing FB was associated with human esophageal cancer in southern Africa (Marasas et al., 1988; Sydenham et al., 1990).

Median fumonisin concentrations usually in corn-based human foods and animal feeds range from 0.1 to 3 µg/g (Hopmans and Murphy, 1993; Murphy et al., 1993; Ross et al., 1991; Sydenham et al., 1991). Fifty milligrams per kilogram of fumonisin B₁ fed to rats from 18 to 24 months was carcinogenic (Gelderblom et al., 1991). However, a no-effect level for fumonisins has not been determined. Human exposure to fumonisins may pose a cancer risk in regions where corn is a dietary staple. Therefore, detoxification of fumonisin-contaminated corn may be necessary. Detoxification of fumonisin-contaminated corn was not accomplished by treatment with 2% ammonia at low pressure for 4 days, a process that successfully decontaminates aflatoxin-contaminated corn (Norred et al., 1991). Although the fumonisin content of ammoniated corn decreased, the hepatotoxicity of the ammoniated corn culture material remained the same as that of nonammoniated fumonisin-contaminated corn. This suggests that other forms of fumonisins or other mycotoxins in the corn can contribute to the toxicity of *F. moniliforme*-contaminated corn. Park et al. (1992) showed decreases in FB₁ levels with high pressure/ambient temperature and low pressure/high temperature ammonia as well as no increases in mutagenic potential due to these processes.




Figure 1. Structures of fumonisin B₁, hydrolyzed fumonisin B₁, and sphingosine.

However, toxicity in animals was not examined. Nixtamalization, the traditional treatment of corn with Ca(OH)₂ and heat, used to produce masa (tortilla flour), has been suggested to decontaminate fumonisin-contaminated corn because almost no fumonisin could be detected in corn products made with masa (Sydenham et al., 1991).

Fumonisin are unusual mycotoxins in that they are relatively water-soluble. The metabolism of fumonisins and their modes of action as carcinogens remain to be fully determined. Fumonisin have been shown to inhibit sphingosine synthesis (Wang et al., 1991) and may promote carcinogenesis by acting as indirect or direct antagonists to sphingosine's cytostatic action (Merrill, 1991), given the partial structural similarity between fumonisins and sphingosine (Figure 1).

* Author to whom correspondence should be addressed.

[†] Iowa State University.

[‡] National Veterinary Services Laboratory.

The dependence of fumonisin toxicity on nutritional status has not been determined, but early studies of fumonisin toxicity and carcinogenicity have examined the effects of feeding *F. moniliforme*-fermented corn (Wilson et al., 1985). Corn has relatively poor nutritional quality. It is deficient in protein and especially lacking in calcium, riboflavin, and lysine (U.S. Department of Agriculture, 1992). The action of many carcinogens has been shown to depend on nutritional status. For example, increasing total caloric intake (Pariza and Boutwell, 1987), fat intake (Carroll and Khor, 1975), or protein intake (Youngman and Campbell, 1991) have been shown to promote experimental carcinogenesis of several types. Protein (Youngman and Campbell, 1991), calorie (Pariza and Boutwell, 1987), or linoleate (Ip et al., 1985) restriction suppresses carcinogenesis. Because nixtamalization improves the nutritive value of corn (Carter and Carpenter, 1982), the hydrolysis of the toxicant may interact with nutritional status to permit or enhance carcinogenic effects of *F. moniliforme* or *F. proliferatum* corn.

This study was performed to determine the effects of nixtamalization and/or nutrient supplementation of *F. proliferatum*-fermented (FP) corn, containing 50 mg/kg fumonisin B₁, on diethylnitrosamine-initiated hepatocarcinogenesis in male F344/N rats. This strain of *Fusarium* was chosen because it is a predominant producer of fumonisin B₁.

MATERIALS AND METHODS

Chemicals. All chemicals were obtained from Sigma Chemical Co., St. Louis, MO, unless otherwise specified.

Animals and Diets. Ten-day-old male F344/N rats were injected intraperitoneally with diethylnitrosamine (15 mg/kg of body weight) in corn oil to initiate carcinogenesis. Eight groups of six rats were fed and provided free access to water for 4 weeks in a room with a 12-h light/dark cycle maintained at 22 °C and 50% humidity. The rats were randomly assigned to one of the following diets: ground corn containing 70 g of dextrose/kg of corn; nixtamalized ground corn containing 70 g of dextrose/kg of corn; corn or nixtamalized corn supplemented with nutrients (per kilogram of corn: casein, 60 g; thiamin, 0.002 g; riboflavin, 0.005 g; pyridoxine, 0.002 g; niacin, 0.010 g; folacin, 0.002 g; CaCO₃, 1.6 g; NaCl, 1.0 g; methionine, 3.0 g; choline, 2.0 g) to approximate an AIN-76 diet (American Institute of Nutrition, 1980) based on the estimated nutrient content of whole-ground unbolted dry corn meal (U.S. Department of Agriculture, 1992). Four of the eight diets were formulated with corn fermented with *F. proliferatum* strain M5991, a strain producing predominantly FB₁ (donated by Dr. Paul Nelson, Pennsylvania State University, University Park, PA), to produce a concentration of approximately 50 mg of fumonisin B₁ (FB₁)/kg of diet.

Corn for fermentation was sterilized by autoclaving 1-kg batches for 40 min at 132 °C. The corn (Variety 636, Dekalb Co.) was determined to be aflatoxin-free by P. F. Ross (Osheim and Ross, 1982), National Veterinary Services Laboratory, Ames, IA. Each batch of sterile corn was inoculated with *F. proliferatum* culture material reconstituted in sterile phosphate buffered saline according to the procedure of Ross et al. (1990). The FP corn was cultured for 3 weeks at 22 °C in the dark, when the corn was frozen and analyzed for FB content (Ross et al., 1991). This fermented corn was dried and mixed with fumonisin-free corn to give a final concentration of 50 mg/kg FB₁ in the diet.

Clean ground corn or FP corn containing 50 mg/kg FB₁ was nixtamalized by heating 300 g of corn with 900 mL water containing 1.2% Ca(OH)₂ at 80–100 °C for 1 h. The mixture was steeped overnight. The nixtamalized [Ca(OH)₂-treated] corn (NFP) was washed with approximately 3 volumes of water and air-dried before nutrient supplementation. Each diet was analyzed in duplicate for FB₁ and hydrolyzed FB₁ (HFB₁) content by analytical HPLC (Wilson et al., 1990).

Animals were weighed and food intakes measured weekly. After euthanasia by CO₂ anesthesia and decapitation, 5 mL of

Table I. Fumonisin B₁ and Hydrolyzed Fumonisin B₁ Contents of Experimental Diets

dietary treatment			FB ₁ , μg/g	HFB ₁ , μg/g
nixtamalized	nutrient-supplemented	<i>F. proliferatum</i> -fermented		
-	-	-	0.3 ± 0.0	ND ^a
-	-	+	48.0 ± 1.3	ND
-	+	-	0.2 ± 0.0	ND
-	+	+	45.2 ± 4.4	ND
+	-	-	0.1 ± 0.1	ND
+	-	+	0.0 ± 0.0	10.7 ± 0.4
+	+	-	0.2 ± 0.0	ND
+	+	+	0.8 ± 0.4	7.6 ± 0.2

^a Not detected at a limit of detection of 25 ng/g. Duplicate samples of each diet were extracted, and each extract was analyzed twice by HPLC with fluorescence detection of the OPA derivative.

rat blood was collected in tubes containing 0.25 mL of Na₂EDTA (40 mg/mL). Plasma was collected after centrifugation of blood for 15 min at 2000 rpm in an IEC low-speed floor-model centrifuge (International Equipment Co., Boston, MA) at room temperature. Liver weight was measured, and a portion of the largest lobe was fixed in formalin, embedded in paraffin, sectioned by microtome to a thickness of 10 μm, stained by standard methods with hematoxylin and eosin (Sheehan and Hrapchak, 1980), and examined microscopically.

Plasma was analyzed for glutamate-pyruvate transaminase (GPT) (Reitman and Frankel, 1957) and total cholesterol (Allain et al., 1981) by colorimetric diagnostic kits which included standards (Sigma). The analyses were performed with a variable-wavelength Gilford 240 spectrophotometer (Gilford Instrument Laboratories, Inc., Oberlin, OH). Cholesterol concentration and GPT activity were determined by extrapolation from standard curves.

All data were analyzed by one-way ANOVA ($p < 0.05$), and least significant differences between treatment means were determined by Duncan's multiple-range tests ($p < 0.05$), using the Statistical Analysis System (Cary, NC) on the Iowa State University mainframe computer.

RESULTS

The *F. proliferatum*-fermented (FP) corn diets contained 45–48 mg/kg FB₁ and no HFB₁ (Table I). Very small amounts of FB₂ and FB₃ were found in the FP corn. Only 0.1–0.3 mg/kg FB₁ and no HFB₁ were detected in uninoculated corn diets. The nixtamalized corn diets fermented with *F. proliferatum* (NFP) contained 0–0.8 mg/kg FB₁ and 8–11 mg/kg HFB₁.

Body weights were greatest in rats fed nixtamalized nutrient-supplemented corn. Nutrient-supplemented rats had significantly greater body weights than did rats fed corn alone (Table II). Body weight was significantly suppressed in rats fed FP corn; however, body weight was most suppressed in rats fed unsupplemented unnixtamalized FP corn. Food intakes did not differ significantly among groups (Table II). Relative liver weight (liver weight as a percentage of body weight) was significantly increased by FP corn only in nutrient-supplemented rats, and especially in rats fed nixtamalized nutrient-supplemented corn (Table II). Rats fed unsupplemented FP corn did not show increased liver weight.

All rats fed FP corn had significantly greater plasma GPT activities than the control groups, with the exception of rats fed unsupplemented nixtamalized FP corn (Table II). In groups fed FP corn, plasma GPT was increased to a greater extent in rats fed nutrient-supplemented corn than in rats fed unsupplemented corn. Compared with groups fed control corn, plasma cholesterol was significantly greater in all groups fed FP corn. The greatest

Table II. Effects of Feeding *F. proliferatum*-Fermented, Nixtamalized, and/or Nutrient-Supplemented Corn on Body Weight, Food Intake, Relative Liver Weight, Plasma Glutamate-Pyruvate Transaminase, and Plasma Cholesterol

dietary treatment			body wt, ^d g	food intake, ^e g	rel liver wt, %	plasma GPT/ SF/mL	plasma cholesterol, mmol/L
FP ^a	nutrient supplement ^b	nixtamal ^c					
-	-	-	74 ± 5 ^c	692 ± 82	3.9 ± 0.5 ^{cd}	55 ± 9 ^c	1.72 ± 0.17 ^c
+	-	-	49 ± 6 ^a	852 ± 182	3.7 ± 0.5 ^{cd}	117 ± 31 ^b	4.57 ± 0.61 ^a
-	+	-	127 ± 10 ^b	811 ± 48	4.0 ± 0.2 ^c	57 ± 7 ^c	2.14 ± 0.20 ^c
+	+	-	75 ± 7 ^c	681 ± 209	5.2 ± 0.7 ^b	139 ± 26 ^a	3.58 ± 0.79 ^b
-	-	+	75 ± 5 ^c	701 ± 59	3.6 ± 0.2 ^{cd}	47 ± 8 ^c	1.74 ± 0.34 ^c
+	-	+	61 ± 4 ^d	726 ± 280	3.3 ± 0.3 ^d	67 ± 5 ^c	4.00 ± 0.57 ^{ab}
-	+	+	162 ± 18 ^a	911 ± 87	4.0 ± 0.3 ^c	51 ± 13 ^c	1.68 ± 0.41 ^c
+	+	+	119 ± 13 ^b	784 ± 215	5.8 ± 0.7 ^a	127 ± 18 ^{ab}	4.61 ± 0.65 ^a

^a *F. proliferatum*-fermented corn. ^b Diets supplemented with adequate protein, vitamins, and minerals to match AIN-76 recommendations. ^c Nixtamalized corn. ^d Body weight gain over 30 days. ^e Food intake over 30 days. / Plasma glutamate-pyruvate transaminase in Sigma-Frankel units. Data are expressed as mean ± standard deviations. Values in a column bearing different superscripts are significantly different, *p* < 0.05.

Table III. Incidence of Hepatic Histopathology^a of Rats Fed Control or *F. proliferatum*-Fermented Diets

dietary treatment			n	% incidence	
nixtamalized	nutrient-supplemented	<i>F. proliferatum</i> -fermented		hepatic adenomas	hepatic cholangiomas
±	±	-	24	0 ^a	0
-	-	+	6	83 ^b	33
-	+	+	6	100 ^b	50
+	-	+	6	14 ^a	33
+	+	+	6	67 ^b	17

^a The microscopic lesions were classified according to the descriptors of Eustis et al. (1990). Each group bearing a letter different from that of the control group is significantly different from the combined controls, as analyzed by chi-square tests using a two-way contingency plan, *p* < 0.05.

increases in plasma cholesterol occurred in rats fed unsupplemented FP corn and in rats fed nutrient-supplemented NFP corn (Table II).

Rats fed nixtamalized or unnixtamalized FP corn developed hepatocellular adenomas, clear cell and hyperplastic foci, and cholangiomas, whereas rats fed clean corn did not (Table III; Figure 2). The photomicrographs are representative of the hepatic histopathological findings from each of the groups fed FP corn. The rats fed FP corn had diffuse mild to severe hyperplastic nodules (Figure 2A). The supplemented FP corn-fed group showed diffuse severe hyperplastic nodules (Figure 2B). The group fed NFP corn showed multiple clear cell foci (Figure 2C), whereas rats fed supplemented NFP corn had severe diffuse hyperplastic nodules (Figure 2D). The rats fed corn without nutrient supplementation showed fatty infiltration of the liver. Rats fed nixtamalized FP corn without nutrient supplementation developed less severe hepatic lesions than did the other groups fed FP diets. The rats fed unsupplemented FP corn also showed less development of neoplasia than the nutrient-supplemented rats fed FP corn.

DISCUSSION

This study demonstrated the hydrolysis of FB₁ during nixtamalization (Figure 1). The hydrolysis of FB₁ by treatment with Ca(OH)₂ and heat did not detoxify FP corn, but rather produced a more toxic product or products; this finding is based on the fact that feeding NFP corn containing 8–11 mg/kg HFB₁ produced some toxicity symptoms as severe as feeding FP corn containing 45–48 mg/kg FB₁, in terms of effects on body weight, relative liver weight, plasma GPT and plasma cholesterol (Table II), and incidence of hepatic adenomas (Table III). The major toxic product of fumonisin hydrolysis during nixtamalization appeared to be HFB₁, but it is possible that other breakdown products, such as calcium-FB complexes and other products more difficult to identify than HFB₁, also played a role in the toxicity of nixtamalized

FP corn. Further studies comparing purified HFB₁ and FB₁ toxicity are needed to prove HFB₁ toxicity.

How likely is it that toxicants in FP-fermented corn, other than FB₁ and its breakdown products, might be responsible for the toxicity and neoplastic responses observed in rats fed FP-fermented corn? Purified FB₁ has been shown to cause male rat hepatocellular carcinomas, when fed at 50 mg/kg in the diet for 18–24 months (Gelderblom et al., 1991), and purified FB₁ has been shown to reproduce the neoplastic and weight-suppressive effects of feeding crude *F. moniliforme* culture to rats. Purified FB₁ has also been shown to cause equine leukoencephalomalacia (Wilson et al., 1992; Kellerman et al., 1990), a disease first associated with feeding *F. moniliforme*-contaminated corn (Wilson and Maronpot, 1971). Although a no-effect level has yet to be established for FB₁, human foods commonly contain 0.2–3.0 mg/kg of FB₁ (Hopmans and Murphy, 1993; Sydenham et al., 1991). The small amounts of FB₁ remaining in NFP corn seem unlikely to be responsible for the severe toxicity signs observed. Fumonisin production by *Fusarium* species was associated with less production of other *Fusarium* mycotoxins (Sydenham et al., 1990). Because crude contamination of human foods by *F. moniliforme* and *F. proliferatum* would be the usual means of exposure to FB₁, identification of potential synergistic agents in *F. moniliforme* may be important. It seems likely that the predominant toxic factor in FP corn was FB₁ (Gelderblom et al., 1988).

The knowledge regarding fumonisin's mechanism of action as a toxicant and carcinogen is limited. Fumonisin are sphingosine synthesis inhibitors (Wang et al., 1991). Sphingolipids have been postulated to be anticarcinogens because they are negative regulators of protein kinase C (Merrill, 1991), a metabolic control point which is activated by numerous carcinogens (Weinstein, 1987). The present results suggested that FB₁ and HFB₁ act in similar ways, because their toxicity signs were similar. Whether these toxicity signs were associated with altered sphingolipid




Figure 2. Photomicrographs of formalin-fixed hematoxylin- and eosin-stained hepatic sections from rats fed *F. proliferatum*-fermented (FP) corn: (A) FP corn only; (B) supplemented FP corn; (C) nixtamalized FP corn (NFP); (D) supplemented nixtamalized FP corn.

metabolism remains to be seen. It seemed reasonable that HFB₁ would also be a sphingosine synthesis inhibitor, bearing more structural similarity to sphingosine than fumonisin itself (Figure 1).

The present study suggested that some fumonisin toxicity signs may be more directly related to carcinogenesis than are other toxicity signs. Plasma GPT levels were significantly elevated in the same groups which show the greatest development of neoplasia (Tables II and III): all rats fed FP corn and rats fed supplemented NFP corn. Hepatomegaly is a common effect of tumor-promoting agents. It has been associated with the most severe neoplasia, which occurred in nutrient-supplemented rats fed FP corn or NFP corn (Table II), but rats fed unsupplemented FP corn also developed neoplasia, without hepatomegaly.

Dramatically increased plasma or serum cholesterol was a sign of *F. moniliforme* toxicity seen in primates (Fincham et al., 1992). In vervets, an African primate species, low-fat diets containing *F. moniliforme* culture material that provided increasing doses of FBs caused a dose/response in plasma cholesterol. This strongly suggested that FBs increased plasma cholesterol in primates. Therefore,

fumonisin was probably responsible for increasing plasma cholesterol in rats in the present study. Other tumor-promoting agents, such as phenobarbital, increased serum cholesterol (Katayama et al., 1991). Perhaps the negative effect of fumonisin on sphingolipids also had the effect of up-regulating cholesterol production, as one of many effects of disrupting basic cell signaling. Increased plasma cholesterol was related to *F. proliferatum*-promoted carcinogenesis, because increased plasma cholesterol occurred in all FP- and NFP-fed rats (Table II). But plasma cholesterol was lower in nutrient-supplemented rats fed FP corn than in rats fed unsupplemented FP corn (Table II). This suggested that when carcinogenesis was more severe (Table III), plasma cholesterol declined. Thus, greater plasma cholesterol may be an indicator of the earlier stages of carcinogenesis. In the rats fed nixtamalized FP corn, unsupplemented and supplemented rats showed similar plasma cholesterol. This may mean either that HFB₁ acted in part by a different mechanism than does FB₁ or that NFP corn containing 11 mg/kg HFB₁ was not quite as potent a carcinogen as FP corn containing 45 mg/kg FB₁, i.e., that hepatocarcinogenesis in the NFP corn-fed rats was not quite as advanced as in the FP corn-fed

rats. The lack of development of neoplastic nodules in unsupplemented rats fed HFB₁ may support either of these two viewpoints (Table III).

This study indicated that the production of hydrolyzed fumonisins in the processing of foods for human consumption may be of significant concern. The further possibility that hydrolyzed fumonisins might be produced *in vivo* seems to be worthy of investigation because of the potential toxicity of such products.

The role of nutritional status in carcinogenesis was also of fundamental concern. The action of fumonisin and its metabolites may depend to some extent on nutritional status. Carcinogenesis by many agents has been shown to be suppressed by nutritional deprivation of varying sorts, including deprivation of calories, protein, and essential fatty acids (Pariza and Boutwell, 1987; Youngman and Campbell, 1991; Ip et al., 1985). For example, diets containing less than 9% protein totally suppressed the promotion of aflatoxin B₁-induced rat hepatocarcinogenesis (Youngman and Campbell, 1991). The corn diet fed in the present study was low in protein and in some other nutrients such as thiamin and niacin for which relative deprivation has not been studied with respect to hepatocarcinogenesis. Rats fed corn alone showed fatty infiltration of the liver (Figure 2), which was a clear sign of malnutrition, and especially of protein malnutrition, due to a lack of the protein needed to synthesize lipoproteins to export fat from the liver (Latham, 1990). Perhaps FB₁ carcinogenesis does not depend on protein status as much as aflatoxin carcinogenesis does. In our study, increased body weight (Table II), due to nutrient supplementation and nixtamalization, was associated with increased plasma GPT and severity of neoplasia (Tables II and III; Figure 2). This was seen in comparing liver from an unsupplemented FP corn-fed rat (Figure 2A) with liver from a supplemented FP corn-fed rat (Figure 2B). Although rats fed NFP corn had about 20% lower body weights than did rats fed supplemented FP corn, only one rat fed NFP corn developed hepatic adenomas, whereas all supplemented FP corn-fed rats had hepatic adenomas (Table III). This suggested that NFP corn containing 11 mg/kg HFB₁ was not as potent a promoter as was FP corn containing 45 mg/kg FB₁. Nixtamalization combined with supplementation of FP corn significantly increased body weights (Table II) and adenomas (Table III). Body weights in this group fed 8 mg/kg HFB₁ were greater than in supplemented FP corn-fed rats (Table II), but the extents of development of neoplasia were similar in both groups (Table III). This suggested that although NFP corn containing 8 mg/kg HFB₁ was not as potent a promoter as was FP corn containing 45 mg/kg FB₁, the better nutritive value of nixtamalized corn (Carter and Carpenter, 1982) made NFP corn almost as hepatocarcinogenic as FP corn. Thus, nixtamalization is probably not a useful strategy for detoxification of *F. moniliforme*- or *F. proliferatum*-contaminated corn in populations of good nutritional status. If fumonisins are nutrient-independent carcinogens, they might be especially significant to carcinogenesis in undernourished populations as well.

ACKNOWLEDGMENT

Journal Paper J-15110, Iowa Agriculture and Home Economics Experiment Station, Ames, IA, Projects 2955, 2844, and 2406, the latter a contributing project to North Central Regional Project NC-129. Additional funds were provided by the Iowa Corn Promotion Board and the ISU/USDA Food Safety Consortium.

LITERATURE CITED

- Allain, C. C.; Poon, L. S.; Chan, C. S. G.; Richmond, W.; Fu, P. C. Enzymatic determination of total serum cholesterol. *Clin. Chem.* 1981, 27, 838-841.
- American Institute of Nutrition. Second report of the ad hoc committee on standards for nutritional studies. *J. Nutr.* 1980, 110, 1726.
- Bezuidenhout, S. C.; Gelderblom, W. C. A.; Gorst-Allman, C. P.; Horak, R. M.; Marasas, W. F. O.; Spiteller, G.; Vleggaar, R. Structure elucidation of the fumonisins, mycotoxins from *Fusarium moniliforme*. *J. Chem. Soc., Chem. Commun.* 1988, 743-745.
- Carroll, K. K.; Khor, H. T. Dietary fat in relation to tumorigenesis. *Prog. Biochem. Pharmacol.* 1975, 10, 308-353.
- Carter, E. G. A.; Carpenter, K. J. The available niacin values of foods for rats and their relation to analytical values. *J. Nutr.* 1982, 112, 2091-2103.
- Eustis, S. L.; Boorman, G. A.; Harada, T.; Popp, J. A. Liver. In *Pathology of the Fischer Rat*; Boorman, G. A., Eustis, S. L., Elwell, M. R., Montgomery, C. A., Jr., MacKenzie, W. F., Eds.; Academic Press: San Diego, CA, 1990; pp 71-94.
- Fincham, J. E.; Marasas, W. F. O.; Taljaard, J. J. F.; Kriek, N. P. J.; Badenhorst, C. J.; Gelderblom, W. C. A.; Seier, J. V.; Smuts, C. M.; Faber, M.; Weight, M. J.; Slazus, W.; Woodroof, C. W.; van Wyk, M. J.; Kruger, M.; Thiel, P. G. Atherogenic effects in a non-human primate of *Fusarium moniliforme* cultures added to a carbohydrate diet. *Atherosclerosis* 1992, 94, 13-25.
- Gelderblom, W. C. A.; Jaskiewicz, K.; Marasas, W. F. O.; Thiel, P. G.; Horak, R. M.; Vleggaar, R.; Kriek, N. P. J. Fumonisins—novel mycotoxins with cancer-promoting activity produced by *Fusarium moniliforme*. *Appl. Environ. Microbiol.* 1988, 54, 1806-1811.
- Gelderblom, W. C. A.; Kriek, N. P. J.; Marasas, W. F. O.; Thiel, P. G. Toxicity and carcinogenicity of the *Fusarium moniliforme* metabolite, fumonisin B₁, in rats. *Carcinogenesis* 1991, 12, 1247-1251.
- Harrison, L. R.; Colvin, B. M.; Greene, J. T.; Newman, L. E.; Cole, J. R. Pulmonary edema and hydrothorax in swine produced by fumonisin B₁, a toxic metabolite of *Fusarium moniliforme*. *J. Lab. Diagn. Invest.* 1990, 2, 217-221.
- Hopmans, E. C.; Murphy, P. A. Detection of fumonisins B₁, B₂, B₃ and hydrolyzed fumonisin B₁ in corn-containing foods. *J. Agric. Food Chem.* 1993, following paper in this issue.
- Ip, C.; Carter, C. A.; Ip, M. M. Requirement of essential fatty acid for mammary tumorigenesis in the rat. *Cancer Res.* 1985, 45, 1997-2001.
- Katayama, T.; Momoto, Y.; Yasutomo, W.; Kato, N. Elevated concentrations of α -tocopherol, ascorbic acid, and serum lipids in rats fed polychlorinated biphenyls, chlorobutanol, or phenobarbital. *J. Nutr. Biochem.* 1991, 2, 92-96.
- Kellerman, T. S.; Marasas, W. F. O.; Thiel, P. G.; Gelderblom, W. C. A.; Cawood, M.; Coetzer, J. A. W. Leukoencephalomalacia in two horses induced by oral dosing of fumonisin B₁. *Onderstepoort J. Vet. Res.* 1990, 57, 269-275.
- Latham, M. C. Protein-energy malnutrition. In *Present Knowledge in Nutrition*, 6th ed.; Brown, M. L., Ed.; International Life Sciences Institute: Washington, DC, 1990; pp 39-46.
- Marasas, W. F. O.; Jaskiewicz, K.; Venter, F. S.; Van Schalkwyk, D. J. *Fusarium moniliforme* contamination of maize in esophageal cancer areas in Transkei. *S. Afr. Med. J.* 1988, 74, 110-114.
- Merrill, A. H., Jr. Cell regulation by sphingosine and more complex sphingolipids. *J. Bioenerg. Biomembr.* 1991, 23, 83-104.
- Murphy, P. A.; Rice, L. D.; Ross, P. F. Fumonisin B₁, B₂, and B₃ content of Iowa, Wisconsin and Illinois corn and corn screenings. *J. Agric. Food Chem.* 1993, 41, 263-266.
- Norred, W. P.; Voss, K. A.; Bacon, C. W.; Riley, R. T. Effectiveness of ammonia treatment in detoxification of fumonisin-contaminated corn. *Food Chem. Toxicol.* 1991, 29, 815-819.
- Osheim, D. L.; Ross, P. F. Cleanup of some mycotoxins by gel permeation chromatography. In *Proceedings of the Midwest AOAC Meeting*; AOAC International: Arlington, VA, 1982; Abstract 23.

- Pariza, M. W.; Boutwell, R. K. Historical perspective: calories and energy expenditure in carcinogenesis. *Am. J. Clin. Nutr.* 1987, 45 (suppl.) 151-156.
- Park, D. L.; Rua, S. M.; Mirocha, C. J.; Abd-Alla, E.-S.; Weng, C. Y. Mutagenic potentials of fumonisin contaminated corn following ammonia decontamination procedure. *Mycopathologia* 1992, 117, 105-108.
- Reitman, S.; Frankel, S. A colorimetric method for the determination of serum glutamic oxaloacetic and glutamic pyruvic transaminases. *Am. J. Clin. Pathol.* 1957, 28, 56-60.
- Ross, P. F.; Nelson, P. E.; Richard, J. L.; Osweiler, G. D.; Rice, L. D.; Plattner, R. D.; Wilson, T. M. Production of fumonisins by *Fusarium moniliforme* and *Fusarium proliferatum* isolates associated with equine leukoencephalomalacia and a pulmonary edema syndrome in swine. *Appl. Environ. Microbiol.* 1990, 56, 3225-3226.
- Ross, P. F.; Rice, L. G.; Plattner, R. D.; Osweiler, G. D.; Wilson, T. M.; Owens, D. L.; Nelson, H. A.; Richard, J. L. Concentrations of fumonisin B₁ in feeds associated with animal health problems. *Mycopathologia* 1991, 114, 129-135.
- Sheehan, D. C.; Hrapchak, B. B. *Theory and Practice of Histotechnology*, 2nd ed.; Mosby: St. Louis, MO, 1980; pp 143-144.
- Sydenham, E. W.; Thiel, P. G.; Marasas, W. F. O.; Shephard, G. S.; Van Schalkwyk, D. J. K.; Koch, K. R. Natural occurrence of some *Fusarium moniliforme* mycotoxins in corn from low and high esophageal cancer prevalence area of the Transkei, Southern Africa. *J. Agric. Food Chem.* 1990, 38, 1900-1903.
- Sydenham, E. W.; Gelderblom, W. C. A.; Thiel, P. G.; Marasas, W. F. O.; Stockenstrom, S. Fumonisin contamination of commercial corn-based human foodstuffs. *J. Agric. Food Chem.* 1991, 39, 2014-2018.
- U.S. Department of Agriculture. *Composition of Foods, raw, processed, prepared*; Agriculture Handbook 8; Washington, DC, 1992.
- Wang, E.; Norred, W. P.; Bacon, C. W.; Riley, R. T.; Merrill, A. H., Jr. Inhibition of sphingosine biosynthesis by fumonisins. Implications for diseases associated with *Fusarium moniliforme*. *J. Biol. Chem.* 1991, 266, 14486-14490.
- Weinstein, I. B. Growth factors, oncogenes, and multistage carcinogenesis. *J. Cell. Biochem.* 1987, 33, 213-224.
- Wilson, B. J.; Maronpot, R. R. Causative fungus agent of leukoencephalomalacia in equine animals. *Vet. Rec.* 1971, 88, 484-486.
- Wilson, T. M.; Nelson, P. E.; Knepp, C. R. Hepatic neoplastic nodules, adenofibrosis, and cholangiocarcinomas in male Fisher 344 rats fed corn naturally contaminated with *Fusarium moniliforme*. *Carcinogenesis* 1985, 6, 1155-1160.
- Wilson, T. M.; Ross, P. F.; Rice, L. D.; Osweiler, G. D.; Nelson, H. A.; Owens, D. L.; Plattner, R. D.; Reggiardo, C.; Noon, T. H.; Pickrell, J. W. Fumonisin B₁ levels associated with an epizootic of equine leukoencephalomalacia. *J. Vet. Diagn. Invest.* 1990, 2, 213-216.
- Wilson, T. M.; Ross, P. F.; Owens, D. L.; Rice, L. G.; Green, S. A.; Jenkins, S. J.; Nelson, H. A. Experimental reproduction of ELEM: a study to determine minimum toxic dose in ponies. *Mycopathologia* 1992, 117, 115-120.
- Youngman, L. D.; Campbell, T. C. High protein intake promotes the growth of hepatic preneoplastic foci in Fischer 344 rats: evidence that early remodeled foci retain the potential for future growth. *J. Nutr.* 1991, 121, 1454-1461.

Received for review March 30, 1993. Revised manuscript received July 2, 1993. Accepted July 19, 1993.*

* Abstract published in *Advance ACS Abstracts*, September 15, 1993.