








## ORIGINAL RESEARCH

# Vegetation state changes in the course of shrub encroachment in an African savanna since about 1850 CE and their potential drivers

Ximena Tabares<sup>1,2</sup>  | Heike Zimmermann<sup>1</sup>  | Elisabeth Dietze<sup>1</sup>  |  
Gregor Ratzmann<sup>3</sup> | Lukas Belz<sup>4</sup> | Andrea Vieth-Hillebrand<sup>5</sup>  | Lydie Dupont<sup>6</sup>  |  
Heinz Wilkes<sup>4</sup> | Benjamin Mapani<sup>7</sup>  | Ulrike Herzschuh<sup>1,2,8</sup> 

<sup>1</sup>Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Potsdam, Germany

<sup>2</sup>Institute of Biochemistry and Biology, Potsdam University, Potsdam, Germany

<sup>3</sup>Institute of Biology, Freie Universität Berlin, Berlin, Germany

<sup>4</sup>Institute for Chemistry and Biology of the Marine Environment, Carl von Ossietzky University of Oldenburg, Oldenburg, Germany

<sup>5</sup>GFZ German Research Centre for Geosciences, Helmholtz Centre Potsdam, Potsdam, Germany

<sup>6</sup>MARUM – Centre for Marine Environmental Sciences, University of Bremen, Bremen, Germany

<sup>7</sup>Department of Geology, University of Namibia, Windhoek, Namibia

<sup>8</sup>Institute of Environmental Science and Geography, Potsdam University, Potsdam, Germany

## Correspondence

Ximena Tabares and Ulrike Herzschuh, Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Telegrafenberg A45, 14473 Potsdam, Germany.

Emails: xtabares@uni-potsdam.de; uherzschuh@awi.de

## Funding information

Bundesministerium für Bildung und Forschung, Grant/Award Number: FKZ:01LL1302A; Ministry of Education, Grant/Award Number: 01LL1302A; Helmholtz Association

## Abstract

1. Shrub encroachment has far-reaching ecological and economic consequences in many ecosystems worldwide. Yet, compositional changes associated with shrub encroachment are often overlooked despite having important effects on ecosystem functioning.
2. We document the compositional change and potential drivers for a northern Namibian *Combretum* woodland transitioning into a *Terminalia* shrubland. We use a multiproxy record (pollen, sedimentary ancient DNA, biomarkers, compound-specific carbon ( $\delta^{13}\text{C}$ ) and deuterium ( $\delta\text{D}$ ) isotopes, bulk carbon isotopes ( $\delta^{13}\text{C}_{\text{org}}$ ), grain size, geochemical properties) from Lake Otjikoto at high taxonomical and temporal resolution.
3. We provide evidence that state changes in semiarid environments may occur on a scale of one century and that transitions between stable states can span around 80 years and are characterized by a unique vegetation composition. We demonstrate that the current grass/woody ratio is exceptional for the last 170 years, as supported by *n*-alkane distributions and the  $\delta^{13}\text{C}$  and  $\delta^{13}\text{C}_{\text{org}}$  records. Comparing vegetation records to environmental proxy data and census data, we infer a complex network of global and local drivers of vegetation change. While our  $\delta\text{D}$  record suggests physiological adaptations of woody species to higher atmospheric  $p\text{CO}_2$  concentration and drought, our vegetation records reflect the impact of broad-scale logging for the mining industry, and the macrocharcoal record suggests a decrease in fire activity associated with the intensification of farming. Impact of selective grazing is reflected by changes in abundance and taxonomical composition of grasses and by an increase of nonpalatable and trampling-resistant taxa. In addition, grain-size and spore records suggest changes in the erodibility of soils because of reduced grass cover.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2019 The Authors. *Ecology and Evolution* published by John Wiley & Sons Ltd.

4. *Synthesis.* We conclude that transitions to an encroached savanna state are supported by gradual environmental changes induced by management strategies, which affected the resilience of savanna ecosystems. In addition, feedback mechanisms that reflect the interplay between management legacies and climate change maintain the encroached state.

#### KEYWORDS

climate change, fossil pollen, land-use change, savanna ecology, sedimentary ancient DNA, state and transition, tree–grass interactions

## 1 | INTRODUCTION

Shrub encroachment is affecting savannas worldwide (Saha, Scanlon, & D'Odorico, 2015; Stevens, Lehmann, Murphy, & Durigan, 2017; Tian, Brandt, Liu, Rasmussen, & Fensholt, 2017). This process is assumed to indicate a savanna state change from open woodlands and grasslands, characterized by dominance of  $C_4$  grasses, to a bush-thickened savanna (Joubert, Rothauge, & Smit, 2008; Meyer, Wiegand, Ward, & Moustakas, 2007). Southern Africa is a hotspot of studies reporting shrub encroachment (O'Connor, Puttick, & Hoffman, 2014; Roques, O'Connor, & Watkinson, 2001; Wigley, Bond, & Hoffman, 2009). This is not only because shrub encroachment is a widespread phenomenon, but also because the area is traditionally used as grazing grounds and encroachment reduces the land-carrying capacity, threatening local economies (De Klerk, 2004; Moyo, O'Keefe, & Sill, 1993).

Shrub encroachment has been described as an alternative stable state occurring several times during the last two millennia in African savannas (Gil-Romera, Lamb, Turton, Sevilla-Callejo, & Umer, 2010). There is also some indication that African savannas are characterized by grassland/woodland phases occurring with a periodicity of 250–600 years (Gillson, 2004; Gil-Romera et al., 2010). Such woodland phases were not always led by the spread of encroaching species (Gil-Romera et al., 2010; Scott, Cooremans, de Wet, & Vogel, 1991), which suggests that a mere increase in woody cover, as reported by several studies since the 20th century (Hoffman, Rohde, & Gillson, 2019; van Rooyen, le Roux, van der Merwe, van Rooyen, & Geldenhuys, 2018; Wiegand, Ward, & Saltz, 2005), does not necessarily mean shrub encroachment. Accordingly, identification of encroached states should be related to the increasing cover of encroacher species and the suppression of perennial grasses (Gil-Romera et al., 2010). However, the low taxonomic resolution of pollen records, particularly regarding Poaceae, constrains the identification of these phases on a long timescale.

It is assumed that resilience of savanna ecosystems is scale dependent. For example, savanna seems to be resilient to changes at a centennial timescale, as the vegetation fluctuates between two stable states of grassland and woodland (Gillson, 2004; Gil-Romera et al., 2010). Conversely, savanna seems to be less resilient at a decadal

scale, as encroachment may occur within a century and without regression to a grassy state even at a patch scale (Rohde & Hoffman, 2012). Similarly, African landscapes are predicted to shift to woodland states, with abrupt transitions at the local scale, but smoother at the continental scale (Higgins & Scheiter, 2012). Transitions have been described as the unstable equilibrium between the “basins of attraction” of stable states (Holling, 1973; Scheffer, Carpenter, Foley, Folke, & Walker, 2001), but surprisingly little is known about the duration of such transitions (Higgins & Scheiter, 2012; Joubert et al., 2008) or the compositional changes they entail (Joubert et al., 2008; Liao, Clark, & DeGloria, 2018).

Different opinions exist as to the causes of shrub encroachment in savannas (Devine, McDonald, Quaipe, & Maclean, 2017; Venter, Cramer, & Hawkins, 2018; D Ward, 2005). Among others, changes in precipitation, atmospheric  $pCO_2$ , vegetation fires, and land use are considered to be triggers and/or drivers (Aleman, Blarquez, & Staver, 2016; Berry & Kulmatiski, 2017; Buitenwerf, Bond, Stevens, & Trollope, 2012; Case & Staver, 2017), which likely interplay at different timescales. For example, while climate shifts are considered the main driver of changes in the grass/woody vegetation ratio on a long timescale (Gil-Romera et al., 2010; Scott et al., 1991), changes in grazing pressure and fire, besides stochastic variations in rainfall, are thought to drive decadal-scale variability (Gillson, 2004; van Rooyen et al., 2018). Interestingly, different studies have shown that precipitation, fire, and herbivory may have different effects on encroacher species, and thus a different effect on savanna stability (Joubert et al., 2008; Joubert, Smit, & Hoffman, 2012, 2013). Hence, disentangling triggers that induce changes of state, as well as understanding stabilizing feedbacks, requires long time series with adequate taxonomic resolution.

In addition, gradual change in environmental conditions is assumed to have little effect on the feedbacks that support a stable state but still alters its resilience (Scheffer et al., 2001). For example, management legacies are assumed to affect the properties of temperate woodland responses to current environmental change up to 200 years after disturbance (Perring et al., 2018). Similarly, shrub encroachment of Namibian savannas has been related to management strategies during colonial administration (Moyo et al., 1993; Rohde & Hoffman, 2012; Verlinden & Kruger, 2007). However, it remains unclear to what extent management may have affected species composition and thus savanna stability in Namibia.

Lake sediments represent unique environmental archives. Traditionally, they are mostly used to resolve millennial-scale environmental changes in savannas (Metwally, Scott, Neumann, Bamford, & Oberhänsli, 2014; Quick et al., 2018). Vegetation reconstruction using pollen deposited in lakes has been demonstrated to be a powerful method to track structural and compositional changes in savannas despite the low pollen taxonomic resolution (Miller & Gosling, 2014; Neumann, Scott, Bousman, & van As, 2010). Recently, sedimentary ancient DNA (*sedaDNA*) using the g-h universal primers (Taberlet et al., 2007) was established as a further proxy. This technique enables the reconstruction of tropical plant diversity at high taxonomic resolution (Boessenkool et al., 2014; Bremond et al., 2017). Further, fire dynamics and its impact on savanna vegetation can be reconstructed from sedimentary charcoal (Colombaroli, van der Plas, Rucina, & Verschuren, 2018; Gillson & Ekblom, 2009), while carbon ( $\delta^{13}\text{C}$ ) and deuterium ( $\delta\text{D}$ ) isotope compositions of plant-wax *n*-alkanes can be used to reconstruct vegetation structure and hydrological variability (Garcin et al., 2018; Miller et al., 2019; Walther & Neumann, 2011). In addition, grain-size analyses of lake sediments are useful to reconstruct sediment mobility in savanna landscapes and to identify soil erosion (Walther & Neumann, 2011). However, high-resolution multiproxy approaches that include the use of *sedaDNA*, and that track vegetation and environmental changes in semiarid savannas at a decadal scale, are still missing.

Lakes are rare in southern Africa. Lake Otjikoto, a sinkhole lake located in the Karstveld area in northern Namibia (Figure 1), represents a unique environmental archive (Scott et al., 1991). Applying a multiproxy approach to lake sediments from Otjikoto including the analyses of pollen, *sedaDNA*, biomarkers, geochemical proxies,

grain size, and macrocharcoal analyses, this study provides the first decadal-scale record of the turnover of open savanna woodlands to shrublands from northern Namibia in the period between about 1,850 and the present day. This enables the discussion of savanna state changes and, in combination with climate and land-use data, the investigation of potential triggers and drivers and related feedbacks of the vegetation.

## 2 | MATERIALS AND METHODS

### 2.1 | Site

The lake basin is formed from a groundwater cave, which developed from soluble dolomite of the Tsumeb subgroup of the Damara sequence (Kamona & Günzel, 2007). Water depth is ~71 m; diameter is ~102 m. Water pH measurements vary between 7.0 and 8.7 (Marchant, 1980). The potential natural vegetation in the vicinity of the lake comprises *Terminalia prunioides*-*Combretum apiculatum* woodlands (Mendelsohn, El Obeid, & Roberts, 2000). Accordingly, a low grass layer and dominance of trees characterize this vegetation type, which is common on shallow sandy-loamy soils. *Terminalia prunioides*, *C. apiculatum*, *C. imberbe*, *Commiphora glandulosa*, *Vachellia reficiens*, *V. hebeclada*, and *Senegalia mellifera* are the most common taxa. The shrub layer comprises *Grewia* sp., *Gymnosporia senegalensis*, *Dichrostachys cinerea*, and *Croton gratissimus*. However, a dense shrubland characterizes the current local vegetation, which is mainly composed of *T. prunioides*, *S. mellifera*, *D. cinerea*, *Ziziphus mucronata*, and *Grewia* sp. Shore vegetation is sparse, with *Sclerocarya birrea* and *Ficus* growing on the steep rock walls




FIGURE 1 Map of the study site in Namibia

surrounding the lake. Crop fields are close to the lake. Mean annual precipitation is 456 mm, falling mainly in the summer months (November–March); mean monthly temperature in summer is 27°C and in winter 15°C (Tsumeb climate station; Harris, Jones, Osborn, & Lister, 2014). Cattle farming is the dominant land use.

## 2.2 | Materials

Two cores—15OJ10 (length 31 cm,  $\varnothing = 6$  cm) and 15OJ06 (length 30 cm,  $\varnothing = 6$  cm)—were collected in March 2015 using UWITEC coring equipment at a water depth of 50 m (19.19467°S, 17.54980°E). The core extraction was carried out with a pulley and without motor support. Divers from Dantica diving club Windhoek assisted with the core extraction from the lake bed. Core 15OJ10 was sampled at a 0.5 cm resolution in the field. We obtained 61 samples for pollen analysis; 30 subsamples were used for sedimentological, geochemical, *n*-alkane, and isotope analyses. Core 15OJ06 was sampled at a 1.5 cm resolution at the Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research. We obtained 21 samples for *sedaDNA* and charcoal analysis. All samples were stored at 4°C.

## 2.3 | Dating

Sixteen subsamples from core 15OJ10 and 21 from core 15OJ06 were analyzed for  $^{210}\text{Pb}$ ,  $^{226}\text{Ra}$ , and  $^{137}\text{Cs}$  by direct gamma assay at the Liverpool University Environmental Radioactivity Laboratory. Measures were obtained using an Ortec HPGe-GWL well-type coaxial low background intrinsic germanium detector (Appleby et al., 1986).  $^{210}\text{Pb}$  dates were calculated using the CRS dating model (Appleby & Oldfield, 1978). Corrections were made using the  $^{137}\text{Cs}$  date 1964 as a reference point (Appleby, 2001).

## 2.4 | Pollen and palynomorph analysis

Pollen extraction was performed using standard procedures (10% HCl, KOH, 40% HF—including 4 hr boiling—and acetolysis) (Faegri & Iversen, 1989; Moore, Webb, & Collison, 1991). We added one tablet of *Lycopodium* spores (batch 1,031) to each sample to calculate pollen concentrations (Stockmarr, 1971).

At least 400 pollen grains were counted per sample using a microscope of x400 magnification. Critical identification was made by x1,000 magnification based on standard literature (Bonnefille & Riollet, 1980; Gosling, Miller, & Livingstone, 2013; Lezine, 2005; Schüler & Hemp, 2016; Scott, 1982), taxonomical studies (Banks & Lewis, 2018; Poston & Nowicke, 1993), and online pollen catalogues (Bremont, 2011). We differentiate the *Acacia* type into *Vachellia* and *Senegalia* by means of columella identification (Kyalangalilwa, Boatwright, Daru, Maurin, & van der Bank, 2013). Spores' identification follows Gelorini, Verbeken, Geel, Cocquyt, and Verschuren (2011). Pollen percentages were calculated based on the total

terrestrial pollen sum per sample. Shore taxa (*Ficus*, Cyperaceae) and aquatic taxa were not included in the pollen sum.

## 2.5 | Sedimentary ancient DNA analysis

The sampling of core 15OJ06 for *sedaDNA* was carried out at +10°C in a climate chamber that was previously cleaned (DNA-ExitusPlus™—AppliChem, Germany; deionized water and ethanol 100%) to prevent contamination. Extraction and polymerase chain reaction (PCR) followed the same procedures as described in Zimmermann et al. (2017). For each sample ( $n = 21$ ) and extraction negative control ( $n = 3$ ), we performed six PCR replicates to ensure the accurate detection of taxa (Ficetola et al., 2015). For each PCR, we used two negative controls and different primer tag combinations leading to a total of 126 PCR products and 30 negative controls. All PCR products were purified using the MinElute PCR Purification Kit (Qiagen, Hilden, Germany) according to the manufacturer's instructions. DNA concentrations were measured with the dsDNA BR Assay Kit (Life Technologies, USA) using 1  $\mu\text{l}$  of the purified amplifications with the Qubit 2.0 fluorometer (Invitrogen, Carlsbad, CA, USA). Purified samples were pooled in equal concentrations and sent to the Fasteris SA sequencing service (Geneva, Switzerland) for library preparation following the MetaFast protocol. Sequencing was realized on an Illumina HiSeq 2,500 platform (2  $\times$  125 bp) (Illumina Inc., San Diego, CA, USA) using 1/10 of a flow cell lane. Obtained data were processed and assigned their taxonomic name using the software package OBITools (Boyer et al., 2016) following the descriptions in Zimmermann et al. (2017). Sequences selected for further analyses needed at least 95% best identity with respect to an entry in the reference database and at least 10 sequence counts in at least two samples in the dataset. Taxa percentages were calculated based on the sum of sequence counts per sample. After sequencing, two replicate batches were excluded from the analysis because they lacked sequencing depth, probably due to errors in the allocation of tags.

## 2.6 | Biomarker analysis

For the analysis of *n*-alkanes, 30 freeze-dried sediment samples were extracted using a dichloromethane containing 1% methanol. Further procedures used for the analysis of *n*-alkanes follow those described by Vogts, Moossen, Rommerskirchen, and Rullkötter (2009) and Badewien, Vogts, Dupont, and Rullkötter (2015).

The carbon ( $\delta^{13}\text{C}$ ) and deuterium ( $\delta\text{D}$ ) isotopic compositions of saturated hydrocarbons were measured using gas chromatography isotope ratio mass spectrometry consisting of a GC-Unit (7890N; Agilent Technology, USA) connected to GC-Isolink and coupled to a Delta V Plus mass spectrometer (Thermo Fisher Scientific, Germany). We used a combustion interface for  $\delta^{13}\text{C}$ , and a pyrolysis furnace for  $\delta\text{D}$  (Supplement 1.A). Isotope measurements were controlled based on *n*-alkane standards with known isotopic composition (Campro Scientific, Germany, and Arndt Schimmelmann, Indiana University, USA).

Isotope measurements of bulk organic carbon ( $\delta^{13}\text{C}_{\text{org}}$ ) were made on decarbonized samples using an elemental analyser (1,108 from Carlo Erba) coupled to a MAT 252 isotope ratio mass spectrometer from Thermo Fisher Scientific (Supplement 1B). All subsamples were analyzed in duplicate.

## 2.7 | Charcoal analysis

Charcoal was extracted from 1 cm<sup>3</sup> subsamples following Mooney and Tinner (2011). Samples were bleached (24 hr) and washed through a sieve (125  $\mu\text{m}$ ) to separate macrocharcoal particles. All charcoal particles (>125  $\mu\text{m}$ ) were counted for each sample using a binocular microscope at  $\times 15$  magnification and a Bogorov counting chamber.

## 2.8 | Geochemical analysis

Total carbon (TC) and total nitrogen (TN) contents were analyzed for 30 freeze-dried subsamples (core 15OJ10) using an Elementar vario EL III (CNS) analyser. An Elementar vario MAX C was used to quantify total organic carbon (TOC). Total inorganic carbon (TIC) was calculated by deducting TOC from TC values.

## 2.9 | Grain-size and end-member modeling analysis (EMMA)

Samples were pretreated with CH<sub>3</sub>COOH (10% for 24 hr) to remove carbonates, and with H<sub>2</sub>O<sub>2</sub> (initial 0.3%, with addition of 10 ml (35%) every second day for up to six weeks) to remove the organic sediment fraction. Grain-size analysis was performed using a Coulter LS 200 Laser Diffraction Particle 30 Analyser.

Robust grain-size end-members (EMs) have been modeled from all grain-size distributions using the EMMA algorithm of Dietze et al. (2012). We used the compact protocol in the R package EMMAgeo (Dietze & Dietze, in press). Definition of the robust EMs considered the main mode classes that were modeled most frequently (see kernel density estimate in Figure S2a), when the number of end-members and weight transformation limit was varied between a minimum and maximum value (after Dietze et al. (2012) and Dietze & Dietze, in press). EM loadings (Figure S2b) represent scaled grain-size class-wise EM contributions, and scores (Figure S2c) represent the contribution of EMs in the samples.

## 3 | RESULTS

### 3.1 | Chronology

The core 15OJ10 covers the period from 1886 to 2015. The results suggest a relatively steady accumulation since the late 1980s with a mean sedimentation rate of 0.34 cm/year. Sedimentation rates in

the 1930s were around 0.21 cm/year, increasing to around 0.31 cm/year by the early 1980s (Figure S3a). Low <sup>210</sup>Pb concentrations in the pre-1930 samples made accurate dating problematic. However, <sup>210</sup>Pb dates suggest even lower sedimentation rates in the late 19th and early 20th century with a mean value of 0.15 cm/year. We applied linear interpolation to obtain the chronology of segments between dated samples.

The core 15OJ06 covers the period from 1849 to 2015. Since the <sup>210</sup>Pb concentrations reach the limit of detection at a depth of 23.25 cm (dated to 1907), we calculated ages for samples below 23 cm by linear extrapolation. The sedimentation rate since the early 1980s has a mean accumulation value of 0.30 cm/year (Figure S3b). The results show lower sedimentation rates increasing from 0.14 cm/year during the first few decades of the 20th century to 0.20 cm/year during the period from 1940 to the 1970s.

### 3.2 | Pollen and nonpollen palynomorph records


We identified 116 pollen and spore taxa (Table S4). The most dominant taxa are Poaceae, Anacardiaceae, and Combretaceae. The pollen diagram captures a signal of grasses decreasing up the core, with values dropping below 60% in the 1970s and from the turn of this century, while the pollen signal of woody vegetation increases (Figure 2).

Based on a visual inspection of all proxies of both cores and guided by the results of the cluster analysis of the pollen record, we defined three zones. The taxa association in zone 1 (1886–1910) comprises Combretaceae, *Dichrostachys*, *Croton*, *Grewia*, *Euclea*, *Senegalia*, and *Petalidium*. Bignoniaceae, *Spirostachys*, and *Colophospermum mopane* yield high percentages at the border to zone 2. The pollen signals of *Dichrostachys* and *Ziziphus* are high in zone 2 (1914–1997), while *Grewia* is lower compared with zone 1. Furthermore, zone 2 is characterized by high abundance of *Myrtaceae*, as well of *Alternanthera*, *Tribulus*, *Pentzia*, *Monechma*, *Opilia*, *Montinia*, *Gymnosporia*, and Euphorbiaceae taxa. Zone 3 (1998–2015) is characterized by high values in *Vachellia*, *Senegalia*, *Grewia*, and *Ziziphus*. This goes together with an increased signal of *Alternanthera* and low values for *Euclea*. The signal of *Prosopis* increases toward the top.

NPP results show a gradual decrease of *Gelasinospora* and *Sordaria* toward the top of the core. *Glomus* reaches highest values in zone 3.

### 3.3 | Sedimentary ancient DNA records

We obtained 4,211,287 sequence counts (sqc) from Illumina sequencing for 21 samples with four replicates. After bioinformatic filtering, 1,840,462 sqc were assigned to 177 taxa (Table S5). Of these, 37 could be identified to family level (905,299 sqc), 58 to genus level (323,335 sqc), and 64 to species level (239,924 sqc). Most dominant taxa are *Moraceae*, *Combretum*, and Anacardiaceae. Overall, the *sedaDNA* record reflects a turnover from taxa characteristic of open savanna woodlands in zone 1 to shrubland taxa in zone 3 (Figure 3).



**FIGURE 2** Percentage abundance diagram of selected pollen and spore taxa. Arrangement of taxa follows their dominance in the pollen zones (separated by dotted lines). Taxa are displayed in subgroups: trees and shrubs (brown), Karroid shrubs (gray), taxa associated with pastures (green), local shore taxa (blue), and spores (purple). Pollen influx, as well as mean abundance percentages of woody, grass, and other pollen taxa, is displayed. Truncated values are indicated on the bars

Zone 1 (1849–1907) is characterized by dominance of *Dichrostachys*, *Combretum*, *Ebenaceae*, and *Catophractes alexandri* and the occurrence of *Gossypium*. A mixed grass layer consisting of perennial grasses such as *Fingerhutia africana*, *Schmidtia*, and *Cenchrus* and annual grasses such as *Brachiaria deflexa*, *Aristida*, and *Tragus* characterizes zones 1 and 2, as well as the occurrence of Karroid shrubs. Zone 2 (1919–1991) is distinguished by a high abundance of taxa associated with pastures such as *Antigonon leptopus*, *Mentzelia*, *Tribuloideae*, *Alternanthera pungens*, *Geigeria*, and *Acanthospermum*, as well as of shrub taxa from the *Euphorbiaceae*, *Celastraceae*, and *Opiliaceae* families. In zone 2, the signals of *Myrtoideae* are high. Zone 3 (1996–2015) is characterized by high signals of *Terminalia*, *Vachellia*, and *Senegalia*, as well as *Grewia*, *Rhamnaceae*, and *Medicago*. The perennial grasses *Hyperthelia dissoluta*, *Panicum*, and *Heteropogon contortus* characterize zone 3. Furthermore, the relative abundance of *Moraceae* increases in the upper part of the core.

### 3.4 | Biogeochemical and macrocharcoal analysis

Concentration of organic and inorganic carbon (TOC, TIC) is highest in zones 1 (5.1%) and 3 (4.9%), respectively (Figure 4). The average chain length (ACL; Poynter, 1989) values for *n*-alkanes with an odd number of carbon atoms ( $C_{27}$ – $C_{33}$ ) range from 31.1 to 30.3 and increase with depth. The carbon preference indices ( $CPI_{25-35}$ ; Marzi, Torkelson, & Olson, 1993) range from 13.6 to 3.4 with an average of 8.4. The C/N ratios range from 18.05 to 13.4 with an average of 16.04. The carbon isotopic compositions of organic matter ( $\delta^{13}C_{org}$ ) range from  $-19.9$  to  $-27.3$  with an average of  $-24.0\text{‰}$  and with the more  $^{13}C$ -depleted values in zone 3. Similarly, the weighted mean  $\delta^{13}C$  values of *n*-alkanes with an odd number of carbon atoms ( $\delta^{13}C_{WMA27-33}$ ) range from  $-24.6\text{‰}$  to  $-32.3\text{‰}$  and have the more  $^{13}C$ -depleted values in zone 3. We also find the more depleted weighted mean  $\delta D$  values ( $\delta D_{WMA27-33}$ ; range:  $-135.7\text{‰}$  to  $-159.6\text{‰}$ )

in this zone. In addition, macrocharcoal flux is highest in zones 1 and 2 and decreases upwards in the core (Figure 3).


### 3.5 | Grain-size end-members

We obtained five rEM (robust grain-size end-members) after calculating similarly-likely EM models (Figure S2). The mean rEMs explain 77% of the mean total variance of the original grain-size dataset. The rEM1 (23% explained variance) has a primary mode between 0.2 and 14.5  $\mu m$  with a maximum at 1.8  $\mu m$ , covering the clay and very fine silt fraction. The rEM2 (13% explained variance) has a broad primary mode between 3 and 66  $\mu m$  covering the whole silt fraction, with a maximum at 18.7  $\mu m$ , which corresponds to medium silt. The rEM3 (14% explained variance) has a mean mode between 16 and 144  $\mu m$ , and a maximum at 51  $\mu m$  (medium silt to fine sand fraction). The rEM4 (23% explained variance) has a very robust mode in the very fine to fine sand between 51 and 310  $\mu m$ , with a maximum at 127  $\mu m$ . The coarsest component rEM5 (26% explained variance) has a robust mode in the fine to medium sand between 86 and 454  $\mu m$ , and a maximum at 211  $\mu m$ .


## 4 | DISCUSSION

### 4.1 | Source of the lake sediments

We consider terrestrial plants to be a major source of organic matter in the sediments of Lake Otjikoto due to their high  $CPI_{25-35}$  and C/N values (Meyers, 1997). Although higher TOC values ( $>4\%$ ) below 25 cm depth, and the oscillating variation of  $\delta^{13}C_{org}$  content below 18 cm, suggest a bacterial and algal contribution to the sediments (cf. Kristen et al., 2007), corresponding C/N values ( $>16$ ) indicate a higher proportion of allochthonous input relative to the autochthonous component of the organic matter (Meyers, 1997).



**FIGURE 3** Diagram of selected *sedaDNA* taxa. Arrangement of taxa follows their dominance in the *sedaDNA* zones (separated by dotted lines). Taxa are displayed in subgroups: trees and shrubs (brown), Karroid shrubs (gray), taxa associated with pastures (green), grasses (yellow), and local shore taxa (blue). Macrocharcoal influx is also displayed



**FIGURE 4** (A) Biogeochemical and sedimentological proxies from core 150J10: Carbon preference indices (CPI<sub>25-35</sub>); average chain length of *n*-alkanes with an odd number of carbon atoms (ACL<sub>27-33</sub>); weighted mean of carbon isotopic compositions of *n*-alkanes ( $\delta^{13}C_{WMA27-33}$ ); carbon isotopic composition of organic matter ( $\delta^{13}C_{org}$ ); weighted mean of deuterium isotopic compositions of *n*-alkanes ( $\delta D_{WMA27-33}$ ); total organic carbon (TOC) content; total inorganic carbon (TIC) content; total organic carbon to nitrogen ratio (C/N); and scores from robust end-member loadings (REM). Dotted lines indicate pollen zones. (B) Historical records: Mean annual precipitation (3-year running mean—MAP) was calculated from climate data (meteorological station Tsumeb), vertical black dotted line corresponds to the long-term MAP, and red dotted lines correspond to the standard deviation (SD); livestock numbers Namibia (data 1915–1959, 1961, 2015; Lange et al., 1998; FAO, 2018b); CO<sub>2</sub> emissions Namibia (FAO, 2018a); documented events (D: drought; F: flood; FMD: foot-and-mouth disease; CBPP: contagious bovine pleuropneumonia (Directorate of Planning, 2011, 2005; Nicholson & Selato, 2000; Schneider, 2012; Sweet, 1998))

## 4.2 | Savanna state change: from open woodland to dense shrubland

### 4.2.1 | Multidecadal shift in the grass/woody ratio

Our multiproxy approach indicates a shift in the dominance among grasses and woody vegetation. In particular, our results show an increase in woody vegetation since about 1915 in the Tsumeb region with a particularly strong increase after about 1998. These results validate vegetation models that suggest an increase in woody cover at low rates since 1,850 in Africa, and an accelerated rate between 1990 and 2010 (Higgins & Scheiter, 2012). Our findings

also support results from remote-sensing analyses of sub-Saharan savannas, which have observed an increase in woody vegetation cover over the last three decades (Mitchard & Flintrop, 2013; Venter et al., 2018).

The ratio of Poaceae/woody pollen (Figure 2) indicates that the current grass/woody ratio is exceptional for the last 170 years and is similar to the ratio found by Scott et al. (1991) in a previous palynological study from Lake Otjikoto. Scott et al. (1991) identified two earlier woodland phases with woody pollen abundances up to 33% and Poaceae pollen below 60% (Figure S6). In our study, we estimate grass pollen abundances to decrease to below 60% in the 1970s and in the 21st century, while woody pollen percentages increase up

to 32% suggesting a woodland phase in zone 3. These results are consistent with a modern pollen-vegetation study conducted in the vicinity of the lake, which found a mean woody pollen abundance of 33% by a mean woody cover of 74% at the study site (Tabares, Mapani, Blaum, & Herzs Schuh, 2018).

Biogeochemical results also support our pollen findings. The relatively constant  $\delta^{13}\text{C}_{\text{WMA27-33}}$  values throughout zones 1 and 2 indicate mixed  $\text{C}_3/\text{C}_4$  vegetation with the  $\text{C}_4$  proportion generally above 50% (Badewien, Vogts, & Rullkötter, 2015). Only in the very recent past (zone 3) does  $\delta^{13}\text{C}_{\text{WMA27-33}}$  shift toward more  $^{13}\text{C}$ -depleted values providing evidence for an increase of the  $\text{C}_3$  proportion up to around 80%, which is also supported by  $\delta^{13}\text{C}_{\text{Org}}$  values around  $-27\%$  (Meyers, 1997). It has been reported that southern African savanna trees show significant variations in their ACL values, which, for example, are higher for *C. apiculatum* and *C. mopane* than for *T. prunioides* and *Ziziphys* spp. (Kristen et al., 2010; Vogts et al., 2009; for Australian trees, see Hoffmann, Kahmen, Cernusak, Arndt, & Sachse, 2013). Assuming a significant contribution of trees to the sedimentary *n*-alkanes, the overall shift of ACL toward lower values throughout the profile is thus in agreement with the vegetation shifts inferred from pollen analyses and *sedDNA* as discussed in detail below.

In addition, grain-size distribution lets us infer changes in landscape openness via the type of sediment transport reflected in the grain-size rEM (Dietze et al., 2014). The two types of sand deposits represented by rEM4 and rEM5 indicate short-term, high energetic sediment input from run-off during rain events or sand storms in an open landscape with reduced grass cover (Dietze et al., 2014; Walther & Neumann, 2011). In particular, within the error of the age model, rEM5 can be associated with the main flood events documented in the region from the 1920s until 2000 (Figure 4). We assume that an increased woody cover (as observed from the 21st century) favors the development of soil crusts on bare patches by constraining trampling (Thomas & Dougill, 2007). Although soil crusts reduce infiltration and promote run-off and erosion, they also increase soil roughness limiting the detachment of soil particles (Bullard, Ockelford, Strong, & Aubault, 2018; Rodríguez-Caballero, Cantón, Chamizo, Afana, & Solé-Benet, 2012; Valentin, 1993). Accordingly, soil crusts may retain the coarser sand fractions, which would explain the low percentages of rEM4 and rEM5 in zone 3. However, prolonged drought (as observed in the 1990s), followed by heavy rains, can damage the crust surfaces (Kidron, Ying, Starinsky, & Herzberg, 2017; Thomas & Dougill, 2007), causing the loss of fine sand-sized particles (100–200  $\mu\text{m}$ ), which require less energy to be detached compared to coarser and finer-sized particles (Bullard et al., 2018; Salles, Poesen, & Govers, 2000). This would explain why only fine sand (rEM4) mobilized during La Niña flood in 2006 (Figure 4).

An increased woody cover would also retain the local dust sedimentation (rEM3), by reducing the near-surface wind energies. Hence, the associated importance of more remote dust (rEM2) from several hundred kilometers away within the detrital sediment fraction is seen during the last decade.

## 4.2.2 | Compositional turnover in the course of shrub encroachment

Our pollen and *sedDNA* results indicate a savanna state change from open *Combretum* woodland to *Terminalia* shrubland during the 20th century at local (i.e., the vicinity of Otjikoto) and probably regional (i.e., the wider Tsumeb area) scales. These results are the first time series with sufficient taxonomic resolution to show compositional change in the course of shrub encroachment in a semiarid savanna at a multidecadal scale. They confirm theoretical assumptions (Joubert et al., 2008; Li, 2002) which were based on compositional observations of only a few decades or on the analyses of spatial gradients (van Rooyen et al., 2018; Strohbach, 2001).

According to our proxies, an open *Combretum* woodland dominated the landscape around Lake Otjikoto until the turn of the 19th to 20th century (i.e., zone 1 in our record). A high grass cover and the dominance of *Combretum* species in the tree layer are characteristic of this vegetation type (Mendelsohn et al., 2000), which is represented by high Poaceae pollen values and up to 50% *Combretum* in the *sedDNA* records. Our results also show that *Grewia*, *D. cinerea*, *Croton*, *Kirkia acuminata*, *C. alexandri* (Bignoniaceae), and Ebenaceae (likely *Euclea* sp.) were common taxa in the tree and shrub layer of this vegetation type. This aligns well with modern observations of this vegetation type preserved on flat areas between dolomite hills (Giess, 1971), particularly on moderately deep soils (Mendelsohn et al., 2000). In addition, our results reflect botanical observations from the early 20th century of the existence of woodlands and lime steppes in the Tsumeb region (Dinter, 1918). A high pollen signal of *Petalidium* (likely *P. eurychlamys*) in the late 19th century and the *sedDNA* record of *Gossypium* reflect early observations regarding both taxa as local indicators of open broad-leaved savanna vegetation (Dinter, 1922; Mildbraed, 1941). We assume that open *Combretum* woodlands represent the potential natural vegetation of the region and a stable savanna state. These results confirm a previous low-resolution pollen record from Lake Otjikoto (Scott et al., 1991) that revealed the presence of open Combretaceae-*Spirostachys* woodlands in the region during the late Holocene.

Our results allow us to infer a transition phase that ranges from 1920 to about 1997 (zone 2 in our record) and provide empirical evidence to previous studies, which assume a transition period of several decades between alternative stable grassland/woodland states (Higgins & Scheiter, 2012; Joubert et al., 2008).

The transition is marked by a high turnover in the vegetation proxy composition. While Poaceae are still abundant during this period, some shrub taxa are also common (*Ziziphys* (Rhamnaceae), *Grewia*, *Phyllanthus/Phyllanthaceae*) particularly during the second half of the 20th century. Furthermore, this transition phase is characterized by unique taxa, particularly those associated with disturbance (e.g., common pasture taxa such as *Antigonon* and *Mentzelia* (López-Olmedo, Meave, & Pérez-García, 2007), and nonpalatable weeds such as *A. pungens*, *Tribulus/Tribuloideae*, and *Geigeria* (Tabares et al., 2018)). This transition is also distinguished by the



propagation of dwarf shrubs (*Monechma*/*Justiciinae*, *Pentzia*), annual grasses (*Tragus*, *B. deflexa*, *Aristida*), and the increase of shrub taxa (*Croton*, *Euphorbia*, *Gymnosporia* (Celastraceae), *Montinia caryophyllacea*, *Opilia*). We assume that our inferred transition phase corresponds to the unstable state described by Joubert et al. (2008), which is distinguished by the above-mentioned characteristics and by the spread of encroacher seedlings (e.g., *S. mellifera*) in a mixed grass sward. The increase of *Senegalia* pollen in zone 3 may therefore indicate the maturity of *S. mellifera* trees established in zone 2.

The vegetation at the local and regional scale around Lake Otjikoto after 1997 (zone 3 in our record) can be characterized as a *T. prunioides* shrubland (*T. prunioides* association in Hüttich et al., 2009; Mendelsohn et al., 2000; *Acacia-T. prunioides* in Strohbach, 2014). This represents a stable encroached savanna state (Joubert et al., 2008) and is supported by the finding that the compositional change within zone 3 is rather small compared to zone 2. The vegetation is characterized by high abundances of *Vachellia*, *Senegalia*, and *Terminalia* and the decline of *Combretum* (Figure 3) in our sedimentary vegetation proxy records, which likely indicates the spread of local encroacher species such as *V. reficiens*, *S. mellifera*, and *T. prunioides* in the region (De Klerk, 2004; Strohbach, 2001). The transition toward *T. prunioides* vegetation type is also reflected by the changes in the shrub layer: The abundances of *Grewia* and *Ziziphus* (Rhamnaceae) increase, while *C. alexandri* (Bignoniaceae) and *Euclea* (Ebenaceae) decrease. Furthermore, a state change toward encroached savanna is indicated by the decline in Poaceae pollen signals suggesting a reduction in grass cover, which is characteristic of *T. prunioides* shrubland (Strohbach, 2014). Interestingly, the increase of Fabaceae (e.g., *Vachellia*, *Senegalia* in pollen and *sedaDNA* values) in zone 3 corresponds with more depleted  $\delta^{13}\text{C}$  values of organic matter. This matches modern studies along spatial gradients that likewise found depleted  $\delta^{13}\text{C}$  values in Fabaceae  $\text{C}_3$  plant matter (Badewien, Vogts, & Rullkötter, 2015; Vogts et al., 2009).

We hypothesize that the current *Terminalia* encroached state may have occurred earlier in the region, as suggested by the compositional changes reflected in an early pollen record of Lake Otjikoto

(Scott et al., 1991). In particular, the early woodland phase (zone O1), which is dominated by Combretaceae, may correspond to a phase of *Terminalia* encroachment, since the abundance of *Spirostachys* is low, while *Grewia* and *Acacia* have high pollen signals (Figure S6). Interestingly, this phase is preceded by high abundances of *Tribulus*, which suggests a transition phase.

Given the changes in species composition and in the grass/woody ratio observed in our study, our results provide empirical evidence to support the hypothesis of Rohde and Hoffman (2012) that transitions from an open woodland into an encroached state in savannas from northern Namibia may occur on a scale of one century and without reversals to a grassy state.

### 4.3 | Triggers and drivers

It is likely that a complex network of global, regional, and local triggers and drivers including land management (forestry, husbandry/cultivation), precipitation, and atmospheric  $p\text{CO}_2$  caused the observed vegetation turnover in the Tsumeb area (Figure 5). However, limited data availability prevents us from doing straightforward multivariate statistical analyses such as decomposition of explained variances to statistically relate driver changes to changes in vegetation (Tian, Herzsuh, Mischke, & Schlütz, 2014); instead, only trends can be compared.

#### 4.3.1 | Forestry in the context of mining

Our results support previous studies that assign the emergence of shrubland in northern Namibia to forest management (Strohbach, 2001). In particular, the expansion of thickets in the Tsumeb region, as indicated in forest reports from the 1950s, has been assigned to selective and broad-scale logging in the course of mining in the early 20th century (Erkkilä, 2001; Lau & Reiner, 1993). *Combretum*, *Spirostachys africana*, and *C. mopane* were cut for props and fuel for the Tsumeb mine (Erkkilä & Siiskonen, 1992), and this is reflected in the decline of these taxa in the Otjikoto




FIGURE 5 Network of triggers and drivers of savanna vegetation change

pollen record. After logging, it is likely that dense thickets of *S. africana* formed because of its ability to form stump shoots and root suckers (Gandiwa, Gandiwa, & Mxozza, 2012; Moyo et al., 1993). The *Spirostachys*/Hippomaneae peaks in the pollen and *sedaDNA* records may indicate resprouting events during the Depression and the Second World War times, in which the mine production came to standstill (Lau & Reiner, 1993). In contrast, other local taxa became reduced because of their low growth rates (e.g., *C. mopane*, Cunningham, 1996) or their weak coppicing ability (e.g., *Combretum*, Strohbach, 2001). In order to supply the mines, *Eucalyptus* was planted in the Tsumeb region from the beginning of the 20th century (Dinter, 1909, 1918) and from 1949 at Lake Otjikoto (Lau & Reiner, 1993), which is reflected in our proxy records (Myrtaceae/Myrtoideae). Later on, gradual environmental changes, such as changes in the tree species composition and the reduction of soil moisture as a consequence of deforestation (Moyo et al., 1993), may have favored the spread of *Vachellia*, *Senegalia*, and *T. prunioides*, which are better adapted to drier conditions.

### 4.3.2 | Husbandry

Land-use records from this study and census information (FAO, 2018b; Lange, Barners, & Motinga, 1998) indicate that the intensity of husbandry increased throughout the 20th century, which likely affected grass species composition and productivity.

The signal of *Sordaria*, an obligate coprophilous fungus spore associated with dung deposits of wild/domestic herbivores (Dietre, Gauthier, & Gillet, 2012; van Geel & Aptroot, 2006; van Geel et al., 2011), suggests the continuous presence of herbivores in the vicinity of the lake. The high values of *Sordaria* in the first half of the 20th century could be the result of herding of cattle at watering points and on productive pastures close to the lake, particularly during drought episodes (as suggested by the fungal peak at about 1946 [Figures 2 and 4]). Surprisingly, the signal of *Sordaria* decreases in the second half of the 20th century although grazing intensity was likely highest during this period—census information on cattle in Namibia indicates an increase from ~700k animals before 1930 to ~2500k after 1950 (FAO, 2018b; Lange et al., 1998).

The decrease in the *Sordaria* signal could be related to a decline of transhumance pastoralism (as similarly observed by Gelorini, Ssemmanda, & Verschuren, 2012] at lakes in Uganda), which is associated with changes in land tenure and management in the region. During the first half of the 20th century, most of the Otjikoto region became freehold land (Mendelsohn, Jarvis, Roberts, & Robertson, 2009), which resulted in the construction of fences, concentration of livestock, and eventually the restriction of traditional transhumance in the surroundings of the lake. From the 1960s, movement of cattle was further restricted because of the outbreak of foot-and-mouth disease (Schneider, 2012). The increase of cattle in the mid-20th century likely originated from the intensification of farming from the 1950s (Lau & Reiner, 1993). Additionally, the introduction

of subsidies for feed and water-hole exploration in response to the prolonged drought in the early 1960s (Figure 4) allowed ranchers to increase their stocks in years with good rain, and to maintain them in the drought years (Schneider, 2012; Sweet, 1998).

As reflected in the *sedaDNA* record, these practices led to the decrease of highly palatable grasses such as *Schmidtia* (likely *S. papophoroides*), *B. deflexa*, and *F. africana*. They were, if at all, replaced by less palatable grasses such as *H. dissoluta* and *H. contortus* which are characteristic of overgrazed areas (Kgosikoma, Mojeremane, & Harvie, 2015). High grazing pressure on pastures during the mid-20th century is also indicated by above-average signals of nonpalatable taxa such as *Geigeria* (Strohbach & Kutuahuripa, 2014) and *Acanthospermum* (Tolsma, Ernst, & Verwey, 1987), as well as by trampling-resistant taxa such as *Alternanthera* and *Tribulus*/Tribuloideae (Tabares et al., 2018). High grazing pressure alongside poor grazing resources would also explain the cultivation of *Medicago* to complement livestock fodder from the 1990s around Lake Otjikoto as indicated in the *sedaDNA* record (Figure 3).

The persistent high grazing pressure on pasture with already low grass density may have lowered the seed production of grasses to such an extent that potential niches could not be (re)invaded (O'Connor & Pickett, 1992; Tessema, de Boer, & Prins, 2016). This might explain why our *sedaDNA* record lacks *F. africana* and *Schmidtia* (likely *S. papophoroides*) toward the present day. According to the state-transition models, the lack of both species may also reflect the vegetation's response to overgrazing when disturbance passed a threshold.

### 4.3.3 | Fire management

Our results show that there were changes in fire management related to husbandry and cultivation near Lake Otjikoto, which further altered the resilience of *Combretum* woodlands and supported shrub establishment in the long-term.

High macrocharcoal flux and high *Gelasinospora* values (a fungus growing preferentially on burned wood [Revelles & van Geel, 2016]) in sediments of the early 20th century point to local forest fires within a radius of about 5 km from the lake (Clark, 1988; Duffin, Gillson, & Willis, 2008). The emergence of cleared areas is reflected in the *sedaDNA* signals of pioneer grasses such as *B. deflexa* and *Tragus*. During the early 20th century, these areas were likely burned prior to crop cultivation as indicated by records of *Sorghum* (Lau & Reiner, 1993). Landscape opening is also reflected by the aeolian input of well-sorted coarse silt (rEM3), which was probably blown in from burned fields (Ravi et al., 2009).

During the mid-20th century, the coincidence of lower charcoal flux and a decrease in *Sorghum* *sedaDNA* suggest that fire was used to manage livestock grazing by fostering the establishment of palatable grasses (Lau & Reiner, 1993; Sweet, 1998). This would explain the rise in *sedaDNA* signals of *Cenchrus* (likely *C. ciliaris* (Gilo & Kelkay, 2017)) and *Panicum* (likely *P. maximum*) in the second half of the 20th century. Both these highly palatable perennial grasses

are common in the *T. prunioides* shrubland (Strohbach, 2014) and are usually planted for hay and pasture (van Oudtshoorn, 2016; Sweet & Burke, 2006). The expansion of monoculture farming in the Tsumeb region, particularly from the 1980s (Lau & Reiner, 1993), is also suggested by decreases in charcoal flux and *Gelasinospora* signals (Figures 2 and 3) toward zone 3, which in turn suggest the suppression of wildfire and/or the reduction of controlled fires to conserve pastures for livestock (Joubert, Smit, & Hoffman, 2012).

An initially reduced grass cover in combination with artificial fire suppression may have affected the grass/shrub competition by reducing mortality of woody seedlings and saplings (Case & Staver, 2017; Joubert et al., 2012). The remaining grass tussocks may have improved the survival of seedlings of encroaching Fabaceae taxa such as *Crotalaria* and *Prosopis*, as the tussocks provide soil moisture and protection against heat stress (De Dios, Weltzin, Sun, Huxman, & Williams, 2014; Wagner, Richter, Joubert, & Fischer, 2018), triggering a vegetation change into an encroached state.

#### 4.3.4 | Precipitation and atmospheric CO<sub>2</sub> changes

Comparing the pollen and *sedaDNA* records of *Vachellia* and *Senegalia* with the mean annual precipitation curve (Figure 4), it seems plausible that drought years from 1980 onwards contributed to the spread of these drought-tolerant taxa. Previous periods with consecutive rainy summers (particularly 1976–1979 and 1954–1956) and reduced grass cover also favored the germination and establishment of their seeds and saplings. These results are in line with several studies, which identified a positive relationship between rainfall, seed production, and seedling survival of encroacher species (Joubert et al., 2008; Joubert, Smit, & Hoffman, 2013; Wagner et al., 2018).

Our records show a die-back of *Senegalia* in the 21st century. This may be related to persistent drought stress, as reported from Botswana in 2013 (Joubert et al., 2013). Persistent drought can also inhibit the recovery of *Cenchrus ciliaris*, *C. apiculatum*, and *D. cinerea* (O'Connor, 1998), which explains the low *sedaDNA* signals of related taxa from the mid-1990s (i.e., zone 3).

Interestingly, the *Vachellia* pollen and *sedaDNA* signals in zone 3 are high when compared with *Senegalia*, which probably reflects differences in drought sensitivity. As observed by Joubert et al. (2013), *S. mellifera* has a lower tolerance for prolonged periods of moisture stress than *Vachellia* species such as *V. reficiens*, *V. erubescens*, and *V. erioloba*. According to the authors, *S. mellifera* mostly relies on an extensive shallow-root system, while *V. reficiens* and *V. erioloba* rely on a deeper tap-root system, which enables water uptake from deep soil layers.

The observed depletion of  $\delta D$  values of plant-derived *n*-alkanes since around 1992 (Figure 4) may also reflect the water uptake of deep-rooted trees from deep soil layers. In arid and semiarid environments, soil water has higher  $\delta D$  values than precipitation because of strong evaporation (Da Silveira Lobo Sternberg, 1988). Consequently, water from soil surface layers is more enriched in D compared to soil water from deep layers (De Deurwaerder et al.,

2018; Ehleringer & Dawson, 1992). Therefore, under dry conditions, xylem water from grasses is expected to have higher  $\delta D$  values compared to trees with a deep root system. Field studies performed in Namibian savannas show that xylem water from deep-rooted species such as *V. erioloba*, *V. hebeclada*, and *Prosopis juliflora* comes from D-depleted deep soil layers (Kanyama, 2017). Our pollen and *sedaDNA* results indicate an increase of *Vachellia* and *Prosopis* in zone 3. However, it is unclear to what extent the isotope composition of xylem water is reflected by leaf lipids. While greenhouse experiments found a correlation between leaf wax and soil water  $\delta D$  composition (Hou, D'Andrea, & Huang, 2008), field studies show that the leaf wax  $\delta D$  signature depends mostly on evaporative enrichment of leaf water during photosynthesis (Feakins & Sessions, 2010).

Considering the influence of photosynthesis on  $\delta D$  values, depletion in zone 3, particularly since the beginning of the 21st century, may also reflect an increase in the water-use efficiency of woody vegetation, which in turn is enhanced through, for example, reduced stomatal conductance and stomata density due to an increasing atmospheric  $pCO_2$  concentration (Figure 4). Studies conducted in tropical environments have shown that trees increase their water-use efficiency at higher  $CO_2$  concentrations, despite high transpiration rates (Brienen, Wanek, & Hietz, 2011; van der Sleen et al., 2014). In contrast, trees in arid and semiarid environments, and under higher  $CO_2$  concentrations, reduce their stomata opening while carbon assimilation increases (Evans, Schortemeyer, McFarlane, & Atkin, 2000; Krishnamurthy & Machavaram, 2000). This improved ratio of carbon assimilation-to-conductance results in an increase of transpiration efficiency (Evans et al., 2000), which in turn leads to  $^1H$  enrichment in leaves and thus a decrease in  $\delta D$  values. Such higher atmospheric  $CO_2$  concentrations may have favored the propagation of *Vachellia* and *Senegalia* toward the present day, because Fabaceae species are expected to benefit from such conditions in nutrient-limited savanna environments (Wagner et al., 2018; Ward, Hoffman, & Collocott, 2014). As observed by Evans et al. (2000), *Acacia* species increase their nitrogen content per unit foliage area when growing under a higher  $CO_2$  concentration. Consequently, the photosynthetic capacity of leaves increases (Evans, 1989), which can then improve the competitiveness of *Acacia* species against grasses.

The interpretation of the leaf wax  $\delta D$  is complicated further as the signature of the water source of plants may have changed as well. The high  $\delta D$  values around 1916 (i.e., in the transition between zones 1 and 2) probably reflect an enrichment of source water. Such an interpretation is supported by the high concentration of inorganic carbon (TIC) and an increase of *Ficus* pollen that inhabits the walls of the sinkhole. A low lake level may result from low rainfall between 1916 and 1922 (Figure 4). The decrease of  $\delta D$  values from the mid-1990s (i.e., at the transition to zone 3) despite high TIC and *Ficus* pollen signals and low precipitation, however, contradicts the interpretation of the leaf wax  $\delta D$  as primarily representing a precipitation signal (Niedermeyer et al., 2016; Sachse et al., 2012) but rather supports our interpretation of changes in the plant water source.

Our results suggest that environmental changes derived from management (i.e., reduced soil moisture, reduced grass cover, changes in species composition and competitiveness, reduced fire intensity) may have affected the resilience of *Combretum* woodlands, making them more susceptible to change into an encroached state by stochastic events such as consecutive years of precipitation and drought, and by high  $p\text{CO}_2$  concentrations. These results are in line with state-transition models, which predict gradual changes in environmental conditions to alter the resilience of stable states, thus enhancing the likelihood of a state change (Scheffer et al., 2001).

#### 4.4 | State-stabilizing feedbacks

Comparing our multiproxy record with historical climate and land-use data allows us to infer the feedback mechanisms that support the stable states observed in our study. This allows us to interpret the observed vegetation changes in the context of state-transition models (Hirota, Holmgren, Van Nes, & Scheffer, 2011; Scheffer et al., 2001).

##### 4.4.1 | Open *Combretum* woodland

Transhumance and intense fires, as reflected by our spore and charcoal records, may have stabilized an open woodland phase. A high grass cover, as observed in phase 1, may produce enough fuel for intense fires, which increase seedling and sapling mortality (Beckett & Bond, 2019; Case & Staver, 2017; Hirota et al., 2011; Staver, Archibald, & Levin, 2011). Since tree seedlings are not as fire-resistant as adult trees and some of them are very palatable (e.g., *Dichrostachys*), a combination of fire and browsing may also have prevented tree seedlings from being recruited to mature age classes (Gillson, 2004; Joubert et al., 2012; Scheffer et al., 2001).

In addition, wetter conditions may have favored competitiveness of *Combretum* and *Spirostachys* against drought-tolerant taxa. An early *Combretum-Spirostachys* woodland phase at Otjikoto was linked to wetter climatic conditions (Scott et al., 1991). Although vegetation models predict a woody state in savannas with increased precipitation (Hirota et al., 2011), our results allow us to infer that under such climatic conditions open *Combretum* woodland is possible, but also *Terminalia* shrubland, as the latter profits from wet years followed by persistent drought.

##### 4.4.2 | *Terminalia* shrubland

Several feedback mechanisms may stabilize the encroached state and inhibit the reverse transition toward an open woodland state:

First, open patches in degraded pastures may have supported the formation of trampling-resistant soil crusts (Dougill & Thomas, 2004) containing nitrogen-fixing cyanobacteria. These soil crusts, in turn, may have favored the establishment of nitrogen-fixing shrubby

Fabaceae, which eventually inhibit grass growth under their canopy (Maron & Connors, 1996). Such a feedback mechanism would explain the decrease in the C/N ratio (Figure 4) in the uppermost sedimentary record due to the allochthonous input of nitrogen-enriched soil materials (Talbot, 2005) in Lake Otjikoto. The latter is also supported by the predominance of fine-sized sediments in zone 3 (rEM1, rEM2), which may also depress the C/N values by uptake of inorganic N, since higher proportions of clay can absorb ammonia well (Meyers, 1997).

Second, the reduction of grass cover toward the present day in Tsumeb coincides with soil erosion, as suggested by *Glomus*, a spore of an arbuscular mycorrhizal fungus associated with eroded soils (Revelles & van Geel, 2016; van Geel et al., 2011) (Figure 2). On bare soil patches with soil crusts, run-off increases and water infiltration decreases, which further reduces the establishment of grass seedlings (Strohbach, 2001). In addition, climate- and/or human-induced damage to soil crusts can lead to soil losses (as suggested by rEM4 in zone 3), which in turn promote rill formation thus reducing the surface run-off and limiting the run-on input to vegetation (i.e., limiting the formation of temporary ponds and soils with water tables (Rodríguez-Caballero et al., 2012; Valentin, 1993)). Our study supports previous studies that find soil erosion in the course of bush encroachment since the second half of the 20th century (De Klerk, 2004; Strohbach, 2001).

Third, the practice of bush removal (indicated for Lake Otjikoto by a slight decrease of shrub pollen in the uppermost core section and observed in the locality of Otjiguinas Farm during fieldwork) enables the establishment of fast-growing species such as *H. dissoluta* (recorded in the uppermost section). This perennial grass spreads across disturbed fields forming dense tall swards, which inhibit the growth of other grasses (Cech, Edwards, & Olde Venterink, 2010; Jordaan, 2017).

Fourth, once established, woody vegetation with a mixed rooting system, such as *S. mellifera* and *Prosopis* sp., can suppress grass production (Smit, 2003), as they can effectively take up soil moisture from the upper soil layer after rain events with their lateral roots and survive dry phases due to water uptake by deep roots (Ansley, Boutton, & Jacoby, 2014; Brown & Archer, 1999). A reduced grass cover in turn reduces the intensity of fire, resulting in the self-propagation of a woody state (Higgins & Scheiter, 2012; Hirota et al., 2011; Scheffer et al., 2001).

Fifth, climate change (i.e., higher atmospheric  $p\text{CO}_2$  concentration and changes in precipitation variability) may induce physiological adaptations (as suggested by our  $\delta\text{D}$  record) enhancing the competitiveness of trees over grasses. Although persistent drought constrains the  $\text{CO}_2$ -related photosynthetic advantages for savanna trees (Nackley et al., 2018), the assimilation rates are still higher for  $\text{C}_3$  plants compared to  $\text{C}_4$  grasses under such conditions (Bellasio, Quirk, & Beerling, 2018). This is because  $\text{C}_4$  grasses under water restriction experience metabolic inhibitions that have a greater effect on leaf assimilation compared to the limited stomatal conductance of savanna trees (Bellasio et al., 2018). Furthermore, under restored watering conditions, photosynthetic rates recover faster when inhibited by stomatal than by metabolic factors, which in turn affects the competitiveness of  $\text{C}_4$

grasses under optimal conditions (Bellasio et al., 2018). Such a feedback mechanism implies that at higher atmospheric  $p\text{CO}_2$  concentration, an increase in the frequency of droughts and heavy rains may favor the expansion of  $C_3$  vegetation (Zhang et al., 2019).

These feedback mechanisms explain the ongoing encroachment processes observed at our study site up to the present, even though management intensity may have adjusted to the local carrying capacity (e.g., decline of cattle numbers from the 1970s (Lange et al., 1998) and a levelling off of cattle populations since 2000) or decreased (e.g., regulation of timber production since 1968 (Erkkilä & Siiskonen, 1992)). The current vegetation may thus represent a management legacy from the first half of the 20th century, as well as the impact of climate change.

## 5 | CONCLUSIONS

We have shown that the expansion of shrubs is a multidecadal trend characterized by the turnover from stable open woodland to a stable encroached state. Our results indicate that such a state change may occur on a scale of a century and without short-term reversions to a grassy state, but with an unstable transition phase, which covers about 80 years and is characterized by a steady turnover of taxa, some of which are restricted to the transition phase.

We demonstrate that the current grass/woody ratio is exceptional for the last 170 years and that such a process is related to gradual changes in soil moisture, erodibility, and species composition and competitiveness. Our inferences largely support prior theoretical considerations about shrub encroachment dynamics in southern African savannas that were mostly based on space-for-time approaches or studies with low taxonomic and temporal resolution.

Our study shows, for the first time, the potential of *sedaDNA* analysis to track vegetation changes in savanna environments, as well as to infer related local and global triggers and drivers. In particular, detailed taxonomical information enabled us to confirm and complement our pollen findings, and thus to reconstruct the turnover from broad-leaved to fine-leaved tree taxa and *Terminalia*.

Taking into account an earlier pollen record of Lake Otjikoto (Scott et al., 1991), we suggest that shrub encroachment is not a new phenomenon and that it possibly occurred in the past in the Tsumeb region, as suggested by changes in taxonomic composition and the grass/woody ratio observed in the previous record. In this sense, the encroached state observed in our study can be interpreted as a successive phase in a context of state transitions and is in line with previous studies that define encroachment as an alternative stable state (Gil-Romera et al., 2010).

Comparing our vegetation proxies to environmental proxies and historical data, we inferred the impact of logging, as well as physiological adaptations related to changes in precipitation regime and high atmospheric  $p\text{CO}_2$  concentrations. Furthermore, detailed information of grass taxonomical composition (which is restricted in the pollen record) made it possible to infer changes in land use, such as crop cultivation, selective grazing, and overgrazing. Combined with

our macrocharcoal record, we deduce changes in fire management associated with the intensification of farming.

Our study helps fill a knowledge gap about the feedback mechanisms supporting shrub encroachment. Such mechanisms reflect the interplay between historical land use and climate change (Perring et al., 2018), enabling us to confirm the far-reaching effects of management legacies on vegetation in savanna ecosystems.

Overall, this study highlights the suitability of multiple lake sedimentary proxies (particularly *sedaDNA*) to resolve complex compositional change and related environmental conditions in semiarid savannas on at least decadal and centennial timescales.

## ACKNOWLEDGMENTS

We are very grateful to Johan Le Roux and Chris Steenkamp from Dantica diving club Windhoek for their assistance in the field. We also thank Sefine Kilinc, Rosemary Shikangalah, Myron Eberle, Sarah Bala, Rabja Popall, and Anke Müllenmeister-Sawall for the laboratory work. Cathy Jenks is thanked for proofreading. We thank the reviewers for the very helpful remarks. The research was funded by the German Federal Ministry of Education and Research (BMBF) within the framework of the SPACES Project OPTIMASS (FKZ:01LL1302A) and the Initiative and Networking Fund of the Helmholtz Association.

## CONFLICT OF INTEREST

None declared.

## AUTHOR'S CONTRIBUTIONS

XT and UH conceived the ideas and designed methodology; XT and LD performed fossil pollen and macrocharcoal analyses; HZ and XT performed *sedaDNA* analyses; XT and ED performed grain-size and end-member modeling analysis; LB, AV, HW, and XT performed biogeochemical and sedimentological analyses. Under supervision of UH, XT wrote a first version of the manuscript that all coauthors commented on. All authors gave final approval for publication.

## DATA AVAILABILITY STATEMENT

The data that support the findings of this study are openly available in PANGAEA repository: Tabares, X., Zimmermann, H., Dietze, E., Ratzmann, G., Belz, L., Vieth-Hillebrand, A., Dupont, L., Wilkes, H., Mapani, B., Herzs Schuh, U. (2019): Palynology, sedimentary ancient DNA, biomarkers, macrocharcoal, and sedimentological data from Lake Otjikoto, Namibia. PANGAEA, <https://doi.org/10.1594/PANGAEA.902988>.


## ORCID

Ximena Tabares  <https://orcid.org/0000-0002-9866-3843>

Heike Zimmermann  <https://orcid.org/0000-0002-0225-5176>

Elisabeth Dietze  <https://orcid.org/0000-0003-4817-8441>

Andrea Vieth-Hillebrand  <https://orcid.org/0000-0002-3688-662X>

Lydie Dupont  <https://orcid.org/0000-0001-9531-6793>

Benjamin Mapani  <https://orcid.org/0000-0002-5354-3397>

Ulrike Herzs Schuh  <https://orcid.org/0000-0003-0999-1261>

## REFERENCES

- Aleman, J. C., Blarquez, O., & Staver, C. A. (2016). Land-use change outweighs projected effects of changing rainfall on tree cover in sub-Saharan Africa. *Global Change Biology*, 22(9), 3013–3025. <https://doi.org/10.1111/gcb.13299>
- Ansley, R. J., Boutton, T. W., & Jacoby, P. W. (2014). Root biomass and distribution patterns in a semi-arid mesquite savanna: Responses to long-term rainfall manipulation. *Rangeland Ecology and Management*, 67(2), 206–218. <https://doi.org/10.2111/REM-D-13-00119.1>
- Appleby, P. G. (2001). Chronostratigraphic techniques in recent sediments. In W. M. Last, & J. P. Smol (Eds.), *Tracking environmental change using lake sediments, vol. 1: Basin analysis, coring, and chronological techniques* (pp. 171–203). Dordrecht: Kluwer Academic Publishers. [https://doi.org/10.1007/0-306-47669-X\\_9](https://doi.org/10.1007/0-306-47669-X_9)
- Appleby, P. G., Nolan, P. J., Gifford, D. W., Godfrey, M. J., Oldfield, F., Anderson, N. J., & Battarbee, R. W. (1986).  $^{210}\text{Pb}$  dating by low background gamma counting. *Hydrobiologia*, 143(1), 21–27. <https://doi.org/10.1007/BF00026640>
- Appleby, P. G., & Oldfield, F. (1978). The calculation of lead-210 dates assuming a constant rate of supply of unsupported  $^{210}\text{Pb}$  to the sediment. *Catena*, 5(1), 1–8. [https://doi.org/10.1016/S0341-8162\(78\)80002-2](https://doi.org/10.1016/S0341-8162(78)80002-2)
- Badewien, T., Vogts, A., Dupont, L., & Rullkötter, J. (2015). Influence of Late Pleistocene and Holocene climate on vegetation distributions in southwest Africa elucidated from sedimentary n-alkanes - differences between 12°S and 20°S. *Quaternary Science Reviews*, 125, 160–171. <https://doi.org/10.1016/j.quascirev.2015.08.004>
- Badewien, T., Vogts, A., & Rullkötter, J. (2015). n-Alkane distribution and carbon stable isotope composition in leaf waxes of C3 and C4 plants from Angola. *Organic Geochemistry*, 89–90, 71–79. <https://doi.org/10.1016/j.orggeochem.2015.09.002>
- Banks, H., & Lewis, G. (2018). Phylogenetically informative pollen structures of “caesalpinoid” pollen (Caesalpinioideae, Cercidoideae, Detarioideae, Dialioideae and Duparquetioideae: Fabaceae). *Botanical Journal of the Linnean Society*, 187, 59–86. <https://doi.org/10.1093/botlinnean/boy005>
- Beckett, H., & Bond, W. J. (2019). Fire refugia facilitate forest and savanna co-existence as alternative stable states. *Journal of Biogeography*, 46(12), 2800–2810. <https://doi.org/10.1111/jbi.13707>
- Bellasio, C., Quirk, J., & Beerling, D. J. (2018). Stomatal and non-stomatal limitations in savanna trees and C4 grasses grown at low, ambient and high atmospheric CO<sub>2</sub>. *Plant Science*, 274, 181–192. <https://doi.org/10.1016/j.plantsci.2018.05.028>
- Berry, R. S., & Kulmatiski, A. (2017). A savanna response to precipitation intensity. *PLoS ONE*, 12(4), e0175402. <https://doi.org/10.1371/journal.pone.0175402>
- Boessenkool, S., Mcglyn, G., Epp, L. S., Taylor, D., Pimentel, M., Gizaw, A., ... Popp, M. (2014). Use of ancient sedimentary DNA as a novel conservation tool for high-altitude tropical biodiversity. *Conservation Biology*, 28(2), 446–455. <https://doi.org/10.1111/cobi.12195>
- Bonnefille, R., & Riollet, G. (1980). *Pollen des savanes d'Afrique orientale*. Paris: Centre National de la Recherche Scientifique.
- Bremond, L. (2011). *Pollen Blade Reference Collection (ISEM UM)*. Retrieved from <https://data.oreme.org/observation/pollen>
- Bremond, L., Favier, C., Ficetola, G. F., Tossou, M. G., Akouégninou, A., Gielly, L., ... Salzmann, U. (2017). Five thousand years of tropical lake sediment DNA records from Benin. *Quaternary Science Reviews*, 170, 203–211. <https://doi.org/10.1016/j.quascirev.2017.06.025>
- Brienen, R. J. W., Wanek, W., & Hietz, P. (2011). Stable carbon isotopes in tree rings indicate improved water use efficiency and drought responses of a tropical dry forest tree species. *Trees*, 25(1), 103–113. <https://doi.org/10.1007/s00468-010-0474-1>
- Brown, J. R., & Archer, S. (1999). Shrub invasion of grassland: Recruitment is continuous and not regulated by herbaceous biomass or density. *Ecology*, 80(7), 2385–2396. <https://doi.org/10.2307/176918>
- Buitenwerf, R., Bond, W. J., Stevens, N., & Trollope, W. S. W. (2012). Increased tree densities in South African savannas: >50 years of data suggests CO<sub>2</sub> as a driver. *Global Change Biology*, 18(2), 675–684. <https://doi.org/10.1111/j.1365-2486.2011.02561.x>
- Bullard, J. E., Ockelford, A., Strong, C., & Aubault, H. (2018). Effects of cyanobacterial soil crusts on surface roughness and splash erosion. *Journal of Geophysical Research: Biogeosciences*, 123(12), 3697–3712. <https://doi.org/10.1029/2018JG004726>
- Case, M. F., & Staver, A. C. (2017). Fire prevents woody encroachment only at higher-than-historical frequencies in a South African savanna. *Journal of Applied Ecology*, 54(3), 955–962. <https://doi.org/10.1111/1365-2664.12805>
- Cech, P. G., Edwards, P. J., & Olde Venterink, H. (2010). Why is abundance of herbaceous legumes low in African savanna? A test with two model species. *Biotropica*, 42(5), 580–589. <https://doi.org/10.1111/j.1744-7429.2009.00622.x>
- Clark, J. S. (1988). Particle motion and the theory of charcoal analysis: Source area, transport, deposition, and sampling. *Quaternary Research*, 30(01), 67–80. [https://doi.org/10.1016/0033-5894\(88\)90088-9](https://doi.org/10.1016/0033-5894(88)90088-9)
- Colombaroli, D., van der Plas, G., Rucina, S., & Verschuren, D. (2018). Determinants of savanna-fire dynamics in the eastern Lake Victoria catchment (western Kenya) during the last 1200 years. *Quaternary International*, 488, 67–80. <https://doi.org/10.1016/j.quaint.2016.06.028>
- Cunningham, P. L. (1996). *Prospects for sustained harvesting of Mopane (Colophospermum mopane) on the Venetia Limpopo Nature Reserve and its implications for browsing ungulates*. MSc thesis, Stellenbosch University.
- Da Silveira Lobo Sternberg, L. (1988). D/H ratios of environmental water recorded by D/H ratios of plant lipids. *Nature*, 333(6168), 59–61. <https://doi.org/10.1038/333059a0>
- De Deurwaerder, H., Hervé-Fernández, P., Stahl, C., Burban, B., Petronelli, P., Hoffman, B., ... Verbeeck, H. (2018). Liana and tree below-ground water competition—evidence for water resource partitioning during the dry season. *Tree Physiology*, 38(7), 1071–1083. <https://doi.org/10.1093/treephys/tpy002>
- De Dios, V. R., Weltzin, J. F., Sun, W., Huxman, T. E., & Williams, D. G. (2014). Transitions from grassland to savanna under drought through passive facilitation by grasses. *Journal of Vegetation Science*, 25(4), 937–946. <https://doi.org/10.1111/jvs.12164>
- De Klerk, J. N. (2004). *Bush encroachment in Namibia: Report on Phase 1 of the Bush Encroachment Research, Monitoring, and Management Project*. Windhoek.
- Devine, A. P., McDonald, R. A., Quaipe, T., & Maclean, I. M. D. (2017). Determinants of woody encroachment and cover in African savannas. *Oecologia*, 183(4), 939–951. <https://doi.org/10.1007/s00442-017-3807-6>
- Dietre, B., Gauthier, É., & Gillet, F. (2012). Modern pollen rain and fungal spore assemblages from pasture woodlands around Lake Saint-Point (France). *Review of Palaeobotany and Palynology*, 186, 69–89. <https://doi.org/10.1016/j.revpalbo.2012.07.002>
- Dietze, E., Hartmann, K., Diekmann, B., Ijmker, J., Lehmkuhl, F., Opitz, S., ... Borchers, A. (2012). An end-member algorithm for deciphering modern detrital processes from lake sediments of Lake Donggi Cona, NE Tibetan Plateau, China. *Sedimentary Geology*, 243–244, 169–180. <https://doi.org/10.1016/j.sedgeo.2011.09.014>
- Dietze, E., Maussion, F., Ahlborn, M., Diekmann, B., Hartmann, K., Henkel, K., ... Haberzettl, T. (2014). Sediment transport processes across the Tibetan Plateau inferred from robust grain-size end members in lake sediments. *Climate of the Past*, 10(1), 91–106. <https://doi.org/10.5194/cp-10-91-2014>
- Dinter, K. (1909). *Deutsch-Südwest-Afrika Flora, Forst und landwirtschaftliche Fragmente*. Leipzig: Theodor Oswald Weigel.
- Dinter, K. (1918). *Botanische Reisen in Deutsch-Südwest-Afrika. Repertorium Specierum Novarum Regni Vegetabilis*, 3, 1–169.

- Dinter, K. (1922). Index der aus Deutsch-Südwestafrika bis zum Jahre 1917 bekannt gewordenen Pflanzenarten X. *Repertorium Specierum Novarum Regni Vegetabilis*, 18, 13–16. <https://doi.org/10.1002/fedr.19220180106>
- Directorate of Planning (2005). *Agricultural statistics bulletin*. Windhoek: Directorate of Planning.
- Directorate of Planning (2011). *Agricultural statistics bulletin*. Windhoek: Directorate of Planning.
- Dougill, A. J., & Thomas, A. D. (2004). Kalahari sand soils: Spatial heterogeneity, biological soil crusts and land degradation. *Land Degradation and Development*, 15(3), 233–242. <https://doi.org/10.1002/ldr.611>
- Duffin, K. I., Gillson, L., & Willis, K. J. (2008). Testing the sensitivity of charcoal as an indicator of fire events in savanna environments: Quantitative predictions of fire proximity, area and intensity. *Holocene*, 18(2), 279–291. <https://doi.org/10.1177/0959683607086766>
- Ehleringer, J.R., & Dawson, T.E. (1992). Water uptake by plants: Perspectives from stable isotope composition. *Plant, Cell and Environment*, 15(9), 1073–1082. <https://doi.org/10.1111/j.1365-3040.1992.tb01657.x>
- Erkkilä, A. (2001). *Living on the land: Change in forest cover in North-Central Namibia 1943–1996*. Joensuu: University of Joensuu.
- Erkkilä, A., & Siiskonen, H. (1992). *Forestry in Namibia 1850–1990*. Joensuu: University of Joensuu.
- Evans, J. R. (1989). Photosynthesis and nitrogen relationships in leaves of C3 plants. *Oecologia*, 78(1), 9–19. <https://doi.org/10.1007/BF00377192>
- Evans, J. R., Schortemeyer, M., McFarlane, N., & Atkin, O. K. (2000). Photosynthetic characteristics of 10 *Acacia* species grown under ambient and elevated atmospheric CO<sub>2</sub>. *Functional Plant Biology*, 27(1), 13–25. <https://doi.org/10.1071/PP99126>
- Faegri, K., & Iversen, J. (1989). *Textbook of pollen analysis*. In K. Faegri, P. Kaland, & K. Krzywinski (Eds.) (4th ed.). Chichester: John Wiley & Sons.
- FAO (2018a). *FAOSTAT agri-environmental indicators - country Indicators*. Retrieved from <http://www.fao.org/faostat/en/#country/147>
- FAO (2018b). *FAOSTAT agri-environmental indicators - livestock patterns*. Retrieved from <http://www.fao.org/faostat/en/#data/EK>
- Feakins, S. J., & Sessions, A. L. (2010). Controls on the D/H ratios of plant leaf waxes in an arid ecosystem. *Geochimica Et Cosmochimica Acta*, 74(7), 2128–2141. <https://doi.org/10.1016/j.gca.2010.01.016>
- Ficetola, G. F., Pansu, J., Bonin, A., Coissac, E., Giguet-Covex, C., De Barba, M., ... Taberlet, P. (2015). Replication levels, false presences and the estimation of the presence/absence from eDNA metabarcoding data. *Molecular Ecology Resources*, 15(3), 543–556. <https://doi.org/10.1111/1755-0998.12338>
- Gandiwa, E., Gandiwa, P., & Mxoxa, T. (2012). Structure and composition of *Spirostachys africana* woodland stands in Gonarezhou National Park, southern Zimbabwe. *International Journal of Environmental Sciences*, 2(4), 2076–2089. <https://doi.org/10.6088/ijes.00202030091>
- Garcin, Y., Deschamps, P., Ménot, G., de Saulieu, G., Schefuβ, E., Sebag, D., ... Sachse, D. (2018). Early anthropogenic impact on Western Central African rainforests 2,600 y ago. *Proceedings of the National Academy of Sciences*, 115(13), 3261–3266. <https://doi.org/10.1073/pnas.1715336115>
- Gelorini, V., Ssemmanda, I., & Verschuren, D. (2012). Validation of non-pollen palynomorphs as paleoenvironmental indicators in tropical Africa: Contrasting ~200-year paleolimnological records of climate change and human impact. *Review of Palaeobotany and Palynology*, 186, 90–101. <https://doi.org/10.1016/j.revpalbo.2012.05.006>
- Gelorini, V., Verbeke, A., van Geel, B., Cocquyt, C., & Verschuren, D. (2011). Modern non-pollen palynomorphs from East African lake sediments. *Review of Palaeobotany and Palynology*, 164(3–4), 143–173. <https://doi.org/10.1016/j.revpalbo.2010.12.002>
- Giess, W. (1971). A preliminary vegetation map of South West Africa. *Dinteria*, 4, 5–114.
- Gillson, L. (2004). Testing non-equilibrium theories in savannas: 1400 years of vegetation change in Tsavo National Park, Kenya. *Ecological Complexity*, 1(4), 281–298. <https://doi.org/10.1016/j.ecocom.2004.06.001>
- Gillson, L., & Ekblom, A. (2009). Untangling anthropogenic and climatic influence on riverine forest in the Kruger National Park. *South Africa. Vegetation History and Archaeobotany*, 18(2), 171–185. <https://doi.org/10.1007/s00334-008-0202-6>
- Gilo, B. N., & Kelkay, T. Z. (2017). Changes in vegetation structure and aboveground biomass in response to traditional rangeland management practices in Borana, southern Ethiopia. *African Journal of Range and Forage Science*, 34(1), 21–31. <https://doi.org/10.2989/10220119.2017.1331934>
- Gil-Romera, G., Lamb, H. F., Turton, D., Sevilla-Callejo, M., & Umer, M. (2010). Long-term resilience, bush encroachment patterns and local knowledge in a Northeast African savanna. *Global Environmental Change*, 20(4), 612–626. <https://doi.org/10.1016/j.gloenvcha.2010.04.008>
- Gosling, W. D., Miller, C. S., & Livingstone, D. A. (2013). Atlas of the tropical West African pollen flora. *Review of Palaeobotany and Palynology*, 199, 1–135. <https://doi.org/10.1016/j.revpalbo.2013.01.003>
- Harris, I., Jones, P. D., Osborn, T. J., & Lister, D. H. (2014). Updated high-resolution grids of monthly climatic observations - the CRU TS3.10 dataset. *International Journal of Climatology*, 34(3), 623–642. <https://doi.org/10.1002/joc.3711>
- Higgins, S. I., & Scheiter, S. (2012). Atmospheric CO<sub>2</sub> forces abrupt vegetation shifts locally, but not globally. *Nature*, 488(7410), 209–212. <https://doi.org/10.1038/nature11238>
- Hirota, M., Holmgren, M., Van Nes, E. H., & Scheffer, M. (2011). Global resilience of tropical forest and savanna to critical transitions. *Science*, 334(6053), 232–235. <https://doi.org/10.1126/science.1210657>
- Hoffman, M. T., Rohde, R. F., & Gillson, L. (2019). Rethinking catastrophe? Historical trajectories and modelled future vegetation change in southern Africa. *Anthropocene*, 25, 100189. <https://doi.org/10.1016/j.ancene.2018.12.003>
- Hoffmann, B., Kahmen, A., Cernusak, L. A., Arndt, S. K., & Sachse, D. (2013). Abundance and distribution of leaf wax n-alkanes in leaves of *Acacia* and *Eucalyptus* trees along a strong humidity gradient in northern Australia. *Organic Geochemistry*, 62, 62–67. <https://doi.org/10.1016/j.orggeochem.2013.07.003>
- Holling, C. S. (1973). Resilience and stability of ecological systems. *Annual Review of Ecology and Systematics*, 4(1), 1–23. <https://doi.org/10.1146/annurev.es.04.110173.000245>
- Hou, J., D'Andrea, W. J., & Huang, Y. (2008). Can sedimentary leaf waxes record D/H ratios of continental precipitation? Field, model, and experimental assessments. *Geochimica Et Cosmochimica Acta*, 72(14), 3503–3517. <https://doi.org/10.1016/j.gca.2008.04.030>
- Hüttich, C., Gessner, U., Herold, M., Strohbach, B. J., Schmidt, M., Keil, M., & Dech, S. (2009). On the suitability of MODIS time series metrics to map vegetation types in dry savanna ecosystems: A case study in the Kalahari of NE Namibia. *Remote Sensing*, 1(4), 620–643. <https://doi.org/10.3390/rs1040620>
- Jordaan, J. J. (2017). Controlling *Hyperthelia dissoluta* (Nees ex. Steud.) Clayton (Yellow thatch grass) through defoliation in southern African rangelands. *African Journal of Ecology*, 55(4), 742–747. <https://doi.org/10.1111/aje.12395>
- Joubert, D. F., Rothauge, A., & Smit, G. N. (2008). A conceptual model of vegetation dynamics in the semiarid Highland savanna of Namibia, with particular reference to bush thickening by *Acacia mellifera*. *Journal of Arid Environments*, 72(12), 2201–2210. <https://doi.org/10.1016/j.jaridenv.2008.07.004>
- Joubert, D. F., Smit, G. N., & Hoffman, M. T. (2012). The role of fire in preventing transitions from a grass dominated state to a bush thickened

- state in arid savannas. *Journal of Arid Environments*, 87, 1–7. <https://doi.org/10.1016/j.jaridenv.2012.06.012>
- Joubert, D. F., Smit, G. N., & Hoffman, M. T. (2013). The influence of rainfall, competition and predation on seed production, germination and establishment of an encroaching *Acacia* in an arid Namibian savanna. *Journal of Arid Environments*, 91, 7–13. <https://doi.org/10.1016/j.jaridenv.2012.11.001>
- Kamona, A. F., & Günzel, A. (2007). Stratigraphy and base metal mineralization in the Otavi Mountain Land, Northern Namibia - a review and regional interpretation. *Gondwana Research*, 11(3), 396–413. <https://doi.org/10.1016/j.gr.2006.04.014>
- Kanyama, C. (2017). *A study of isotopic composition of xylem water of woody vegetation and groundwater along a precipitation gradient in Namibia*. MSc thesis. University of Namibia.
- Kgosikoma, O. E., Mojeremane, W., & Harvie, B. (2015). The impact of livestock grazing management systems on soil and vegetation characteristics across savanna ecosystems in Botswana. *African Journal of Range and Forage Science*, 32(4), 271–278. <https://doi.org/10.2989/10220119.2015.1008042>
- Kidron, G. J., Ying, W., Starinsky, A., & Herzberg, M. (2017). Drought effect on biocrust resilience: High-speed winds result in crust burial and crust rupture and flaking. *Science of the Total Environment*, 579, 848–859. <https://doi.org/10.1016/j.scitotenv.2016.11.016>
- Krishnamurthy, R. V., & Machavaram, M. (2000). Is there a stable isotope evidence for the CO<sub>2</sub> fertilization effect? *Journal of Earth System Science*, 109(1), 141–144. <https://doi.org/10.1007/BF02719157>
- Kristen, I., Fuhrmann, A., Thorpe, J., Rohl, U., Wilkes, H., & Oberhänsli, H. (2007). Hydrological changes in southern Africa over the last 200 ka as recorded in lake sediments from the Tswaing impact crater. *South African Journal of Geology*, 110(2–3), 311–326. <https://doi.org/10.2113/gssajg.110.2-3.311>
- Kristen, I., Wilkes, H., Vieth, A., Zink, K.-G., Plessen, B., Thorpe, J., ... Oberhänsli, H. (2010). Biomarker and stable carbon isotope analyses of sedimentary organic matter from Lake Tswaing: Evidence for deglacial wetness and early Holocene drought from South Africa. *Journal of Paleolimnology*, 44(1), 143–160. <https://doi.org/10.1007/s10933-009-9393-9>
- Kyalangalilwa, B., Boatwright, J. S., Daru, B. H., Maurin, O., & van der Bank, M. (2013). Phylogenetic position and revised classification of *Acacia* s.l. (Fabaceae: Mimosoideae) in Africa, including new combinations in *Vachellia* and *Senegalia*. *Botanical Journal of the Linnean Society*, 172(4), 500–523. <https://doi.org/10.1111/boj.12047>
- Lange, G. M., Barners, J. I., & Motinga, D. J. (1998). Cattle numbers, biomass, productivity and land degradation in the commercial farming sector of Namibia, 1915–95. *Development Southern Africa*, 15(4), 555–572. <https://doi.org/10.1080/03768359808440031>
- Lau, B., & Reiner, P. (1993). *100 years of agricultural development in colonial Namibia: A historical overview of visions and experiments*. Windhoek: National Archives of Namibia.
- Lezine, A. M. (2005). African pollen database. Retrieved from <http://apd.sedoo.fr/accueil.htm>
- Li, B.-L. (2002). A theoretical framework of ecological phase transitions for characterizing tree-grass dynamics. *Acta Biotheoretica*, 50, 141–154. <https://doi.org/10.1023/a:1016562208170>
- Liao, C., Clark, P. E., & DeGloria, S. D. (2018). Bush encroachment dynamics and rangeland management implications in southern Ethiopia. *Ecology and Evolution*, 8(23), 11694–11703. <https://doi.org/10.1002/ece3.4621>
- López-Olmedo, L. I., Meave, J. A., & Pérez-García, E. A. (2007). Floristic and structural contrasts between natural savannas and anthropogenic pastures in a tropical dry landscape. *Rangeland Journal*, 29, 181–190. <https://doi.org/10.1071/RJ07007>
- Marchant, J. W. (1980). *Hydrogeochemical exploration at Tsumeb*. PhD thesis. University of Cape Town.
- Maron, J. L., & Connors, P. G. (1996). A native nitrogen-fixing shrub facilitates weed invasion. *Oecologia*, 105(3), 302–312. <https://doi.org/10.1007/BF00328732>
- Marzi, R., Torkelson, B. E., & Olson, R. K. (1993). A revised carbon preference index. *Organic Geochemistry*, 20(8), 1303–1306. [https://doi.org/10.1016/0146-6380\(93\)90016-5](https://doi.org/10.1016/0146-6380(93)90016-5)
- Mendelsohn, J., El Obeid, S., & Roberts, C. (2000). *A profile of North-Central Namibia*. Windhoek: Gamsberg Macmillan.
- Mendelsohn, J., Jarvis, A., Roberts, C., & Robertson, T. (2009). *Atlas of Namibia. A portrait of the land and its people* (3rd ed.). Cape Town: Sunbird Publishers.
- Metwally, A. A., Scott, L., Neumann, F. H., Bamford, M. K., & Oberhänsli, H. (2014). Holocene palynology and palaeoenvironments in the savanna biome at Tswaing crater, central South Africa. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 402, 125–135. <https://doi.org/10.1016/j.palaeo.2014.03.019>
- Meyer, K. M., Wiegand, K., Ward, D., & Moustakas, A. (2007). The rhythm of savanna patch dynamics. *Journal of Ecology*, 95(6), 1306–1315. <https://doi.org/10.1111/j.1365-2745.2007.01289.x>
- Meyers, P. A. (1997). Organic geochemical proxies of paleoceanographic, paleolimnologic, and paleoclimatic processes. *Organic Geochemistry*, 27(5–6), 213–250. [https://doi.org/10.1016/S0146-6380\(97\)00049-1](https://doi.org/10.1016/S0146-6380(97)00049-1)
- Mildbraed, J. (1941). Beitrage zur Flora von Deutsch-Sudwestafrika. *Notizblatt Des Königl. Botanischen Gartens Und Museums Zu Berlin*, 15(3), 448. <https://doi.org/10.2307/3995082>
- Miller, C., Finch, J., Hill, T., Peterse, F., Humphries, M., Zabel, M., & Schefuß, E. (2019). The drivers of late Quaternary climate variability in eastern South Africa. *Climate of the Past Discussions*, 1–34. <https://doi.org/10.5194/cp-2019-4>
- Miller, C. S., & Gosling, W. D. (2014). Quaternary forest associations in lowland tropical West Africa. *Quaternary Science Reviews*, 84, 7–25. <https://doi.org/10.1016/j.quascirev.2013.10.027>
- Mitchard, E. T. A., & Flintrop, C. M. (2013). Woody encroachment and forest degradation in sub-Saharan Africa's woodlands and savannas 1982–2006. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 1982–2006. <https://doi.org/10.1098/rstb.2012.0406>
- Mooney, S., & Tinner, W. (2011). The analysis of charcoal in peat and organic sediments. *Mires and Peat*, 7, 1–18.
- Moore, P., Webb, J., & Collison, M. (1991). *Pollen analysis*. Oxford: Wiley-Blackwell.
- Moyo, S., O'Keefe, P., & Sill, M. (1993). *The Southern African environment: Profiles of the SADC countries*. London: Routledge.
- Nackley, L. L., Betzelberger, A., Skowno, A., G. West, A., Ripley, B. S., Bond, W. J., & Midgley, G. F. (2018). CO<sub>2</sub> enrichment does not entirely ameliorate *Vachellia* karroo drought inhibition: A missing mechanism explaining savanna bush encroachment. *Environmental and Experimental Botany*, 155, 98–106. <https://doi.org/10.1016/j.envexpbot.2018.06.018>
- Neumann, F. H., Scott, L., Bousman, C. B., & van As, L. (2010). A Holocene sequence of vegetation change at Lake Eteza, coastal KwaZulu-Natal, South Africa. *Review of Palaeobotany and Palynology*, 162(1), 39–53. <https://doi.org/10.1016/j.revpalbo.2010.05.001>
- Nicholson, S. E., & Selato, J. C. (2000). The influence of La Nina on African rainfall. *International Journal of Climatology*, 20(14), 1761–1776. [https://doi.org/10.1002/1097-0088\(20001130\)20:14<1761:AID-JOC580>3.0.CO;2-W](https://doi.org/10.1002/1097-0088(20001130)20:14<1761:AID-JOC580>3.0.CO;2-W)
- Niedermeyer, E. M., Forrest, M., Beckmann, B., Sessions, A. L., Mulch, A., & Schefuß, E. (2016). The stable hydrogen isotopic composition of sedimentary plant waxes as quantitative proxy for rainfall in the West African Sahel. *Geochimica Et Cosmochimica Acta*, 184, 55–70. <https://doi.org/10.1016/j.gca.2016.03.034>
- O'Connor, T. G. (1998). Impact of sustained drought on a semi-arid *Colophospermum mopane* savanna. *African Journal of Range and Forage Science*, 15(3), 83–91. <https://doi.org/10.1080/10220119.1998.9647948>



- O'Connor, T. G., & Pickett, G. A. (1992). The influence of grazing on seed production and seed banks of some African savanna grasslands. *The Journal of Applied Ecology*, 29(1), 247–260. <https://doi.org/10.2307/2404367>
- O'Connor, T. G., Puttick, J. R., & Hoffman, M. T. (2014). Bush encroachment in southern Africa: Changes and causes. *African Journal of Range and Forage Science*, 31(2), 67–88. <https://doi.org/10.2989/10220119.2014.939996>
- Perring, M. P., Bernhardt-Römermann, M., Baeten, L., Midolo, G., Blondeel, H., Depauw, L., ... Verheyen, K. (2018). Global environmental change effects on plant community composition trajectories depend upon management legacies. *Global Change Biology*, 24(4), 1722–1740. <https://doi.org/10.1111/gcb.14030>
- Poston, M. E., & Nowicke, J. W. (1993). Pollen morphology, Trichome types, and relationships of the Gronovioideae (Loasaceae). *American Journal of Botany*, 80(6), 689. <https://doi.org/10.2307/2445439>
- Poynter, J. G. (1989). *Molecular stratigraphy: The recognition of palaeoclimatic signals in organic geochemical data*. PhD thesis, University of Bristol.
- Quick, L. J., Chase, B. M., Wündsche, M., Kirsten, K. L., Chevalier, M., Mäusbacher, R., ... Haberzettl, T. (2018). A high-resolution record of Holocene climate and vegetation dynamics from the southern Cape coast of South Africa: Pollen and microcharcoal evidence from Eilandvlei. *Journal of Quaternary Science*, 33(5), 487–500. <https://doi.org/10.1002/jqs.3028>
- Ravi, S., D'Odorico, P., Wang, L., White, C. S., Okin, G. S., Macko, S. A., & Collins, S. L. (2009). Post-fire resource redistribution in desert grasslands: A possible negative feedback on land degradation. *Ecosystems*, 12(3), 434–444. <https://doi.org/10.1007/s10021-009-9233-9>
- Revelles, J., & van Geel, B. (2016). Human impact and ecological changes in lakeshore environments. The contribution of non-pollen palynomorphs in Lake Banyoles (NE Iberia). *Review of Palaeobotany and Palynology*, 232, 81–97. <https://doi.org/10.1016/j.revpalbo.2016.05.004>
- Rodríguez-Caballero, E., Cantón, Y., Chamizo, S., Afana, A., & Solé-Benet, A. (2012). Effects of biological soil crusts on surface roughness and implications for runoff and erosion. *Geomorphology*, 145–146, 81–89. <https://doi.org/10.1016/j.geomorph.2011.12.042>
- Rohde, R. F., & Hoffman, M. T. (2012). The historical ecology of Namibian rangelands: Vegetation change since 1876 in response to local and global drivers. *Science of the Total Environment*, 416, 276–288. <https://doi.org/10.1016/j.scitotenv.2011.10.067>
- Roques, K. G., O'Connor, T. G., & Watkinson, A. R. (2001). Dynamics of shrub encroachment in an African savanna: Relative influences of fire, herbivory, rainfall and density dependence. *Journal of Applied Ecology*, 38(2), 268–280. <https://doi.org/10.1046/j.1365-2664.2001.00567.x>
- Sachse, D., Billault, I., Bowen, G. J., Chikaraishi, Y., Dawson, T. E., Feakins, S. J., ... Kahmen, A. (2012). Molecular paleohydrology: interpreting the hydrogen-isotopic composition of lipid biomarkers from photosynthesizing organisms. *Annual Review of Earth and Planetary Sciences*, 40(1), 221–249. <https://doi.org/10.1146/annurev-earth-042711-105535>
- Saha, M. V., Scanlon, T. M., & D'Odorico, P. (2015). Examining the linkage between shrub encroachment and recent greening in water-limited southern Africa. *Ecosphere*, 6(9), 1–16. <https://doi.org/10.1890/ES15-00098.1>
- Salles, C., Poesen, J., & Govers, G. (2000). Statistical and physical analysis of soil detachment by raindrop impact: Rain erosivity indices and threshold energy. *Water Resources Research*, 36(9), 2721–2729. <https://doi.org/10.1029/2000WR900024>
- Scheffer, M., Carpenter, S., Foley, J. A., Folke, C., & Walker, B. (2001). Catastrophic shifts in ecosystems. *Nature*, 413(6856), 591–596. <https://doi.org/10.1038/35098000>
- Schneider, H. P. (2012). The history of veterinary medicine in Namibia. *Journal of the South African Veterinary Association*, 83(1), a4. <https://doi.org/10.4102/jsava.v83i1.4>
- Schüler, L., & Hemp, A. (2016). Atlas of pollen and spores and their parent taxa of Mt Kilimanjaro and tropical East Africa. *Quaternary International*, 425, 301–386. <https://doi.org/10.1016/j.quaint.2016.07.038>
- Scott, L. (1982). Late Quaternary fossil pollen grains from the Transvaal, South Africa. *Review of Palaeobotany and Palynology*, 36(3–4), 241–278. [https://doi.org/10.1016/0034-6667\(82\)90022-7](https://doi.org/10.1016/0034-6667(82)90022-7)
- Scott, L., Cooremans, B., de Wet, J. S., & Vogel, J. C. (1991). Holocene environmental changes in Namibia inferred from pollen analysis of swamp and lake deposits. *Holocene*, 1(1), 8–13. <https://doi.org/10.1177/095968369100100103>
- Smit, G. N. (2003). The importance of ecosystem dynamics in managing the bush encroachment problem in southern Africa (pp. 14–22). In *Proceedings of the Seventh International Rangelands Congress*, Durban.
- Staver, A. C., Archibald, S., & Levin, S. A. (2011). The global extent and determinants of savanna and forest as alternative biome states. *Science*, 334(6053), 230–232. <https://doi.org/10.1126/science.1210465>
- Stevens, N., Lehmann, C. E. R., Murphy, B. P., & Durigan, G. (2017). Savanna woody encroachment is widespread across three continents. *Global Change Biology*, 23(1), 235–244. <https://doi.org/10.1111/gcb.13409>
- Stockmarr, J. (1971). Tablets with spores used in absolute pollen analysis. *Pollen Et Spores*, 13, 615–621.
- Strohbach, B. J. (2001). Vegetation degradation in Namibia. *Journal of Namibia Scientific Society*, 48, 127–156.
- Strohbach, B. J. (2014). Vegetation of the eastern communal conservancies in Namibia: I. Phytosociological descriptions. *Koedoe*, 56(1), 1–18. <https://doi.org/10.4102/koedoe.v56i1.1116>
- Strohbach, B. J., & Kutuahuripa, J. T. (2014). Vegetation of the eastern communal conservancies in Namibia: II. Environmental drivers. *Koedoe*, 56(1), a1117. <https://doi.org/10.4102/koedoe.v56i1.1117>
- Sweet, J. (1998). "Livestock: Coping with Drought". Namibia: FAO/AGAP. [http://www.fao.org/documents/pub\\_det.asp?xml:lang=en&pub\\_xml:id=154104](http://www.fao.org/documents/pub_det.asp?xml:lang=en&pub_xml:id=154104)
- Sweet, J., & Burke, A. (2006). *Country pasture/forage resource profiles*. Namibia: FAO.
- Tabares, X., Mapani, B., Blaum, N., & Herzs Schuh, U. (2018). Composition and diversity of vegetation and pollen spectra along gradients of grazing intensity and precipitation in southern Africa. *Review of Palaeobotany and Palynology*, 253, 88–100. <https://doi.org/10.1016/j.revpalbo.2018.04.004>
- Taberlet, P., Coissac, E., Pompanon, F., Gielly, L., Miquel, C., Valentini, A., ... Willerslev, E. (2007). Power and limitations of the chloroplast trnL (UAA) intron for plant DNA barcoding. *Nucleic Acids Research*, 35(3), e14. <https://doi.org/10.1093/nar/gkl938>
- Talbot, M. R. (2005). Nitrogen isotopes in palaeolimnology. In W. M. Last & J. P. Smol (Eds.), *Tracking environmental change using lake sediments* (pp. 401–439). Dordrecht: Kluwer Academic Publishers. [https://doi.org/10.1007/0-306-47670-3\\_15](https://doi.org/10.1007/0-306-47670-3_15)
- Tessema, Z. K., de Boer, W. F., & Prins, H. H. T. (2016). Changes in grass plant populations and temporal soil seed bank dynamics in a semi-arid African savanna: Implications for restoration. *Journal of Environmental Management*, 182, 166–175. <https://doi.org/10.1016/j.jenvman.2016.07.057>
- Thomas, A. D., & Dougill, A. J. (2007). Spatial and temporal distribution of cyanobacterial soil crusts in the Kalahari: Implications for soil surface properties. *Geomorphology*, 85(1–2), 17–29. <https://doi.org/10.1016/j.geomorph.2006.03.029>
- Tian, F., Brandt, M., Liu, Y. Y., Rasmussen, K., & Fensholt, R. (2017). Mapping gains and losses in woody vegetation across global tropical drylands. *Global Change Biology*, 23(4), 1748–1760. <https://doi.org/10.1111/gcb.13464>
- Tian, F., Herzs Schuh, U., Mischke, S., & Schlüt, F. (2014). What drives the recent intensified vegetation degradation in Mongolia – Climate change or human activity? *The Holocene*, 24(10), 1206–1215. <https://doi.org/10.1177/0959683614540958>

- Tolsma, D. J., Ernst, W. H. O., & Verwey, R. A. (1987). Nutrients in soil and vegetation around two artificial waterpoints in eastern Botswana. *The Journal of Applied Ecology*, 24(3), 991–1000. <https://doi.org/10.2307/2403995>
- Valentin, C. (1993). Soil crusting and sealing in West Africa and possible approaches to improved management. In Food and Agriculture Organization of the United Nations. Soil Resources, Management, and Conservation Service (Ed.), *Soil tillage in Africa: Needs and challenges* (pp. 95–128). Rome: FAO Soils Bulletin. <http://www.fao.org/3/T1696E/T1696e00.htm>
- van der Sleen, P., Groenendijk, P., Vlam, M., Anten, N. P. R., Boom, A., Bongers, F., ... Zuidema, P. A. (2014). No growth stimulation of tropical trees by 150 years of CO<sub>2</sub> fertilization but water-use efficiency increased. *Nature Geoscience*, 8(1), 24–28. <https://doi.org/10.1038/ngeo2313>
- van Geel, B., & Aptroot, A. (2006). Fossil ascomycetes in Quaternary deposits. *Nova Hedwigia*, 82(3), 313–329. <https://doi.org/10.1127/0029-5035/2006/0082-0313>
- van Geel, B., Gelorini, V., Lyaruu, A., Aptroot, A., Rucina, S., Marchant, R., ... Verschuren, D. (2011). Diversity and ecology of tropical African fungal spores from a 25,000-year palaeoenvironmental record in southeastern Kenya. *Review of Palaeobotany and Palynology*, 164(3–4), 174–190. <https://doi.org/10.1016/j.revpalbo.2011.01.002>
- van Oudtshoorn, F. (2016). Identification guide to Southern African grasses: An identification manual with keys, descriptions and distributions. *African Journal of Range & Forage Science*, 33(4), 279–280. <https://doi.org/10.2989/10220119.2016.1264482>
- van Rooyen, M. W., le Roux, A., van der Merwe, H., van Rooyen, N., & Geldenhuys, C. (2018). Long-term vegetation change (>20 years) in the plains habitat on the Goegap Nature Reserve, Succulent Karoo, South Africa. *African Journal of Range & Forage Science*, 35(3–4), 289–302. <https://doi.org/10.2989/10220119.2018.1498802>
- Venter, Z. S., Cramer, M. D., & Hawkins, H. J. (2018). Drivers of woody plant encroachment over Africa. *Nature Communications*, 9(1), <https://doi.org/10.1038/s41467-018-04616-8>
- Verlinden, A., & Kruger, A. S. (2007). Changing grazing systems in central north Namibia. *Land Degradation & Development*, 18(2), 179–197. <https://doi.org/10.1002/ldr.769>
- Vogts, A., Moossen, H., Rommerskirchen, F., & Rullkötter, J. (2009). Distribution patterns and stable carbon isotopic composition of alkanes and alkan-1-ols from plant waxes of African rain forest and savanna C3 species. *Organic Geochemistry*, 40(10), 1037–1054. <https://doi.org/10.1016/j.orggeochem.2009.07.011>
- Wagner, T. C., Richter, J., Joubert, D. F., & Fischer, C. (2018). A dominance shift in arid savanna: An herbaceous legume outcompetes local C4 grasses. *Ecology and Evolution*, 8(13), 6779–6787. <https://doi.org/10.1002/ece3.4188>
- Walther, S. C., & Neumann, F. H. (2011). Sedimentology, isotopes and palynology of late Holocene cores from Lake Sibaya and the Kosi Bay system (KwaZulu-Natal, South Africa). *South African Geographical Journal*, 93(2), 133–153. <https://doi.org/10.1080/03736245.2011.591982>
- Ward, D. (2005). Do we understand the causes of bush encroachment in African savannas? *African Journal of Range & Forage Science*, 22(2), 101–105. <https://doi.org/10.2989/10220110509485867>
- Ward, D., Hoffman, M. T., & Collocott, S. J. (2014). A century of woody plant encroachment in the dry Kimberley savanna of South Africa. *African Journal of Range and Forage Science*, 31(2), 107–121. <https://doi.org/10.2989/10220119.2014.914974>
- Wiegand, K., Ward, D., & Saltz, D. (2005). Multi-scale patterns and bush encroachment in an arid savanna with a shallow soil layer. *Journal of Vegetation Science*, 16(3), 311–320. <https://doi.org/10.1111/j.1654-1103.2005.tb02369.x>
- Wigley, B. J., Bond, W. J., & Hoffman, M. T. (2009). Bush encroachment under three contrasting land-use practices in a mesic South African savanna. *African Journal of Ecology*, 47(1), 62–70. <https://doi.org/10.1111/j.1365-2028.2008.01051.x>
- Zhang, W., Brandt, M., Penuelas, J., Guichard, F., Tong, X., Tian, F., & Fensholt, R. (2019). Ecosystem structural changes controlled by altered rainfall climatology in tropical savannas. *Nature Communications*, 10(671), <https://doi.org/10.1038/s41467-019-08602-6>
- Zimmermann, H. H., Raschke, E., Epp, L. S., Stoof-Leichsenring, K. R., Schwamborn, G., Schirrmeister, L., ... Herzschuh, U. (2017). Sedimentary ancient DNA and pollen reveal the composition of plant organic matter in Late Quaternary permafrost sediments of the Buor Khaya Peninsula (north-eastern Siberia). *Biogeosciences*, 14(3), 575–596. <https://doi.org/10.5194/bg-14-575-2017>

## SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

**How to cite this article:** Tabares X, Zimmermann H, Dietze E, et al. Vegetation state changes in the course of shrub encroachment in an African savanna since about 1850 CE and their potential drivers. *Ecol Evol*. 2020;10:962–979. <https://doi.org/10.1002/ece3.5955>

© 2020. This work is published under <http://creativecommons.org/licenses/by/4.0/>(the “License”). Notwithstanding the ProQuest Terms and Conditions, you may use this content in accordance with the terms of the License.