

KAT1 is not essential for stomatal opening

Alexander Szyroki^{*†}, Natalya Ivashikina^{†‡}, Petra Dietrich[‡], M. Rob G. Roelfsema[‡], Peter Ache[‡], Birgit Reintanz^{*}, Rosalia Deeken[‡], Matthias Godde^{*}, Hubert Felle[§], Ralf Steinmeyer[‡], Klaus Palme^{*}, and Rainer Hedrich^{*†¶}

^{*}Max-Delbrück-Laboratorium, Carl-von-Linné-Weg 10, 50829 Cologne, Germany; [‡]Julius-von-Sachs-Institut, Lehrstuhl für Molekulare Pflanzenphysiologie und Biophysik, Julius-von-Sachs-Platz 2, 97082 Würzburg, Germany; and [§]Botanisches Institut I, Senckenbergstrasse 17, 35392 Giessen, Germany

Communicated by Jozef S. Schell, Max Planck Institute for Plant Breeding Research, Cologne, Germany, December 26, 2000 (received for review December 10, 2000)

It is generally accepted that K⁺ uptake into guard cells via inward-rectifying K⁺ channels is required for stomatal opening. To test whether the guard cell K⁺ channel KAT1 is essential for stomatal opening, a knockout mutant, *KAT1::En-1*, was isolated from an *En-1* mutagenized *Arabidopsis thaliana* population. Stomatal action and K⁺ uptake, however, were not impaired in KAT1-deficient plants. Reverse transcription-PCR experiments with isolated guard cell protoplasts showed that in addition to KAT1, the K⁺ channels AKT1, AKT2/3, AtKC1, and KAT2 were expressed in this cell type. In impalement measurements, intact guard cells exhibited inward-rectifying K⁺ currents across the plasma membrane of both wild-type and *KAT1::En-1* plants. This study demonstrates that multiple K⁺ channel transcripts exist in guard cells and that KAT1 is not essential for stomatal action.

Guard cells represent the best characterized plant cell type with respect to ion transport and signal transduction. Opening of stomata in response to various stimuli such as light, low CO₂ concentrations, or the phytohormone auxin is supposed to require K⁺ uptake through inward-rectifying K⁺ channels in the plasma membrane (1–8). Likewise, stomatal closure caused by the wilting hormone abscisic acid (ABA), or high CO₂, is suggested to involve signal cascades that down-regulate the K⁺ inward-rectifier and promote ion efflux (9–13). Even though the inward-rectifying K⁺ channel KAT1 has been cloned (14), characterized in oocytes (15–18), and shown to express in guard cells (19), its impact for the functional K⁺ channel in the plasma membrane of guard cells is still unknown (20). To analyze the physiological role of the KAT1 channel *in planta*, we have isolated a mutant in the *KAT1* gene by using transposon mutagenesis (21). We show that disruption of the *KAT1* gene does not affect stomatal opening, although underlying potassium currents were altered.

Materials and Methods

Arabidopsis thaliana wild-type and *KAT1::En-1* plants were grown in a growth chamber or in the greenhouse under a dark/light regime of 16:8 h. Plants with fully developed leaves were used (growth period 5–8 weeks).

Reverse Genetic Screening and Reverse Transcription (RT)-PCR Experiments. For functional analysis of the *KAT1* gene, we screened for a knockout mutant in a collection of *Arabidopsis* plants mutagenized by the maize transposon *En-1* (21). In a reverse genetic approach (22) employing a combination of *KAT1*- and *En-1*-specific PCR-primers, we identified the mutant *KAT1::En-1* 7AAD31 (Fig. 1A). *KAT1*-primers were K1–5 (5'-TAG ACG CTG AGT ATT TCC CAC CAA A-3') and K1–4 (5'-TCC ATC AAC GTA GAC AGT GAA GTC C-3'). *En-1*-specific primers were En-205 (5'-AGA AGC ACG ACG GCT GTA GAA TAG GA-3') and En-8130 (5'-GAG CGT CGG TCC CCA CAC TTC TAT AC-3'). For RT-PCR experiments, guard cell protoplasts were isolated as described below and mRNA was purified twice with the Dynabeads mRNA Direct kit (Dyna, Oslo) to minimize DNA contaminations. First strand cDNA was prepared by using Superscript RT (GIBCO/BRL) and diluted for RT-PCR 20-fold in water. PCR was performed in a LightCycler (ROCHE) with

the LightCycler-FastStart DNA Master SYBR Green I Kit (ROCHE). The following K⁺ channel-specific primers were used: KAT1fwd (5'-ACT TCC GAC ACT GC-3'), KAT1rev (5'-CCC AAA TGA CAT CTA A-3'), KAT2fwd (5'-ATA TTG ATA TGG GGT CA-3'), KAT2rev (5'-ATC TAT TTC TGC GTT TT-3'), AKT1fwd (5'-CCA ACT GTT GCG TAT-3'), AKT1rev (5'-CTG CGT GGT ACT CC-3'), AKT2/3fwd (5'-AAA ATG GCG AAA ACA C-3'), AKT2/3rev (5'-CGC TGC TTC ACA TAG AA-3'), AKT5fwd (5'-AGG CCA CAG TTG TTC-3'), AKT5rev (5'-CGC CAT TTT CTG ATA A-3'), AKT6fwd (5'-GCC AGT GCG GTT AC-3'), AKT6rev (5'-GAC TCA ATC GCT TGG TA-3'), AtKC1fwd (5'-ATA TTG CGA TAC ACA AG-3'), AtKC1rev (5'-GAC CTA ACT TCG CTA AT-3'), GORKfwd (5'-CCT CCT TTA ATT TAG AAG-3'), GORKrev (5'-GCT CCA TCC GAT AG-3'), SKORfwd (5'-TGA CCC GAA TAA GAC AG-3'), and SKORrev (5'-TGT GTT TCC CCA TCT G-3'). The GenBank accession numbers are as follows: KAT1 (X93022), KAT2 (CAA16801), AKT1 (X62907), AKT2/3 (U40154/U44745), AKT5 (AJ249479), AKT6 (AC006053), AtKC1 (U81239), AtGORK (AJ279009), SKOR (AJ223358), and *Arabidopsis* actins (cf. ref. 23). cDNA quantities were calculated by using LIGHTCYCLER 3.1 (ROCHE). All quantifications were normalized to actin cDNA fragments amplified by ACTfwd (5'-GGT GAT GGT GTG TCT-3') and ACTrev (5'-ACT GAG CAC AAT GTT AC-3'). To enable detection of contaminating genomic DNA, the primers for KAT2, AKT1, AKT2/3, AKT5, AKT6, and GORK were selected to flank up to three introns. All kits were used according to the manufacturer's protocols.

Patch-Clamp Experiments. Guard cell protoplasts were isolated as described (24) with the enzyme solution adjusted to 560 mosmol kg⁻¹ by using D-Sorbitol. Current measurements were performed by using an EPC-7 patch-clamp amplifier (HEKA, Lambrecht, Germany) and low-pass filtered with an eight-pole Bessel filter at a cutoff frequency of 2 kHz. Data were sampled at 5 kHz, digitized (ITC-16, Instrutech, Elmont, NY), stored on hard disk, and analyzed with PULSE software (HEKA). Patch pipettes were prepared from Kimax-51 glass (Kimble, Vineland, NY) and coated with silicone (Sylgard 184 silicone elastomer kit, Dow Corning). To determine membrane potentials, the command voltages were corrected off-line for liquid junction potentials (25). The standard pipette solution (cytoplasm) contained 150 mM potassium gluconate, 2 mM MgCl₂, 10 mM EGTA, 2 mM MgATP, and 10 mM Hepes/Tris (pH 7.4). Sealing solution contained 20 mM CaCl₂ and 30 mM potassium gluconate (pH 5.6). The standard bathing medium was composed of 10 mM Mes/Tris (pH 5.6) in addition to 1 mM CaCl₂ and 30 mM

Abbreviations: ABA, abscisic acid; RT, reverse transcription.

[†]A.S. and N.I. contributed equally to this work.

[¶]To whom reprint requests should be addressed. E-mail: hedrich@botanik.uni-wuerzburg.de.

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. §1734 solely to indicate this fact.




Fig. 1. Characterization of the plant line tagged by insertion of the *En-1* in the *KAT1* gene. (A) Position of the *En-1* transposon insertion. The diagram depicts the genomic organization of the *KAT1* gene (accession no. X93022) with introns (lines) and exons (black boxes). The nucleotide sequence (below) flanking both ends of the transposon insertion (gray box, designated *En-1*) in the plant line *KAT1::En-1* 7AAD31 shows the duplication of codon 374 (AAC). Above, the positions of the *KAT1*-specific primers K1-4 and K1-5 are indicated. The directions of the *En-1*-specific primers En-205 and En-8130 are depicted above the *En-1* box. (B) Southern blot analysis of *KAT1::En-1* plants. The Southern blot on the left side shows two representative F₁ (1 and 3) and the wild-type (wt) plants, digested with *Pst*I and hybridized with a probe specific to the 3' region of *KAT1*. The transposon-tagged allele of *KAT1* shows a signal of 2.4 kb, and somatic reversion events are visible as a faint band of the expected wild-type size (\approx 12 kb). On the right side, the *KAT1* (black box) and the inserted *En-1* (gray box) are shown and the relevant *Pst*I sites (designated P) are indicated.

potassium gluconate. All solutions were adjusted to 560 mosmol g^{-1} by using D-sorbitol. Chemicals were obtained from Sigma.

Impalement Measurements. For the impalement of guard cells with microelectrodes (26), the abaxial epidermis was peeled and attached to a microscope slide by using Medical Adhesive (VM 355, Ulrich AG, St. Gallen, Switzerland). Opening of stomata was provoked by keeping the epidermal strips in the following solution: 50 mM KCl, 0.1 mM CaCl_2 , and 5 mM Mes/Bis-Tris propane (pH 6.5), which was aerated with CO_2 -free air and illuminated at a photon flux density of 250 $\mu\text{mol m}^{-2}$ per s (halogen lamp Type 6423, Philips). Before impalement, the epidermal strips were transferred to the following bath solution: 50 mM KCl and 1 mM $\text{Ca}(\text{OH})_2$, buffered to pH 6.0 with Mes. Where indicated, the Ca^{2+} concentration was increased to 20 mM with CaCl_2 .

Guard cells were impaled on an upright microscope (Axioskop 2FS, Carl Zeiss), at an angle of 40°. Double barreled electrodes were pulled from borosilicate glass capillaries (GC100F-10, Clark Electromedical Instruments, Pangbourne, U.K.), two capillaries were aligned, heated, and twisted 360° on a customized electrode puller (L/M-3P-A, List Medical Electronic Darmstadt, Germany) and the tip was pulled on a laser puller (P-2000, Sutter Instrument Co., Novato, CA). The electrodes were filled with 300 mM potassium acetate (pH 7.5) and had a tip resistance ranging from 100 to 240 M Ω .

The electrodes were connected via 300 mM KCl bridges and AgCl/Cl half cells to a double microelectrode amplifier (VF-102,

Bio-Logic, Claix, France) equipped with headstages of 10^{11} Ω input impedance. Voltage step protocols were applied via an ITC-16 interface (Instrutech Corp., Elmont, NY) under control of Pulse software (HEKA). The test voltages were fed into a differential amplifier (CA-100, Bio-Logic) connected to the VF-102 amplifier. The data were low pass filtered at 300 Hz with an 8-pole Bessel filter (type 902, Frequency Devices, Haverhill, MA) and sampled at 1 kHz.

Measurements of Apoplastic K^+ Concentrations. K^+ concentrations were recorded by K^+ -sensitive microelectrodes as described (27). Briefly, the measuring electrode was inserted through a half-open stoma into the apoplastic cavity below. The voltage reference was positioned in a neighboring stoma. Both electrodes were connected to a high-impedance amplifier (FD 223; WP-Instruments, Sarasota, FL) that simultaneously measured and subtracted the signals coming from the K^+ electrode and the voltage electrode. Single pipettes were pulled on a vertical puller (List Instruments, Darmstadt, Germany) and silanized internally by using a 0.2% tributylsilane/chloroform solution. After heat stabilization at 200°C for 1 h, the cooled pipettes were backfilled with the sensor mixture (Fluka, no. 60398) dissolved in a mixture of polyvinylchloride/tetrahydrofuran (40 mg/ml) at a ratio of 30:70 (vol/vol). After evaporation of the tetrahydrofuran, the remaining firm gel was topped with the undiluted sensor mixture followed by the reference solution, which consisted of 100 mM Mes/Tris mixed to pH 6 in 0.5 mM KCl.

Stomatal Assay. Leaves were harvested at 8 a.m. from darkened plants. Epidermal strips were peeled off under dim red light, washed for 1 min in a solution without K^+ (pH 4.5), and transferred to the test solutions containing 0.1 mM CaCl_2 and 5 mM Mes/BTP (pH 5.6) and KCl at the desired concentration. Epidermal peels were kept either in the dark or light (400 $\mu\text{mol m}^{-2}$ per s) for 3 h. To take stomatal images, the strips were pasted onto coverslips by using silicon glue, covered with the respective solution, and mounted on the microscope (Axiovert 100, Zeiss). The stomatal aperture was defined as the maximal width between the inner cuticular lips. No differences in the length of the stomatal complex were observed between wild-type and knockout plants. For each condition, >100 stomata were analyzed in three different plants.

Gas Exchange Measurements. A section of the predarkened *Arabidopsis* leaf was enclosed in a sandwich-type cuvette (diameter 2.1 cm) with two windows on the upper and lower side. Gas flow through the cuvette was set to a constant rate of 0.5 liter min^{-1} . Relative humidity was 40% and the temperature was 25°C. Transpiration rates were measured by infrared gas analysis technique by using the Binos instrument (Heraeus, Hanau, Germany). Experiments were started at 8 a.m.

Results and Discussion

Isolation of a *KAT1* Transposon Insertion Mutant from *Arabidopsis thaliana*. In a reverse genetic approach (22) employing a combination of *KAT1*- and *En-1*-specific PCR-primers, we identified the mutant *KAT1::En-1* 7AAD31. Sequence analysis revealed that *En-1* was inserted into the sixth exon of the *KAT1* gene, thereby disrupting the ORF. This insertion caused a duplication of the nucleotides 1,764–1,766, which code for Asn374 (Fig. 1A). Southern blot analysis of six F₁ plants revealed that the parental line 7AAD31 was homozygous for the *En-1* insertion (data not shown). In the F₁ plants, we observed a low degree of somatic reversion as indicated by a faint band on the Southern blot at the size of the *KAT1* wild-type signal (\approx 12 kb, Fig. 1B). This signal constituted 12–17% of the overall signal, an amount frequently observed for somatic reversion of *En-1* in *Arabidopsis* (28).




Fig. 2. Stomatal performance of wild-type and *KAT1::En-1* plants. (A) Light-induced stomatal opening from wild-type (open bars) and *KAT1::En-1* (filled bars) plants. Stomatal apertures were measured after a 3-h illumination period of epidermal peels incubated in concentrations of KCl as indicated. Mean values (\pm standard errors) are shown. (B) Stomatal movement in wild-type and *KAT1::En-1* leaves in response to CO₂ and ABA. Leaves from predarkened plants were placed in a gas exchange chamber and superfused continuously with CO₂-free air for stomatal opening. Application of either 340 ppm of CO₂ or 100 μ M ABA (arrows) induced stomatal closure, whereas removal of CO₂ led to reopening of stomata (arrows). Transpiration rates are given in arbitrary units. Bars represent 20 min and 0.5 mmol m⁻² per s, respectively. (C) Apoplastic changes in K⁺ activity (pK) during light-dependent stomatal opening of representative wild-type and *KAT1::En-1* plants. Following the dark-conditioning characterized by stable K⁺ activities the application of continuous white light (200 μ mol m⁻² per s) caused a decrease in the K⁺ activity (increase in pK) reaching a stable value after \approx 20 min.

Stomatal Movement in *KAT1::En-1* Mutants Is Identical to That in Wild-Type Plants. Wild-type and mutant plants were indistinguishable with respect to plant morphology, guard cell size (radius $r = 4.1 \pm 1.9 \mu\text{m}$ for the wild type and $r = 4.2 \pm 2.1 \mu\text{m}$ for *KAT1::En-1*, calculated from the whole-cell capacitance of 18 and 16 guard cells, respectively), and number of chloroplasts per guard cell (8 ± 1 , $n = 14$, for wild type and 8 ± 1 , $n = 16$, for *KAT1::En-1*). Furthermore, no significant differences between the photosynthetic capacity of guard cell chloroplasts from wild-type and *KAT1::En-1* plants could be detected in Microscopy-PAM chlorophyll fluorescence measurements (data not shown; ref. 29).

In search for the phenotype of a *KAT1* insertion mutant, we measured the stomatal aperture in epidermal peels of predarkened wild-type and mutant plants in response to a 3-h light period. Stomatal opening assays were started at 8 a.m. because recent experiments demonstrated that in the morning stomatal opening is based on K⁺ uptake, whereas in the afternoon it is also driven by sugar accumulation (30, 31). Unexpectedly, light-dependent stomatal apertures of *KAT1::En-1* plants at different K⁺ concentrations were similar to those of wild-type plants (Fig. 2A). Even in the presence of 10 μ M fusicoccin, a compound that hyperactivates the H⁺-ATPase and thereby induces stomatal




Fig. 3. Quantitative RT-PCR. (A) Gel electrophoresis of RT-PCR products of either wild-type (left lanes) or *KAT1::En-1* (right lanes) guard cell protoplast mRNA (positive signals only); representative of $n = 3$ experiments. Fragments were cloned and sequenced for verification. Notice that *KAT1* transcripts are also detectable in the *KAT1::En-1*. Fragment length are for *KAT1* = 379 bp, *KAT2* = 392 bp, *AKT1* = 347 bp, *AKT2/3* = 353 bp, *AtKC1* = 373 bp, and *AtGORK* = 496 bp. (B) Quantification of K⁺ channel transcripts by external standards relative to actin in different guard cell protoplast preparations of wild-type (WT) and *KAT1::En-1* (EN; $n = 3 \pm$ SD). The figure shows the calculated amount of cDNA molecules in the individual probes. *KAT1* expression in the *KAT1::En-1* compared with the wild type is about 6%.

opening (32), stomatal behavior of mutant and wild-type plants was indistinguishable (data not shown).

To verify that stomata in intact leaves behave similarly to those in isolated epidermal strips, we mounted *Arabidopsis* wild-type and mutant leaves into a gas exchange chamber. Using infrared gas analysis, we monitored changes in transpiration, a measure for stomatal movement (33). Predarkened wild-type and mutant plants opened their stomata in response to CO₂-free air, but they closed upon increase of CO₂ to 340 ppm (Fig. 2B). Feeding the wilting hormone ABA via the petiole to leaves from wild-type and knockout plants induced the rapid closure of reopened stomata. Using K⁺-selective electrodes, we tested whether stomatal opening in wild-type and knockout plants is accompanied by K⁺ changes in the apoplast. Following the light stimulus, the K⁺ concentration in the apoplastic cleft between the guard cells and their neighboring cells decreased from $3.97 \pm 0.92 \text{ mM}$ to $2.25 \pm 0.94 \text{ mM}$ ($n = 7$) in wild-type plants and from $4.55 \pm 1.21 \text{ mM}$ to $3.17 \pm 1.12 \text{ mM}$ ($n = 11$) in *KAT1::En-1* plants (Fig. 2C). These values corresponded to a decrease in apoplastic K⁺ concentration of $1.72 \pm 0.08 \text{ mM K}^+$ in the wild type compared with $1.38 \pm 0.31 \text{ mM}$ in *KAT1::En-1*. These results show that K⁺ uptake into guard cells from *KAT1::En-1* is not impaired by the mutation and indicates that K⁺ uptake in the mutant is rescued by other K⁺ transporters.

Multiple K⁺ Channel Genes Are Transcribed in Guard Cells of *Arabidopsis*. To identify other transcripts for K⁺ channels that could influence K⁺ uptake, we performed RT-PCR experiments on mRNA isolated from preparations of guard cell protoplasts of both wild-type and *KAT1::En-1* plants. As expected for somatic reversion, we detected low levels of *KAT1* mRNA in guard cell protoplasts from *KAT1::En-1* (\approx 6% of wild-type level, Fig. 3, cf.




Fig. 4. Inward-rectifying K^+ currents in intact guard cells from wild-type and *KAT1* knockout plants. (A) Hyperpolarization-induced K^+ currents in the presence of 1 mM Ca^{2+} and 50 mM K^+ in guard cells from wild-type (Left) and *KAT1::En1* (Right). Cells were clamped from a holding potential of -100 mV to test pulses ranging from -100 mV to -220 mV in 20-mV decrements. (B) Activation of K^+ channels in wild-type (Left) and *KAT1::En1* (Right) using the pulse protocol as in A. Currents were recorded in the presence of 20 mM Ca^{2+} and 50 mM K^+ . (C) Steady-state current-voltage relation of K^+ currents in wild type (filled symbols) and *KAT1::En1* (open symbols) in the presence of 1 mM (Left) and 20 mM Ca^{2+} (Right). Data points represent mean values \pm SEM, $n = 8$ (wild type, 1 mM Ca^{2+}), 7 (*KAT1::En1*, 1 mM Ca^{2+}), and 6 (wild type and *KAT1::En1*, 20 mM Ca^{2+}). Note the higher Ca^{2+} sensitivity of inward K^+ channels in *KAT1::En1* compared with the wild type. Inward K^+ currents were observed in 20 out of 32 (wild type) and 21 out of 44 measurements (*KAT1::En1*).

Fig. 1B). Using gene-specific primers in RT-PCR experiments, we probed for the presence of *KAT1*-related channel transcripts. Among them, *KAT1* (15), *KAT2* (34), and *AKT1* (35) encode inward-rectifying, *GORK* (36) and *SKOR* (37) encode outward-rectifying, and *AKT2/3* (38) encodes largely voltage-independent K^+ channels. We analyzed the transcript levels of *AtKCL1*, *AKT5*, and *AKT6*, channels that have not been functionally expressed yet. In addition to *KAT1*, we found *KAT2*, *AKT1*, *AtKCL1*, *AKT2/3*, and *GORK* to be expressed in wild-type guard cells (Fig. 3). In contrast, *AKT5*, *AKT6*, and *SKOR* mRNA could not be detected. The finding that guard cells express multiple K^+ channels is in line with staining of guard cells expressing GUS under the control of the individual promoters for *KAT2* (34), *AKT1* (39), *AKT2/3* (40), or *GORK* (data not shown, cf. ref. 36). Despite the strong reduction in *KAT1* transcripts, expression




Fig. 5. K^+ channel activities of guard cell protoplasts from wild-type and *KAT1::En1* plants. (A) Wild-type inward K^+ currents in the whole-cell configuration of the patch-clamp technique upon hyperpolarization of the guard cell plasma membrane. Test pulses to voltages between -9 and -189 mV were applied in 20-mV decrements starting from a holding potential of -49 mV. Absence (B) or reduction (C) of inward K^+ currents in response to the same voltage protocol as used in A in two different guard cell protoplasts from *KAT1::En1* mutants. The bath contained 30 mM K^+ and 1 mM Ca^{2+} (pH 5.6). (D) Histogram of steady-state K^+ current (I_{ss}) amplitudes at -189 mV of 18 cells from wild type (open bars) and 95 cells from *KAT1::En1* plants (filled bars). (E) Steady-state current-voltage relations for K^+ channels from wild-type (open circles, \pm SEM, $n = 18$) and *KAT1::En1* (filled circles, \pm SEM, $n = 16$) plants.

levels of the other K^+ channel transcripts in mutant guard cells remained largely unaffected (Fig. 3B).

Inward-Rectifying K^+ Currents in Intact Guard Cells from Wild-Type and *KAT1::En1* Plants.

To compare the properties of the K^+ inward rectifier between the wild type and mutant, we used two different approaches: impalement of double barreled microelectrodes into intact guard cells and patch-clamp analysis on enzymatically isolated guard cell protoplasts. Upon hyperpolarization of guard cells impaled by microelectrodes, time- and voltage-dependent inward K^+ currents were elicited in both wild-type and mutant plants (Fig. 4A). At 1 mM extracellular Ca^{2+} , K^+ current amplitudes were not significantly smaller in *KAT1::En1* compared with wild type (Fig. 4C). In the presence of 20 mM Ca^{2+} , however, a pronounced difference in current amplitudes between wild type and mutant plants appeared (Fig. 4B and C). The decrease of K^+ currents at strong hyperpolarization results from a voltage-dependent Ca^{2+} block, characteristic for the guard cell inward rectifier (41–43). Apparently, K^+ channels in the *KAT1::En1* exhibit a higher Ca^{2+} sensitivity compared with wild type. When heterologously expressed, *KAT1* (42), *KAT2* (our own observations), and *AKT1* (A. Bertl, personal communication) are Ca^{2+} -insensitive. Because *AKT2/3* represents the only *Arabidopsis* K^+ channel known to be susceptible to block by Ca^{2+} ions (38), this channel could possibly account for the Ca^{2+} -sensitive component of the guard cell inward rectifier. The pronounced Ca^{2+} sensitivity of mutant guard cells compared with wild type therefore may result from a relative increase in *AKT2/3* channels with respect to the entire K^+ channel pool.

In line with the impalement studies, patch-clamp experiments on guard cell protoplasts from *Arabidopsis* wild-type plants revealed the presence of inward-rectifying K⁺ currents (Fig. 5A) (20, 42, 44, 45). In contrast, however, the major fraction of *KATI::En-1* protoplasts (79%) lacked this K⁺ conductance (Fig. 5B and D). In the remaining 21% of cells, hyperpolarization elicited time-dependent inward K⁺ currents of reduced amplitude (Fig. 5C and E). Based on their voltage-dependence (Fig. 5E), these residual currents largely resembled those of the guard cell inward rectifier observed in wild-type *Arabidopsis*. The strong reduction in K⁺ currents of mutant guard cell protoplasts compared with wild-type cells correlates with the decrease of *KATI* mRNA levels. This coincidence indicates that *KATI* represents the dominant inward K⁺ channel recorded in patch-clamp measurements on enzymatically isolated guard cell protoplasts. In intact guard cells, however, qualitative rather than quantitative differences between the mutant and wild type were observed. The differences in K⁺ currents between intact guard cells and protoplasts may reflect, for example, alterations in signaling chains induced by elicitors released by cell wall degradation or changes in the turgor.

The presence of *KAT2*, *AKT1*, *AtKCI1*, and *AKT2/3* transcripts in guard cells together with the K⁺ current fingerprint of *KATI* knockout plants documents that multiple K⁺ channels coexpress in this cell type. We thus conclude that inward-rectifying K⁺ currents in wild-type *Arabidopsis* guard cells are not based just on *KAT1* homomers. K⁺ currents observed in *KATI::En-1* may therefore be mediated by Ca²⁺-insensitive K⁺ channels (e.g., *AKT1*, *KAT2*) and Ca²⁺-sensitive (*AKT2/3*) K⁺ channels. Whether the different channels form homo- or heteromeric K⁺ channels is under current investigation (46, 47). In conclusion, we predict that in the absence of *KAT1* guard cell K⁺ channel homeostasis guarantees stomatal function.

We are grateful to Petra Spormaker for help and support during the early stage of the project. We also thank Michaela Lehnen, Petra Tänzler, and Spidola Neimanis for excellent technical assistance as well as Susanne Michel and the ADIS group for sequence analysis. We are also grateful to Chang-Hyo Goh for chlorophyll fluorescence measurements. This work was funded by the European Community's BIOTECH and INCO-Copernicus programs and the Deutsche Forschungsgemeinschaft to R.H. and K.P.

1. Assmann, S. M. & Haubrick, L. L. (1996) *Curr. Opin. Cell Biol.* **8**, 458–467.
2. Hedrich, R., Hoth, S., Becker, D., Dreyer, I. & Dietrich, P. (1998) in *NATO ASI Series*, eds. Lo Schiavo, F., Last, R. L., Morelli, G. & Reikhel, N. V. (Springer, Berlin), Vol. H 104, pp. 35–45.
3. MacRobbie, E. A. C. (1998) *Philos. Trans. R. Soc. London B Biol. Sci.* **353**, 1475–1488.
4. Thiel, G. & Wolf, A. H. (1997) *Trends Plant Sci.* **2**, 339–345.
5. Ward, J. M. & Schroeder, J. I. (1997) in *Molecular and Cell Biology Updates: Signal Transduction in Plants*, ed. Aducci, P. (Birkhauser, Boston), pp. 1–22.
6. Willmer, C. & Fricker, M. (1996) *Stomata* (Chapman and Hall, London).
7. Zimmermann, S., Ehrhardt, T., Plesch, G. & Müller-Röber, B. (1999) *Cell. Mol. Life Sci.* **55**, 183–203.
8. Grabov, A. & Blatt, M. R. (1998) *J. Exp. Bot.* **49**, 351–360.
9. Li, J., Wang, X. Q., Watson, M. B. & Assmann, S. M. (2000) *Science* **287**, 300–303.
10. Armstrong, F., Leung, J., Grabov, A., Brearley, J., Giraudat, J. & Blatt, M. R. (1995) *Proc. Natl. Acad. Sci. USA* **92**, 9520–9524.
11. Grabov, A. & Blatt, M. R. (1998) *Proc. Natl. Acad. Sci. USA* **95**, 4778–4783.
12. Jacob, T., Ritchie, S., Assmann, S. M. & Gilroy, S. (1999) *Proc. Natl. Acad. Sci. USA* **96**, 12192–12197.
13. Grabov, A. & Blatt, M. R. (1999) *Plant Physiol.* **119**, 277–288.
14. Anderson, J. A., Huprikar, S. S., Kochian, L. V., Lucas, W. J. & Gaber, R. F. (1992) *Proc. Natl. Acad. Sci. USA* **89**, 3736–3740.
15. Schachtman, D. P., Schroeder, J. I., Lucas, W. J., Anderson, J. A. & Gaber, R. F. (1992) *Science* **258**, 1654–1658.
16. Véry, A. A., Gaymard, F., Bosseux, C., Sentenac, H. & Thibaud, J. B. (1995) *Plant J.* **7**, 321–332.
17. Hoshi, T. (1995) *J. Gen. Physiol.* **105**, 309–328.
18. Hedrich, R., Moran, O., Conti, F., Busch, H., Becker, D., Gambale, F., Dreyer, I., Kuch, A., Neuwinger, K. & Palme, K. (1995) *Eur. Biophys. J.* **24**, 107–115.
19. Nakamura, R. L., McKendree, W. L., Jr., Hirsch, R. E., Sedbrook, J. C., Gaber, R. F. & Sussman, M. R. (1995) *Plant Physiol.* **109**, 371–374.
20. Ichida, A. M., Pei, Z. M., Baizabal-Aguirre, V. M., Turner, K. J. & Schroeder, J. I. (1997) *Plant Cell* **9**, 1843–1857.
21. Wisman, E., Guillermo, H. C., Franz, P. & Seadler, H. (1998) *Plant Mol. Biol.* **37**, 989–999.
22. Baumann, E., Lewald, J., Saedler, H., Schulz, B. & Wisman, E. (1998) *Theor. Appl. Genet.* **97**, 729–734.
23. An, Y. Q., McDowell, J. M., Huang, S., McKinney, E. C., Chambliss, S. & Meagher, R. B. (1996) *Plant J.* **10**, 107–121.
24. Hedrich, R., Busch, H. & Raschke, K. (1990) *EMBO J.* **9**, 3889–3892.
25. Neher, E. (1992) *Methods Enzymol.* **207**, 123–131.
26. Roelfsema, M. R. & Prins, H. B. (1997) *Planta* **202**, 18–27.
27. Felle, H. H., Hanstein, S., Steinmeyer, R. & Hedrich, R. (2000) *Plant J.* **24**, 297–304.
28. Cardon, G. H., Frey, M., Saedler, H. & Gierl, A. (1993) *Plant J.* **3**, 773–784.
29. Goh, C.-H., Schreiber, U. & Hedrich, R. (1999) *Plant Cell Environ.* **22**, 1057–1070.
30. Talbot, L.-D. & Zeiger, E. (1996) *Plant Physiol.* **111**, 1051–1057.
31. Lu, P., Zhang, S. Q., Outlaw, W. H., Jr., & Riddle, K. A. (1995) *FEBS Lett.* **362**, 180–184.
32. Baunsgaard, L., Fuglsang, A. T., Jahn, T., Korthout, H. A., de Boer, A. H. & Palmgren, M. G. (1998) *Plant J.* **13**, 661–671.
33. Raschke, K. (1979) in *Encyclopedia of Plant Physiology, New Series, Physiology of Movements*, ed. Feinleib, H. W. A. M. E. (Springer, New York), Vol. 7, pp. 383–441.
34. Pilot, G., Lacombe, B., Gaymard, F., Cherel, I., Boucherez, J., Thibaud, J. B. & Sentenac, H. (2000) *J. Biol. Chem.*, in press.
35. Bertl, A., Anderson, J. A., Slayman, C. L., Sentenac, H. & Gaber, R. F. (1994) *Folia Microbiol.* **39**, 507–509.
36. Ache, P., Becker, D., Ivashikina, N., Dietrich, P., Roelfsema, M. R. G. & Hedrich, R. (2000) *FEBS Lett.* **486**, 93–98.
37. Gaymard, F., Pilot, G., Lacombe, B., Bouchez, D., Bruneau, D., Boucherez, J., Michaux-Ferriere, N., Thibaud, J. B. & Sentenac, H. (1998) *Cell* **94**, 647–655.
38. Marten, I., Hoth, S., Deeken, R., Ache, P., Ketchum, K. A., Hoshi, T. & Hedrich, R. (1999) *Proc. Natl. Acad. Sci. USA* **96**, 7581–7586.
39. Lagarde, D., Basset, M., Lepetit, M., Conejero, G., Gaymard, F., Astruc, S. & Grignon, C. (1996) *Plant J.* **9**, 195–203.
40. Lacombe, B., Pilot, G., Michard, E., Gaymard, F., Sentenac, H. & Thibaud, J. B. (2000) *Plant Cell* **12**, 837–851.
41. Dietrich, P., Dreyer, I., Wiesner, P. & Hedrich, R. (1998) *Planta* **205**, 277–287.
42. Brüggemann, L., Dietrich, P., Dreyer, I. & Hedrich, R. (1999) *Planta* **207**, 370–376.
43. Fairley-Grenot, K. A. & Assmann, S. M. (1992) *J. Membr. Biol.* **128**, 103–113.
44. Brüggemann, L., Dietrich, P., Becker, D., Dreyer, I., Palme, K. & Hedrich, R. (1999) *Proc. Natl. Acad. Sci. USA* **96**, 3298–3302.
45. Pei, Z. M., Kuchitsu, K., Ward, J. M., Schwarz, M. & Schroeder, J. I. (1997) *Plant Cell* **9**, 409–423.
46. Dreyer, I., Becker, D., Bregante, M., Gambale, F., Lehnen, M., Palme, K. & Hedrich, R. (1998) *FEBS Lett.* **430**, 370–376.
47. Urbach, S., Cherel, I., Sentenac, H. & Gaymard, F. (2000) *Plant J.* **23**, 527–538.