### DOCUMENT RESUME

ED 352 745 EC 301 107

AUTHOR

Haas, Julie; Popowicz, Louanne

TITLE

Toddler-Parent Playgroups: Empowering Parents in

Language Intervention.

PUB DATE

Nov 91

NOTE

14p.; Paper presented at the Annual Convention of the

American Speech-Language-Hearing Association

(Atlanta, GA, November 22-25, 1991). Pages 2-7 will not reproduce well due to extremely small and/or

broken print.

PUB TYPE

Speeches/Conference Papers (150) -- Guides -

Non-Classroom Use (055)

EDRS PRICE

MF01/PC01 Plus Postage.

DESCRIPTORS

Case Studies; \*Communication Skills; \*Early Intervention; Interaction; \*Language Handicaps; Models; Parent Education; \*Parent Participation; Play; Preschool Education; Program Implementation;

Skill Development; Speech Therapy; \*Toddlers

**IDENTIFIERS** 

\*Developmental Delays

### **ABSTRACT**

In an attempt to provide a model of language intervention tailored to toddler needs, this poster session gives an overview of a playgroup program implemented by a speech-language pathologist . . an early intervention specialist. The program's aim is to improve communicative abilities while maintaining the integrity of the child-caregiver interaction. Caregivers are active participants in therapy, and children are never separated from the caregivers and expected to interact with the therapist. The speech-language pathologist and behavior/family specialist act as facilitators, targeting intervention strategies and modelling techniques. The program is designed to avoid directive intervention that often leads to nonverbal behavior or fewer communicative attempts. Another element of the program that is designed to enable improved interactions is structure--the same set of toys is used during each session, the same set of rules is followed, and responses to behaviors are consistent. Four case histories of toddlers with developmental delays are provided, including target language goals and frequency of parents' use of specific language intervention techniques over the course of the playgroup intervention. (JDD)



<sup>\*</sup> Reproductions supplied by EDRS are the best that can be made

### .DR. GERTRUDE A. BARBER CENTER



U.S. DEPARTMENT OF EDUCATION
Office of Educational Research and Improvement EDUCATIONAL RESOURCES INFORMATION CENTER (ERIC)

- CENTER (ERIO)

  This document has been reproduced as received from the person or organization originating it
- Minor changes have been made to improve reproduction quality
- Points of view or opinions stated in this docu-ment do not necessarily represent official OERI position or policy

4 ~ N 10

70

### POSTER SESSION

American Speech/Language/Hearing Association Convention November 22-25, 1991 Atlanta, GA

Title:

Toddler-Parent Play Groups: Empowering Parents in Language

vention

Presentors:

Louanne Popowicz, M.Ed.

Parent Education Specialist

Julie Haas, M.A., CCC-SLP Speech/Language Pathologist

Address:

136 East Avenue

Erie, PA 16507

Phone:

(814) 453-7661, extension 244

PERMISSION TO REPRODUCE THIS MATERIAL HAS BEEN GRANTED BY

プスペピー

BEST COPY AVAILABLE

TO THE EDUCATIONAL RESOURCES INFORMATION CENTER (ERIC)



TDD (814) 459-4211

### TODDLER-PARENT PLAY GROUPS: Empowering Parents in Language Intervention

### Issue:

Providing language intervention for toddler-age children requires an understanding of the special needs of toddlers and parents if successful communicative interactions are to be fostered. Increasing numbers of speech/language pathologists who have not received specialized training in working with toddlers are now beginning to provide direct service to this population and their families. Because service providers do not always understand toddler development, programs often mirror intervention for preschool or school-age children. In an attempt to provide a model of language intervention tailored to toddler needs, the poster session will give an overview of a program implemented by a speech/language pathologist and a family specialist.

Objective: To illustrate the purpose and structure of an innovative toddler-parent playgroup as a model demonstration for language intervention.

### Program Philosophy:

The premise of the program is to improve communicative abilities while maintaining the integrity of the child-caregiver interaction. Three essential components include:

- 1) The caregiver remains the child's primary communicative partner.
- 2) The speech/language pathologist and family specialist act as participant observers.
- 3) Intervention techniques include incidental teaching modeled by therapists.

### Objectives of the Toddler-Parent Play Group:

- 1) To provide a nonthreatening child-parent centered versus adult-directed form of intervention.
- 2) To enable primary caregivers as experts in their child's development.
- 3) To provide transdisciplinary intervention from a family specialist and speech/language pathologist.
- 4) To increase child's frequency of nonverbal and verbal communicative behaviors through the use of incidental teaching methods.

### Procedures:

Participants include 2-5 parent-toddler dyads and two early intervention specialists. Parents are active participants in therapy, not observers. The child is never separated from the caregiver and expected to interact with the therapist. The primary caregiver-child dyad remains the most valuable and central focus. The speech/language pathologist and family specialist act as facilitators, targeting intervention strategies and modeling techniques. The therapists do not tell the parent how to interact unless the parent outwardly asks, but rather model an interaction and encourage parents to analyze their own interactions. Incidental language techniques modeled by the therapist will be outlined in the poster session.



Programs which use more adult-directed intervention often seem to increase "terrible two" behavior such as excessive tantruming, crying, or other behaviors evident in a power struggle between the adult and toddler. It has been the experience of the authors that directive intervention often leads to nonverbal behavior or less communicative attempts. Toddlers are particularly perceptive of adult expectations placed on them and feel that pressure. A child needs to feel safe, knowing and trusting the adult before he/she will venture to interact. Reluctance in interacting can be a by-product when a child does not feel secure.

Another element of this program that enables improved interactions is structure. Toddles respond and interact when they can know what to expect. For this reason the same basic set of toys are used each session, the same set of rules are followed, and behavior are consistently responded to in the same manner. The poster session will include a typical schedule for an hour session.

In summary, this poster session includes a child-parent centered method of language intervention which allows parents to equally participate in programming. Examples of four parent-toddler dyads will be presented. Although long term data has not yet been collected, this intervention approach provides possibility for further research.

/af



# TODDLER - PARENT PLAYGROUPS: EMPOWERING PARENTS IN LANGUAGE INTERVENTION

Dr. Gertrude A. Barber Center Early Intervention Program Louanne Popowicz, M.E.D., Counseling Iulie Haas, M.A. CCC-SI.P

SESSION # SUOB - PSOSA

To illustrate the purpose and structure of a toddler-parent playgroup as a model for Statement of Purpose: language intervention

Language Stimulation Planned Environment Adult Support Young Child anguage and Developing Self Esteem Children's ncidental Hour of

The Dr. Gertrude A. Barber Center is a nonprofit educachildren and adults with disabilities from birth to adult. tional, residential and rehabilitation facility serving

136 East Avenue Erie, PA 16507

## PHILOSOPHY

## **OBJECTIVES**

- form of intervention. To increase parents'

## Therapists should have an

- psychosocial development in communicative interactions. Use of incidental language order to foster successful understanding of child
  - of psychosocial development techniques supports theories The toddler parent dyad is relationship to foster the most umportant
- facilitated when toddlers feel Progress in therapy is communication

## STAGES OF PSYCHOSOCIAL DEVELOPMENT

Consistency, continuity and negative care may arouse BIRTH to 1 YEAR Irust ve. Mistrust

Opportunities to try out skills at

Autonomy vs. Doubt

2 to 3 YEARS

her own way lead to autonomy.

Overprotection or lack of

child's own pace and in his /

## **PHILOSOPHY**

- To provide a nonthreatening child-parent vs adult directed
- frequency of using incidental
- themselves as experts in their To enable parents to see language techniques.
  - umplemented by a family transdisciplinary model child's development. counselor and SLP. To provide

# **ACCORDING TO ERIKSON**

trust. Inadequate inconsistent or sammes of experience lead to mistrust

## **PARTICIPANTS**

2-5 toddler-caregiver/parent dyads 1 coddler with developmentally age appropriate abilities (non-delayed

WHICH FOSTERS A SENSE

OF TRUST

CONSISTENT ROUTINE

CONSTRUCTA

**PROCEDURES** 

Targeted toddlers exhibit a range of communication-language-speech

Snack

Toddlers exhibit a range of social emotional developmental needs (behavioral issues, parent-child attachment issues.)

Story-Music

## CONSTRUCT AN EMOTIONAL ENVIRONMENT WHICH FOSTERS A SENSE OF AUTONOMY $2 \times 5$ wall murror, Toddier table and child chaus, Child-size book shelf, Child-size toy self, water table

MATERIALS

Therapist respects the parent as the "home-base" for the child's Therapist models positive, accepting emotional tones. emotional refueiting. TOYS

doctor lat

sets of multi-cultural family dolls baby dolls and bottles/blankets Fisher Price School / Discovery Basket of farm/zoo animals oddler books Purse set

support may lead to doubt about

ability to control self or

N

Therapiet assists the parent in

establishing harits and rules.

acknowledge child's feelings

Therapist uses words to

## Therapist assists child in age appropriate negotiation skills. Therapist models empathac

Crayons, paper, playdough

### ENVIRONMENT WHICH FOSTERS A SENSE OF SUCCESS **CONSTRUCT A**

- (continuity of people-toys). Provide opportunities for child

Provide a predictable environment

Provide toys which permit exploration and foster communication. Choice

## CONSTRUCT A COMMUNICATIVELANGUAGE BASED ENVIRONMENT WHICH FOSTERS A SENSE OF

- USE OF INCIDENTAL LANGUAGE TECHNIQUES)
- Therapist comments on child's behaviors or vocalizations as CODTENUES (TVE.
- Therapist models use of expension. Therapiet models use of parallel
- Therapist waits for child
- Therapist acts as participant observer. responses

# BEST COPY AUGULANIE

CASE HISTORY: Sara

D.O.B.: 3/3/89

C.A.: 2 years, 7 months

Diagnosis: Developmental Delay

(Genetic testing being completed)

### Early Intervention Services

Sara ttends the playgroup with her mother and also attends individual physical therapy.

The following scores were assessed at 2 years 6 months of age using the <u>Battelle Developmental Inventory</u>, (Newborg 1988) and the <u>Sequenced Inventory of Communication Development</u>, (Hedrick, Prather and Tobin, 1984):

Cognition: 12 months Receptive Language: 12 months Self-Help/Adaptive: 10 months Expressive language: 12 months Social/Emotion: 14 months Fine Motor: 9-11 months

Gross Motor: 12 months

### Speech-Language-Communication Abilities

Sara communicates primarily through intentional communication (using reaching-vocalizing in conjunction with eye gaze to request attention or an object). She has begun to use gross gestures in an imitative context to request "eat" and "more". Consonant use is limited to vowel sounds and an occasional /m/.

### Case History

Sara was born full term and "small for gestation age". She was a failure to thrive infant and underwent extensive chromosomal testing, specifically for Cornella DeLange Syndrome. Her gross/fine motor, oral motor and cognitive intervention continue to be an important part of her overall programming for communication.

### Targeted Communication Goals Within the Playgroup

- 1. Increase frequency of parent use of child-centered intervention techniques (following child's lead, providing choices to elicit responses).
- 2. Increase frequency of child's communicative attempts, specifically gross gestures.



Frequency of parent-use of child-centered language techniques.

\* recorded for 30 minutes of 1 hour sessions.

SESSION #	FREQUENCY OF PARENT-USE OF TECHNIQUES
Baseline i	0
Baseline 2	0
Baseline 3	0
Intervention 1	0
Intervention 2	2
Intervention 3	0
Intervention 4	5
Intervention 5	7
Intervention 6	5

### CASE HISTORY - Jennifer

D.O.B.: 1/13/90

C.A.: 1 year, 10 months

<u>Diagnosis:</u> Developmental Delay

### Early Intervention Services:

Jennifer attends the playgroup with her adoptive mother and attends individual physical/occupational therapy.

The following scores were assessed at 1 year, 9 months of age using the <u>Battelle Developmental Inventory</u> (Newborg, 1988) and the <u>Sequenced</u> <u>Inventory of Communication Development</u>, (Hedrick, Prather. and Tobin, 1984):

Cognition: 12 months Receptive Language: 16 months
Social/Emotional 14 months Expressive Language: 12 months
Fine Motor: 12 months Self Help/Adaptive: 12 months
Gross Motor: 10 months

### <u> Speech/Language - Communication Abilities:</u>

Jennifer demonstrates use of intentional communication (use of vocalizations in conjuction with eye gaze to gain adult assistance or attention). She frequently uses labeling and greeting, but rarely initiates requests for objects or attention. Sound play has recently increased in frequency and types of sounds used. Increased sound use has developed along with improved oral-motor and feeding skills.

### Case History

Jennifer has been in her adoptive home for the last 10 months. She was born with a heart defect and cogenital anomalies of the hands and feet. She has had a history of feeding problems, including esophageal reflex.

### Targeted Communication Goals Within the Playgroup

- 1. Increase frequency of parent use of child-centered intervention techniques (specifically providing choices, use of expansion and commenting on the child's behaviors as communicative).
- 2. Increase frequency of child's communicative attempts, specifically use of vocalization in a requesting context.



Frequency of parent-use of child -centered language techniques.

\* recorded for 30 minutes of 1 hour session.

SESSION #	FREQUENCY OF	PARENT-USE	OF TECHNIQUES
Baseline 1	10		
Baseline 2	12		
Baseline 3	18		
Baseline 4	12		
Intervention 1	25		
Intervention 2	32		
Intervention 3	41		
Intervention 4	25		
Intervention 5	23		
Intervention 6	26		

CASE HISTORY: Alyssa

D.O.B.: 9/14/89

C.A.: 2 years, 2 months (26 months)

Diagnosis: \* Repaired cleft of the soft palate

\* Pierre Robin Syndrome (small mandible)

\* Delay in expressive language

### Early Intervention Services:

Alyssa attends 1/2 hour of individual speech-language therapy and a 1 hour playgroup per week with her mother and a peer with developmentally age appropriate abilities.

### Developmental Scores

According to the <u>Battelle Developmental Inventory</u>, (Newb**0**rg, 1988) administered at 24 months of age:

Cognition: 24-29 months Personal-Social: 25 months
Gross Motor: 24 months Receptive Language: 28 months
Fine Motor: 24 months Expressive Language: 24 months

Adaptive: 24 months

### Speech-Language Scores

According to the <u>Sequenced Inventory of Communication Development</u> (Hedrick, Prather and Tobin, 1984), The <u>Arizona Articulation Proficiency Scale</u> (Fudula and Reynolds, 1986) and a speech language sample at 24 months of age.

Receptive language age: 28 months Expressive language age: 28 months

Speech intelligibility: total AAPS scores 48 - severe articulation delay.

Eepressive and receptive language abilities are age appropriate. Speech is characterized by hypernasality and glottal stops. There is no use of bilabials. Speech pattern consists of omissions and /k,g,n/ substitutions.

Communicative attempts are infrequent and almost always directed towards Alyssa's mother.

### Case History

Alyssa was born following an uncomplicated, full term pregnancy. She experienced respiratory distress at birth requiring intubation to clear her airways. She had difficulties in nipple feeding for the first few months of life. She received OT/PT services for a period of 6 months (14-20 months) for motor delays.

arganization and a first of

### Target Language Goals Within the Playgroup

- 1. Frequency and maintenance of parent use of child-center intervention techniques.
- 2. Frequency of child's communicative attempts.

Frequency of parent-use of child-centered language techniques

\* recorded for 30 minutes of 1 hour sessions.

SESSION #		FREQUENCY	OF PARENT-USE	OF TECHNIQUES
Baseline 1		10		
Baseline 2		11		
Baseline 3		6		
Intervention	1	15		
Intervention	2	21		
Intervention	3	25		
Intervention	4	0		
Intervention	5	26		
Intervention	6	28		

3/28/89 D.O.B.:

C.A.: 2 years, 7 months (31 months)

Repaired bilateral complete cleft of lip and palate. Diagnosis:

Frequent middle ear infections and fluctuating conductive

loss. Has bilateral myringotomy tubes.

Developmental Delay with severe deficit in expressive

speech and language.

### Early Intervention Services:

Brandon attends a 1/2 hour of individual speech-language therapy and a 1hour playgroup per week with his father.

### Developmental Scores

According to the Battelle Developmental Inventory (Newborg, 1984) administered at 31 months of age:

Cognition: Gross Motor: 32 months

Personal-Social:

26 months

Fine Motor:

31-34 months

Adaptive:

25 months

Receptive Language: 30-31 months

38 months

Expressive Language: 22 months

Total developmental age:

28 months

### Speech-Language Scores:

According to the Sequenced Inventory of Communication Development, (Hedrick, Prather, and Tobin, 1984) and a speech-language sample at 31 months of age:

Expressive Communication Age: 24 months Receptive Communication Age: 32 months

Brandon uses 2-4 word phrases that are completely unintelligible to most listeners, even when the referent is known. Most single words are intelligible to primary caregivers. Communicative attempts remain frequent even though his messages are not clearly received. Brandon, however does not frequently demonstrate frustration when his communicative attempts are not understood by the listener. He regularly makes second attempts to be understood, complete with pointing and gestures.

Speech is characterized by hypernasality, several omissions of initial and final consonants, frequent glottal stops, substitutions of Igl - Idl, Ikl - Itl, Ihl - fricatives, Iml - Ip, bl.



### Case History

Brandon was born with a low birth weight after a complicated pregnancy (blood pressure, toxemia, bleeding). Prior to his second birthday he underwent 6 surgeries which included repair of the right and left lip and palate and insertion of myringotomy tubes. He was tube fed for the first four months of life. Nipple feedings remained lengthy and difficult due to his inability to suck. Excessive drooling and poor proprioceptive sensory awareness was observed until age 2 years 4 months.

### Targeted Language Goals Within the Playgroup

- 1. Focus remains on the child's communicative attempts. Increase frequency of parent use of child-centered intervention techniques.
- 2. Increase frequency and maintenance of child's communicative attempts.

Frequency of parent-use of child-centered language techniques

\* recorded for 30 minutes of 1 hour sessions.

SESSION #	FREQUENCY OF	PARENT-USE	OF TECHNIQUES
Baseline 1	0		
Baseline 2	1		
Baseline 3	0		
Intervention 1	0		
Intervention 2	0		
Intervention 3	1		
Intervention 4	4		

