

DOCUMENT RESUME

ED 076 537

SP 006 442

TITLE Strand V: Education for Survival. First Aid and Survival Education. Grades 4, 5, 6.
INSTITUTION New York State Education Dept., Albany. Bureau of Elementary Curriculum Development.
PUB DATE [70]
NOTE 54p.
EDRS PRICE MF-\$0.65 HC-\$3.29
DESCRIPTORS Accidents; Course Descriptions; *Elementary Grades; *First Aid; Grade 4; Grade 5; Grade 6; *Health Education; Heart Rate; *Injuries; *Instructional Materials

ABSTRACT

GRADES OR AGES: Grades 4-6. SUBJECT MATTER: First aid and survival education. ORGANIZATION AND PHYSICAL APPEARANCE: The guide is divided into seven sections: introduction to first aid; wounds and control of minor bleeding; respiratory emergencies and resuscitation; poisoning; traumatic shock; and injuries from abnormal conditions. The publication format of four columns gives the outline of content, the major understandings and fundamental concepts, suggested teaching aids and learning activities, and supplementary information for teachers. The course objectives are presented in the introduction. The guide is soft covered. OBJECTIVES AND ACTIVITIES: Each subsection contains questions and topics for discussion. The supplementary information provides teachers with further discussion material. INSTRUCTIONAL MATERIALS: Lists of multimedia resources are presented for teachers and students. Information on the procurement of audiovisual aids, mannequins, teaching kits, and pamphlets is also included. STUDENT ASSESSMENT: No provision is made. OPTIONS: The guide is suggestive only. (BRB)

THE UNIVERSITY OF THE STATE OF NEW YORK

Regents of the University (with years when terms expire)

1984 Joseph W. McGovern, A.B., LL.B., L.H.D., LL.D., D.C.L.,
Chancellor ----- New York
1985 Everett J. Penny, B.C.S., D.C.S.,
Vice Chancellor ----- White Plains
1978 Alexander J. Allan, Jr., LL.D., Litt.D. ----- Troy
1973 Charles W. Millard, Jr., A.B., LL.D., L.H.D. ----- Buffalo
1972 Carl H. Pforzheimer, Jr., A.B., M.B.A., D.C.S., H.H.D. ----- Purchase
1975 Edward M.M. Warburg, B.S., L.H.D. ----- New York
1977 Joseph T. King, LL.B. ----- Queens
1974 Joseph C. Indelicato, M.D. ----- Brooklyn
1976 Mrs. Helen B. Power, A.B., Litt.D., L.H.D. ----- Rochester
1979 Francis W. McGinley, B.S., LL.B., LL.D. ----- Glens Falls
1980 Max J. Rubin, LL.B., L.H.D. ----- New York
1971 Kenneth B. Clark, A.B., M.S., Ph.D., Litt.D. ----- Hastings on Hudson
1982 Stephen K. Bailey, A.B., B.A., M.A., Ph.D., LL.D. ----- Syracuse
1983 Harold E. Newcomb, B.A. ----- Owego
1981 Theodore M. Black, A.B. ----- Sands Point

President of the University and Commissioner of Education
Ewald B. Nyquist

Executive Deputy Commissioner of Education
Gordon M. Ambach

Deputy Commissioner for Elementary and Secondary Education
Herbert F. Johnson

Associate Commissioner for Instructional Services
Philip B. Langworthy

Assistant Commissioner for Instructional Services (General Education)
Bernard F. Haake

Director, Curriculum Development Center
William E. Young

Chief, Bureau of Elementary Curriculum Development
Robert H. Johnstone

Director, Division of General Education
Ted T. Grenda

Chief, Bureau of School Health Education
John S. Sinacore

SP 006 442

ED 076537

STRAND V


EDUCATION FOR SURVIVAL


FIRST AID AND SURVIVAL EDUCATION

Grades 4, 5, 6

U.S. DEPARTMENT OF HEALTH,
EDUCATION & WELFARE
NATIONAL INSTITUTE OF
EDUCATION
THIS DOCUMENT HAS BEEN REPRO-
DUCED EXACTLY AS RECEIVED FROM
THE PERSON OR ORGANIZATION ORIGIN-
AL. POINTS OF VIEW OR OPINIONS
STATED DO NOT NECESSARILY REPRESENT
OFFICIAL NATIONAL INSTITUTE OF
EDUCATION POSITION OR POLICY

The University of the State of New York/The State Education Department
Bureau of Elementary Curriculum Development/Albany, New York 12224





STRAND V

EDUCATION FOR SURVIVAL

FIRST AID AND SURVIVAL EDUCATION

GRADES 4, 5, 6

OVERVIEW

First aid education is essential for every American. The knowledge of first aid may someday make it possible for a person to save a life. This life may be that of a stranger or of a person who is very dear. In some cases the life he saves may be his own.

First aid education makes it possible for the individual to deal with the many health emergencies that occur in everyday living. It helps him to meet these health emergencies without confusion or panic, until professional aid becomes available.

All teachers involved with this course should hold a standard and advanced Red Cross first aid card or its equivalent. The practical work in the Red Cross first aid courses should be of particular value to the teacher in deciding how to handle class practice and demonstrations.

STRAND V

EDUCATION FOR SURVIVAL

FIRST AID AND SURVIVAL EDUCATION

GRADES 4, 5, 6

OBJECTIVES

Pupils in grades 4, 5, and 6 should:

- develop an appreciation of first aid measures as a means of saving lives and minimizing disease and injury.
- develop an awareness that immediate action in calling for professional help can be vital.
- know which person or agency is equipped to provide the help needed and how to obtain it.
- develop a feeling of confidence rather than fear in administering first aid.
- develop competency in using first aid kits and equipment.
- know about respiratory emergencies and develop competency in the administration of the various methods of artificial respiration.
- know about the various kinds of wounds and develop competency in controlling bleeding.
- know about the various kinds of shock and the first aid procedures to follow for each type.
- know about the various kinds of poisons that affect the body and the first aid procedures to follow.
- understand how elementary school children can prevent accidents from occurring.

OUTLINE OF CONTENT

MAJOR UNDERSTANDINGS AND FUNDAMENTAL CONCEPTS

SUGGESTED TEACHING AIDS AND LEARNING ACTIVITIES

SUPPLEMENTARY INFORMATION FOR TEACHERS

I. Introduction to First Aid

First aid involves immediate temporary care and not treatment.

What is first aid? What is the difference between first aid and treatment?

A. Definition

First aid is the immediate and temporary care given to an injured or sick person until the services of a physician can be obtained.

B. Values of first aid training

Having knowledge of first aid will give the student a feeling of some competency to care for himself and others and will serve to make him aware of preventable safety problems.

The pupil can care for his own injuries and the injuries of others. He is helped to improve his safety consciousness which aids in the prevention of accidents. First aid can be extremely valuable during catastrophes, such as wars, floods, hurricanes, earthquakes, and tornadoes.

C. General procedures to follow in first aid emergencies

The first aider should function in a logical and systematized way rather than in a state of panic and disorganization.

Learn the meaning of panic.

Describe several accident situations and discuss with pupils the general first aid procedures that should be followed in each case.

Example: A man has just been hit by a car. He is lying in the street and a large crowd is beginning to form around him. As a first aider, what would you do to help

Factors that should be considered include:

- determining the victim's injuries
- determining what injuries should be dealt with first:
- getting the victim's name and address
- carrying out the determined procedures
- securing necessary help
- protecting victim from further harm

OUTLINE OF CONTENT

MAJOR UNDERSTANDINGS AND
FUNDAMENTAL CONCEPTS

SUGGESTED TEACHING AIDS
AND LEARNING ACTIVITIES

SUPPLEMENTARY INFORMATION
FOR TEACHERS

this man?

Point out that securing necessary help is vital. Call for professional help as soon as possible.

Show McGraw-Hill film-strip Your Responsibilities in First Aid. An introductory film to first aid also may be shown, such as the American Red Cross films, Checking for Injuries and The How and Why of First Aid.

Discuss the consequences that might occur if it were proven that a first aider did something that further injured the victim.

Panic can be worse or as bad as an injury. Keep yourself and the injured person calm. Encourage others to do the same. Discuss possible injurious results of panic. Act out in simple skits.

If an injured person is found unconscious keep him lying on his side so

- psychological reassurance for victim
- dealing with external factors of weather, traffic, crowds, etc.

The following bibliographical sources may be checked for further information: Cole, pp. 1-16, and Henderson, pp. 34-56

It is best to summon medical help immediately and keep the victim comfortable.

A first aider might find himself in a lawsuit if he negligently and carelessly administered first aid which resulted in injuries or aggravation of the original injury.

See bibliography:

Gold and Gold: "First Aid and Legal Responsibility," Jan., 1963.

OUTLINE OF CONTENT

MAJOR UNDERSTANDINGS AND FUNDAMENTAL CONCEPTS

SUGGESTED TEACHING AIDS AND LEARNING ACTIVITIES

SUPPLEMENTARY INFORMATION FOR TEACHERS

D. First aid kits and materials

First aid kits should be available in situations where accidents are likely to occur.

that his tongue does not fall back in his mouth to choke him. Move him as little as possible. Call for professional help.

Have several kits available to show to pupils. Show and discuss the contents of each kit.

Contents of kits usually include adhesive compresses, bandage compresses, plain gauze pads, gauze roller bandages, eye dressing

Very often the first aider must improvise with available materials.

Have pupils report on the first aid supplies they have in their homes.

packets, plain absorbent cotten, triangular bandages, tweezers, and scissors, antiseptic soap (containing hexachlorophene), aspirin, elastic bandages 3" wide, and safety pins.

Have the pupils make first aid kits for the home, car, and one for camping trips. Wooden boxes 3" x 5" x 2" may be constructed to house these kits.

Make a list of possible kit items (at right), and discuss this list with your school nurse teacher.

Discuss materials that might be available when there is no first aid kit for supplies.

II. Wounds and the Control of Minor Bleeding

It is vital to keep wounds clean. Wash with soap and water if possible.

First aid for wounds varies according to the nature and possible results of each.

OUTLINE OF CONTENT

MAJOR UNDERSTANDINGS AND
FUNDAMENTAL CONCEPTS

SUGGESTED TEACHING AIDS
AND LEARNING ACTIVITIES

SUPPLEMENTARY INFORMATION
FOR TEACHERS

Show filmstrip and record The History of Nursing - Florence Nightingale, Metropolitan Life, to show the importance of cleanliness.

Restate that it is most important to get professional help.

At home have a telephone list posted including:

- . family doctor
- . nearest poison control center
- . pharmacist's night number
- . police
- . fire

You could make this list on oak tag as a present for your parents.

Elicit children's definitions of a wound.

A. Wound defined
A wound is a break in the skin or in the mucous membrane lining one of the body cavities.

B. Kinds of wounds
There are several different kinds of wounds. Each type has specific characteristics with which the first aider should be familiar.

OUTLINE OF CONTENT

MAJOR UNDERSTANDINGS AND FUNDAMENTAL CONCEPTS

SUGGESTED TEACHING AIDS AND LEARNING ACTIVITIES

SUPPLEMENTARY INFORMATION FOR TEACHERS

1. abrasion

An abrasion is an irregular, superficial, open wound of the skin in which the outer layers are scraped off.

Discuss the kinds of accidents that produce this kind of injury.

There is relatively little bleeding. Scraping of the skin on mats, floors, or concrete would produce an abrasion.

2. incised wound

An incised wound is a clean, sharp cut in the skin which may involve capillaries, veins, or arteries.

Discuss the kinds of accidents that produce this kind of injury.

These wounds are produced by objects such as glass, razors, and knives. They usually appear as straight lines and may bleed profusely.

3. puncture wound

A puncture wound is a deep penetration of the skin which is caused by such objects as bullets, knives, nails, and ice picks.

Discuss the kinds of accidents that produce this kind of injury.

Large, irregularly shaped or deeply imbedded splinters are an example of a puncture wound.

A major danger of this kind of wound is tetanus which is a severe form of infection. Other forms of infection are also a danger.

4. laceration

A laceration is a wound in which there is a jagged, ragged tearing of the tissues.

Discuss the kinds of accidents that produce this kind of injury.

This type of wound is usually produced by tearing the skin on a nail or piece of metal, or being involved in a car accident. They also occur quite frequently in industry.

Bleeding and/or infection are dangers associated with lacerations.

OUTLINE OF CONTENT

MAJOR UNDERSTANDINGS AND FUNDAMENTAL CONCEPTS

SUGGESTED TEACHING AIDS AND LEARNING ACTIVITIES

SUPPLEMENTARY INFORMATION FOR TEACHERS

5. contusion

A contusion is a wound which is commonly referred to as a bruise and occurs when tiny blood vessels are broken by a hard blow or fall.

Examples include black and blue marks on the body and the black eye. The clot changes in color due to the new blood coming into the area and the blood clot gradually being absorbed.

C. Dangers of wounds

The dangers of wounds include loss of blood, infection, and tetanus.

Use the American Red Cross film, Wounds.

The general first aid procedures for all wounds are:

1. prevent contamination
2. provide protection
3. control bleeding, if necessary

1. infection

All wounds are subject to infection because broken skin permits entry of microorganisms.

Soap and water cleansing is best. If soap contains hexachlorophene, it can be helpful if use is repeated. Wash AROUND wound. Alcohol or iodine can be used to cleanse AROUND wound to prevent bacteria from multiplying. Cover with clean gauze. If wound is serious, get professional help. Some bleeding helps to clean the wound. The body will provide help for its own healing if we cooperate. Strong antiseptics on wounds kill tissue. This area is then a place for bacteria to multiply, for it is defenseless with no

Bacteria found on the skin or the object that caused the wound may cause the infection to develop.

Signs of infection include redness, swelling of the area, pains, pus, fever, and enlarged glands (in the neck, under the armpit, or in the groin, depending on where the infection is located).

First aid for infected wounds includes sterile wet compresses and inactivity of the part.

Cleanse the area thoroughly using soap and water and apply a sterile dressing or a Band-Aid. In an emergency situation, any type of clean cloth may be used for a dressing.

OUTLINE OF CONTENT

MAJOR UNDERSTANDINGS AND FUNDAMENTAL CONCEPTS

SUGGESTED TEACHING AIDS AND LEARNING ACTIVITIES

SUPPLEMENTARY INFORMATION FOR TEACHERS

help from body cells.

Physical activity favors
the spread of infection.

The loss of a first tooth
can leave a kind of wound
in the mouth. A certain
amount of bleeding will
occur. Rinse mouth with
clean water. Use
slightly warm water with
a little salt added if
you can. Keep your hands
out of your mouth.

Infected wounds should have
medical care.

A scab forms to help keep
out infection. Pulling
it loose can allow in-
fection to enter the
wound. If bandage is
stuck, soak it with hydro-
gen peroxide (H_2O_2 Solu-
tion 3%) to soften scab
for less damage to
wounded area.

Almost all children have
had antitetanus shots.
This is not always true
for adults.

Report a wound from rusty
metal objects to an adult
in charge immediately.
Nails, splinters, and
gardening tools can be a
dangerous source of
tetanus or lockjaw be-
cause of the presence of

Tetanus or lockjaw if
caused by the organism
Clostridium Tetani. It
multiplies where there is
little or no oxygen. The
toxins produced by the
organism affect the central
nervous system producing
muscle spasms and convul-
sion. The victim has diffi-
culty opening his jaws and
spasm of the facial muscles
often causes a fixed smile

2. tetanus
Tetanus must be considered
a danger in all wounds.

OUTLINE OF CONTENT

MAJOR UNDERSTANDINGS AND FUNDAMENTAL CONCEPTS

SUGGESTED TEACHING AIDS AND LEARNING ACTIVITIES

SUPPLEMENTARY INFORMATION FOR TEACHERS

manure in soil.

and elevated eyebrows.
During convulsions, the rigidity of the chest wall interferes with respiration causing asphyxia, which may lead to death. Immunization against tetanus is the best method of protection.

3. severe bleeding

Control of severe bleeding commands urgent attention.

The first aid objective is to stop the bleeding at once.

Application of pressure is recommended as the safest and most effective method for controlling bleeding.

Demonstrate pressure methods of controlling bleeding.

1. Direct pressure - preferably using a sterile or clean folded cloth over the wound, apply firm and steady pressure with the hand.
2. Pressure dressing - Folds of clean cloth may be held firmly against the wound with a bandage.
3. Pressure points - Press with fingers or heel of the hand to hold the supplying blood vessel against

Rapid or extensive loss of blood can severely affect all body functions; therefore, the control of severe bleeding is one of the most urgent procedures of first aid.

Adults can readily overcome the loss of a pint of blood but the loss of a quart may be a serious matter.

The American Red Cross recommends only two points on each side of the body to use as pressure points. They are the brachial artery on the inner half of the arm, midway between the elbow and armpit. Pressure against the main vessel and bone there diminishes bleeding in the upper extremity below the point of pressure. Pressure against the femoral artery

OUTLINE OF CONTENT

MAJOR UNDERSTANDINGS AND
FUNDAMENTAL CONCEPTS

SUGGESTED TEACHING AIDS
AND LEARNING ACTIVITIES

SUPPLEMENTARY INFORMATION
FOR TEACHERS

An improperly used tourniquet can cause serious injury.

the underlying bone. If bleeding is excessive, call for professional help as soon as you can.

and bone just below the groin on the front, inner half of the thigh diminishes bleeding in the extremity below the point of pressure.

A tourniquet should be used only as a last resort.

This decision should be made by a professional person only.

The tourniquet should be used only when the decision has been made that it may be necessary to sacrifice a limb to save a life. Tissue damage resulting in gangrene may result from indiscriminate use of a tourniquet, and fatal shock has been attributed to effects resulting from release of the tourniquet. A tourniquet should be used only by persons who are aware of the dangers associated with its use.

D. First aid for wounds requiring special attention

1. animal bites

There is always the danger of infection and rabies whenever an animal bite is involved.

Discuss types of wounds that may be associated with animal bites; i.e., puncture, laceration, bruise, etc.

Any warm-blooded animal that is infected with rabies can transmit the disease to humans through any break in the skin. Animals that frequently bite humans include cats, dogs, rats, rabbits, bats, squirrels, and raccoons.

OUTLINE OF CONTENT

MAJOR UNDERSTANDINGS AND FUNDAMENTAL CONCEPTS

SUGGESTED TEACHING AIDS AND LEARNING ACTIVITIES

SUPPLEMENTARY INFORMATION FOR TEACHERS

Assign library projects to determine

- what hydrophobia (rabies) means
- how it affects the human body
- treatment
- whom to contact following animal bites (police and local health department)

Show filmstrip and record, Louis Pasteur, The Germ Theory (Metropolitan Life). Have group do research as to what Pasteur did in helping rabies victims.

Medical attention should always be sought following animal bites.

2. nosebleeds

Nosebleeds are usually not serious unless they are profuse or prolonged.

Most minor nosebleeds can be controlled by having the victim sit down, tilt his head backward, and breathe through his mouth.

3. eye injuries

The eyes are very delicate organs and any injury to them must be brought to the attention of a physician immediately.

Rabies is almost invariably fatal to humans unless there is prompt treatment.

Since the bites of animals always present the danger of rabies, an animal involved in biting a person should never be killed. Authorities should catch the animal and retain it for observation. Untrained or inexperienced people should not attempt to capture a dog that has bitten someone.

The wound should be cleansed thoroughly and covered as soon as possible.

Nosebleeds may occur spontaneously or as a result of injury. If a nosebleed is not brought under control within a reasonable period of time, a physician should be consulted.

Automobile accidents and athletic injuries frequently involve the eye. Children are sometimes hit in the eye with rocks or poked in the eye with pencils and

OUTLINE OF CONTENT

MAJOR UNDERSTANDINGS AND
FUNDAMENTAL CONCEPTS

SUGGESTED TEACHING AIDS
AND LEARNING ACTIVITIES

SUPPLEMENTARY INFORMATION
FOR TEACHERS

Prior to receiving medical attention, the injured eye should be covered with a sterile or clean cloth patch.

Discuss special areas, such as shops, science laboratories, and physical education spaces, where accidents might, most likely occur.

Discuss safety procedures to follow to avoid such accidents.

other sharp objects.

The eye should be covered with a loose sterile dressing and the victim should receive medical care as soon as possible. Cover both eyes to prevent movement of injured as well as uninjured eye.

III. Respiratory Emergencies and Resuscitation

Respiratory emergencies must be dealt with immediately in order to save the person's life.

Have some students report on the basic anatomy and physiology of the respiratory system and the basic principles of the respiratory process.

For information on the respiratory system read:

Henderson, op. cit., pp. 7-9.

American Red Cross. Textbook for Juniors. op. cit., Chapter 6.

Respiratory emergencies involve respiratory obstruction, the cessation of breathing, and the mechanical interference with respiration. Factors that will cause a stoppage of breathing include objects becoming stuck in the wind pipe, electric shock, carbon monoxide, and other gas poisoning, compression of the chest by a cave-in, and prolonged exposure to the cold.



OUTLINE OF CONTENT

MAJOR UNDERSTANDINGS AND FUNDAMENTAL CONCEPTS

A. Common first aid emergencies involving the stoppage of breathing

Discuss some of the objects that might cause respiratory obstruction - toys, food, pins, etc.

A fish or chicken bone lodged in the throat can be a serious problem. Have the person eat dry, unbuttered bread. Call for professional help immediately.

B. Symptoms of an insufficient supply of oxygen

The major symptom associated with an insufficient supply of oxygen is blueness of the skin.

C. Objectives of first aid for respiratory emergencies

The two major objectives of first aid for respiratory emergencies are:
1. to maintain open air passages
2. to get air into the lungs

SUGGESTED TEACHING AIDS AND LEARNING ACTIVITIES

Stoppage of breathing may result from a concussion of the brain, a fracture of the skull, certain types of neck fractures, electric shock, certain drugs (especially the narcotics and sleeping pills composed of barbiturates), lack of oxygen in the air, toxic gases (carbon monoxide), and heart disease.

The victim's head is thrown back, the eyes bulge, and his face becomes bluish-red in color. The person struggles to get air.

Achievement of the major objectives is attempted through:
1. checking the nose, mouth, and throat for foreign objects
2. positioning the patient with neck hyperextended to maintain an open airway
3. forcing a flow of air into and from the lungs by mechanical action that stimulates the function of the respiratory muscles.

SUPPLEMENTARY INFORMATION FOR TEACHERS

OUTLINE OF CONTENT

MAJOR UNDERSTANDINGS AND FUNDAMENTAL CONCEPTS

SUGGESTED TEACHING AIDS AND LEARNING ACTIVITIES

SUPPLEMENTARY INFORMATION FOR TEACHERS

D. Mouth-to-mouth breathing

The mouth-to-mouth method of rescue breathing is recommended by the American Red Cross because it has many advantages over other methods.

Show how to clear a person's mouth of foreign objects before starting artificial respiration.

Demonstrate the mouth-to-mouth method of giving artificial respiration on the Resusci-Anne or Tammi inflatable mannequin.

Consult film listing at end of the unit for films on mouth-to-mouth resuscitation.

This technique of artificial respiration is the most practical method for emergency ventilation. Regardless of the cause of the cessation of breathing, it may be used on an individual of any age who has stopped breathing in the absence of equipment or of help from a second person.

See New York State Health Department pamphlet "Rescue Breathing" for procedure.

E. Manual methods of artificial respiration

Manual methods of artificial respiration require considerable strength and endurance when applied for long periods.

Pupils in grades 4, 5, and 6 should not be expected to use these methods as easily or as skillfully as they can employ rescue breathing techniques.

IV. Poisoning

A. Poison defined

A poison is any substance which, when taken into the body, will produce an injurious or deadly effect.

Poisoning can be intentional or accidental.

Have students list some products commonly found around the home that

Poisoning can result from taking an overdose of medicine; being bitten by snakes, insects, and marine life; inhaling poisonous gases; eating toxic foods; and attempting suicide.

Many useful common products contain poisonous substances.

OUTLINE OF CONTENT

MAJOR UNDERSTANDINGS AND FUNDAMENTAL CONCEPTS

SUGGESTED TEACHING AIDS AND LEARNING ACTIVITIES

SUPPLEMENTARY INFORMATION FOR TEACHERS

B. Effects of poisons

The effects of poisons upon the body depend upon the type, the quantity consumed, and the age and physical condition of the victim.

The two dangerous general effects of oral poisons are that they burn the tissues of the digestive organs and are absorbed into the system from the digestive tract.

contain poisonous substances. Places to consider are:

1. the medicine cabinet
2. under the sink
3. the garage
4. outside the house

Show a film on poisoning, such as Poisons in the House, which is produced by Association Films, or Children at Play with Poison, which is produced by Creative Arts Studio, Inc., or Poisoning from the American Red Cross.

Show slides Keep Out of the Reach of Children by Administration.

Do a report on the responsibility of older children in regard to preventing accidental poisoning of younger children.

Refer to Alcohol and Drug Education in Strand II.

Poisons can be grouped, depending on their physiological action, into four groups including corrosives, irritants, neurotoxins, and hemotoxins.

Corrosives include strong acids and alkalis. They cause a local destruction of the tissues with which they come in contact.

Irritants are those which produce congestion of the organ with which they come in contact. The effect that the poison ivy leaf has on the skin is a good example of an irritant.

Neurotoxins include the narcotics, barbiturates, alcohols, and anesthetics. They affect the nerves or some of the basic processes within the cell.

OUTLINE OF CONTENT

MAJOR UNDERSTANDINGS AND
FUNDAMENTAL CONCEPTS

SUGGESTED TEACHING AIDS
AND LEARNING ACTIVITIES

SUPPLEMENTARY INFORMATION
FOR TEACHERS

C. General signs and
symptoms of
poisoning

The symptoms and signs of poisoning vary greatly, depending upon the amount of poison ingested and the time elapsed since ingestion.

Hemotoxins, such as carbon monoxide and hydrogen cyanide, combine with the blood and prevent oxygen from forming hemoglobin.

Many poisons cause no symptoms until absorbed into the system. General signs to look for include the odor of poison on the person's breath, a discoloration of the lips and mouth, bottles or packages of drugs or poisonous chemicals near the victim, pain or burning sensation in throat, puncture wounds resembling a snake bite or other kind of bite, abdominal pain, vomiting, visual disturbances, headache, mental confusion, hallucinations, convulsions or a deep sleep, and evidence in the mouth of eating wild berries or leaves.

D. First aid for
poisoning due to
chemicals

First aid for oral poisoning must be immediate to prevent absorption of the poison into the blood stream.

1. general objectives
of first aid for
poisoning

The general objectives of first aid procedures are:
1. dilute the poison

Water is usually the most readily available substance for dilution. Several

OUTLINE OF CONTENT

MAJOR UNDERSTANDINGS AND FUNDAMENTAL CONCEPTS

2. eliminate the poison (except as noted)
3. neutralize it if elimination is not possible

SUGGESTED TEACHING AIDS AND LEARNING ACTIVITIES

SUPPLEMENTARY INFORMATION FOR TEACHERS

glasses of it should be given in the dilution procedure. An important procedure in many cases is to induce vomiting to eliminate the substance. When the poisonous substance was strong enough to burn the tissues going down, it will burn again coming up and vomiting should not be induced in such cases! An attempt should be made to neutralize such substances.

2. special procedures

Some poisons require special first aid procedures.

Discuss how to determine what substance has been taken; i.e., lips and mouth, the container, sudden onset, victim might say so, etc.

Discuss the incidence of childhood poisoning by: aspirin, lead products, insecticides.

Elicit several rules for prevention of poisoning.

1. Keep poisons out of reach of children.
2. Label products carefully and correctly.
3. Read labels carefully.
4. Separate poisonous substances from others.

A strong acid should be neutralized with an alkali; milk of magnesia or baking powder solution.

A strong alkali should be neutralized with an acetic substance such as lemon juice or vinegar.

Other cases when vomiting should not be induced include poisoning by strychnine (used in rat poisons) kerosene, or if the victim is unconscious. In such cases, keep the patient warm and get medical help immediately.

OUTLINE OF CONTENT

MAJOR UNDERSTANDINGS AND
FUNDAMENTAL CONCEPTS

SUGGESTED TEACHING AIDS
AND LEARNING ACTIVITIES

SUPPLEMENTARY INFORMATION
FOR TEACHERS

3. the container

The container of the poisonous substance can provide valuable clues.

Learn the meaning of antidote.

5. Discard unused medicines.
6. Never leave medicine at the bedside of a sick person.
7. Put transparent tape over labels so the writing stays clear.

Ask students to check the information listed on products commonly found around the home. Ask about types of ingredients in the product, precautions for use, and directions for first aid.

Many products list an antidote to be given in case of accidental poisoning.

The universal antidote can be given when a specific one is not known. It consists of one part each of strong tea and milk of magnesia combined with two parts of burnt toast.

Substances that coat the linings of the digestive organs to reduce intestinal absorption include milk, olive oil, salad oil, mineral oil, and egg white.

4. poison control centers

Poison control centers are available in many areas.

Have students find out where poison control centers in their area are located.

Poison control centers list many commonly used products and can recommend the specific antidote to give for each.

OUTLINE OF CONTENT

MAJOR UNDERSTANDINGS AND FUNDAMENTAL CONCEPTS

SUGGESTED TEACHING AIDS AND LEARNING ACTIVITIES

SUPPLEMENTARY INFORMATION FOR TEACHERS

E. Emergencies caused by poisonous bites or stings

Many types of organisms produce venom or other irritating substances that may create an emergency requiring first aid.

Assign special research projects on poisonous snakes, insects, and marine life. Ask for distinguishing characteristics of species which are hazardous or toxic to humans.

Bites of snakes, bees, wasps, hornets, spiders, scorpions, caterpillars, moths, stingrays, salt-water catfish, and jellyfish may produce varying degrees of poisoning of the body.

1. snake bites

The fang and tooth marks of poisonous and non-poisonous snakes are distinguishable.

Show pictures of different kinds of snakes.

The wound caused by a venomous snake is distinguished by one or two distinct punctures by the fangs.

venomous and nonvenomous snakes. Have a committee report on poisonous varieties and where they are located in the United States. Show on a map with key as to areas where poisonous snakes are found. Do the same with a world map if it is appropriate for group.

venomous nonvenomous



a. nonvenomous snakes

The bite of a nonvenomous snake produces a physical reaction in the area of the bite.

The physical reaction to a nonvenomous snake bite is localized slight burning and swelling at the site of the wound. No general reaction usually develops except, perhaps, an emotional one. However, you should consult your family doctor.

OUTLINE OF CONTENT

MAJOR UNDERSTANDINGS AND FUNDAMENTAL CONCEPTS

SUGGESTED TEACHING AIDS AND LEARNING ACTIVITIES

SUPPLEMENTARY INFORMATION FOR TEACHERS

b. venomous snakes

The venom of poisonous snakes spreads throughout the body by the circulatory system and affects the body's vital processes.

Discuss the symptoms that occur with interference with the body's systems: shortness of breath, nausea, and vomiting, weak and rapid pulse, dimness of vision, and unconsciousness.

There are four kinds of poisonous snakes in the United States including the rattlesnake, copperhead, cottonmouth moccasin, and the coral snake. The first three are known as pit vipers and have a pit on each side of the head between the eyes and the nostrils. Their venom affects the circulatory system. The venom from the coral snake is extremely toxic and affects the nervous system. Most snake bite fatalities in this country are caused by rattlesnakes since man has greater chance of contact with this snake.

c. first aid for snake bites

1. nonvenomous snakes

Nonvenomous snake bites should be treated as ordinary puncture wounds.

Review the objectives for treating simple wounds.

First aid involves having the person stop physical activity immediately and rest. Put a constriction band around the extremity just above the bite tight enough to slow the flow of blood in the surface

2. venomous snakes

The objective of first aid in case of a bite by a venomous snake is to slow down the absorption of the venom into the system until medical help can be secured.

Discuss with students steps that can be taken to prevent being bitten by snakes.

OUTLINE OF CONTENT

MAJOR UNDERSTANDINGS AND FUNDAMENTAL CONCEPTS

SUGGESTED TEACHING AIDS AND LEARNING ACTIVITIES

SUPPLEMENTARY INFORMATION FOR TEACHERS

2. Insect bites

Bites and stings of insects can be dangerous to individuals who are sensitive to the injected poison.

Discuss prevention of insect bites through the use of insect repellants and other precautionary measures.

Individual pupils may be interested in researching and reporting on poisoning by the:

Black widow spider
Brown recluse spider
Jellyfish
Portugese Man O'War
Scorpion
Stingray (Whip ray)
Tarantula

These reports may be assigned to pupils whose families vacation in areas where these creatures are found.

vessels (veins). The victim should be taken to a hospital as soon as possible so that he can get the needed treatment.

In a nonsensitive individual, a bee sting may produce painful swelling with redness, aching, and itching. Severe hives and generalized swelling of the tissues may result if the person receives several bites at one time. In the severely allergic individual, a bee sting may make the person desperately ill requiring emergency treatment to save his life.

According to the American Medical Association, the venom of these stinging insects causes more deaths each year than are caused from bites of rattlesnakes. Death occurs from respiratory obstruction caused by swelling of the vocal cords or by severe constriction and congestion of the bronchial tubes resulting in shock.

OUTLINE OF CONTENT

MAJOR UNDERSTANDINGS AND FUNDAMENTAL CONCEPTS

SUGGESTED TEACHING AIDS AND LEARNING ACTIVITIES

SUPPLEMENTARY INFORMATION FOR TEACHERS

a. first aid for persons not sensitive to insect bites and stings

First aid measures for insect bites for non-sensitive persons consists of relieving itching and pain.

Wash area very well with soap and warm water. If sensitivity is suspected, call for medical help.

Application of a paste of baking soda, cold cream, calamine, or ice water relieves pain and itching.

b. first aid for persons sensitive to insect bites and stings

A person known to be sensitive to insect bites and stings should secure preventive medical advice.

Use pamphlets The Fly and The Mosquito (N. Y. State Department of Health), to discuss insect elimination by cleanliness.

Use filmstrip and record Walter Reed (Met. Life) for the history of the scientific method in the control of yellow fever.

Do a report on the anopheles mosquito and malaria.

The American Medical Association has suggested that persons known to be sensitive to insect bites wear an identification tag to inform others in case they are bitten or stung. Pending medical care, absorption of the poisonous substance can be retarded by a tight constricting band if the wound is on an extremity. Ice applied to the wound and physical inactivity are other first aid measures to use.

F. Poisoning resulting from contact with toxic plants

There are numerous plants which cause a marked reaction on contact with human skin and may produce severe symptoms if eaten.

Show pictures and discuss some plants that have toxic effects upon humans. These could include: poison ivy, poison oak, poison sumac, poison parsley, buttercup,

It is unwise to believe that a person is immune to a given plant toxin. Individuals who have always been relatively insensitive to plant poisons (e.g., poison ivy) may suddenly become very

OUTLINE OF CONTENT	MAJOR UNDERSTANDINGS AND FUNDAMENTAL CONCEPTS	SUGGESTED TEACHING AIDS AND LEARNING ACTIVITIES	SUPPLEMENTARY INFORMATION FOR TEACHERS
1. first aid for contact with toxic plants	The objective of first aid for contact with toxic plants is to relieve the discomfort of the symptoms.	crowfoot, primrose, rue arnica, death camas, western monkshood, rosary pea, blue night- shade, black nightshade, and others.	sensitive to it.
		Discuss symptoms that might occur from contact with toxic plants. These might include redness, swelling, pain, fever, blisters, itching, etc.	Washing the exposed area with soap and water helps remove the toxic substance. Soothing and cooling lotions help to relieve the discomfort.
		Use pamphlet, N. Y. State Department of Health, <u>Poison IVY</u> .	
		Poison ivy is a vine and usually grows among other plants and shrubs or on trees. It may be hard to identify. Here's what to look for: Three little leaflets make up one leaf, each leaflet is about three inches long, edges may be smooth or have little notches or teeth. The green leaves of summer turn red in the fall. In spring, small greenish-white flowers bloom in clusters. Clusters of white berries are borne each fall. Each berry	

OUTLINE OF CONTENT

MAJOR UNDERSTANDINGS AND
FUNDAMENTAL CONCEPTS

SUGGESTED TEACHING AIDS
AND LEARNING ACTIVITIES

SUPPLEMENTARY INFORMATION
FOR TEACHERS

is about the size of a
raisin.

An old superstition
used to be that eating
poison ivy would keep
you from getting it.
This is NOT true; you
could become very ill
if you do this.

Keep the infected
area very clean with
soap and water.

Never burn poison ivy
to get rid of it. The
smoke inhaled can be
extremely dangerous. It
must be destroyed by
chemical means.

If poison ivy cannot
be cured in a reasonable
time by bathing the ef-
fected area, etc., see
your doctor.

Have a child cut out a
model of a poison ivy
plant. Put it on the
bulletin board.

OUTLINE OF CONTENT

MAJOR UNDERSTANDINGS AND FUNDAMENTAL CONCEPTS

SUGGESTED TEACHING AIDS AND LEARNING ACTIVITIES

SUPPLEMENTARY INFORMATION FOR TEACHERS

G. Poisoning by toxic food substances

Food poisoning may result from eating plants or marine life that contain toxic substances or from foods that become contaminated through improper preservation.

1. mushroom poisoning

Mushroom poisoning can be avoided by using only those obtained through commercial sources.

Discuss with pupils the differences between edible and poisonous mushrooms. Show pictures of the Amanita phalloides and Amanita muscaria mushrooms. See bibliography for textbooks containing pictures.

Since mushroom poisoning is often very severe, people who gather their own mushrooms in the woods must be aware of the danger involved.

There is no sure way in which poisonous varieties can be identified.

There is no really sure way to identify mushrooms for eating purposes. Leave that to the experts who grow them.

The Amanita phalloides contains a toxin that liquefies tissue cells all over the body, especially the red blood cells. Fifty percent of the victims die.

The Amanita muscaria contains the toxic drug muscarine which has a strong action on the nervous system. This plant is dangerous but causes death less frequently than the Amanita phalloides.

OUTLINE OF CONTENT	MAJOR UNDERSTANDINGS AND FUNDAMENTAL CONCEPTS	SUGGESTED TEACHING AIDS AND LEARNING ACTIVITIES	SUPPLEMENTARY INFORMATION FOR TEACHERS
a. symptoms	Mushroom poisoning causes severe digestive and systemic reactions.		There is acute abdominal pain, diarrhea, and vomiting. After 24 hours, the person may go into shock, have convulsions, and become comatose. Death occurs after several days.
b. emergency care	In case of mushroom poisoning, the physician should be called immediately.		First aid involves absolute bed rest, treatment for shock, and immediate summoning of a physician.
2. poisoning by fish and seafood	Ingestion of contaminated fish and seafood will produce serious digestive and systemic symptoms.	Discuss the symptoms that would be likely to occur from ingestion of contaminated fish products. These include: intense gastrointestinal symptoms, skin reactions, nervous system signs of headache and dizziness, and respiratory problems of swallowing and breathing.	Fish and seafood may become contaminated by bacterial contamination or putrefaction as the result of improper storage. Bacterial contamination and putrefaction are primary causes of poisoning. Mussel, clam, and oyster poisoning which result from the ingestion of poisonous plankton may produce paralysis from respiratory failure. The poison secreted by plankton is known as a mytilotoxin which is not destroyed by cooking. The problem occurs primarily on the Pacific coast between the months of June to October.

OUTLINE OF CONTENT

**MAJOR UNDERSTANDINGS AND
FUNDAMENTAL CONCEPTS**

**SUGGESTED TEACHING AIDS
AND LEARNING ACTIVITIES**

**SUPPLEMENTARY INFORMATION
FOR TEACHERS**

If there is any doubt at all about the edibility of a particular fish, it should not be eaten under any circumstances.

Emergency care consists of securing the physician immediately.

Under certain conditions specific bacteria invade foods causing them to be poisonous to humans.

Staphylococcus is the most common type of food poisoning.

Food handlers with staphylococcal skin infections are mainly responsible for its spread. The bacteria multiply in the food, producing an enterotoxin responsible for the symptoms of poisoning. The staphylococcus organism grows chiefly in custards, cream-filled pastry, milk, processed meat, and fish.

Discuss the symptoms which occur: nausea, vomiting, abdominal cramps, diarrhea, headache, fever, and shock, if the reaction is severe.

Foods kept at room temperatures or in large containers in a refrigerator are frequently the cause of outbreaks of staphylococcus poisoning. Discuss.

Fatalities are rare and symptoms usually subside in a day.

It is due to the ingestion of foods contaminated with Salmonella organisms. The greatest source of human infection are poultry

Discuss the symptoms which include chills, fever, vomiting, diarrhea, and abdominal pain.

Salmonellosis is a very serious infectious food poisoning that usually occurs in epidemics.

3. poisoning by bacterial infection of food products

a. staphylococcus food poisoning

b. Salmonella poisoning

OUTLINE OF CONTENT

MAJOR UNDERSTANDINGS AND FUNDAMENTAL CONCEPTS

SUGGESTED TEACHING AIDS AND LEARNING ACTIVITIES

SUPPLEMENTARY INFORMATION FOR TEACHERS

Discuss the prevention of *Salmonella* food poisoning. Discuss why it is important for food handlers to be free of bacteriological infections.

Eggs, which are a perishable food item, must always be refrigerated.

products, including chickens, ducks, turkeys, and eggs. Meats, cold cuts, frankfurters, fish, dairy products, and many vegetables, both cooked and uncooked may also be contaminated. Contamination occurs from the handling of the food by a carrier of the bacteria, who himself may not show any symptoms.

c. botulism

Botulism is a form of food poisoning that can be very serious.

Discuss the symptoms of botulism.

Symptoms include stomach upset, dimness of vision followed by double vision, difficulty in talking and swallowing, and paralysis of the throat muscles resulting in the individual choking when he tries to swallow.

The person should be kept at rest and given fluids as soon as nausea and vomiting subside. A physician should see the victim.

Botulism results from the ingestion of canned foods contaminated by botulinus spores. Home canned vegetables are especially likely to become contaminated unless they are sterilized in a pressure cooker. Canned foods which are softened, show bubbles, or have a rancid odor should not be used. More than 50 percent of the victims of botulism die. Death occurs from cardiac and respiratory failure.

OUTLINE OF CONTENT

MAJOR UNDERSTANDINGS AND FUNDAMENTAL CONCEPTS

SUGGESTED TEACHING AIDS AND LEARNING ACTIVITIES

SUPPLEMENTARY INFORMATION FOR TEACHERS

Prompt medical care is essential in cases of botulism.

Discuss the prevention of botulism. Discuss the proper methods of canning foods. Have pupils write reports on such topics as "The Commercial Methods Used in the Canning of Foods," and "Laws Regulating the Preserving of Foods."

Botulism affects the respiratory, circulatory, and nervous systems. It causes death within three to seven days after infection.

The person should be taken to a hospital where antitoxin can be given and artificial respiration can be continued as long as signs of life remain.

V. Traumatic Shock

Traumatic shock occurs as a state of depression of vital body functions resulting in circulatory failure.

Traumatic shock should always be considered as a possible consequence of

Mention other types of shock:

1. electric shock: the concern of the first aider should be to rescue the victim without harm to himself. (See American Red Cross First Aid

Some distinction should be made regarding other types of shock; however, the first aider of this age group should not be expected to make the diagnosis required to provide certain types of first aid. The first aid

OUTLINE OF CONTENTS

MAJOR UNDERSTANDINGS AND FUNDAMENTAL CONCEPTS

any serious accident or
emotional event.

SUGGESTED TEACHING AIDS AND LEARNING ACTIVITIES

Manual.) Following
rescue, artificial
respiration may be
necessary.
2. emotional shock:
procedures would be
the same as for
traumatic shock.
3. anaphylactic shock:
requires immediate
medical attention.
4. insulin shock:
results from a
diabetic's need for
sugar. If the con-
dition is positively
known, sugar could be
administered. If
there is any doubt
whatsoever, the first
aider should wait for
medical advice.

SUPPLEMENTARY INFORMATION FOR TEACHERS

procedures for traumatic
shock should suffice for
most emergencies that
this age group would be
expected to cope with.

Signs of traumatic shock
may not be immediately
apparent.

The first aider should be
alert for symptoms of shock.

Immediately following an
accident or emotional
event, the person may appear
to be all right physically
and emotionally only to
have the impact of the
event affect him seriously
within a few minutes or
hours.

Major symptoms of shock
are:
1. skin - cold, clammy,
moist, pale

OUTLINE OF CONTENT

MAJOR UNDERSTANDINGS AND
FUNDAMENTAL CONCEPTS

The first aid objectives for prevention and reduction of shock are the same.

Discuss the first aid procedures for traumatic shock. Show a film or filmstrip on shock, such as the United States Department of Health, Education and Welfare Medical Self-Help Training filmstrip on Shock.

Keep the person still, as comfortable as possible, and warm. Do not give any alcohol, once considered helpful.

It is better not to give the victim any liquid or solids since this procedure can be very dangerous. Call for professional help as soon as possible.

SUPPLEMENTARY INFORMATION
FOR TEACHERS

2. eyes - dilated pupils, glaring or staring
3. pulse - may vary, rapid and weak or absent
4. nausea - may be present

First aid procedures for preventing and reducing the effects of shock are:

1. position - Keep the patient prone to favor circulation to the head and chest unless such a position makes breathing difficult.
2. preserve body heat - Do not add heat.
3. give fluids sparingly if there is no possibility that the patient may soon have to be operated upon.
4. carefully check for injuries that could contribute to the shock condition
5. do not give stimulants
6. manipulate the victim as little as possible

OUTLINE OF CONTENT

MAJOR UNDERSTANDINGS AND
FUNDAMENTAL CONCEPTS

SUGGESTED TEACHING AIDS
AND LEARNING ACTIVITIES

SUPPLEMENTARY INFORMATION
FOR TEACHERS

VI. Burns and Other
Conditions Caused
by Abnormal
Temperatures

A. Burns

Burns are tissue injuries that may be mild or severe.

The American Red Cross film First Aid for Burns may be used.

Burns are classified by degree according to depth or severity.

Smoke inhalation can be as dangerous as a bad burn. Call for help so the person can be brought to a fresh air supply as soon as possible.

Fire is not a toy. It is a help to man and a danger....only to be used wisely and with great care and respect for its power. Look up and find out how fire changed the life of man. Put camp-fires out completely.

Characteristics of burns:
1. First degree - skin surface reddens
2. Second degree - blisters form
3. Third degree - all layers of the skin are charred and burned. Scars will result.

1. causes

Burns may result from thermal heat, sunburn, or chemicals.

Thermal burns result from exposure to excessive heat; i.e., stove water.

Chemical burns may result from contact with strong acids, alkalis, or other corrosives.

Cold water may be used to alleviate pain of first degree burns.

2. objectives

First aid objectives for burns are directed at relieving pain, preventing shock, and preventing infection.

Severely burned persons are in danger of going into shock for several weeks following the accident.

OUTLINE OF CONTENT

MAJOR UNDERSTANDINGS AND FUNDAMENTAL CONCEPTS

SUGGESTED TEACHING AIDS AND LEARNING ACTIVITIES

SUPPLEMENTARY INFORMATION FOR TEACHERS

3. procedures

First aid procedure for burns consist of covering the areas with sterile cloth to prevent air from reaching the burn.

Exposure of burned skin surfaces to air causes severe pain and increases the chances of shock. Layers of cloth exclude the air.

The cloth should be as sterile as possible to prevent the introduction of germs into the exposed surfaces. Vaseline or other greasy substances should never be applied because the doctor must only remove them later (unless the burn is very small)

Slight burns are best treated quickly with cold water to reduce skin temperature. If serious, call for help. Sterile gauze with a generous supply of petroleum jelly can be helpful for first degree burns only. People used to put butter on burns, but since much of our butter contains salt this only produces further irritation.

Cold creams or lotions may be helpful in alleviating first degree sunburn.

Whenever blisters form, whether from burns or other causes, first aid should simply include covering them with dry, sterile dressings.

First aid for burns of the eyes consists of washing the eyes with large volumes of water and seeking medical care.

OUTLINE OF CONTENT

MAJOR UNDERSTANDINGS AND FUNDAMENTAL CONCEPTS

SUGGESTED TEACHING AIDS AND LEARNING ACTIVITIES

SUPPLEMENTARY INFORMATION FOR TEACHERS

B. Frostbite

Frostbite occurs as the result of freezing of a body part.

Discuss symptoms of frostbite:

1. flushed skin before it occurs
2. white to yellowish gray glossy skin as it occurs
3. lack of pain
4. intense coldness and numbness

In cases of frostbite, it is important to avoid further tissue damage that could be caused by rubbing or massaging. The consequences could be tissue death (gangrene).

First aid for frostbite consists of careful handling and gradual rewarming.

Discuss the danger of the old remedy of rubbing the frostbitten part with snow.

The frostbitten part should be rewarmed gradually by covering it in woolen material or immersing it in water at body temperature (90° - 100° F.)

C. Problems caused by excessive heat

Heat exhaustion and heat stroke are caused by exposure to excessive heat conditions.

The two conditions are different in symptoms and severity.

Have students prepare charts to distinguish the symptoms and first aid for heat exhaustion and heat stroke.

Try to remove person from heat or to provide shade. Call for professional help.

Heat stroke is extremely dangerous. The patient becomes flushed, the temperature rises to 106° and above, the skin is hot and dry, and unconsciousness may ensue.

OUTLINE OF CONTENT	MAJOR UNDERSTANDINGS AND FUNDAMENTAL CONCEPTS	SUGGESTED TEACHING AIDS AND LEARNING ACTIVITIES	SUPPLEMENTARY INFORMATION FOR TEACHERS
<p>VII. Injuries to Bones, Muscles, and Joints</p> <p>A. Fracture</p>	<p>A fracture is a break in a bone.</p>	<p>Discuss causes and prevention of fractures.</p> <p>Discuss types of fractures and the first aid problems involved with each type.</p> <ol style="list-style-type: none"> 1. simple or closed - bone is bent or cracked; no open wound 2. compound or open - bone ends protrude through break in the skin 	<p>While awaiting medical care, the first aider should try to bring down the temperature by bathing the patient with alcohol or water. If conscious, the patient would profit by having fluids.</p> <p>General symptoms would include pain and tenderness at the site of the break, loss of use of the extremity containing the break, swelling and deformity of the broken part, a grating sensation in the broken area, discoloration of the skin, shortening of the limb, and the presence of a wound with a bone visible or projecting through.</p> <p>Some types of simple fractures may have no obvious symptoms.</p> <p>The major objective with a fracture is to immobilize the broken ends of</p>
<p>Objectives of first aid are determined by the nature of the fracture.</p>	<p>The McGraw-Hill First Aid Series filmstrip on <u>Bone, Muscle and Joint</u></p>		

OUTLINE OF CONTENT

MAJOR UNDERSTANDINGS AND
FUNDAMENTAL CONCEPTS

SUGGESTED TEACHING AIDS
AND LEARNING ACTIVITIES

SUPPLEMENTARY INFORMATION
FOR TEACHERS

Injuries may be shown at the beginning of this unit. The American Red Cross film, First Aid for Injuries to Bones, Joints and Muscles, may be used.

the bone and the adjacent joints. If a wound is associated with the fracture, additional consideration must be given to controlling bleeding and preventing infection.

Discuss items that could be used in an emergency to immobilize a fracture (magazines, pillows, blankets).

Immobilization can be achieved with many types of articles commonly found around the home.

Keep person as warm and comfortable as possible, and call for professional help immediately.

A splint consists of material that can be applied to a broken limb to make it rigid. The splint should immobilize the joint above and below the broken bone. Proper splinting relieves pain and often lessens shock. Splints also help to prevent further injury. The splint should be used by an experienced person only.

B. Sprain

A sprain is an injury where there is damage done to the ligaments surrounding a joint without the displacement of bones.

Discuss the causes of sprains and the kinds of accidents that result in this kind of injury. Have pupils tell about their experiences with sprains. Discuss the symptoms of a sprain.

It is caused by a sudden twist or wrench of a joint, as when the foot is unexpectedly caught in an obstruction while walking or running.

OUTLINE OF CONTENT

MAJOR UNDERSTANDINGS AND FUNDAMENTAL CONCEPTS

SUGGESTED TEACHING AIDS AND LEARNING ACTIVITIES

SUPPLEMENTARY INFORMATION FOR TEACHERS

Symptoms include pain at the site of the sprain, swelling, loss of function in the affected part, black and blue discoloration, and tenderness in the area of the sprain.

The possibility of a fracture should be suspected in all sprains since a sprain cannot be distinguished from a fracture except by X-ray.

These injuries frequently are associated with small chip fractures.

The objective of first aid for a sprain is to reduce the pain caused by swelling of the tissues.

C. Strain

A strain is an overstretching or tearing of a muscle or a tendon.

Cold application should be applied the first 24 hours. The sprained part should be immobilized and rested.

They are frequently caused by lifting heavy objects improperly. Violent, unexpected movements also produce strains. The back muscles are frequently involved.

Discuss the causes of strains and the kinds of accidents that produce this kind of injury. Have pupils tell about their experiences with strains.

Pain on movement is the primary symptom. Movement is limited due to the pain.

A strain can be greatly helped by a good hot bath or shower.

OUTLINE OF CONTENT

MAJOR UNDERSTANDINGS AND FUNDAMENTAL CONCEPTS

SUGGESTED TEACHING AIDS AND LEARNING ACTIVITIES

SUPPLEMENTARY INFORMATION FOR TEACHERS

The objective of first aid for a strain is to allow the injured part to repair through rest.

Discuss prevention of strains through proper lifting, carrying, etc.

Stiffness may also be present.

The injured part should be kept at rest and heat should be applied.

Show film How to Avoid Muscle Strains.

Lifting should be done with the muscles of the legs and not the back. The lift should be done with the back straight, the feet firmly planted on the ground, and the knees bent. The lift is made by straightening the legs so that the weight is distributed evenly along the strong, bony structures of the vertebral column and pelvis, rather than borne by the back muscles alone.

D. Dislocations

A dislocation is an injury where there is damage done to the ligaments surrounding a joint with a temporary removal of a bone from its normal position.

Discuss the causes of dislocations and examples of accidents that would cause this kind of injury. Discuss the role of athletics in producing dislocations.

After completing the materials on fractures, sprains, and dislocations, devise accident problems involving these injuries

Dislocations occur as a result of a twisting force applied to a bone near a joint. Falls, blows, and sudden spastic muscular contractions frequently produce dislocations. There is normally an unnatural shape of the affected part, severe pain in the joint, rigidity and loss of

OUTLINE OF CONTENT

MAJOR UNDERSTANDINGS AND
FUNDAMENTAL CONCEPTS

SUGGESTED TEACHING AIDS
AND LEARNING ACTIVITIES

and have the pupils provide the needed first aid.

Example:

An elderly man is found lying on the ground in his yard. His left leg is twisted under his right leg at an odd angle. Blood is seeping through his left trouser leg. A ladder with a broken rung is found nearby. What would you do as a first aider?

The injured part should be properly immobilized and the person transported to a hospital.

Who do you call in an emergency?

SUPPLEMENTARY INFORMATION
FOR TEACHERS

function, and marked swelling.

Keep person warm. Do not move injured person. If he is unconscious, try to keep his head on the side so that his tongue does not fall back in his mouth to choke him. Call for professional help immediately.

Make a list as a review including:

1. parent or nearest dependable adult
2. family doctor
3. police
4. fire department
5. poison control center
6. other, due to area and appropriateness to group

OUTLINE OF CONTENT

MAJOR UNDERSTANDINGS AND
FUNDAMENTAL CONCEPTS

SUGGESTED TEACHING AIDS
AND LEARNING ACTIVITIES

SUPPLEMENTARY INFORMATION
FOR TEACHERS

Plan for a visit from or
to one of these agencies
to discuss what its role
is in emergency care.

These supplementary aids have not been evaluated. The list is appended for teacher convenience only and teachers in the field are requested to critically evaluate the materials and to forward their comments to the Curriculum Development Center.

FIRST AID AND SURVIVAL EDUCATION

Grades 4--6

Multimedia Resources

TEACHER REFERENCES

BOOKS

- American Academy of Pediatrics, Committee on Accident Prevention. Accidental poisoning in childhood. Evanston. American Academy of Pediatrics. 1956.
- American College of Surgeons, Committee on Trauma. Emergency care of the sick and injured. Edited by R. H. Kennedy. Philadelphia. W. B. Saunders. 1966.
- American Medical Association. First aid manual. Chicago. American Medical Association. 1967.
- _____. Today's health guide. Bauer, W. W. ed. 1965.
- American National Red Cross. First aid textbook. 4th ed. Washington, D. C. Doubleday and Co. 1957.
- _____. First aid textbook for juniors. 2d ed. rev. 1966.
- _____. A supplement on artificial respiration to accompany Red Cross textbooks.
First aid
First aid textbook for juniors
Life saving and water safety
Teaching Johnny to swim. ARC. 1068B. 1959.
- _____. Mother's aid course instructor's guide. ARC. 1672. 1966.

- ____ Standard and advanced courses first aid instructor's manual. ARC. 1090. 1966.
- Armstrong, Richard. Cold hazard. Boston. Houghton Mifflin Co. 1961.
- Bellios, A. D. Handbook of first aid and bandaging. 5th ed. Baltimore. Williams and Wilkins Co., Inc. 1963.
- Board of Education the City of New York. Health teaching in elementary schools. pp. 54-60. Curriculum Bulletin 1955-56 Series No. 5. New York. Board of Education the City of New York. 1956.
- ____ Health teaching in secondary schools. pp. 81-85. Curriculum Bulletin 1960-61 Series No. 6. 1961.
- Bolton, W. W. What to do until the doctor comes. Chicago. Reilly and Lee and Co. 1960.
- Brennan, W. T. & Ludwig, D. T. Guide to problems and practices in first aid and civil defense. Dubuque. W. C. Brown and Co. 1962.
- Byrd, O. E. Health today and tomorrow. Teacher's ed. pp. 357-383. Summit. Laidlaw Brothers Publishers. 1966.
- Clark, R. L. & Cumley, R. W. eds. The book of health. 2d ed. pp. 633-673. Princeton. D. Van Nostrand Co. 1962.
- Cole, W. H. & Puestow, C. B. First aid: Diagnosis and management. 6th ed. New York. Appleton-Century Crofts Co., Inc. 1965.
- Consumer Reports. eds. The medicine show. rev. Mt. Vernon. Consumers Union. 1963. \$1.50.
- Cramer Chemical Company. First aid digest. Charlotte. Gardner Publishers. 1967.
- Equitable Life Assurance Society. Home health emergencies. New York. Equitable Life Assurance Society. 1956.
- Fishbein, M. & Irwin, L. W. First aid training. Chicago. Lyons and Carnahan Co., Inc. 1961.
- Halstead, B. W. Dangerous marine animals. Cambridge. Cornell Maritime Press. 1959.
- Henderson, John. Emergency medical guide. 2d ed. New York. Blakiston Division, McGraw-Hill Book Co. 1969.
- Hoffman, F. P. Artificial respiration. New York. Queens College. undated.
- Igel, B. H. First aid. A programmed text teacher's manual. New York. Instructional Materials Laboratories, Inc. 1965.
- John Hancock Life Insurance Company. When the unexpected happens. Boston. John Hancock Life Insurance Co. 1965.

- Lyght, C. E. ed. The Merck manual of diagnosis and therapy. 11th ed. Rahway, Merck, Sharp and Dohme Research Laboratories. 1966.
- Metropolitan Life Insurance Company, First aid. New York. Metropolitan Life Insurance Company. 1960.
- Miller, L. M. Handbook of first aid. Pleasantville. Reader's Digest. 1960.
- Nassau Tuberculosis and Health Association. Curriculum guide for health and safety kindergarten through grade twelve. Roslyn. Nassau Tuberculosis and Health Association. 1966.
- ____ Part I: The beginning years. K-6. pp. 1-4.
- ____ Part II: The middle years. 7-9. pp. 9-10.
- ____ Part III: The senior years. 10-12. pp. 14-15.
- National Safety Council. Accident facts. Chicago. National Safety Council.
- New York State Department of Health and Civil Defense Commission. Guide for medical aides in aid stations. rev. New York. New York State Department of Health and Civil Defense Commission. 1964.
- ____ Guide for self-help and neighbor help for the injured. 1951.
- ____ Instructor's handbook for self-help and neighbor help for the injured. 1951.
- ____ Instructor's handbook for psychological first aid. 1958.
- ____ Rescue breathing: improved techniques of expired air resuscitation. 1961.
- North Carolina State Department of Public Instruction. Health education. pp. 319-322. Publication No. 287. Raleigh. North Carolina State Department of Public Instruction. 1953.
- Sharpe, J. C. & Marx, F. D. eds. Management of medical emergencies. 2d ed. New York. McGraw-Hill Book Co. 1969.
- Stephenson, H. E. Emergency care and first aid. Boston. Little Brown and Co. 1964.
- Suchman, E. A. & Scherzer, A. L. Current research in childhood accidents. New York. The Association for the Aid of Crippled Children. 1960.
- Thorndike, A. Athletic injuries: prevention, diagnosis and treatment. 5th ed. Philadelphia. Lea and Febiger Co., Inc. 1962.

United States Department of Defense and United States Department of Health, Education and Welfare. Family guide, emergency health care. Washington, D. C. United States Government Printing Office. 1965.

Accidents and children. Children's Bureau.

United States Public Health Service, Division of Accident Prevention. Teaching poison prevention in kindergarten and primary grades. Washington, D. C. United States Government Printing Office. 1965.

PERIODICAL ARTICLES

American Academy of Pediatrics.

New hazards of chemicals and drugs to children. News letter supplement. January 15, 1968.

Bulletin of the New York Academy of Medicine.

Present status of the treatment of burns. Vol. 43 52-55, 89-90, 92. January 1962.

Consultant.

Treatment of frostbite today. pp. 44-47. January 1967.

First Aider.

The L-C treatment of venomous bites and stings. Vol. XXV. No. 13. 236-239. April 18, 1966.

Instructor.

Six graders practical rescue breathing. Vol. 77:124. March 1968.

Journal of Health, Physical Education and Recreation.

Accidents among elementary children. Vol. XXIX. No. 4. 20-21. April 1958.

Colorful first aid teaching. Vol. XXXVI. No. 4. 31, 47. April 1965.

First aid and legal liability. Vol. XXXIV. No. 1. 42-43. January 1963.

Home made mannequin for first aid classes. Vol. XXXVIII. No. 2. 65. February 1966.

How our schools teach home nursing and first aid. Vol. XXIII. No. 3. 3052. March 1952.

Medical self-help in the curriculum. Vol. XXXVI. No. 8. 71. October 1965.

Rescue breathing pilot program. Vol. XXXIV. No. 3. 84. March 1963.

Teaching oral resuscitation. Vol. XXI. No. 5. 34-35, 44. May-June 1960.

Journal of School Health

Childhood accidents. Vol. XXVIII. No. 10. 350. November 1956.

Nature and frequency of accidents among school children in New York State. Vol. XXVII. No. 10. 343.

December 1958.

Journal of the American Medical Association.

- Cardio-pulmonary resuscitation. 198: 372-379. October 24, 1966.
International comments: When is a patient dead? Vol. 197. No. 8. 153. August 22, 1966.
Personal protection from blood-sucking arthropods. Vol. 196. 236-239. April 18, 1966.
Treatment of snakebite. Vol. 193. No. 5. 336-338. August 2, 1965.
When death occurs: Some practical aspects. Vol. 197. No. 12. 329-330. September 19, 1966.

New England Journal of Medicine.

- Accidental poisoning. Vol. 277. No. 21. 1135-1137. November 23, 1967.
Electric shock. Vol. 268. No. 20. 1127-1128. May 16, 1963.

Today's Health.

- Artificial respiration. Vol. XL. No. 4. 6-7, 85-87. April 1962.
Can your child handle an emergency? Vol. XXXIX. No. 1. 54-55, 79-81. January 1961.
First aid for athletic injuries. Vol. XLII. No. 10. 49. October 1964.
How to make yourself fall safe. Vol. XLII. No. 4. 72-76. April 1964.
Lead poisoning: New look at an old hazard. Vol. XL. No. 1. 52-53, 89-90, 92. January 1962.
A medical I. D. tag can save your life. Vol. XLI. No. 7. 8-13. July 1963.
New help for the burn injured child. Vol. XLIII. No. 43-45, 70, 73-75. July 1965.
Ordeal by sting. Vol. XXXVIII. No. 7. 30-31. 69-72. July 1960.
Suddenly you're involved. Vol. XLII. No. 6. 26-29, 65-66. June 1964.
Survival begins at home: part 4: artificial respiration. Vol. XL. No. 4. 6-7, 84-87. April 1962.
Part 5: shock. Vol. XL. No. 5. 24, 27, 79-80. May 1962.
Part 6: bleeding and bandaging. Vol. XL. No. 6. 32-33, 58-63. June 1962.
Part 7: fractures and splinting. Vol. XL. No. 7. 12-13, 41-43. July 1962.
Part 9: burns. Vol. XL. No. 9. 56-59. September 1962.
Tragedy of needless drowning deaths. Vol. XLIII. No. 7. 46-48, 66. July 1965.
When the heart stops. Vol. XXXVIII. No. 11. 14, 80-81. November 1960.
When lightning strikes. Vol. XLII. No. 7. 14-17, 72. July 1964.
Why people faint. Vol. XLI. No. 8. 18-19, 53, 55-58. August 1963.
You can prevent poison accidents. Vol. XXIX. No. 10. 36-37, 86, 88. October 1961.

STUDENT REFERENCES

BOOKS

- American Medical Association. First aid manual. rev. ed. Chicago. American Medical Association. 1967.
- American National Red Cross. First aid textbook. 4th ed. Washington, D. C. Doubleday and Co. 1957.
- _____ First aid textbook for juniors. 2d ed. rev. 1966.
- _____ A supplement on artificial respiration to accompany Red Cross textbooks:
First aid.
First aid textbook for juniors.
Life saving and water safety.
Teaching Johnny to swim. ARC 1086B. 1959.
- Equitable Life Assurance Society. Home health emergencies. New York. Equitable Life Assurance Society. 1956.
- Forest Service, United States Department of Agriculture. First aid guide. Washington, D. C. United States Department of Agriculture. \$.25 for campers.
- Igel, B. H. First aid. A programmed text. New York. Instructional Materials Laboratories, Inc. 1965
- John Hancock Life Insurance Company. When the unexpected happens. Boston. John Hancock Life Insurance Company. 1964.
- Johnson and Johnson. First aid guide. New Brunswick. Johnson and Johnson Co. 1967.
- New York State Department of Health and Civil Defense Commission. Rescue breathing: improved techniques of expired air resuscitation. New York. New York State Department of Health and Civil Defense Commission. 1961.
- Reader's Digest. Handbook of first aid. Pleasantville. Reader's Digest. September 1960.

United States Department of Health, Education and Welfare. Dennis the Menace takes a poke at poison. A comic book. Washington, D. C. United States Government Printing Office. 1961.

United States Office of Civilian Mobilization. First aid. Washington, D. C. Supt. of Documents. leaflet \$.05 1961.

AUDIOVISUAL AIDS

FILMS

American National Red Cross. (consult your local chapter for these films)

Artificial respiration

Checking for injuries

First aid for bones, joints and muscles

First aid for burns

First aid for shock

First aid: Part I and II

Poisoning

Target: babies and children

The how and why of first aid

Wounds

Your breath can save a life

Breath of life. Adams Productions, Inc., Box 1048. Santa Monica, California. 90406. 27 min. color.

Films (Available from Film Library, New York State Department of Health, 84 Holland Avenue, Albany, New York 12208.

Accidentally yours. 15 min. color.

Breath of life. 16 min. color.

Children at play with poison. 10 min. color.

Danger--high voltage. 15 min. color.

Donald's fire survival plan. 11 min. color.

Electrical safety in the home. 13 min. color.

Film library, New York State Department of Health, 84 Holland Avenue, Albany, New York 12208.

Fires and wires. 21 min. color

First aid now. 26 min. color.

First aid on the spot. 10 min. black and white.

Hands of action. 42 min. black and white.

How to avoid muscle strain. 15 min. black and white.

Life in your hands. 11 min. black and white.
One day's poison. 27 min. black and white.
Poison in the house. 10 min. color.
Pulse of life. 27 min. color.
Rescue breathing. 22 min. black and white.
To save a life. 14 min. color.
Water rescue. 12 min. color.

FILMSTRIPS (Available from McGraw-Hill Films. First aid series, Text Film Department, 330 W. 42nd Street, New York 36, New York)

Artificial respiration.
First aid.
First aid - bleeding and shock.
First aid - bone, muscle and joint injuries.
First aid - wounds.
Five basic steps in first aid.
Your responsibility in first aid.

FILMSTRIPS - SOUND (Available from Metropolitan Life Insurance Company)

Louis Pasteur. The germ theory and the history of immunization.
The history of nursing. Florence Nightingale and the importance of cleanliness.
Walter Reed. History of work with yellow fever, experimentation and control.

SLIDES

Keep out of the reach of children. Food and Drug Administration, Visual Service Branch. Bureau of Education and Voluntary Compliance, Washington, D. C. 20204.

CHARTS AND POSTERS

Flip Chart. Self-help and neighbor help for the injured course. 18 1/2 x 28 1/2 in. 102pp. color. English and Spanish available. New York State Department of Health. New York State Civil Defense Department. 1963.

Flip Chart. Training course for medical aides in aid stations. 18 1/2 x 28 1/2 in. 100pp. color. New York State Department of Health. 84 Holland Avenue. Albany, New York 12208. 1964.

Safety guides for upper elementary grades. ARC 1447. 8 1/2 x 11 in. set of nine. Consult your local American National Red Cross.

MANNEQUINS

Resusci Anne & Resusci Andy. Life size for use in resuscitation training. Guardian Safety Equipment Company, 37 East 21st Street, Linden, New Jersey 07037. Hygienic face shields. Available from Laerdal Medical Corporation, 136 Marbledale Road, Tuckahoe, New York 10707.

Respiration. Half-bodied mannequin for use in resuscitation training. Model F., including the external cardiac resuscitation unit, shirt, fibre carrying case, 100 face shields. Additional face shields available. Alderson Research Laboratories, Inc. 729. Canal Street, Stamford, Connecticut. Approx. \$198.00.

Resuse - Kate. 24 in. full-bodied mannequin of a child for use in resuscitation training. No. 110. Simulaid. Woodstock, New York 12498. Approx. \$22.50.

Resusci Baby. Lifelike normal sized baby mannequin for training in mouth-to-mouth resuscitation and cardiopulmonary resuscitation of infants. Further information write: Laerdal Medical Corporation, 136 Marbledale Road, Tuckahoe, New York 10707.

Tammi. Life size mannequin for use in teaching mouth-to-mouth resuscitation and external cardiac compression. Further information write: Uni/Flex Medical Supply Company, Rockford, Illinois 61101

Anatomic Anne. Training mannequin that visually shows circulation of blood and inflation of lungs during the correct performance of cardiopulmonary resuscitation. Further information write: Laerdal Medical Corporation, 136 Marbledale Road, Tuckahoe, New York 10707.

TEACHING KITS

Medical self-help instructors kit. United States Department of Defense and United States Department of Health, Education and Welfare.

Health survival instructor's guide.

Course introduction booklet.

11 lesson plan books.

Family guide-emergency health care manual.

11 35 mm filmstrips.

Examination books.

Grading templates.

11 16 mm sound-color films.
Film. When disaster strikes. narrated by Danny Thomas. 13 1/2 min. color.

Consult New York State Department of Health, 84 Holland Avenue, Albany, New York 12208.
No charge for kit or student supplies.

PAMPHLETS (Available from New York State Department of Health, 84 Holland Avenue, Albany, New York 12208.)

Poison Ivy. 1952.

The fly. rev. 1959.

The mosquito. 1954.